



HAL
open science

Statistical analysis of the precision of the Match method

R. Lehmann, P. von Der Gathen, M. Rex, M. Streibel

► **To cite this version:**

R. Lehmann, P. von Der Gathen, M. Rex, M. Streibel. Statistical analysis of the precision of the Match method. Atmospheric Chemistry and Physics Discussions, 2005, 5 (3), pp.3225-3268. hal-00301455

HAL Id: hal-00301455

<https://hal.science/hal-00301455>

Submitted on 18 Jun 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Precision of the
Match method**

R. Lehmann et al.

Statistical analysis of the precision of the Match method

R. Lehmann, P. von der Gathen, M. Rex, and M. Streibel

Alfred Wegener Institute for Polar and Marine Research, Research Unit Potsdam,
Telegrafenberg A43, D-14473 Potsdam, Germany

Received: 2 February 2005 – Accepted: 15 April 2005 – Published: 24 May 2005

Correspondence to: R. Lehmann (rlehmann@awi-potsdam.de)

© 2005 Author(s). This work is licensed under a Creative Commons License.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

Abstract

The Match method quantifies chemical ozone loss in the polar stratosphere. The basic idea consists in calculating the forward trajectory of an air parcel that has been probed by an ozone measurement (e.g., by an ozone sonde or satellite) and finding a second ozone measurement close to this trajectory. Such an event is called a “match”. A rate of chemical ozone destruction can be obtained by a statistical analysis of several tens of such match events. Information on the uncertainty of the calculated rate can be inferred from the scatter of the ozone mixing ratio difference (second measurement minus first measurement) associated with individual matches. A standard analysis would assume that the errors of these differences are statistically independent. However, this assumption may be violated because different matches can share a common ozone measurement, so that the errors associated with these match events become statistically dependent. Taking this effect into account, we present an analysis of the uncertainty of the final Match result. It has been applied to Match data from the Arctic winters 1995, 1996, 2000, and 2003. For these ozone-sonde Match studies the effect of the error correlation on the uncertainty estimates is rather small: compared to a standard error analysis, the uncertainty estimates increase by 15% on average. However, the effect is more pronounced for typical satellite Match analyses: for an Antarctic satellite Match study (2003), the uncertainty estimates increase by 60% on average.

1. Introduction

The Match method was developed to quantify chemical ozone loss in the Arctic stratosphere (von der Gathen et al., 1995; Rex et al., 1998, 1999). The basic idea is the following: After an air parcel has been probed by an ozone sonde, its forward trajectory is calculated. If a second ozone sonde comes close to this trajectory, i.e. its distance from the trajectory is smaller than a pre-defined threshold, then the measurements of the two ozone sondes form a “match”. This can happen by chance or may be attained

Precision of the Match method

R. Lehmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Precision of the Match method

R. Lehmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

by launching the second sonde intentionally so that it approaches the trajectory at the appropriate time. In the final analysis, backward trajectories may also be applied to determine matches. It is possible that an ozone measurement forms matches with more than one other ozone measurement.

5 A variant of the original Match method uses satellite data instead of ozone sonde measurements (Sasano et al., 2000; Terao et al., 2002; and Sects. 8.4 and 8.5 of the present paper). Although all formulae derived in the present paper are also applicable to satellite or other data, we will prefer the terminology of the original Match method (e.g., “first sonde” and “second sonde” of a match).

10 Under ideal circumstances (no measurement errors, no trajectory error, zero distance of the second sonde from the trajectory), the difference of the ozone mixing ratios measured by the first and second sonde would be equal to the change of the ozone mixing ratio along the trajectory. Under the assumption that mixing can be neglected, this is equal to the chemical ozone loss in the corresponding air parcel.

15 As the above-listed errors are non-zero in reality, it is not possible to draw conclusions from a single pair of ozone observations. However, if several tens of such pairs are available, then an ozone loss rate can be obtained by statistical methods. For this, the differences between the ozone mixing ratio of the first and second measurement of several pairs of “matching” sondes are plotted versus a variable that is expected to correlate with ozone destruction, usually the time that the corresponding trajectories spent in sunlight. Then a linear fit, which is forced through the origin (0,0) of the coordinate system, is performed. The slope of that line yields an estimate of the mean ozone loss rate, e.g., ozone loss per hour of sunlight. An example is given in Fig. 1.

20 A standard assumption of a linear regression analysis is the statistical independence of the errors of the quantities entering the analysis. In the case of the Match method, these quantities are the ozone mixing ratio differences of several matches. As one ozone measurement may enter several matches, the errors of those matches will not be uncorrelated. The influence of this effect on the Match analysis will be investigated in this paper. It turns out that the linear regression still yields an unbiased estimate of

Precision of the Match method

R. Lehmann et al.

the mean ozone loss rate (Sect. 5), but the calculation of the corresponding uncertainty (“error bars”) is affected.

First, the effect of the use of correlated ozone data on the uncertainty of the Match results is illustrated with the help of a highly simplified example in Sect. 2. Then the uncertainties associated with the Match method are reviewed in Sect. 3. The formulae for the exact numerical treatment of this effect are derived in Sects. 4–6 and summarised in algorithmic form in Sect. 7. Finally, results of the application of the new formulae to data from five Match campaigns are presented in Sect. 8.

2. Illustration of the effect of correlated matches

As mentioned in Sect. 1, the ozone mixing ratio differences (second measurement minus first measurement) associated with individual match events, like in Fig. 1, may be correlated, because one ozone measurement may be part of several match events. This will influence the estimation of the precision of the slope of the regression line (cf. Fig. 1), i.e. of the ozone loss rate. In order to illustrate this effect, we consider a highly simplified example: We assume that only two matches occurred and that there was no ozone loss along the corresponding two trajectories. Furthermore, the sunlit times associated with the two matches are assumed to be equal and will be denoted by t_0 . Then the ozone loss rate determined by the linear regression is equal to the mean of the ozone loss rates calculated for the individual match events. In order to simplify the situation further, we assume that only one ozone measurement per match has a measurement error and that this error can only assume two values ($+\delta$ and $-\delta$) of equal probability. All other errors associated with the Match method are assumed to vanish. Now we consider three cases:

1) The two matches are independent: If we assume that the measurements of the second sondes of these matches are associated with an error, then the ozone mixing ratio differences measured in the two match events, d_1 and d_2 , may assume the following

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

**Precision of the
Match method**

 R. Lehmann et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Print Version](#)
[Interactive Discussion](#)

EGU

values (with equal probabilities):

$$d_1 = +\delta, d_2 = +\delta \text{ or}$$

$$d_1 = +\delta, d_2 = -\delta \text{ or}$$

$$d_1 = -\delta, d_2 = +\delta \text{ or}$$

$$5 \quad d_1 = -\delta, d_2 = -\delta .$$

The mean ozone destruction rate will be $\frac{\delta}{t_0}$ in the first case, 0 in the second and third case, and $-\frac{\delta}{t_0}$ in the fourth case. This is illustrated in Fig. 2.

2) The two matches have a common second sonde: If we assume that this sonde is associated with an error, then the ozone mixing ratio differences measured may assume the following values (with equal probabilities):

$$d_1 = +\delta, d_2 = +\delta \text{ or}$$

$$d_1 = -\delta, d_2 = -\delta .$$

The mean ozone destruction rate will be $\frac{\delta}{t_0}$ in the first case, and $-\frac{\delta}{t_0}$ in the second case. This is illustrated in Fig. 3. As the extreme values of the ozone destruction rate ($\frac{\delta}{t_0}, -\frac{\delta}{t_0}$) of case 1 still occur, but the middle value (0) does not appear, the variance of the estimated ozone loss rate has increased compared to case 1.

3) The two matches have a common sonde, which is second sonde for the first match and first sonde for the second match: If we assume that this sonde is associated with an error, then the ozone mixing ratio differences measured may assume the following values (with equal probabilities):

$$d_1 = +\delta, d_2 = -\delta \text{ or}$$

$$d_1 = -\delta, d_2 = +\delta .$$

The mean ozone destruction rate will be 0 in both cases. This is illustrated in Fig. 4. In this example the variance of the ozone loss rate is zero, i.e. clearly smaller than in the first case.

The results of these three very simple examples are summarised in Fig. 5. They illustrate that pairs of matches with a common second (or first) sonde will increase the variance of the estimated ozone loss rate, compared to the case of independent matches. Pairs of matches with a common sonde, that is first sonde for one of the matches and second sonde for the other one, will decrease the variance of the estimated ozone loss rate. These simple considerations will be confirmed by the calculations in Sect. 6.

3. Uncertainties associated with the Match method

3.1. Random errors versus systematic errors

In general, measurement errors can be classified as systematic (“accuracy”) or random errors (“precision”).

Systematic errors are inherent in the measurement technique and can only be estimated by an investigation of this technique itself. Systematic errors of the Match method can be caused, e.g., by approximations in the code that calculates the vertical position of trajectories. For a discussion of systematic errors see Rex et al. (1998), Morris et al. (2004).

Random errors can be estimated by statistical methods from the scatter of measurements. The errors of the Match method contain a significant random component, arising, e.g., from random errors of the ozone measurements and of the meteorological data that enter the trajectory calculations. Information on the magnitude of the random errors associated with individual match events can be obtained from the scatter of the corresponding ozone mixing ratio differences around the regression line describing the mean ozone loss (cf. Fig. 1). Then straight-forward statistical methods allow to esti-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Precision of the Match method

R. Lehmann et al.

mate the random error of the slope of the regression line itself, i.e. of the ozone loss rate determined by the Match method.

This paper will deal exclusively with random errors, i.e. with the precision of the Match method. The term “uncertainty” will be used in this sense.

Precision may be quantified by the standard deviation of the random variable describing the quantity of interest. This standard deviation is usually unknown, but an estimate may be obtained from realisations of the random variable. Error bars are a graphical representation of the precision, in the above-cited Match publications they represent one standard deviation. Similarly, here we will use the term “precision estimate”, or alternatively “error bar”, for denoting an estimate of the standard deviation of the random errors of the Match results.

3.2. Sources of uncertainties associated with the Match method

The uncertainties associated with the Match method have the following sources, which are schematically depicted in Fig. 6:

- a) measurement error of the first and second ozone sonde of a match; this includes the error of the ozone measurement itself and the errors of the pressure and temperature measurements, which are translated into an error of the potential temperature level to which an ozone measurement is assigned; furthermore, the ozone mixing ratio at the point of the sonde measurement may deviate from the mean mixing ratio in a larger air parcel, e.g. within the match radius, due to small-scale ozone variations;
- b) trajectory error (horizontal and vertical), which results in an ozone measurement displaced from the required position;
- c) match radius, i.e. the distance of the measurement of the second sonde from the trajectory at the time of the measurement; this also results in an ozone measurement displaced from the required position;

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

- d) deviation of the ozone loss rate on an individual trajectory from the mean loss rate in a region of interest.

The Match method attempts to limit the above-listed error sources if possible (Rex et al., 1998, 1999):

- 5 a) Standard operation procedures for participating ozone sonde stations have been worked out. However, certain measurement errors are inevitable, unless improved ozone sondes become available.
- b) The Match method includes procedures to limit the effect of the trajectory error: A cluster of trajectories around the trajectory of interest is calculated. If these trajectories diverge significantly, it is assumed that the main trajectory is more error-prone. In this case it is discarded, i.e. not used for establishing a match. In order to ensure that vertical trajectory errors are not translated into large errors of the ozone mixing ratio, only ozone profiles with vertical gradients below a certain threshold are used.
- 10
- 15 c) In order to limit the error of the ozone mixing ratio resulting from a non-zero match radius, a suitable maximum match radius is applied.
- d) The variability of the ozone loss rate on different trajectories is determined by the inhomogeneity of the chlorine activation in the region of interest and also by differences of the solar zenith angle during solar illumination along the trajectories. It is thus objectively present, but it may, to some degree, be influenced by the choice of the region of interest (whole polar vortex, vortex core etc.).
- 20

Error a) is an instrumental error, whereas errors b)–d) are related to the technique of Match. We will denote the combined effect of b)–d) as “net match error” (“net”, because the measurement errors a) are not included).

25 The exact knowledge of the sources of errors will not be of relevance for the derivations in the subsequent sections. However, this work will be based on the fact that

Precision of the Match method

R. Lehmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Precision of the Match method

R. Lehmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

there are two categories of random errors: those associated with the ozone measurements, which may be shared by several matches, and individual net errors of each match. All of these errors will be assumed to be statistically independent from each other. This assumption needs some discussion: Two match events having a common first sonde also share a common trajectory segment, starting at the position of the first sonde measurement. That is why their trajectory errors can be correlated, which may lead to correlated net match errors. If this effect was significant, then it would be reflected in the covariance (of the ozone mixing ratio difference) of pairs of matches sharing a common first sonde. It would have a larger absolute value than the corresponding covariance for pairs of matches that share a common ozone sonde, but no common trajectory segment, i.e. for the case “second sonde of first match = first sonde of second match”. However, for the ozone-sonde data analysed in Sect. 8, this was not the case, so that we can conclude that the correlation between the net match errors is negligible.

4. Statistical description of the uncertainties associated with the Match method

As a basis for estimating the uncertainty of the final Match result, we are going to provide a statistical description of the uncertainties discussed in Sect. 3. First we introduce the following notations:

n = number of ozone observations,

m = number of match events,

\bar{r} = mean ozone loss rate (loss per sunlit time) for the atmospheric region probed by the matches under consideration [ppb/h] (to be estimated),

c_k = ozone mixing ratio determined by the k -th sonde measurement, $k = 1, \dots, n$,

d_i = difference of the ozone measurements of the i -th match, $i = 1, \dots, m$,

t_i = sunlit time of the i -th match, $i = 1, \dots, m$,

Precision of the Match method

R. Lehmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

δ_k = error of the k -th ozone measurement, $k = 1, \dots, n$,

Δ_i = net match error of the i -th match event, resulting from the combined effect of the trajectory error, the non-zero match radius, and the deviation of the ozone loss rate on the i -th trajectory from the mean ozone loss rate \bar{r} , $i = 1, \dots, m$,

ϵ_i = total error associated with the i -th match, $i = 1, \dots, m$,

$k_1(i)$ = index of the first sonde measurement of the i -th match, $i = 1, \dots, m$,

5 $k_2(i)$ = index of the second sonde measurement of the i -th match, $i = 1, \dots, m$.

In order to simplify the notation, we define the following vectors:

$$\mathbf{c} = (c_1, \dots, c_n)^T, \quad \mathbf{d} = (d_1, \dots, d_m)^T, \quad \mathbf{t} = (t_1, \dots, t_m)^T,$$

$$\boldsymbol{\delta} = (\delta_1, \dots, \delta_n)^T, \quad \boldsymbol{\Delta} = (\Delta_1, \dots, \Delta_m)^T, \quad \boldsymbol{\epsilon} = (\epsilon_1, \dots, \epsilon_m)^T,$$

where T denotes the transpose of a vector (or matrix).

10 For storing the information which sonde measurement contributes to which match events, we define the following $m \times n$ matrix \mathbf{M} (“Match matrix”), each row of which corresponds to a match event and each column of which corresponds to an ozone observation:

$$m_{ik} = \begin{cases} 1 & \text{if the second sonde of the } i\text{-th match is sonde } k, \\ -1 & \text{if the first sonde of the } i\text{-th match is sonde } k, \\ 0 & \text{if the } i\text{-th match does not use sonde } k. \end{cases} \quad (1)$$

15 For example, the matrix describing a first match between the ozone observations number 1 and 2 and a second match between the ozone observations number 2 and 3 is

$$\mathbf{M} = \begin{pmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{pmatrix}. \quad (2)$$

Precision of the Match method

R. Lehmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

The ozone mixing ratio difference d_i associated with the i -th match event is the difference between two ozone mixing ratios measured by the second and by the first sonde of this match event:

$$d_i = c_{k_2(i)} - c_{k_1(i)} \quad (3)$$

5 or, in matrix form,

$$\mathbf{d} = \mathbf{M} \cdot \mathbf{c} .$$

The ozone mixing ratio difference d_i deviates from the ozone loss $\bar{r} \cdot t_i$ expected from the mean loss rate by the total match error e_i :

$$d_i = \bar{r} \cdot t_i + e_i . \quad (4)$$

10 The total match error e_i may be expressed by the errors of the ozone sonde measurements and the net match error of the i -th match as follows:

$$e_i = \delta_{k_2(i)} - \delta_{k_1(i)} + \Delta_i . \quad (5)$$

Taking into account the definition of the Match matrix \mathbf{M} , we may write Eqs. (4) and (5) in vector form:

$$15 \quad \mathbf{d} = \bar{r} \cdot \mathbf{t} + \mathbf{e} , \quad (6)$$

$$\mathbf{e} = \mathbf{M} \cdot \boldsymbol{\delta} + \boldsymbol{\Delta} . \quad (7)$$

We assume that the errors δ_k of all individual ozone observations and the net match errors Δ_j are unbiased, i.e. do not comprise a systematic error:

$$\mathbf{E}(\boldsymbol{\delta}) = \mathbf{0} , \quad (8)$$

$$20 \quad \mathbf{E}(\boldsymbol{\Delta}) = \mathbf{0} , \quad (9)$$

where $\mathbf{E}(\cdot)$ denotes the expected value of a random vector, and $\mathbf{0}$ is the null vector of appropriate size. From these two equations and Eq. (7) it follows that the total match errors are also unbiased:

$$\mathbf{E}(\mathbf{e}) = \mathbf{0} . \quad (10)$$

Precision of the Match method

R. Lehmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

We assume that the errors δ_k of all individual ozone observations are statistically independent and their variances are identical, namely σ_δ^2 . Analogously, the net match errors Δ_i are assumed to be statistically independent of each other and independent of the measurement errors δ_k (cf. Sect. 3.2); their variance, which is assumed to be independent of i , is denoted by σ_Δ^2 . Then the corresponding covariance matrices can be written as:

$$\text{Cov}(\boldsymbol{\delta}) = \sigma_\delta^2 \cdot \mathbf{I}, \quad (11)$$

$$\text{Cov}(\boldsymbol{\Delta}) = \sigma_\Delta^2 \cdot \mathbf{I}, \quad (12)$$

where the matrices \mathbf{I} are identity matrices of appropriate size ($n \times n$ or $m \times m$). From these two equations and the statistical independence of the errors δ_k , $k = 1, \dots, n$, and Δ_i , $i = 1, \dots, m$, we can obtain an expression for the covariance matrix of the total match errors:

$$\begin{aligned} \text{Cov}(\mathbf{e}) &= \mathbf{M} \cdot \text{Cov}(\boldsymbol{\delta}) \cdot \mathbf{M}^T + \text{Cov}(\boldsymbol{\Delta}), && \text{because of Eq. (7),} \\ &= \mathbf{M} \cdot \sigma_\delta^2 \cdot \mathbf{I} \cdot \mathbf{M}^T + \sigma_\Delta^2 \cdot \mathbf{I}, && \text{because of Eqs. (11) and (12),} \\ &= \sigma_\delta^2 \cdot \mathbf{M} \cdot \mathbf{M}^T + \sigma_\Delta^2 \cdot \mathbf{I}. \end{aligned} \quad (13)$$

The elements μ_{ij} of the matrix $\mathbf{M} \cdot \mathbf{M}^T$ are:

$$\mu_{ij} = \begin{cases} 2 & \text{if } i = j, \\ 1 & \text{if the matches } i \text{ and } j \text{ (} j \neq i \text{) have a common first or a common second sonde,} \\ -1 & \text{if the second sonde of match } i \text{ is the first sonde of match } j \text{ (} j \neq i \text{) or vice versa,} \\ 0 & \text{else.} \end{cases}$$

For the example in Eq. (2) we obtain

$$\mathbf{M} \cdot \mathbf{M}^T = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}. \quad (14)$$

In order to show the effect of the off-diagonal elements of $\mathbf{M} \cdot \mathbf{M}^T$ more clearly, we split this matrix into two matrices containing the diagonal and the off-diagonal elements of $\mathbf{M} \cdot \mathbf{M}^T$, respectively:

$$\mathbf{M} \cdot \mathbf{M}^T = 2 \cdot \mathbf{I} + \mathbf{\Omega} , \quad (15)$$

5 where the elements ω_{ij} of the matrix $\mathbf{\Omega}$ are:

$$\omega_{ij} = \begin{cases} 1 & \text{if the matches } i \text{ and } j \text{ (} j \neq i \text{) have a common first or a common} \\ & \text{second sonde,} \\ -1 & \text{if the second sonde of match } i \text{ is the first sonde of match } j \text{ (} j \neq i \text{)} \\ & \text{or vice versa,} \\ 0 & \text{else.} \end{cases} \quad (16)$$

From this definition it is evident that $\mathbf{\Omega}$ is symmetric. Replacing $\mathbf{M} \cdot \mathbf{M}^T$ in Eq. (13) by the expression in Eq. (15), we obtain:

$$10 \text{ Cov}(\boldsymbol{\epsilon}) = (\sigma_{\Delta}^2 + 2\sigma_{\delta}^2) \cdot \mathbf{I} + \sigma_{\delta}^2 \cdot \mathbf{\Omega} . \quad (17)$$

After defining

$$\sigma^2 = \sigma_{\Delta}^2 + 2\sigma_{\delta}^2 , \quad (18)$$

we can write Eq. (17) in the form

$$\text{Cov}(\boldsymbol{\epsilon}) = \sigma^2 \cdot \mathbf{I} + \sigma_{\delta}^2 \cdot \mathbf{\Omega} . \quad (19)$$

15 5. The linear regression for calculating the ozone loss rate

In order to obtain an estimate \hat{r} of the mean ozone loss rate \bar{r} , a linear regression of the ozone mixing ratio differences d_i on the corresponding sunlit times t_i is performed. The fit is forced to pass through the origin (0, 0) of the coordinate system, because in

Precision of the Match method

R. Lehmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

a time interval of zero length no ozone destruction can occur. This linear regression analysis determines the value of r that minimizes the expression

$$\sum_{i=1}^m (d_i - r \cdot t_i)^2 = (\mathbf{d} - r \cdot \mathbf{t})^T (\mathbf{d} - r \cdot \mathbf{t}) . \quad (20)$$

The solution \hat{r} of this minimization problem is

$$\hat{r} = \frac{\sum_{i=1}^m t_i \cdot d_i}{\sum_{i=1}^m t_i^2} = \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t}^T \mathbf{d} . \quad (21)$$

Together with Eq. (6) this yields

$$\begin{aligned} \hat{r} &= \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t}^T \cdot (\bar{r} \cdot \mathbf{t} + \boldsymbol{\epsilon}) \\ &= \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t}^T \mathbf{t} \cdot \bar{r} + \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t}^T \cdot \boldsymbol{\epsilon} \\ &= \bar{r} + \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t}^T \cdot \boldsymbol{\epsilon} . \end{aligned} \quad (22)$$

This proves that the estimate \hat{r} is unbiased, because $E(\boldsymbol{\epsilon}) = \mathbf{0}$ according to Eq. (10). Furthermore, it leads to the following equation for the variance of the estimate \hat{r} :

$$\begin{aligned} \sigma_{\hat{r}}^2 &= \left(\frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t}^T \right) \cdot \text{Cov}(\boldsymbol{\epsilon}) \cdot \left(\frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t}^T \right)^T \\ &= \left(\frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t}^T \right) \cdot \left(\sigma^2 \cdot \mathbf{I} + \sigma_{\delta}^2 \cdot \boldsymbol{\Omega} \right) \cdot \left(\frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t}^T \right)^T , \quad \text{because of Eq. (19),} \end{aligned}$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

$$= \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \sigma^2 + \frac{\mathbf{t}^T \boldsymbol{\Omega} \mathbf{t}}{(\mathbf{t}^T \mathbf{t})^2} \cdot \sigma_\delta^2. \quad (23)$$

After defining

$$\omega = \frac{\mathbf{t}^T \boldsymbol{\Omega} \mathbf{t}}{\mathbf{t}^T \mathbf{t}} = \frac{\sum_{ij} \omega_{ij} \cdot t_i \cdot t_j}{\sum_{i=1}^m t_i^2}, \quad (24)$$

we can rewrite Eq. (23) in the form

$$\sigma_f^2 = \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot (\sigma^2 + \omega \cdot \sigma_\delta^2). \quad (25)$$

6. Estimation of uncertainties

The variances $\sigma^2 = \sigma_\Delta^2 + 2 \cdot \sigma_\delta^2$ and σ_δ^2 in Eq. (25) are unknown. In this section we are going to derive estimates for them.

6.1. Information on σ^2

Taking into account that the diagonal elements of $\boldsymbol{\Omega}$ are zero, we see from Eq. (19) that the diagonal elements of $\text{Cov}(\boldsymbol{\epsilon})$ are σ^2 , i.e. together with Eq. (10) we obtain

$$E(e_i^2) = \sigma^2, \quad (26)$$

and thus, because of Eq. (4),

$$E\left\{(d_i - \bar{r} \cdot t_i)^2\right\} = \sigma^2. \quad (27)$$

This means that $(d_i - \bar{r} \cdot t_i)^2$, $i = 1, \dots, m$, are unbiased estimates of σ^2 . Consequently, the expression $\frac{1}{m} \cdot \sum_{i=1}^m (d_i - \bar{r} \cdot t_i)^2$ is also an unbiased estimate of σ^2 , but with a

smaller variance than the individual terms. As \bar{r} is unknown, we have to replace it by the estimate \hat{r} . This has the consequence that the arising estimate for σ^2 will no longer be unbiased. However, it should still contain much information on σ^2 . That is why we consider the following sum and calculate its expected value:

$$s_1 = \sum_{i=1}^m (d_i - \hat{r} \cdot t_i)^2 . \quad (28)$$

The sum s_1 is also known as “chi-square” (χ^2). After recalling the definition of ω in Eq. (24), we can write the expected value of s_1 as follows (see Appendix B):

$$E(s_1) = (m - 1) \cdot \sigma^2 - \omega \cdot \sigma_\delta^2 . \quad (29)$$

For the data analysed in Sects. 8.1–8.3, the mean values of $m - 1$ and ω are 41.3 and 1.1, respectively. This means that $E(s_1)$ is dominated by the term containing σ^2 .

6.2. Special case: $\sigma_\delta = 0$

If $\sigma_\delta = 0$, i.e. if the sonde measurement errors vanish and thus the (total) match errors are uncorrelated, then Eq. (29) reduces to

$$E(s_1) = (m - 1) \cdot \sigma^2 . \quad (30)$$

This means that an unbiased estimate \hat{s}^2 of σ^2 can be obtained from

$$\hat{s}^2 = \frac{s_1}{m - 1} . \quad (31)$$

Substituting this value for σ^2 in Eq. (25), we obtain the “classical” error estimate

$$\hat{s}_{\hat{r}}^2 = \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \frac{s_1}{m - 1} = \frac{1}{m - 1} \cdot \frac{\sum_{i=1}^m (d_i - \hat{r} \cdot t_i)^2}{\sum_{i=1}^m t_i^2} . \quad (32)$$

Precision of the Match method

R. Lehmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

6.3. Special case: $\sigma_{\Delta} = 0$

If $\sigma_{\Delta} = 0$, i.e. if the net match errors vanish, then $\sigma^2 = 2 \cdot \sigma_{\delta}^2$, and Eq. (29) reduces to

$$E(s_1) = \{2 \cdot (m - 1) - \omega\} \cdot \sigma_{\delta}^2. \quad (33)$$

Thus an unbiased estimate $\hat{\sigma}_{\delta}^2$ of σ_{δ}^2 can be obtained from

$$\hat{\sigma}_{\delta}^2 = \frac{s_1}{2 \cdot (m - 1) - \omega}. \quad (34)$$

Substituting this value for σ_{δ}^2 in Eq. (25), we obtain the error estimate

$$\begin{aligned} \hat{\sigma}_f^2 &= \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot (2 + \omega) \cdot \frac{s_1}{2 \cdot (m - 1) - \omega} \\ &= \frac{1 + \frac{\omega}{2}}{m - 1 - \frac{\omega}{2}} \cdot \frac{\sum_{i=1}^m (d_i - \hat{r} \cdot t_i)^2}{\sum_{i=1}^m t_i^2}, \end{aligned} \quad (35)$$

This is identical to Eq. (32) if $\omega = 0$, e.g. if the covariance matrix $\text{Cov}(\mathbf{e})$ is diagonal and hence the match errors are uncorrelated.

For the derivation of Eq. (35) the effect of the correlation between the ozone data used in different match events, due to the multiple use of the same ozone measurements in several matches, has been taken into account. As the estimate $\hat{\sigma}_f^2$ in Eq. (35) has been derived under the assumption $\sigma_{\Delta}^2 = 0$, i.e. the ozone measurement errors δ_i alone determine the total match error, it may provide an upper bound of the effect of taking into account the above-mentioned correlation. For the data analysed in Sects. 8.1–8.3, the error bar $\hat{\sigma}_f$ from Eq. (35) is, on average, 24% larger than the “classical” estimate according to Eq. (32).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

6.4. Information on σ_δ^2

If we compare the off-diagonal elements in Eq. (19), we obtain for $i \neq j$:

$$E(\epsilon_i \cdot \epsilon_j) = \omega_{ij} \cdot \sigma_\delta^2, \quad (36)$$

and thus, because of Eq. (4),

$$E\{(d_i - \bar{r} \cdot t_i) \cdot (d_j - \bar{r} \cdot t_j)\} = \omega_{ij} \cdot \sigma_\delta^2. \quad (37)$$

If $\omega_{ij} \neq 0$, then $\omega_{ij} = 1$ or $\omega_{ij} = -1$ and consequently $\omega_{ij}^2 = 1$, so that we obtain from Eq. (37) by multiplication by ω_{ij} :

$$E\{\omega_{ij} \cdot (d_i - \bar{r} \cdot t_i) \cdot (d_j - \bar{r} \cdot t_j)\} = \sigma_\delta^2. \quad (38)$$

Now we proceed as in Sect. 6.1: Eq. (38) means that $\omega_{ij} \cdot (d_i - \bar{r} \cdot t_i) \cdot (d_j - \bar{r} \cdot t_j)$, with i, j such that $\omega_{ij} \neq 0$, are unbiased estimates of σ_δ^2 . Consequently, the arithmetic mean of these expressions is also an unbiased estimate of σ_δ^2 , but with a smaller variance than the individual terms. As \bar{r} is unknown, we have to replace it by the estimate \hat{r} . This has the consequence that the arising estimate for σ_δ^2 will no longer be unbiased. However, it should still contain much information on σ_δ^2 . That is why we consider the following sum and calculate its expected value:

$$s_2 = \sum_{i=1}^m \sum_{j=1(\omega_{ij} \neq 0)}^m \omega_{ij} \cdot (d_i - \hat{r} \cdot t_i) \cdot (d_j - \hat{r} \cdot t_j). \quad (39)$$

As adding the zero terms corresponding to $\omega_{ij} = 0$ does not alter this sum, we can write s_2 also in the form

$$s_2 = \sum_{i=1}^m \sum_{j=1}^m \omega_{ij} \cdot (d_i - \hat{r} \cdot t_i) \cdot (d_j - \hat{r} \cdot t_j). \quad (40)$$

After defining

$$\omega_1 = \sum_{i=1}^m \sum_{j=1}^m \omega_{ij}^2, \quad (41)$$

$$\omega_2 = \frac{(\mathbf{\Omega} \cdot \mathbf{t})^T \cdot (\mathbf{\Omega} \cdot \mathbf{t})}{\mathbf{t}^T \mathbf{t}}, \quad (42)$$

we can obtain the following expression for the expected value of s_2 (see Appendix C):

$$E(s_2) = -\omega \cdot \sigma^2 + (\omega_1 - 2 \cdot \omega_2 + \omega^2) \cdot \sigma_\delta^2. \quad (43)$$

As $\omega_{ij} = 0$ or $\omega_{ij} = \pm 1$, $i, j = 1, \dots, m$, the value ω_1 is equal to the number of non-zero elements of the matrix $\mathbf{\Omega}$. For the data analysed in Sects. 8.1–8.3, the mean values of ω_1 , ω_2 , and ω , are 179, 5.5, and 1.1, respectively. This means that $E(s_2)$ is dominated by the term containing σ_δ^2 .

6.5. Estimates for σ^2 and σ_δ^2

If we define the 2×2 matrix \mathbf{A} by

$$\mathbf{A} = \begin{pmatrix} m-1 & -\omega \\ -\omega & \omega_1 - 2 \cdot \omega_2 + \omega^2 \end{pmatrix}, \quad (44)$$

then Eqs. (29) and (43) may be written as

$$E \begin{pmatrix} s_1 \\ s_2 \end{pmatrix} = \mathbf{A} \cdot \begin{pmatrix} \sigma^2 \\ \sigma_\delta^2 \end{pmatrix}. \quad (45)$$

Consequently,

$$E \left\{ \mathbf{A}^{-1} \cdot \begin{pmatrix} s_1 \\ s_2 \end{pmatrix} \right\} = \mathbf{A}^{-1} \cdot \mathbf{A} \cdot \begin{pmatrix} \sigma^2 \\ \sigma_\delta^2 \end{pmatrix} = \begin{pmatrix} \sigma^2 \\ \sigma_\delta^2 \end{pmatrix}. \quad (46)$$

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

This means that \hat{s}^2 and \hat{s}_δ^2 given by

$$\begin{pmatrix} \hat{s}^2 \\ \hat{s}_\delta^2 \end{pmatrix} = \mathbf{A}^{-1} \cdot \begin{pmatrix} s_1 \\ s_2 \end{pmatrix} \quad (47)$$

are unbiased estimates of σ^2 and σ_δ^2 , respectively. The inverse of the matrix \mathbf{A} is

$$\mathbf{A}^{-1} = \frac{1}{D} \cdot \begin{pmatrix} \omega_1 - 2 \cdot \omega_2 + \omega^2 & \omega \\ \omega & m - 1 \end{pmatrix} \quad (48)$$

5 with

$$D = (m - 1) \cdot (\omega_1 - 2 \cdot \omega_2 + \omega^2) - \omega^2 = (m - 1) \cdot (\omega_1 - 2 \cdot \omega_2) + (m - 2) \cdot \omega^2 \quad (49)$$

Thus we finally obtain from Eq. (47):

$$\hat{s}^2 = \frac{1}{D} \cdot \left\{ (\omega_1 - 2 \cdot \omega_2 + \omega^2) \cdot s_1 + \omega \cdot s_2 \right\}, \quad (50)$$

$$\hat{s}_\delta^2 = \frac{1}{D} \cdot \left\{ \omega \cdot s_1 + (m - 1) \cdot s_2 \right\}. \quad (51)$$

10 6.6. Estimate for σ_f^2

If we substitute the estimates \hat{s}^2 for σ^2 and \hat{s}_δ^2 for σ_δ^2 from Eqs. (50), (51) in Eq. (25), then we obtain the desired estimate \hat{s}_f^2 of σ_f^2 :

$$\begin{aligned} \hat{s}_f^2 &= \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \left(\hat{s}^2 + \omega \cdot \hat{s}_\delta^2 \right) \\ &= \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \frac{1}{D} \cdot \left\{ (\omega_1 - 2 \cdot \omega_2 + \omega^2) \cdot s_1 + \omega \cdot s_2 + \omega^2 \cdot s_1 + (m - 1) \cdot \omega \cdot s_2 \right\} \\ &= \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \frac{1}{D} \cdot \left\{ (\omega_1 - 2 \cdot \omega_2 + 2 \cdot \omega^2) \cdot s_1 + m \cdot \omega \cdot s_2 \right\}. \end{aligned} \quad (52)$$

15

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

It might be desirable to calculate the standard deviation of this estimate, i.e. the “error bars of the error bars”, and investigate whether it is a minimum-variance estimate. However, this remains beyond the scope of the present paper.

6.7. Avoidance of unfeasible values

5 It cannot be excluded that the estimate \hat{s}_δ^2 of the variance σ_δ^2 becomes negative. As a negative variance is unrealistic, this case will be treated by assuming $\sigma_\delta^2 = 0$ and applying Eq. (32). If on the other hand, $\hat{s}_\delta^2 > \hat{s}^2$, which corresponds to a negative estimate for the variance σ_Δ^2 , then it will be assumed that $\sigma_\Delta^2 = 0$ and Eq. (35) will be applied. These modifications might, in principle, destroy the unbiasedness of the final estimate \hat{s}_f^2 . However, for the 96 match ensembles investigated in Sects. 8.1–8.3, the introduction of the above-described sign-restrictions changed the mean value of the estimate \hat{s}_f^2 by only 0.04%.

15 In order to prevent a division by zero in Eqs. (50) and (51), the complete algorithm in the next section will check whether D according to Eq. (49) is zero. If this is the case, then the “classical” error estimate Eq. (32) will be applied. This might occur, e.g., if $\Omega = 0$, i.e. if all match events use independent sonde measurements, in which case the application of Eq. (32) yields the correct estimate. However, in the examples of Sect. 8 this never occurred.

7. Complete formulae for estimating the precision of Match

20 Summarising the results of the previous sections, we obtain the following algorithm for the calculation of estimates for the precision of the Match results. For these estimates we will also use the more illustrative denotation “error bars”.

**Precision of the
Match method**

R. Lehmann et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Print Version](#)
[Interactive Discussion](#)
Input: m = number of match events, d_i = difference of the ozone mixing ratio (second minus first sonde measurement) of the i -th match, $i = 1, \dots, m$, t_i = sunlit time of the i -th match, $i = 1, \dots, m$,5 $k_1(i)$ = identifier of the first sonde of the i -th match, $i = 1, \dots, m$, $k_2(i)$ = identifier of the second sonde of the i -th match, $i = 1, \dots, m$,Output: \hat{r} = estimated ozone loss rate, \hat{s}_r^2 = estimated variance of \hat{r} .10 Algorithm:

1. Calculate an estimate of the ozone loss rate by linear regression:

$$\hat{r} = \frac{\sum_{i=1}^m t_i \cdot d_i}{\sum_{i=1}^m t_i^2}.$$

15 2. Set up the matrix Ω that stores the information on ozone sondes shared by two or more match events:

$$\omega_{ij} = \begin{cases} 1 & \text{if } k_1(i) = k_1(j) \text{ or } k_2(i) = k_2(j), \text{ and } i \neq j, \\ -1 & \text{if } k_2(i) = k_1(j) \text{ or } k_1(i) = k_2(j), \text{ and } i \neq j, \\ 0 & \text{else.} \end{cases}$$

3. Calculate the auxiliary expressions s_1 , s_2 , ω , ω_1 , ω_2 , and D :

$$s_1 = \sum_{i=1}^m (d_i - \hat{r} \cdot t_i)^2, \quad (53)$$

$$s_2 = \sum_{i=1}^m \sum_{j=1}^m \omega_{ij} \cdot (d_i - \hat{r} \cdot t_i) \cdot (d_j - \hat{r} \cdot t_j), \quad (54)$$

$$\omega = \frac{\sum_{ij} \omega_{ij} \cdot t_i \cdot t_j}{\sum_{i=1}^m t_i^2}, \quad (55)$$

$$\omega_1 = \sum_{i=1}^m \sum_{j=1}^m \omega_{ij}^2, \quad (56)$$

$$\omega_2 = \frac{(\mathbf{\Omega} \cdot \mathbf{t})^T \cdot (\mathbf{\Omega} \cdot \mathbf{t})}{\mathbf{t}^T \mathbf{t}}, \text{ where } \mathbf{t} = (t_1, \dots, t_m)^T, \quad (57)$$

$$5 \quad D = (m - 1) \cdot (\omega_1 - 2 \cdot \omega_2) + (m - 2) \cdot \omega^2. \quad (58)$$

4. If $D = 0$, then use the “classical” error bars:
If $D = 0$, then goto step 8.

5. Calculate estimates \hat{s}^2 for σ^2 and \hat{s}_δ^2 for σ_δ^2 :

$$\hat{s}^2 = \frac{1}{D} \cdot \left\{ (\omega_1 - 2 \cdot \omega_2 + \omega^2) \cdot s_1 + \omega \cdot s_2 \right\}, \quad (59)$$

$$10 \quad \hat{s}_\delta^2 = \frac{1}{D} \cdot \left\{ \omega \cdot s_1 + (m - 1) \cdot s_2 \right\}. \quad (60)$$

6. Sign check for \hat{s}_δ^2 and $\hat{s}_\Delta^2 = \hat{s}^2 - 2 \cdot \hat{s}_\delta^2$:

If $\hat{s}_\delta^2 < 0$, then goto step 8.

If $\hat{s}_\delta^2 > \frac{1}{2} \cdot \hat{s}^2$, then goto step 9.

7. Calculate error bar:

$$15 \quad \hat{s}_r^2 = \frac{1}{\sum_{i=1}^m t_i^2} \cdot \left(\hat{s}^2 + \omega \cdot \hat{s}_\delta^2 \right). \quad (61)$$

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

stop.

8. "Classical" error bar (zero sonde measurement errors):

$$\hat{s}_f^2 = \frac{1}{m-1} \cdot \frac{\sum_{i=1}^m (d_i - \hat{r} \cdot t_i)^2}{\sum_{i=1}^m t_i^2} \quad (62)$$

stop.

5 9. Error bar for zero net match errors, i.e. only sonde measurement errors occur:

$$\hat{s}_f^2 = \frac{1 + \frac{\omega}{2}}{m-1 - \frac{\omega}{2}} \cdot \frac{\sum_{i=1}^m (d_i - \hat{r} \cdot t_i)^2}{\sum_{i=1}^m t_i^2} \quad (63)$$

stop.

8. Examples

10 8.1. Data: ozone-sonde Match campaigns

The analysis in the following Sects. 8.2 and 8.3 is based on all data from the Arctic Match campaigns of the winters 1994/95 (potential temperature levels 450 K, 475 K, 500 K; Rex et al., 1999), 1995/96 (475 K; Rex et al., 1997), 1999/2000 (450 K, 475 K, 500 K; Rex et al., 2002), and 2002/03 (475 K; Streibel et al., 2005). These data correspond to 96 match ensembles, i.e. the calculation of 96 ozone loss rates by the application of a linear regression like in Fig. 1.

8.2. Estimates of the measurement errors and the net match errors

For the data introduced in the previous subsection, the mean values of the estimates \hat{s}_σ^2 and \hat{s}_Δ^2 are $2.4 \cdot 10^4 \text{ ppb}^2$ and $2.7 \cdot 10^4 \text{ ppb}^2$, respectively. This means that the errors of the ozone measurements and the net match errors are of the same order of magnitude.

20

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Precision of the Match method

R. Lehmann et al.

Extracting the square root from the mean value of $\hat{\delta}_\delta^2$ yields $\hat{\delta}_\delta = 156$ ppb, which corresponds to 6% of the mean ozone mixing ratio (of all sonde measurements entering the mentioned matches). This is consistent with the precision of ozone sonde measurements of 5.7%, obtained during the Jülich Ozone Sonde Intercomparison Experiment (quadratic mean of the precision values for SPC-6A and ENSCI sondes with a 1% KI solution in the height range of 15–25 km given by Smit and Straeter, 2004a, Table 9, and Smit and Straeter, 2004b, Table 14).

8.3. Comparison of new and old error bars

Up to now the “classical” error bars, according to Eq. (62), have been used in the Match analysis. Figure 7 shows both the new error bars, according to $\hat{\delta}_r$ in Eq. (61), and the old ones. It can be seen that the new error bars are slightly larger on average. For ozone loss rates greater than approximately 2 ppb/h, the loss rates are greater than the old error bars, so that the ozone loss can be considered significant. This does not change when the new error bars are used instead of the old ones.

Figure 8 displays the ratio of the new error bars to the old ones. It varies between 0.96 and 1.68. The 90%-quantile is 1.32, i.e. for 90% of the data points the ratio is less than 1.32. The mean value of the ratio is 1.15.

On average every ozone sonde measurement was used in slightly more than 2 match events (see triangular arrowhead in Fig. 8). In order to express statements like the latter one more concisely, we introduce the following term:

oversampling rate := average number of matches to which an ozone measurement contributes.

The new error bars have been constructed, in order to account for the multiple use of ozone sonde measurements in several match events. It can be expected that, on average, the new error bars deviate more from the old ones if the oversampling rate increases. This is indeed the case, as can be seen from the regression line added in Fig. 8. The slope of this line is 0.12, i.e. an increase of the oversampling rate by 1 results in an average increase of the ratio “new error bar/old error bar” by 12%. The

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

regression line almost crosses the point (1,1). This means that the old and new error bars coincide if each sonde measurement is used in only one match event, i.e. if all sondes used in the matches are different from each other. This is also an expected result.

5 8.4. Data: satellite Match study

In order to test the formulae of Sect. 7 for larger oversampling rates, i.e. in order to extend Fig. 8 to the right, we consider an additional Match study, which is based on satellite data. The analysis in the following Sect. 8.5 uses data from an Antarctic Match study based on ozone observations by the Polar Ozone and Aerosol Measurement
10 III (POAM III) instrument in 2003 (potential temperature level 475 K). These data correspond to 15 match ensembles, i.e. the calculation of 15 ozone loss rates by the application of a linear regression like in Fig. 1.

8.5. Comparison of new and old error bars

For the satellite Match study, the oversampling rate is approximately 5 (cf. Fig. 9), i.e.
15 it is significantly larger than for the ozone-sonde Match campaigns. An extrapolation of the results of Fig. 8 suggests that this leads to larger ratios of the new errors bars to the old ones. As can be seen in Fig. 9, this is indeed the case. The slope of the regression line added in Fig. 9 is 0.14, i.e. an increase of the oversampling rate by 1 results in an average increase of the ratio “new error bar/old error bar” by 14%, which
20 is rather similar to the corresponding value for the ozone-sonde Match campaigns in Sect. 8.3. For the satellite Match study, the ratio of the new error bars to the old ones varies between 1.1 and 2.8, the mean value is 1.6.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

9. Conclusions

A detailed analysis of the random errors of the ozone loss rate calculated by the Match method has been presented. It differs from a standard analysis by taking into account that the same ozone sonde measurement may be used in several matches, so that the ozone mixing ratio differences (second minus first sonde measurement) of these matches become statistically dependent. For four Arctic ozone-sonde Match campaigns, this effect leads to changes of the error bars between -4% and $+68\%$. On average, the error bars increase by 15% . This does not change the conclusions about the statistical significance of the ozone loss rates observed. For an Antarctic satellite Match study, the error bars increase by 10% to 180% , on average by 60% .

Appendix A General matrix identities

Let us assume that $\mathbf{e}_i \in R^m$ denotes the i -th unit vector, i.e. its i -th element is 1, all other elements are zero. Then we obtain:

$$\sum_{i=1}^m \mathbf{e}_i \cdot \mathbf{e}_i^T = \mathbf{I} . \quad (64)$$

This can be easily proven, because $\mathbf{e}_i \cdot \mathbf{e}_i^T$ is an $m \times m$ matrix, the i -th diagonal element of which is 1, all other elements are zero.

Let us assume that \mathbf{A} is an $m \times m$ matrix, $\mathbf{a}, \mathbf{b} \in R^m$ are vectors. Then we obtain:

$$\begin{aligned} \sum_{i=1}^m \mathbf{e}_i^T \cdot \mathbf{A} \cdot \mathbf{a} \cdot \mathbf{b}^T \cdot \mathbf{e}_i &= \sum_{i=1}^m \mathbf{b}^T \cdot \mathbf{e}_i \cdot \mathbf{e}_i^T \cdot \mathbf{A} \cdot \mathbf{a} , \text{ because } \mathbf{b}^T \cdot \mathbf{e}_i \text{ is a real number,} \\ &= \mathbf{b}^T \cdot \left(\sum_{i=1}^m \mathbf{e}_i \cdot \mathbf{e}_i^T \right) \cdot \mathbf{A} \cdot \mathbf{a} \end{aligned}$$

3251

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

$$= \mathbf{b}^T \cdot \mathbf{A} \cdot \mathbf{a} , \text{ because of Eq. (64).} \quad (65)$$

Further we obtain:

$$\begin{aligned} \sum_{i=1}^m \mathbf{e}_i^T \cdot \mathbf{A} \cdot \mathbf{a} \cdot \mathbf{b}^T \cdot \mathbf{A} \cdot \mathbf{e}_i &= \sum_{i=1}^m \mathbf{e}_i^T \cdot \mathbf{A} \cdot \mathbf{a} \cdot (\mathbf{A}^T \cdot \mathbf{b})^T \cdot \mathbf{e}_i \\ &= (\mathbf{A}^T \cdot \mathbf{b})^T \cdot \mathbf{A} \cdot \mathbf{a} , \text{ because of Eq. (65) with } \mathbf{b} \hat{=} \mathbf{A}^T \cdot \mathbf{b} , \\ &= \mathbf{b}^T \cdot \mathbf{A}^2 \cdot \mathbf{a} . \end{aligned} \quad (66)$$

Appendix B Calculation of $E(s_1)$

The sum s_1 defined in Eq. (28) can be written in vector notation as

$$s_1 = (\mathbf{d} - \hat{\mathbf{r}} \cdot \mathbf{t})^T \cdot (\mathbf{d} - \hat{\mathbf{r}} \cdot \mathbf{t}) . \quad (67)$$

The term $(\mathbf{d} - \hat{\mathbf{r}} \cdot \mathbf{t})$ occurring in this expression may be transformed as follows:

$$\begin{aligned} \mathbf{d} - \hat{\mathbf{r}} \cdot \mathbf{t} &= \mathbf{d} - \mathbf{t} \cdot \hat{\mathbf{r}} \\ &= \mathbf{d} - \mathbf{t} \cdot \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t}^T \mathbf{d} , \text{ because of Eq. (21),} \\ &= \left(\mathbf{I} - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t} \cdot \mathbf{t}^T \right) \cdot \mathbf{d} . \end{aligned} \quad (68)$$

Let us define the matrix

$$\mathbf{J} = \mathbf{I} - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t} \cdot \mathbf{t}^T . \quad (69)$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

It is symmetric and fulfills the following equations:

$$\mathbf{J} \cdot \mathbf{t} = \left(\mathbf{I} - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t} \cdot \mathbf{t}^T \right) \cdot \mathbf{t} = \mathbf{t} - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t} \cdot \mathbf{t}^T \mathbf{t} = \mathbf{t} - \mathbf{t} = \mathbf{0} , \quad (70)$$

$$\mathbf{J}^T \cdot \mathbf{J} = \mathbf{J} \cdot \mathbf{J} = \mathbf{J} \cdot \left(\mathbf{I} - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t} \cdot \mathbf{t}^T \right) = \mathbf{J} - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot (\mathbf{J} \cdot \mathbf{t}) \cdot \mathbf{t}^T = \mathbf{J} . \quad (71)$$

Now Eq. (68) can be written as

$$\begin{aligned} 5 \quad \mathbf{d} - \hat{\mathbf{r}} \cdot \mathbf{t} &= \mathbf{J} \cdot \mathbf{d} \\ &= \mathbf{J} \cdot (\bar{\mathbf{r}} \cdot \mathbf{t} + \boldsymbol{\epsilon}) , \text{ because of Eq. (6),} \\ &= \mathbf{J} \cdot \boldsymbol{\epsilon} , \text{ because of Eq. (70).} \end{aligned} \quad (72)$$

Then the expected value of s_1 defined in Eq. (67) can be calculated:

$$\begin{aligned} 10 \quad E(s_1) &= E \left((\mathbf{d} - \hat{\mathbf{r}} \cdot \mathbf{t})^T \cdot (\mathbf{d} - \hat{\mathbf{r}} \cdot \mathbf{t}) \right) \\ &= E \left(\boldsymbol{\epsilon}^T \cdot \mathbf{J}^T \cdot \mathbf{J} \cdot \boldsymbol{\epsilon} \right) , \text{ because of Eq. (72),} \\ &= E \left(\boldsymbol{\epsilon}^T \cdot \mathbf{J} \cdot \boldsymbol{\epsilon} \right) , \text{ because of Eq. (71),} \\ &= E \left\{ \boldsymbol{\epsilon}^T \cdot \left(\mathbf{I} - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t} \cdot \mathbf{t}^T \right) \cdot \boldsymbol{\epsilon} \right\} \\ &= E \left(\boldsymbol{\epsilon}^T \boldsymbol{\epsilon} \right) - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot E \left(\boldsymbol{\epsilon}^T \cdot \mathbf{t} \cdot \mathbf{t}^T \cdot \boldsymbol{\epsilon} \right) \\ 15 \quad &= E \left(\sum_{i=1}^m \epsilon_i^2 \right) - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot E \left(\mathbf{t}^T \cdot \boldsymbol{\epsilon} \cdot \boldsymbol{\epsilon}^T \cdot \mathbf{t} \right) \quad \text{because } \boldsymbol{\epsilon}^T \cdot \mathbf{t} = \mathbf{t}^T \cdot \boldsymbol{\epsilon} \\ &= \left(\sum_{i=1}^m E \left(\epsilon_i^2 \right) \right) - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t}^T \cdot E \left(\boldsymbol{\epsilon} \cdot \boldsymbol{\epsilon}^T \right) \cdot \mathbf{t} \end{aligned}$$

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Print Version](#)
[Interactive Discussion](#)

$$\begin{aligned}
 &= \left(\sum_{i=1}^m \text{Var}(\epsilon_i) \right) - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t}^T \cdot \text{Cov}(\boldsymbol{\epsilon}) \cdot \mathbf{t}, \quad \text{because of } E(\boldsymbol{\epsilon}) = \mathbf{0}, \\
 &= \left(\sum_{i=1}^m \sigma^2 \right) - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t}^T \cdot \left(\sigma^2 \cdot \mathbf{I} + \sigma_\delta^2 \cdot \boldsymbol{\Omega} \right) \cdot \mathbf{t}, \quad \text{because of Eq. (19),} \\
 &= m \cdot \sigma^2 - \frac{\mathbf{t}^T \mathbf{t}}{\mathbf{t}^T \mathbf{t}} \cdot \sigma^2 - \frac{\mathbf{t}^T \boldsymbol{\Omega} \mathbf{t}}{\mathbf{t}^T \mathbf{t}} \cdot \sigma_\delta^2 \\
 &= (m - 1) \cdot \sigma^2 - \frac{\mathbf{t}^T \boldsymbol{\Omega} \mathbf{t}}{\mathbf{t}^T \mathbf{t}} \cdot \sigma_\delta^2. \tag{73}
 \end{aligned}$$

Together with the definition of ω in Eq. (24) we thus obtain

$$E(s_1) = (m - 1) \cdot \sigma^2 - \omega \cdot \sigma_\delta^2. \tag{74}$$

Appendix C Calculation of $E(s_2)$

The sum s_2 defined in Eq. (40) can be transformed to vector notation as follows:

$$\begin{aligned}
 s_2 &= \sum_{i=1}^m \sum_{j=1}^m \omega_{ij} \cdot (d_i - \hat{r} \cdot t_i) \cdot (d_j - \hat{r} \cdot t_j) \\
 &= \sum_{i=1}^m \sum_{j=1}^m \mathbf{e}_i^T \boldsymbol{\Omega} \mathbf{e}_j \cdot \mathbf{e}_i^T (\mathbf{d} - \hat{r} \cdot \mathbf{t}) \cdot \mathbf{e}_j^T (\mathbf{d} - \hat{r} \cdot \mathbf{t}) \\
 &= \sum_{i=1}^m \sum_{j=1}^m \mathbf{e}_i^T \boldsymbol{\Omega} \mathbf{e}_j \cdot \mathbf{e}_j^T (\mathbf{d} - \hat{r} \cdot \mathbf{t}) \cdot (\mathbf{d} - \hat{r} \cdot \mathbf{t})^T \mathbf{e}_i
 \end{aligned}$$

$$\begin{aligned}
 &= \sum_{i=1}^m \mathbf{e}_i^T \boldsymbol{\Omega} \cdot \left(\sum_{j=1}^m \mathbf{e}_j \cdot \mathbf{e}_j^T \right) \cdot (\mathbf{d} - \hat{\mathbf{r}} \cdot \mathbf{t}) \cdot (\mathbf{d} - \hat{\mathbf{r}} \cdot \mathbf{t})^T \mathbf{e}_i \\
 &= \sum_{i=1}^m \mathbf{e}_i^T \boldsymbol{\Omega} \cdot (\mathbf{d} - \hat{\mathbf{r}} \cdot \mathbf{t}) \cdot (\mathbf{d} - \hat{\mathbf{r}} \cdot \mathbf{t})^T \mathbf{e}_i, \text{ because of Eq. (64)}.
 \end{aligned}$$

Thus we obtain:

$$\begin{aligned}
 E(S_2) &= \sum_{i=1}^m \mathbf{e}_i^T \boldsymbol{\Omega} \cdot E \left\{ (\mathbf{d} - \hat{\mathbf{r}} \cdot \mathbf{t}) \cdot (\mathbf{d} - \hat{\mathbf{r}} \cdot \mathbf{t})^T \right\} \cdot \mathbf{e}_i \\
 &= \sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot E \left(\mathbf{J} \cdot \boldsymbol{\varepsilon} \cdot \boldsymbol{\varepsilon}^T \cdot \mathbf{J}^T \right) \cdot \mathbf{e}_i, \text{ because of Eq. (72),} \\
 &= \sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \mathbf{J} \cdot \text{Cov}(\boldsymbol{\varepsilon}) \cdot \mathbf{J} \cdot \mathbf{e}_i, \text{ because of } E(\boldsymbol{\varepsilon}) = \mathbf{0} \text{ and the symmetry of } \mathbf{J}, \\
 &= \sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \mathbf{J} \cdot (\sigma^2 \cdot \mathbf{I} + \sigma_\delta^2 \cdot \boldsymbol{\Omega}) \cdot \mathbf{J} \cdot \mathbf{e}_i, \text{ because of Eq. (19),} \\
 &= \left(\sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \mathbf{J} \cdot \mathbf{J} \cdot \mathbf{e}_i \right) \cdot \sigma^2 + \left(\sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \mathbf{J} \cdot \boldsymbol{\Omega} \cdot \mathbf{J} \cdot \mathbf{e}_i \right) \cdot \sigma_\delta^2. \tag{75}
 \end{aligned}$$

The expressions in front of σ^2 and σ_δ^2 are evaluated separately:

$$\begin{aligned}
 &\sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \mathbf{J} \cdot \mathbf{J} \cdot \mathbf{e}_i \\
 &= \sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \mathbf{J} \cdot \mathbf{e}_i, \text{ because of Eq. (71),}
 \end{aligned}$$

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

$$\begin{aligned}
 &= \sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \left(\mathbf{I} - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t} \cdot \mathbf{t}^T \right) \cdot \mathbf{e}_i, \text{ because of the definition of } \mathbf{J} \text{ in Eq. (69),} \\
 &= \sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \mathbf{e}_i - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \mathbf{t} \cdot \mathbf{t}^T \cdot \mathbf{e}_i \\
 &= \sum_{i=1}^m \omega_{ii} - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t}^T \cdot \boldsymbol{\Omega} \cdot \mathbf{t}, \text{ because of Eq. (65),} \\
 &= -\frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t}^T \cdot \boldsymbol{\Omega} \cdot \mathbf{t}, \text{ because all diagonal elements of } \boldsymbol{\Omega} \text{ are zero, cf. Eq. (16),} \\
 5 \quad &= -\omega, \text{ because of the definition of } \omega \text{ in Eq. (24).} \tag{76}
 \end{aligned}$$

The expression in front of σ_{δ}^2 in Eq. (75) is

$$\begin{aligned}
 &\sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \mathbf{J} \cdot \boldsymbol{\Omega} \cdot \mathbf{J} \cdot \mathbf{e}_i \\
 &= \sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \left(\mathbf{I} - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t} \cdot \mathbf{t}^T \right) \cdot \boldsymbol{\Omega} \cdot \left(\mathbf{I} - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t} \cdot \mathbf{t}^T \right) \cdot \mathbf{e}_i, \\
 &\quad \text{because of the definition of } \mathbf{J} \text{ in Eq. (69),} \\
 10 \quad &= \sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \boldsymbol{\Omega} \cdot \mathbf{e}_i - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \boldsymbol{\Omega} \cdot \mathbf{t} \cdot \mathbf{t}^T \cdot \mathbf{e}_i \\
 &\quad - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \mathbf{t} \cdot \mathbf{t}^T \cdot \boldsymbol{\Omega} \cdot \mathbf{e}_i + \frac{1}{(\mathbf{t}^T \mathbf{t})^2} \cdot \sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \mathbf{t} \cdot \mathbf{t}^T \cdot \boldsymbol{\Omega} \cdot \mathbf{t} \cdot \mathbf{t}^T \cdot \mathbf{e}_i. \tag{77}
 \end{aligned}$$

We are going to evaluate the four sums occurring in Eq. (77) separately:

$$\sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \boldsymbol{\Omega} \cdot \mathbf{e}_i = \sum_{j=1}^m (\boldsymbol{\Omega} \cdot \mathbf{e}_j)^T \cdot \boldsymbol{\Omega} \cdot \mathbf{e}_j, \text{ because } \boldsymbol{\Omega} \text{ is symmetric,}$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

$$\begin{aligned}
 &= \sum_{j=1}^m \begin{pmatrix} \omega_{1j} \\ \dots \\ \omega_{mj} \end{pmatrix}^T \cdot \begin{pmatrix} \omega_{1j} \\ \dots \\ \omega_{mj} \end{pmatrix} \\
 &= \sum_{i=1}^m \sum_{j=1}^m \omega_{ij}^2.
 \end{aligned} \tag{78}$$

$$\sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \boldsymbol{\Omega} \cdot \mathbf{t} \cdot \mathbf{t}^T \cdot \mathbf{e}_i = \mathbf{t}^T \cdot \boldsymbol{\Omega}^2 \cdot \mathbf{t}, \quad \text{because of Eq. (65)}. \tag{79}$$

$$\sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \mathbf{t} \cdot \mathbf{t}^T \cdot \boldsymbol{\Omega} \cdot \mathbf{e}_i = \mathbf{t}^T \cdot \boldsymbol{\Omega}^2 \cdot \mathbf{t}, \quad \text{because of Eq. (66)}. \tag{80}$$

$$\begin{aligned}
 \sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \mathbf{t} \cdot \mathbf{t}^T \cdot \boldsymbol{\Omega} \cdot \mathbf{t} \cdot \mathbf{t}^T \cdot \mathbf{e}_i &= \mathbf{t}^T \cdot \boldsymbol{\Omega} \cdot \mathbf{t} \cdot \mathbf{t}^T \cdot \boldsymbol{\Omega} \cdot \mathbf{t}, \quad \text{because of Eq. (65)}, \\
 &= \left(\mathbf{t}^T \cdot \boldsymbol{\Omega} \cdot \mathbf{t} \right)^2.
 \end{aligned} \tag{81}$$

10 By substituting the expressions of Eqs. (78)–(81) in Eq. (77) we obtain:

$$\begin{aligned}
 &\sum_{i=1}^m \mathbf{e}_i^T \cdot \boldsymbol{\Omega} \cdot \mathbf{J} \cdot \boldsymbol{\Omega} \cdot \mathbf{J} \cdot \mathbf{e}_i \\
 &= \sum_{i=1}^m \sum_{j=1}^m \omega_{ij}^2 - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t}^T \cdot \boldsymbol{\Omega}^2 \cdot \mathbf{t} - \frac{1}{\mathbf{t}^T \mathbf{t}} \cdot \mathbf{t}^T \cdot \boldsymbol{\Omega}^2 \cdot \mathbf{t} + \frac{1}{(\mathbf{t}^T \mathbf{t})^2} \cdot \left(\mathbf{t}^T \cdot \boldsymbol{\Omega} \cdot \mathbf{t} \right)^2 \\
 &= \sum_{i=1}^m \sum_{j=1}^m \omega_{ij}^2 - 2 \cdot \frac{\mathbf{t}^T \cdot \boldsymbol{\Omega}^2 \cdot \mathbf{t}}{\mathbf{t}^T \mathbf{t}} + \left(\frac{\mathbf{t}^T \cdot \boldsymbol{\Omega} \cdot \mathbf{t}}{\mathbf{t}^T \mathbf{t}} \right)^2
 \end{aligned}$$

Precision of the Match method

R. Lehmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

$$= \sum_{i=1}^m \sum_{j=1}^m \omega_{ij}^2 - 2 \cdot \frac{(\boldsymbol{\Omega} \cdot \mathbf{t})^T \cdot \boldsymbol{\Omega} \cdot \mathbf{t}}{\mathbf{t}^T \mathbf{t}} + \left(\frac{\mathbf{t}^T \cdot \boldsymbol{\Omega} \cdot \mathbf{t}}{\mathbf{t}^T \mathbf{t}} \right)^2, \quad \text{because } \boldsymbol{\Omega} \text{ is symmetric,}$$

$$= \omega_1 - 2 \cdot \omega_2 + \omega^2, \quad \text{because of the definition of } \omega_1, \omega_2, \text{ and } \omega \text{ in Eqs. (41), (42),}$$
(24).
(82)

By substituting the expressions of Eqs. (76) and (82) in Eq. (75) we obtain:

$$E(s_2) = -\omega \cdot \sigma^2 + (\omega_1 - 2 \cdot \omega_2 + \omega^2) \cdot \sigma_\delta^2. \quad (83)$$

Acknowledgement. We thank K. Hoppel (Naval Research Laboratory, Washington) for providing the POAM III data on which the analysis in Sects. 8.4–8.5 is based. This work was partly supported by the QUOBI project of the EC DG Research under the contract EVK2-2001-00129.

References

- Morris, G. A., Bojkov, B. R., Lait, L. R., and Schoeberl, M. R.: A review of the Match technique as applied to AASE-2/EASOE and SOLVE/THESEO 2000, Atmos. Chem. Phys. Discuss., 4, 4665–4717, 2004,
[SRef-ID: 1680-7375/acpd/2004-4-4665](#).
- Rex, M., Harris, N. R. P., von der Gathen, P., et al.: Prolonged stratospheric ozone loss in the 1995–96 Arctic winter, Nature, 389, 835–838, 1997.
- Rex, M., von der Gathen, P., Harris, N. R. P., et al.: In situ measurements of stratospheric ozone depletion rates in the Arctic winter 1991/1992: A Lagrangian approach, J. Geophys. Res., 103, 5843–5853, 1998.
- Rex, M., von der Gathen, P., Braathen, G. O., et al.: Chemical ozone loss in the Arctic winter 1994/95 as determined by the Match technique, J. Atmos. Chem., 32, 35–59, 1999.
- Rex, M., Salawitch, R. J., Harris, N. R. P., et al.: Chemical depletion of Arctic ozone in winter 1999/2000, J. Geophys. Res., 107, 8276, doi:10.1029/2001JD000533, 2002.
- Sasano, Y., Terao, Y., Tanaka, L., Yasunari, T., Kanzawa, H., Nakajima, H., Yokota, T., Nakane, H., Hayashida, S., and Saitoh, N.: ILAS observations of chemical ozone loss in the Arctic vortex during early spring 1997, Geophys. Res. Lett., 27, 213–216, 2000.

**Precision of the
Match method**R. Lehmann et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

Smit, H. G. J. and Straeter, W.: JOSIE-1998, Performance of ECC Ozone Sondes of SPC-6A and ENSCI-Z Type, WMO Global Atmosphere Watch report series, No. 157 (Technical Document No. 1218), World Meteorological Organization, Geneva, 2004a.

5 Smit, H. G. J. and Straeter, W.: JOSIE-2000, Jülich Ozone Sonde Intercomparison Experiment 2000, The 2000 WMO international intercomparison of operating procedures for ECC-ozone sondes at the environmental simulation facility at Jülich, WMO Global Atmosphere Watch report series, No. 158 (Technical Document No. 1225), World Meteorological Organization, Geneva, 2004b.

10 Streibel, M., Rex, M., von der Gathen, P., et al.: Chemical ozone loss in the Arctic winter 2002/2003 determined with Match, Atmos. Chem. Phys. Discuss., accepted, 2005.

Terao, Y., Sasano, Y., Nakajima, H., Tanaka, L., and Yasunari, T.: Stratospheric ozone loss in the 1996/1997 Arctic winter: Evaluation based on multiple trajectory analysis for double-sounded air parcels by ILAS, J. Geophys. Res., 107, 8210, doi:10.1029/2001JD000615, 2002.

15 von der Gathen, P., Rex, M., Harris, N. R. P., et al.: Observational evidence for chemical ozone depletion over the Arctic in winter 1991–92, Nature, 375, 131–134, 1995.

**Precision of the
Match method**

R. Lehmann et al.




Fig. 1. Difference ΔO_3 of the ozone mixing ratio determined by the second and first sonde of matches in dependence on the time t_s that the corresponding trajectory spent in sunlight. The slope of the regression line is the mean ozone loss rate, expressed as ozone loss per sunlit time. The data represent all Arctic match events of 14–28 January 1995, 475 K.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

Precision of the Match method

R. Lehmann et al.




Fig. 2. Illustration of the uncertainty of the slope of the linear regression line in the case of two independent matches. Full circles denote the ozone mixing ratio at the position of the first and second sonde of the corresponding match. Empty circles indicate the ozone mixing ratio obtained by sonde measurements, having an error $\pm\delta$. The ozone loss rate derived from the measurements is represented by the slope of the straight line in each panel. t_0 denotes the time t_s that the corresponding trajectory spent in sunlight.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

Precision of the Match method

R. Lehmann et al.




Fig. 3. Illustration of the uncertainty of the slope of the linear regression line in the case of two matches having a common second sonde. Symbols are as in Fig. 2.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

**Precision of the
Match method**

R. Lehmann et al.




Fig. 4. Illustration of the uncertainty of the slope of the linear regression line in the case of two matches sharing one sonde that is first sonde for one of the matches and second sonde for the other match. Symbols are as in Fig. 2.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

Precision of the Match method

R. Lehmann et al.




Fig. 5. Illustration of the uncertainty of the slope of the linear regression line for the three cases displayed in Figs. 2–4.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

Precision of the Match method

R. Lehmann et al.




Fig. 6. Schematic representation of the uncertainties associated with the Match method.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Print Version](#)
[Interactive Discussion](#)

EGU

Precision of the Match method

R. Lehmann et al.




Fig. 7. The new (crosses) and old (circles) estimate of the precision of the ozone loss rate obtained by Match, expressed as one standard deviation, versus the corresponding ozone loss rate. Regression lines for the new and old results have been added (dotted lines). Moreover, the vertical line corresponding to zero ozone loss and the bisecting line (uncertainty = ozone loss) have been highlighted. The data represent all match ensembles of the Arctic winters 1994/95, 1995/96, 1999/2000, 2002/03.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

Precision of the Match method

R. Lehmann et al.




Fig. 8. The ratio “new precision estimate/old precision estimate” versus the oversampling rate (= the average number of matches to which an ozone measurement contributes). A regression line and the horizontal line corresponding to a ratio of 1 (new error bars = old error bars) have been added. Moreover, the mean values of the abscissae and the ordinates of the data points have been highlighted by triangular arrowheads. The same data as in Fig. 7 have been used.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU

Precision of the Match method

R. Lehmann et al.




Fig. 9. The ratio “new precision estimate/old precision estimate” versus the oversampling rate, as in Fig. 8. The data represent match ensembles of an Antarctic satellite (POAM III) Match study in 2003.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

EGU