



HAL
open science

Characterization of positive air ions in boreal forest air at the Hyytiälä SMEAR station

U. Hörrak, P. P. Aalto, J. Salm, K. Komsaare, H. Tammet, J. M. Mäkelä, L.
Laakso, M. Kulmala

► **To cite this version:**

U. Hörrak, P. P. Aalto, J. Salm, K. Komsaare, H. Tammet, et al.. Characterization of positive air ions in boreal forest air at the Hyytiälä SMEAR station. *Atmospheric Chemistry and Physics Discussions*, 2007, 7 (4), pp.9465-9517. hal-00302933

HAL Id: hal-00302933

<https://hal.science/hal-00302933>

Submitted on 18 Jun 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Positive air ions in
boreal forest air**

U. Hörrak et al.

Characterization of positive air ions in boreal forest air at the Hyytiälä SMEAR station

U. Hörrak¹, P. P. Aalto², J. Salm¹, K. Komsaare¹, H. Tammet¹, J. M. Mäkelä³, L. Laakso², and M. Kulmala²

¹Institute of Physics, University of Tartu, 18 Ülikooli St., 50090 Tartu, Estonia

²Department of Physical Sciences, Division of Atmospheric Sciences P.O. Box 64, 00014 University of Helsinki, Finland

³Tampere University of Technology, Institute of Physics, P.O. Box 692, 33101, Tampere, Finland

Received: 8 June 2007 – Accepted: 18 June 2007 – Published: 3 July 2007

Correspondence to: U. Hörrak (urmas.horrak@ut.ee)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

Abstract

The behavior of the concentration of positive small (or cluster) air ions and naturally charged nanometer aerosol particles (aerosol ions) has been studied on the basis of measurements carried out in a boreal forest at the Hyytiälä SMEAR station, Finland, during the BIOFOR III campaign in spring 1999. Statistical characteristics of the concentrations of cluster ions, two classes of aerosol ions of the sizes of 2.5–8 nm and 8–ca. 20 nm and the quantities that determine the balance of small ions in the atmosphere have been given for the nucleation event days and non-event days. The dependence of small ion concentration on the ion loss (sink) due to aerosol particles was investigated applying a model of bipolar diffusion charging of particles by small ions. The small ion concentration and the ion sink were closely correlated (correlation coefficient -87%) when the fog events and the hours of high relative humidity (above 95%), as well as nocturnal calms and weak wind (wind speed $<0.6 \text{ m s}^{-1}$) had been excluded. However, an extra ion loss term presumably due to small ion deposition on coniferous forest with a magnitude equal to the average ion loss to pre-existing particles is needed to explain the observations. Also the hygroscopic growth correction of measured aerosol particle size distributions was found to be necessary for proper estimation of the ion sink. In the case of nucleation burst events, variations in the concentration of small positive ions were in accordance with the changes caused by the ion sink due to aerosols; no clear indication of positive ion depletion by ion-induced nucleation was found. The estimated average ionization rate of the air at the Hyytiälä station in early spring, when the ground was partly covered with snow, was about $6 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$. The study of the charging state of nanometer aerosol particles (2.5–8 nm) revealed a strong correlation (correlation coefficient 88%) between the concentrations of particles and positively charged particles (positive air ions) during nucleation bursts. The estimated charged fraction of particles, which varied from 3% to 6% considering various nucleation event days, confirms that these particles are almost quasi-steady state charged. Also the particles and air ions in the size range of 8–ca. 20 nm showed a good qualitative con-

ACPD

7, 9465–9517, 2007

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

sistency; the correlation coefficient was 92%.

1 Introduction

The formation and growth of ultrafine aerosol particles in the atmosphere have been studied during the last decade at many different locations around the world because of their possible impact on the radiation balance and thereby on the climate of the Earth (Kulmala et al., 2004a; Birmili et al., 2003; Tunved et al., 2003; Iida et al., 2006). Various theories have been elaborated to explain the mechanisms of new particle generation (e.g. Raes and Janssens, 1985; Yu and Turco, 2000, 2001; Laakso et al., 2002; Korhonen et al., 1999, Kulmala et al., 2000; Napari et al., 2002). Despite the frequently observed particle formation events, the microphysical mechanisms of nucleation have remained unclear because of the very small size of newly born particles (about 1–2 nm). This size range is not within the scale of modern aerosol instruments using artificial charging of particles. Air ion spectrometers are suitable to measure naturally charged clusters and particles down to about 0.4 nm, but the charging state of newly born particles is not known in the atmosphere (due to different particle formation mechanisms). Experimental study of new particle formation in the atmosphere is complicated also because the size range of new particles coincides with the size range of cluster ions (0.4–1.6 nm). The knowledge about the behavior of ion clusters, particles and their charged fraction (air ions) during nucleation events could contribute to elucidate the role of different nucleation mechanisms.

The mobility spectrum of cluster ions is sensitive to the content of many trace species (gases or vapors at concentrations below 1 part per billion) in the air giving rise to certain specific ion families with characteristic mobilities (Nagato and Ogawa, 1998; Parts and Luts, 2004). The positive and negative cluster ions exhibit very different chemical composition (Eisele and Tanner, 1990; Beig and Brasseur, 2000; Luts and Parts, 2002). Typically positive core ions consist of hydronium (H_3O^+), ammonium (NH_4^+) and a variety of organic compounds (e.g. pyridine), while the dominant negative

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

core ions besides parent species NO_3^- and HSO_4^- include also strong acids (nitric acid and sulfuric acid). However, the differences in physical properties of ions (mass, size and mobility) are not large, e.g. the mean mobility of negative cluster ions near the ground is about 13% higher than that of positive ions (Hörrak et al., 2000).

5 The importance of air ions in the particle formation processes has been shown in several laboratory experiments (Ramamurthi et al., 1993; Kim et al., 1997, 1998; Han-
son and Eisele, 2002; Froyd and Lovejoy, 2003a, b; Wilhelm et al., 2004). Compre-
hensive overviews about the charging and charge distribution on aerosol particles in a
bipolar ion environment can be found in (Hoppel and Frick, 1986; Reischl et al., 1996).
10 According to laboratory experiments and theoretical considerations, the steady state
charge distribution on large aerosol particles determined by the diffusion of small ions
to particles is close to the Boltzmann distribution. Corrections have been found to be
necessary in the case of ultrafine particles below 100 nm (Fuchs and Sutugin, 1971;
Hoppel and Frick, 1986; Tammet, 1991; Reischl et al., 1996). However, only few ex-
15 perimental studies on the subject have been carried out in the real atmosphere. So
far, the problems of the bursts of intermediate ions (generation of charged nanome-
ter aerosol particles) and of the charge distribution on freshly nucleated atmospheric
nanoparticles have been briefly discussed (Hörrak et al., 1998a, b, 2003a, b; Mäkelä
et al., 2001a, 2003; Tamm et al., 2001). An extensive study of air ions and aerosol
20 particles was recently conducted in a boreal forest at the Hyytiälä SMEAR station, Fin-
land, during the QUEST campaign (21 March–10 April 2003) (Kulmala et al., 2004b;
Laakso et al., 2004a, b). One of the important results of the QUEST 2 campaign was
a charge asymmetry on intermediate air ions (1.6–5 nm) with an excess of negative
charges and overcharging of nanometer aerosol particles (compared to the steady
25 state bipolar charging probability) in the size range of 3–5 nm during weak and mod-
erate nucleation bursts (Hörrak et al., 2004a, b; Laakso et al., 2004a). The formation
of negatively charged nanometer particles by ion-induced nucleation is in accordance
with the results of laboratory experiments (Froyd and Lovejoy, 2003b; Wilhelm et al.,
2004).

**Positive air ions in
boreal forest air**

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The objective of this work was a comparative study of positive air ions (naturally charged small or cluster ions and aerosol particles) and fine aerosol particles with the main focus on the balance between small ions and aerosols, as well as on the charging state of nanometer particles during nucleation events. Simultaneous measurements of the concentration of cluster ions and the size distribution of aerosol particles enable to estimate also the ionization rate of the air (or ion production rate), which is a key factor in ion-induced nucleation.

2 Measurements

Atmospheric aerosols and the concentration of positive air ion mobility classes were measured in a boreal forest at the Hyytiälä SMEAR II station, Finland, during the BIOFOR III campaign (31 March–29 April 1999) (Kulmala et al., 2001a; Aalto et al., 2001). The SMEAR II (Station for Measuring Forest Ecosystem-Atmosphere Relations) is located on a flat terrain at the Hyytiälä Forestry Field Station of the University of Helsinki about 220 km north-west of Helsinki. The station is surrounded by pine-dominated forests of middle density covering extended areas. The terrain is a subject to modest height variation (up to about 40 m). The oblong Lake Kuivajärvi (141 m a.s.l.) is located about 600 m south-west of the station. The ground in the vicinity is igneous and metamorphic bedrock (notably granites) partly exposed or covered by a thin layer of soil (maximum up to about 2 m). The nearest largest city, Tampere (204 000 inhabitants), is located about 60 km south-west of the station. A detailed description of the station has been given by Vesala et al. (1998) and by Hari and Kulmala (2005).

Aerosol particle size distribution was measured by means of a dual DMPS (Differential Mobility Particle Sizer) system covering the sizes of 3–500 nm (Aalto et al., 2001). The devices had a Hauke-type differential mobility analyzer (DMA), a closed-loop sheath flow arrangement and a Krypton-85 aerosol neutralizer. For large particles, an aerodynamical particle sizer (APS, TSI model 3320) was used to measure the size distribution from 0.7 μm up to 20 μm . This instrument was located in a truck about

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

40 m from the main cottage of the SMEAR station where the DMPS system had been installed. Both instruments had sampling lines of the 2-m level inside the forest. The sample flow and sheath flow of the DMPS, as well as the sample flow of the APS were not dried. However, the size distributions of particles measured by means of both devices can be considered as “dry size distributions” due to low relative humidity inside the instruments (especially in the APS). Additional information about the accumulation mode and coarse mode particles in the diameter range of 0.2–32 μm was obtained using in-situ optical particle counters: a Particle Measuring System’s (PMS) Active Scattering Aerosol Spectrometer Probe (ASASP-300), measuring in the size range 0.2–3 μm , and a Classical Scattering Aerosol Spectrometer Probe (CSASP-100), covering the sizes above 1 μm (Kulmala et al., 2001a; Hämeri et al., 2001). The optical counters sampled wet-ambient aerosol particles on a tower at a height of about 18 m above the ground and about 30 m from the main cottage.

In order to estimate the concentration of positive air ions in different mobility classes, four integral ion counters UT-8401 were applied (Aalto et al., 2001). Negative air ions were not measured during the BIOFOR III campaign. The ion counters were the Ger-dien type aspiration counters in which the naturally charged particles (cluster ions and aerosol ions) deposited from the airflow by the electric field in the mobility analyzer according to their electrical mobilities and the electric current was measured by electrometrical amplifier. The limiting mobilities of four ion counters, determined by the voltage, flow rate and effective capacity of the mobility analyzer, were set to 0.02, 0.063, 0.2, and 0.64 $\text{cm}^2 \text{V}^{-1} \text{s}^{-1}$, respectively. Due to the nature of the integral ion counter, the concentrations of air ion mobility classes can be calculated by means of a special inversion algorithm (Tammet, 1970; Israël, 1970). The method is based on the measurements at several different limiting mobilities to determine the dependence of the ion concentration on the reciprocal of the limiting mobility (or current-voltage characteristic curve). At a certain mobility, the intercept of the characteristic curve tangent on the concentration axis yields the concentration of ions with mobilities higher than this specific mobility, and the difference between the intercepts yields the ion concentration in a

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Positive air ions in
boreal forest air**

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

mobility class. In practice, a piecewise linear approximation of the characteristic curve was used and the specific mobilities of 0.355 and $0.0355 \text{ cm}^2 \text{ V}^{-1} \text{ s}^{-1}$ were selected as boundaries between the mobility classes. The lower boundary of the lowest mobility class was conventionally set to $0.005 \text{ cm}^2 \text{ V}^{-1} \text{ s}^{-1}$ (Aalto et al., 2001). The concentrations of positive small or cluster ions (Stokes-Millikan mobility equivalent diameter less than 2.5 nm) and two classes of aerosol ions – charged fine nanometer particles (2.5 – 8 nm) and coarse nanometer particles (8 –ca. 20 nm) – were derived. Air ions were measured as naturally charged in the bipolar atmospheric air. The upper boundary of 20 nm cannot be exactly determined because of the contribution of charged aerosol particles of larger sizes: the larger the particle, the smaller is its contribution to signal. The sampling height of ion counters was 2 m above the ground; the data were recorded every 10 min . The distance between the measurement locations of air ions and aerosols was about 450 m . For details about the instrumentation, see (Kulmala et al., 2001a; Aalto et al., 2001).

The average temperature in the first half of April was 2.2°C and in the second half 7.0°C . The depth of the snow cover was about 40 cm at the beginning and 20 cm in the middle of April; the snow cover melted quickly during the week after 15 April 1999.

3 Theoretical background

3.1 The balance of ions in the atmosphere

Small positive and negative ions are generated in the atmosphere mainly by cosmic radiation and natural radioactivity. Two concurrent processes, ion-ion recombination and adsorption of small ions by aerosol particles, are responsible for the loss of small ions. The simplified balance equation of small ions in a bipolar environment, considering equal numbers of positive and negative small ions and symmetrical charging of

aerosol particles, can be written as (Hoppel and Frick, 1986)

$$\frac{dn}{dt} = q - \alpha n^2 - \beta_{\text{eff}} N_{\text{tot}} n, \quad (1)$$

where q is the ionization rate, α is the recombination coefficient, n is the concentration of small ions, N_{tot} is the total concentration of aerosol particles, and β_{eff} is the effective ion-aerosol attachment coefficient for polydisperse aerosol particles. The time constant of the transfer to the steady state condition ($dn/dt=0$) is about 1 min in the continental boundary layer air. This balance equation does not take into account the loss of small ions by ion-induced nucleation due to its complicated nature. The simplified balance equation of small ions (Eq. 1) has an analytical steady state solution:

$$n = \frac{2q}{\beta_{\text{eff}} N_{\text{tot}} + \sqrt{\beta_{\text{eff}}^2 N_{\text{tot}}^2 + 4\alpha q}}. \quad (2)$$

It can also be presented in the form: $n=q\tau(q)$, where the parameter $\tau(q)$ has the dimension of time and can therefore be interpreted as the “mean lifetime” of small ions in the steady state conditions (Israël, 1970). The mean lifetime of small ions is inversely proportional to the total ion loss caused by ion-aerosol attachment and by the recombination of small ions.

The independent parameters of the balance equation have the following average values often cited in publications. The recombination coefficient α , which depends on the nature of small ions and the properties of the environment (Israël, 1970; Hoppel and Frick, 1986), has an average value of about $1.5 \times 10^{-6} \text{ cm}^3 \text{ s}^{-1}$. In continental areas, the parameter β_{eff} varies in the range of $(1-2) \times 10^{-6} \text{ cm}^3 \text{ s}^{-1}$, depending on the size distribution of aerosol particles (Hoppel, 1985). The average ionization rate near the ground of about $10 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$ (caused by radioactive substances in the ground and in the air, and by cosmic radiation) is considered as an average for continental areas.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The *ion sink* due to aerosols is given by the equation (Hoppel and Frick, 1986; Tammet, 1991):

$$S_a = \beta_{\text{eff}} N_{\text{tot}} = \int_r \sum_{i=-\infty}^{\infty} \beta_i(r) p_i(r) N(r) dr, \quad (3)$$

were $\beta_i(r)$ is the ion-aerosol attachment coefficient between small positive (or negative) ions and particles with radius r and with i (or $-i$) elementary charges, $p_i(r)$ is the probability of carrying i elementary charges on a particle and $N(r)$ is the size distribution of aerosol particles.

The attachment of small ions to aerosol particles gives rise to a size-dependent statistical charge distribution on the aerosol particles (Hoppel and Frick, 1986; Reischl et al., 1996). The time in which the aerosol particle population approaches charge equilibrium depends on the collision rate between small ions and aerosol particles, and also on the particle formation mechanism (e.g. homogeneous nucleation, ion-induced nucleation). Considering typical atmospheric conditions and initially neutral ultrafine aerosol particles, the time required to reach charge equilibrium is typically in the range of about 0.5–1 h (Hoppel, 1985). For rural areas, away from direct sources of aerosol particles, the assumption of charge equilibrium is presumed to be fairly good. However, the charge distribution on freshly nucleated nanometer particles needs further study.

3.2 Calculation of the ion sink

The effect of atmospheric aerosols on the small ion concentration was estimated applying a simplified model of bipolar diffusion charging of aerosols (by small ions) and assuming the steady state conditions and symmetrical charging (Tammet, 1991; Hörrak et al., 1998b). The calculation of the ion sink Eq. (3) is based on the model of ion-aerosol attachment (Hoppel and Frick, 1986; 1990). The charging probability $p_i(r)$ for the steady state is calculated using the algorithm given by Tammet (1991) with improvements described in Hörrak et al. (1998b). The model of charge distribution is

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

based on the expression for the attachment coefficient of a small (cluster) ion to an aerosol particle:

$$\beta_i(r) = \frac{4\pi r k T Z}{e} \frac{x}{\exp(x) - 1} \sqrt{1 - \frac{2}{2 + i(i-1) + r/(5\text{nm})}}, \quad (4)$$

where Z is the mean mobility of cluster ions, $x = ie^2/(4\pi r \epsilon_0 k T)$, ϵ_0 is the electric constant, k is the Boltzmann constant, and T is the temperature. The Eq. (4) approximates the tabulated results of Hoppel and Frick (1990) and is also in accordance with the experimental data of charging probabilities of nanometer aerosol particles (Reischl et al., 1996).

The function $\rho_i(r)$ is defined by the equation:

$$\rho_i(r) = \frac{N_i(r)}{N(r)} \frac{N_i(r)}{N_0(r) + 2 \cdot \sum_{j=1}^{\infty} N_j(r)}, \quad (5)$$

where $N_i(r)$ or $N_j(r)$ is the fraction concentration of particles at the radius r with i or j elementary charges, $N(r)$ is the total concentration and $N_0(r)$ is the concentration of neutral particles at the radius r . In the steady state $N_i(r)/N_0(r) = \prod_{j=1}^i \beta_{j-1}(r)/\beta_{-j}(r)$.

The equivalent, or the single size ion-aerosol attachment coefficient

$$w(r) = \sum_{i=-\infty}^{\infty} \beta_i(r) \rho_i(r) \quad (6)$$

in Eq. (3) is almost proportional to the particle diameter and, therefore, the ion sink S_a can be interpreted as the diameter concentration of aerosol particles (Hoppel and Frick, 1986). A simple approximation function originates from Tammet (1991):

$$w(r) = \sqrt{\frac{2r - 1 \text{ nm}}{2r + 5 \text{ nm}}} \frac{2r}{40 \text{ nm}} 10^{-6}, \quad (7)$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

where the particle radius r is in nanometers (nm) and $w(r)$ is in $\text{cm}^3 \text{s}^{-1}$. Equation (7) was improved compared to (Tammet 1991): the square root was added, which provides a better agreement with exact calculations by means of Eqs. (4), (5) and (6). The dependence of the single size ion-aerosol attachment coefficient on the particles of different sizes (2 nm–10 μm) is shown in Fig. 1. The mean mobility of positive and negative cluster ions of $1.45 \text{ cm}^2 \text{ V}^{-1} \text{ s}^{-1}$, standard temperature (20°C) and air pressure (1013 mb) have been used in these calculations.

The depletion rate of small air ions due to aerosols (ion sink) S_a is found by integrating over the size distribution of aerosol particles of 3–500 nm measured by means of the dual DMPS system. The effect of large particles above 500 nm was studied using the data of the Aerodynamic Particle Sizer (APS) and optical particles counters. The particle size distributions in the diameter range of 0.72–20 μm , measured with the APS, were converted to the mobility diameter by dividing the aerodynamic diameter by the square root of the estimated particle density of 1.9 g cm^{-3} (Aalto et al., 2001). The ion sink is closely correlated with the condensation sink (condensation rate of vapors on aerosol particles) (Kulmala et al., 2001b) because both factors are mainly depending on the diameter concentration of aerosol particles (first moment of the size distribution).

4 Results and discussion

4.1 Statistical characterization of small air ions and ion loss

Statistical characteristics of the positive small (cluster) ion concentration, the sink of small ions on aerosol particles of 3–500 nm measured by the DMPS, as well as the factors of the small ion balance Eq. (2) are presented in Table 1, separately for the nucleation event days and non-event days. The days have been classified as *nucleation event days* or *non-event days* according to the occurrence of new particle formation followed by the particle size distribution measurements. The classification given in Kulmala et al. (2001a) was used in the present study with one exception: the first half of 21

April, when polluted air masses were present, was excluded from the list of nucleation event days.

Considering all the measurements, the ion sink on aerosols S_a varied in the range of $3.9 \times 10^{-4} - 5.8 \times 10^{-2} \text{ s}^{-1}$, commonly $1 \times 10^{-3} - 1.2 \times 10^{-2} \text{ s}^{-1}$. The smallest values were recorded after the inflow of clean maritime air masses, while the highest values correspond to the polluted continental air masses, when the accumulation mode aerosol particles are mainly responsible for the ion sink. During the nucleation event days (13 days) the average ion sink was about 1.8 times smaller compared to non-event days, and the mean concentration of small ions was higher: 530 cm^{-3} versus 424 cm^{-3} , respectively. The highest concentrations of small ions up to about 1200 cm^{-3} were recorded during nighttime calms. The high concentrations were probably caused by the accumulation of radon near the ground during temperature inversions (Porstendörfer, 1994; Penttinen et al., 2003), which can significantly increase the ionization rate of the air. In the nucleation event days, the average diurnal variation of the small ion concentration displayed the maximum of about 625 cm^{-3} during nighttime at 22:00–23:00 LST, and the minimum of 460 cm^{-3} in the morning at 07:00–08:00 LST. The secondary maximum of 580 cm^{-3} , which appeared at noon (11:30 LST), was correlated with the minimum in the ion sink of $2.6 \times 10^{-3} \text{ s}^{-1}$ (the maximum of ion sink of $5.2 \times 10^{-3} \text{ s}^{-1}$ appeared at 24 LST). The secondary maximum of small ions took place before the intensive generation of nanometer particles; it preceded the maximum of the average diurnal variation of 3 nm particles by about 1 h. In general, the small ions measured at Hyytiälä showed similar regularities as at Tahkuse, but the concentrations were about two times higher than at the Tahkuse Observatory, Estonia (Hörrak et al., 2000, 2003b).

Extra information about negative cluster ions was gained from the mobility distribution measurements in the range of $0.5 - 3.2 \text{ cm}^2 \text{ V}^{-1} \text{ s}^{-1}$ at the same place during the QUEST 2 campaign in spring 2003. The average concentration of negative cluster ions and their standard deviation was $575 \pm 114 \text{ cm}^{-3}$. This is about 1.14 times lower than the concentration of positive cluster ions ($655 \pm 118 \text{ cm}^{-3}$) at Hyytiälä in early spring (Hirsikko et al., 2005). The concentrations of positive and negative ions were closely

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

correlated; the correlation coefficient was 93.6%. The mean mobilities of negative and positive cluster ions were $1.69 \text{ cm}^2 \text{ V}^{-1} \text{ s}^{-1}$ and $1.45 \text{ cm}^2 \text{ V}^{-1} \text{ s}^{-1}$, respectively (Laakso et al., 2004b), and the correlation between the quantities was about 60%.

It is worth to point out that during the BIOFOR III campaign at the Hyytiälä station, the loss of small ions due to ion-ion recombination (assuming the recombination coefficient $1.5 \times 10^{-6} \text{ cm}^3 \text{ s}^{-1}$) was about 16% of the total ion loss due to small ion recombination and attachment to the aerosol particles, in average. Typically, it is less than 5% for continental areas (Tammet, 1991). In the case of very clean air, e.g. after the Arctic cold air advection from the Northern Atlantic, the small ion recombination was responsible for about 70% of ion losses (Table 1). Here we neglected the effect of ion-induced nucleation on the loss of small ions. We also assumed equal concentrations of positive and negative small air ions; in general, the ratio of positive to negative ions is about 1.12 (Hörrak et al., 2000). The peak concentrations of small ions at Hyytiälä, recorded in the very clean air (about 800 cm^{-3}), are still considerably smaller than the concentrations (up to 1200 cm^{-3}) recorded during nocturnal calms (radon accumulation effect).

The sink of small ions at the Hyytiälä station was mainly determined by the concentration of particles in the size range of 3–500 nm measured by the DMPS; the contribution of larger particles measured by the APS (mobility equivalent diameter 0.52– $15 \mu\text{m}$) was about two orders of magnitude smaller (see Fig. 2). For example, during the nucleation event days (non-event days), an average value of an ion sink measured by the APS was $4.88 \times 10^{-5} \text{ s}^{-1}$ ($11.47 \times 10^{-5} \text{ s}^{-1}$) and maximum was $14.5 \times 10^{-5} \text{ s}^{-1}$ ($49.6 \times 10^{-5} \text{ s}^{-1}$), respectively. The percentage contribution of particles in the size range of the APS was about 1.5% in average, and always less than 8%. Therefore, if high accuracy is not essential when calculating the ion sink, then we can take into account only the DMPS measurements. However, the estimated ion sinks are probably to a certain extent underestimated because we did not take into account the hygroscopic growth of particle size due the effect of ambient humidity. During the BIOFOR campaign, Hämeri et al. (2001) studied the hygroscopic growth of aerosol particles in the dry particle diameter range of 0.01– $2 \mu\text{m}$ when taken from the dry state (relative humidity $\text{RH} < 5\%$)

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

to RH=90%. The growth factors ranged between 1.0 and 1.6 considering particles of different sizes, and a clear diurnal variation of the growth factors of nucleation, Aitken and accumulation mode particles was found during nucleation event days.

The effective attachment coefficient (β_{eff}), which is proportional to the mean diameter of aerosol particles, varied in the range of $2 \times 10^{-7} - 3.85 \times 10^{-6} \text{ cm}^3 \text{ s}^{-1}$. These changes in the remote environment are affected by the long-range transport of aerosol pollutants and/or clean air advection, by the transformation of aerosol size distribution due to the evolution of boundary layer and new particle generation (Nilsson et al., 2001a, b). During the nucleation burst days the effective attachment coefficient showed a considerable diurnal variation; on the average, it varied from about $6 \times 10^{-7} \text{ cm}^3 \text{ s}^{-1}$ up to $1.8 \times 10^{-6} \text{ cm}^3 \text{ s}^{-1}$, displaying a nighttime maximum at about 04:00–05:00 LST and a minimum in the afternoon at 16:00 LST. During the peak time of nanometer particle generation by photochemical nucleation around noon, the effective attachment coefficient was almost always less than $6 \times 10^{-7} \text{ cm}^3 \text{ s}^{-1}$. The effective attachment coefficients found by Hoppel (1985) at the Naval Research Laboratory in urban Washington, D.C. were 1.15×10^{-6} and $1.97 \times 10^{-6} \text{ cm}^3 \text{ s}^{-1}$, and the ion sinks calculated from those measurements were 1.6×10^{-2} and $4.64 \times 10^{-2} \text{ s}^{-1}$ for the periods of good and poor visibility, respectively. Measurements in the marine environment near the east coast of the United States revealed the attachment coefficients from $0.89 \times 10^{-6} \text{ cm}^3 \text{ s}^{-1}$ up to $2.63 \times 10^{-6} \text{ cm}^3 \text{ s}^{-1}$ and in the remote Atlantic as high as $3.18 \times 10^{-6} \text{ cm}^3 \text{ s}^{-1}$. The corresponding ion sinks were in the range of $1.8 \times 10^{-2} - 2.4 \times 10^{-2} \text{ s}^{-1}$ and $6.46 \times 10^{-4} \text{ s}^{-1}$, respectively. The effective attachment coefficients in considerably different environments are comparable with each other and with average values found in the boreal forest at Hyytiälä. The ion sink influenced by local air pollution, anthropogenic and natural sources of aerosols is a subject to considerable differences.

The average ionization rate of about $2.8 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$, calculated from the balance Eq. (1) considering the steady state conditions, is too small for continental stations. It is probably underestimated by a factor of 2. The average ionization rate at 1 m height from the ground about $10 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$ is considered as an average

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

for continental areas (Israël, 1970, 1973; Chalmers, 1967). Three main ionizing agencies (factors) have the following contribution: the ionization rate caused by radioactive substances in the ground is about $4.6 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$ and in the air (radon, radioactive aerosol) $4 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$, and by cosmic radiation $1.5\text{--}1.8 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$.

5 Considering different locations, the ionization rate could exhibit considerable variations depending on the content of radioactive substances in the ground, on the soil properties and due to local orography. The measurements below 1 m strongly depend on the ionization profile. The temporal variations in the ionization rate (annual and diurnal variation) are mostly due to variations in radon and thoron concentration in the air
10 (Porstendörfer, 1994; Hoppel et al., 1986). Our estimates of the ionization rate did not take into account the loss of small ions by the deposition on the coniferous forest, which could affect the measurements at the Hyytiälä station, as well as the differences in the sizes of wet ambient particles and those sampled by the DMPS. Presumably, the uncertainties of ion loss caused by these factors can explain the small values of
15 ionization rate. The exceptionally high ionization rates found on 21 April with the nighttime maximum of $14.6 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$ (Table 1b) are probably the artifacts caused by local air pollution, which did not affect the ion measurements. The most reasonable maximum value of the ionization rate estimated from the density distribution (excluding data of 21 April) is equal to that found for the nucleation event days, about 7.6
20 ion pairs $\text{cm}^{-3} \text{ s}^{-1}$. The average diurnal variation of the ionization rate displayed a moderate variation with the nighttime maximum of about $3.4 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$ and the minimum of $2 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$ in the afternoon.

Laakso et al. (2004a) applied the same approach to the calculation of the ionization rate based on the measurements of the QUEST 2 campaign at the Hyytiälä station in
25 spring 2003. The effect of the hygroscopic growth of the measured dry aerosol particle size distributions due to ambient humidity was also taken into account, but the average ionization rate stayed at a low value of $2.6 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$.

The measurements at Hyytiälä show that the atmosphere always contains small air ions that can act as condensation centers (of about 1 nm size) in the nucleation pro-

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

cess. During intensive generation of nanometer aerosol particles the concentration of small ions was in the range of $380\text{--}780\text{ cm}^{-3}$, considering different event days. Therefore, when the homogeneous nucleation of new aerosol particles takes place in the atmosphere, it should be accompanied by ion-induced nucleation (see e.g. Hoppel et al., 1994; Raes and Van Dingenen, 1992). To maintain the small ion population, the production rate of aerosols by ion-induced nucleation at the Hyytiälä station should be limited by the ionization rate (about $4\text{--}6\text{ ion pairs cm}^{-3}\text{ s}^{-1}$) (Laakso et al., 2004a).

4.2 Correlation between the concentration of small ions and ion sink

The correlation between small ion concentration and the ion sink due to aerosol particles of the sizes of $3\text{--}500\text{ nm}$ is depicted in Fig. 3. The overall correlation is weak owing to the nighttime calms and weak wind, and due to high humidity periods (probably fog events), when the concentration of small ions varied independently of the ion sink. The fog is known to affect the conductivity of atmospheric air, reducing the concentration of small ions (Dolezalek, 1963; Hoppel et al., 1986). Contrary to that, the nighttime calms and weak wind favor the accumulation of radon near the ground, resulting in an increase in the ionization rate and, therefore, in the concentration of small ions up to about 1200 cm^{-3} . Unfortunately, the radon concentration was not measured during the BIOFOR campaign. The accumulation of radon is confirmed by the measurements carried out at Hyytiälä during 2000–2003 (Penttinen et al., 2003). An alternative explanation of the occurrence of nighttime high concentrations proceeds from the hypothesis that the deposition rate of small ions on the coniferous forest is reduced in the case of weak wind and stable boundary layer, while the turbulent mixing can effectively enhance the deposition (Tammets and Kimmel, 1998; Tammets et al., 2001).

The fog events were identified comparing the data of the aerodynamical particle sizer (APS) and the optical aerosol spectrometer probes (ASASP-300 and CSASP-100). The APS measures dry aerosol size distribution, while the optical probes sample wet-ambient particles in-situ at a high flow rate, thus providing measurements of hydrated aerosol size distributions affected by the ambient relative humidity. During high

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

humidity (above 99%), the size spectra were often completely different above the diameter of $0.5\ \mu\text{m}$, probably due to the presence of fog droplets, which are supposed to reside in the size range of $2\text{--}40\ \mu\text{m}$. We found an indication of fog during the nighttime and/or early morning on 1, 2, 7, 15, 17 and 23 April. The effect of five fog events, when the ion counters were working properly and providing correct results, can be clearly seen in Fig. 3. The concentration of small ions had undergone a considerable decrease (deviation from the regression line), reaching values less than about $150\ \text{cm}^{-3}$. These situations were correlated with fog events.

Another factor that could affect the concentration of small ions is precipitation (snow, sleet, rain). Precipitation can decrease the ionization rate due to the attenuation of gamma and beta radiation from the ground and due to the block of radon exhalation by the wet ground surface. Contrary to that, precipitation can also cause an increase in the ionization rate close to the ground due to the wet deposition of short-lived radioactive radon daughters attached to aerosol particles (Hatakka et al., 1998). The effect of precipitation (snow, sleet, rain) was studied, but no clear direct or indirect effect on the concentration of small ions was found.

To find out a clear relationship between the concentration of small ions and the ion sink, we excluded the presumable fog events and also the hours of high humidity (more than 95%), as well as the periods of weak wind (wind speed less than $0.6\ \text{m s}^{-1}$) from the dataset. The results are depicted in Fig. 3b. In the case of the smallest ion sink of $3.9 \times 10^{-4}\ \text{s}^{-1}$, recorded after the inflow of clean maritime air masses from the North Atlantic (Mäkelä et al., 2001b), the concentration of small ions rose up to $800\ \text{cm}^{-3}$, while during the polluted continental air masses (coming from Russia over St. Petersburg) with a considerably higher ion sink of $2.6 \times 10^{-2}\ \text{s}^{-1}$ the small ion concentration was diminishing down to $150\ \text{cm}^{-3}$.

In accordance with the balance equation of small ions (2), if the ion sink is below about $10^{-3}\ \text{s}^{-1}$, the small ion concentration becomes nearly independent of that of aerosol particles approaching its limit in the case of aerosol free atmosphere. The concentration limits are about 1030, 1830 or $2580\ \text{cm}^{-3}$ if the ionization rate is 1.6, 5

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

or 10 ion pairs $\text{cm}^{-3} \text{s}^{-1}$, respectively. The concentration limit of 1030cm^{-3} best suits our data, but the regression line in Fig. 3 is still far from the relationship predicted by Eq. (2), displaying significantly higher ion concentrations considering the ion sinks above 10^{-3}s^{-1} . The fact that the ionization rate of the air depends on the content of radioactive aerosols might only partially explain the discrepancy.

The correlation between the small ion concentration and the ion sink was stronger during the nucleation event days, because there were no fog events (except one, on 2 April) and relative humidity was considerably lower (average 62%), compared with the non-event days (average 82%). If to take into account only the hours when the atmospheric boundary layer was well mixed (wind speed higher than 1m s^{-1}), then the nonlinear correlation coefficient was -92% .

4.3 Estimation of the ionization rate

As discussed in the Sect. 4.1, the ionization rate derived from the balance equation of small ions (Eq. 1) has unlikely low values for continental stations: the average is about $2.8 \text{ion pairs cm}^{-3} \text{s}^{-1}$. We assume that the balance equation is valid in general, but the ion loss has been underestimated. However, the concentration of small ions (n) and the ion sink (S_a) were closely correlated (see Fig. 3b). Therefore, we are looking for a constant extra ion loss, the addition of which can solve the discrepancy between the experimental data and the results derived from the balance equation. We applied a method of least squares to find out statistically grounded estimates of the free parameters of the balance equation: the ionization rate and a constant term of ion loss. To minimize the influence of local effects and fog events on the results, the periods of well-mixed atmospheric boundary layer were selected (wind speed $>1 \text{m s}^{-1}$, relative humidity $<95\%$) for the data analysis. The results of the regression analysis are depicted in Fig. 4. The statistical estimate of the ionization rate is $4.8 \text{ion pairs cm}^{-3} \text{s}^{-1}$ and that of the additional constant ion loss is $5.2 \times 10^{-3} \text{s}^{-1}$. Thus, the improved balance

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

equation of small ions in steady state conditions can be written as

$$q = \alpha n^2 + (S_a + S_e) n \quad (8)$$

This extra ion loss (S_e), which is approximately equal to the mean ion sink induced by aerosol particles of the sizes of 3–500 nm, could be caused by the enhanced deposition of small ions on the coniferous forest (Tamm et al., 2006). The ionization rate of 4.8 ion pairs $\text{cm}^{-3} \text{s}^{-1}$ is typical for the continental stations of high latitude during the wintertime when the snow cover is present. The same results were obtained when the hours with the wind speed higher than 0.6 m s^{-1} and relative humidity less than 95% were taken into account.

The estimated ionization rate of $4.8 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$ fairly coincides with the value of $4.5 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$ calculated for the same place from the external radiation and radon concentration measurements carried out during the QUEST 2 measurement campaign in spring 2003 (Laakso et al., 2004a). Based on the results of the regression analysis, we can conclude that the ionization rate in Table 1 is systematically underestimated by about $2 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$ and the ion loss of about $5.2 \times 10^{-3} \text{ s}^{-1}$. Also, the extra ion loss reduces significantly (up to about 4 times) the percentage of small ion loss due to the ion-ion recombination given in Table 1. The corrected mean value of the percentage contribution is about 7% and the maximum is 18%.

4.4 Charging state of nanometer aerosol particles

Simultaneous measurements of the concentrations of aerosol particles and naturally charged particles (air ions) in the atmosphere in the same size range enable to estimate the charged fraction or charging probability of particles. Considering the accuracy of nanometer particle measurements, it is possible to study the process only during the nucleation event days, when the enhanced concentrations of nanometer particles (3–10 nm) up to about $12\,000 \text{ cm}^{-3}$ (bursts) are observable in the size spectra in fine weather conditions during the daytime.

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

To find the aerosol particle concentration in the size range coinciding with that of aerosol ions m (2.5–8 nm), we extrapolated the size distribution below 3 nm. The upper boundary of the aerosol ion size class N (8–ca. 20 nm) is not exactly determined because of the applied method of air ion measurements by means of integral ion counters. Despite all, the analysis of the correlation between the concentrations of nanometer aerosol particles and their charged fraction (positive air ions) showed a good qualitative consistency in the case of the nucleation event days (Fig. 5). During the non-event days, the concentrations of particles and air ions in the size class of 2.5–8 nm were close to low background (see Table 2), and the correlation was poor in general, while the particles and ions in the size class of 8–ca. 20 nm could show considerably higher concentrations and, therefore, also a stronger correlation (e.g. on 9 April, the correlation coefficient was 95%).

During the nucleation event days, the enhancement of particle concentration in the size range of 2.5–8 nm precedes that in the size range of 8–20 nm as a consequence of particle growth toward large sizes. The average diurnal variation in the concentration of particles (and air ions) in the size class of 2.5–8 nm showed the maximum of 1910 cm^{-3} (120 cm^{-3}) at noon (12:00–13:00 LST), and in the size class of 8–20 nm about 3600 cm^{-3} (170 cm^{-3}) four hours later at 16–17 LST. A wide minimum of particles (and ions) in the size class of 2.5–8 nm about $20\text{--}50\text{ cm}^{-3}$ ($45\text{--}50\text{ cm}^{-3}$) was recorded during the nighttime and early morning hours at 22:00–07:00 LST; the minimum of larger particles (and ions) about $300\text{--}400\text{ cm}^{-3}$ ($45\text{--}50\text{ cm}^{-3}$) took place in the early morning at 04:00–09:00 LST.

The correlation between the concentrations of aerosol particles and air ions in Fig. 5a is substantially affected by the nucleation burst on 5 April, when the concentrations of nanometer particles (2.5–8 nm) up to $12\,030\text{ cm}^{-3}$ were recorded. Taking into account only the core of the scatterplot (data points below the particle concentration of about 5000 cm^{-3}), the correlation coefficient decreases slightly from 87.6% down to 84.6%, but the relationship (correlation curve) remains nearly the same. The coarse nanometer particles (8–20 nm) and their charged fraction (positive air ions) in Fig. 5b are even

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

more strongly correlated compared to nanometer particles; the correlation coefficient is 92%. Detailed information about the correlation coefficients for various nucleation days is presented in Table 3. The particle formation events in Table 3 have been classified into three classes according to the amount of generated new nanometer particles and the clearness of the subsequent growth of particles from the nucleation mode up to the Aitken mode size range (Kulmala et al., 2001a; Mäkelä et al., 2000). The best ones were classified as class 1. The events with the high concentration of background aerosol or non-continuous growth characteristics were classified as classes 2 or 3. We found that the correlation between the concentrations of nanometer particles and air ions was always linear during the nucleation bursts, considering different generation rates (followed by the maximum concentration of particles) and the growth rates of nanometer particles (Table 3). The event class 1 displayed the strongest correlation for the size class of 2.5–8 nm, because the maximum concentrations of air ions are then well above the background (about 40 cm^{-3}). Based on these results we can conclude that the air ion measurements can give essential information about the basic regularities of the behavior of nanometer aerosol particles.

Since the size fractions of particles measured by means of ion counters and the DMPS were related one-to-one, the scatterplot between the nanometer aerosol particles and their charged fraction (air ions) could be fitted with a line of the linear regression function, the intercept of which should be zero, and the slope giving the mean charging probability for the fraction. The results of the regression analysis showed a linear correlation between the nanometer aerosol particles and their charged fraction, but with a non-zero intercept of about $20\text{--}60 \text{ cm}^{-3}$ (Fig. 5). This step at the zero value of the aerosol concentration (intercept) is probably due to methodological and instrumental uncertainties in the air ion fraction concentration estimation by means of integral ion counters (Tammet, 1970). This step in air ion concentration (average about $40\text{--}50 \text{ cm}^{-3}$) was comparable, considering the size classes of 2.5–8 nm and 8–ca.20 nm during nucleation events or the non-event days. In this study we did not take into account the non-zero intercept and interpreted the slope of the linear regression

Positive air ions in boreal forest air

U. Hörrak et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

line as an estimate of the mean charged fraction of particles (or experimental charging probability). The estimates of the positively charged fraction of nanometer aerosol particles (2.5–8 nm) given in Table 3 varied in the range of 3–6%. The average of the most pronounced events of class 1 is about 4%. These estimates are close to the steady state bipolar charging probability of 2.6% for the mean size of the fraction of 5.3 nm (Reischl et al., 1996). Considering the measurement accuracy, we cannot state that the excess of positively charged nanometer particles indicates an overcharge of atmospheric aerosols due to the ion-induced nucleation. The charged fractions of coarse nanometer aerosol particles (8–ca. 20 nm) given in Table 3 are only the order of magnitude estimates because of the uncertainty in the upper boundary of air ion size class. Despite that, it showed a nearly constant value, average about 4%. The steady state bipolar charging probability for the mean size of the fraction of 12.7 nm should be about 7.2%.

Recent measurements during the QUEST 2 campaign showed that the charge distribution on nanometer aerosol particles was often significantly asymmetric with an excess of negative charges below the particle diameter of about 5 nm, decreasing as a function of particle size. The charging probability for the smallest nanometer particles (3–4 nm) varied during different nucleation event days in the range of about 2–6% and 2–15% for positively and negatively charged particles, respectively (Hörrak et al., 2004a, b).

4.5 Case studies

Four nucleation event days were selected for the case studies to characterize the behavior of positive air ions of different classes: small ions and charged nanometer particles (Fig. 6). Some complementary information about the behavior of air ions and aerosol particles during these nucleation event days can be found in (Hörrak et al., 2003a).

The diurnal variation in the concentration of small ions was different on different nucleation event days. On 2 April 1999, the data showed gradual increase in the

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

concentration of small ions before the onset of the generation of nanometer aerosol particles at 12:30 LST. This is in general accordance with the decrease in the ion sink due to aerosol particles (3–500 nm) from $(5–6) \times 10^{-3} \text{ s}^{-1}$ during the nighttime down to the minimum of $1.2 \times 10^{-3} \text{ s}^{-1}$ in the afternoon (Fig. 7a). The ion sink stayed relatively constant (about $1.4 \times 10^{-3} \text{ s}^{-1}$) during the intensive generation of nanometer particles. The increase in the ion sink caused by nanometer particles was compensated by the decrease in the concentration of the Aitken and accumulation mode particles. Therefore, no considerable changes in the concentration of small ions during the generation of nanometer aerosol particles can be found. The decrease followed the growth of new particles towards large sizes in the afternoon, when the particles in the size class of 8–20 nm displayed the concentration maximum. On 2 April, the concentrations of nanometer aerosol particles (2.5–8 nm) and their charged fraction (positive aerosol ions) showed only a modest correlation (the correlation coefficient was 84%), because a rise in the concentration of aerosol ions precedes (about 10 min, or one measurement cycle) that of aerosol particles measured by the DMPS, probably due to spatial inhomogeneity of the nucleation process.

A peculiarity of this day (2 April) was the fog, which was detected by means of optical spectrometers (ASASP-300, CSASP-100) during 02:00–07:00 LST in the morning with the maximum intensity around 04:50 a.m. The effect of the fog on the concentration of small ions is clearly visible in Fig. 7a, where the small ion concentration decreases independently of the ion sink (based on DMPS data) down to about 150 cm^{-3} around 04:50 a.m. The scatterplot between the small ion concentration and the ion sink in Fig. 9a revealed a separate group below the concentration of about 300 cm^{-3} . Presumably the fog affects all the data below the small ion concentration of about 300 cm^{-3} . Another separate group of data with a relatively high concentration of small ions up to 710 cm^{-3} , which corresponds to the nighttime weak winds less than 0.2 m s^{-1} , is probably due to the effect of radon on the ionization rate of the air. The nucleation event on 2 April, 1999, was analyzed in detail by Aalto et al. (2001).

The next example in Fig. 6c demonstrates the effect of clean air inflow on 10 April

**Positive air ions in
boreal forest air**

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1999. The increase in the concentration of small ions during the nighttime and early morning (up to 800 cm^{-3}) followed the decrease in the ion sink from $5.5 \times 10^{-3} \text{ s}^{-1}$ down to $3.9 \times 10^{-4} \text{ s}^{-1}$ at 06:30 LST. The ion sink was decreasing due to the inflow of clean maritime air mass from the North Atlantic (Mäkelä et al., 2001b). The gradual increase in the ion sink after 06:30 LST up to $4.1 \times 10^{-3} \text{ s}^{-1}$ at 24:00 LST was caused by the local production of aerosols (Fig. 7b). The rise in the concentration of small ions in the evening after 18:00 LST was probably due to the accumulation of radon favored during the nighttime calm and weak wind; the wind speed was below 0.5 ms^{-1} and the temperature inversion developed after 19:00 LST. This separate group with the high concentration of small ions is clearly seen in the scatterplot of the small ion concentration versus the ion sink in Fig. 9c.

A characteristic feature of 10 April was that the concentration of small ions in the clean air (very small ion sink on aerosols) was mainly determined by small ion recombination. Ion-ion recombination was responsible for 75% of total ion loss (due to small ion recombination and attachment to the aerosol particles) at 06:30 LST. In the beginning of the burst of nanometer aerosol particles, the ion-ion recombination was responsible for 40–55% of total ion loss (estimated by means of Eq. (1) considering the steady state condition and neglecting the effect of ion-induced nucleation). During the other days selected for the case studies (on 2, 5 and 14 April) the maximum percentage of ion loss caused by small ion recombination was 44%, 30% and 20%, respectively. Just before the burst of nanometer particles, these percentages were 27%, 25% and 20%, respectively. The hypothetical ion loss due to the enhanced deposition of small ions on the coniferous forest discussed in Sect. 4.3 was not taken into account in these estimates. Taking into account also the extra ion loss, the maximum percentage of small ion losses due to ion-ion recombination decreased about 4 times down to 17% on 10 April. On the other days, on 2, 5 and 14 April this percentage was 13%, 11% and 9%, respectively.

5 and 14 April 1999, were the only days during the BIOFOR III campaign when the concentration of small (or cluster) ions was considerably decreasing during the nu-

5 cleation burst (Figs. 6b, d). These days are of particular interest to the study of the hypothetical role of ion-induced nucleation on the decrease of small ion concentration. The nucleation burst on 5 April was the most intensive in April, with the peak concentration of nanometer particles (2.5–8 nm) of about $12\,000\text{ cm}^{-3}$, which is about two times
10 higher than on other days. On 5 April, the minimum of the ion sink (about $2 \times 10^{-3}\text{ s}^{-1}$) was recorded before the burst of nanometer particles at 09:00 LST (Fig. 8a). The decrease in the concentration of small ions occurred simultaneously with a rapid increase in the concentration of aerosol ions (2.5–8 nm), as well as the concentration of nanometer aerosol particles recorded by the DMPS. The oscillation in the concentration of small ions after the nucleation burst was correlated with the changes in the
15 ion sink induced by the concentration of coarse nanometer particles (8–ca. 20 nm) (Figs. 6b and 8a).

Also on 14 April, the ion sink displayed the minimum ($2.9 \times 10^{-3}\text{ s}^{-1}$) before the onset of the generation of nanometer particles at noon (12:20 LST) (Fig. 8b); the decrease of small ions followed after that. Similarly to the previous case (on 5 April), the concentrations of fine nanometer particles (2.5–8 nm) and coarse nanometer particles (8–ca. 20 nm) started to increase almost simultaneously, afterwards displaying a delay in the peak concentrations about 30 min (Figs. 6b and d). In general, the delay between the concentration maxima of two ion (or particle) classes, affected by the
20 particle growth towards large sizes after the generation, was many times longer (about 3.5 h). The growth rate of nanometer particles (in the size range of 3–10 nm) on 5 and 14 April was comparatively high: 6.7 and 5.4 nm h^{-1} , respectively (Table 3). The study of the evolution of the distribution of the ion sink on particle sizes on 5 and 14 April (Fig. 8) confirms that the decrease in the concentration of small ions was caused by an intensive generation of nanometer aerosol particles followed by a rapid growth of particles towards large sizes. The scavenging efficiency of small ions by aerosol particles in the size class of 8–20 nm is about 3 times higher compared to the particles in the size class of 2.5–8 nm. Thus, on both days the decrease in the small ion concentrations can well be explained by the effect of aerosols (Figs. 9b, d).
25

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The nucleation events on both 5 and 14 April, with a rapid increase in the concentration of charged fine nanometer particles (2.5–8 nm) and coarse nanometer particles (8–ca. 20 nm), showed a good correlation between the nanometer aerosol particles and their charged fraction (positive air ions) (Table 3). Kulmala et al. (2001b) applied an analytical model and found that the formation rate of 3 nm particles was $1.6 \text{ cm}^{-3} \text{ s}^{-1}$ on 5 April and $1.1 \text{ cm}^{-3} \text{ s}^{-1}$ on 14 April. The estimated nucleation rates (generation rate of 1 nm particles) were $10\text{--}100 \text{ cm}^{-3} \text{ s}^{-1}$ and $50\text{--}80 \text{ cm}^{-3} \text{ s}^{-1}$, respectively.

As a result of the case studies, we can conclude that both the ion sink on aerosols and small ion recombination are essential factors of small ion loss in the comparatively clean air at Hyytiälä. If the nucleation burst of nanometer particles is able to cause changes in the ion sink by about $2 \times 10^{-3} \text{ s}^{-1}$, then the effect is clearly observable in the concentration of small ions. This change in the ion sink is significant considering the extent of its variation during nucleation event days (Fig. 9).

Examination of the nucleation events selected for the case studies, as well as other events during the BIOFOR III campaign at Hyytiälä, did not show any clear indication of the ion-induced nucleation in the atmosphere – the scavenging of small ions before or during intensive generation of nanometer particles. In general, it is accepted that the nucleation on small (or cluster) ions is energetically more favored compared to particle formation by homogeneous nucleation from pure vapor(s) (Hoppel et al., 1994; Raes and Van Dingenen, 1992; Nadykto and Yu, 2003). In some cases, the barrierless nucleation can occur, e.g. nucleation of H_2SO_4 on negative cluster ions (Lowejoy et al., 2004). On the other hand, the analysis of the nucleation events by Kulmala et al. (2001b) revealed that the nucleation rate (or the formation rate of 1 nm particles) should be in the range of $10\text{--}100 \text{ cm}^{-3} \text{ s}^{-1}$ to explain the formation rate of 3 nm particles of about $1 \text{ cm}^{-3} \text{ s}^{-1}$ during the BIOFOR III campaign at Hyytiälä. Such high nucleation rates (compared to the ionization rate of about 4–6 ion pairs $\text{cm}^{-3} \text{ s}^{-1}$) should reduce the concentration of cluster ions assuming that the cluster ions are involved in the nucleation process. However, the measurements did not exhibit significant decrease in the concentration of positive cluster ions as the indication of ion-induced

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

nucleation during nucleation bursts. If the concentration decreased, then it correlated well with the ion sink due to aerosol particles. The recent QUEST 2 campaign measurements at Hyytiälä in spring 2003 indicated that also the concentration of negative cluster ions behaved similarly during the nucleation bursts; even the excess of negative intermediate ions (1.6–5 nm) and the negative overcharge on nanometer aerosol particles (3–5 nm) was found (Hörrak et al., 2004a, b; Laakso et al., 2004b). The latter is indeed a clear indication of ion-induced nucleation on negative cluster ions. As a result, we can conclude that at Hyytiälä, the positive cluster ions less likely nucleate (probably due to thermodynamic properties proposed by Froyd and Lovejoy (2003a, b)) and the ion-induced nucleation rate is always less than the ionization rate. Therefore, the cluster ion concentration does not reflect the changes caused by ion-induced nucleation, probably because its variability due to other factors (e.g. ionization rate, ion sink on aerosols, turbulent mixing of air) is commonly considerably higher.

4.6 Study of the effect of the hygroscopic growth of aerosol particles on the ion sink and on the ionization rate

The dependence of the hygroscopic growth factor (GF) of submicron particles ($d < 0.53 \mu\text{m}$) on ambient relative humidity (RH) was estimated according to the parameterization given by Laakso et al. (2004b)

$$GF = \frac{d_{\text{wet}}}{d_{\text{dry}}} = \left(1 - \frac{\text{RH}}{100}\right)^{-\frac{d_{\text{dry}}}{3215\text{nm}} - 0.0847} \quad (9)$$

Equation 9 is based on a simple model by Zhou et al. (2001), wherein the exponent as a function of particle size was found from the particle GF measurements in the size range of 10–264 nm, carried out at Hyytiälä during the BIOFOR III campaign (Hämeri et al., 2001). The model gives an estimate of the average GF for externally mixed aerosols of different hygroscopic growth modes weighted by their corresponding fractions. In general, Eq. (9) is valid in the case of ambient relative humidity $\text{RH} \leq 90\%$ and

the particle sizes below 270 nm. Above these limits, the model has not been verified due to the absence of corresponding particle GF measurements at Hyytiälä. In accordance with the recent results of the particle wet and dry size distribution measurements at Hyytiälä during May and July 2004 (Birmili et al., 2006), we have limited the GF of particles larger than 280 nm with that found for 280 nm size particles. The latter is argued, since by Birmili et al. (2006) the particles of 200–400 nm showed nearly the same hygroscopic properties. Thus, with certain reservations, the GF model (Eq. 9) can be applied for the particles up to about 500 nm and for ambient RHs up to about 95%. In the case of high RH of 96–99%, where the uncertainty of the RH measurements is about 3% and the particle GF can change by a factor of 1.5–2 (Wex et al., 2005; Svenningsson et al., 2005), the exploitation of the model is the most critical.

Statistical description of the parameters given in the Table 1, corrected by the ambient relative humidity, is presented in Table 4, together with the air temperature and relative humidity. Here we have limited the ambient relative humidity by 99% in calculations. In general, the changes in the mean and median values of recalculated parameters due to the effect of hygroscopic growth of aerosol particles (3–500 nm) are in the range of 10%–33%. The changes are about 1.5–2.3 times smaller in the case of nucleation event days if compared to non-event days owing to different meteorological conditions. Thus, the aerosol data correction by the hygroscopic growth factor accounts for about 18% and 28% increase in the mean ion sink due to aerosol particles in the case of nucleation event days and non-event days, respectively. Correspondingly, the mean ionization rate also increases by about 13% and 23% (see Tables 1 and 4). The average diurnal variations in the ion sink, effective attachment coefficient and ionization rate discussed in Sect. 4.1 showed about 25%, 33% and 12% increase in amplitude, respectively, due to higher nighttime maxima of the parameters induced by the hygroscopic GF correction of aerosol data.

The limitation of the GF for particles larger than 280 nm results in a small decrease less than 4% in the ion sink due to particles, since the ion sink at Hyytiälä was mainly caused by the accumulation mode particles. The dependence of the ion sink, found

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Positive air ions in boreal forest air

U. Hörrak et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

from the DMPS measurements applying the hygroscopic growth correction (Eq. 9), on ambient relative humidity can well be approximated by the same power function (Eq. 9) with the particle dry diameter of 154 nm in the exponent. The nonlinear correlation coefficient between the relative humidity and the ratio of ion sink corrected by the ambient humidity to that based on original data is 98.8%. The ratio increases moderately from the value of 1.07 at RH=40% up to 1.24 at RH=80% and henceforth, rises exponentially from 1.36 at RH=90% up to 1.84 at RH =99%. Nearly similar humidity dependence of the ratio of the calculated ionization rates (ratio of corrected by RH to the original) was found with the ratios of 1.06, 1.19, 1.31 and 1.84 corresponding to the relative humidities of 40%, 80%, 90% and 99%, respectively. The correlation coefficient between the ratio and RH was 97.9%. This demonstrates the importance of the GF correction of the DMPS measurements, when estimating the ion sink (especially in the case of high relative humidity) and the necessity for appropriate GF models.

In general, the contribution of wet coarse particles (0.52–32 μm) measured by the optical particle counters (ASASP-300 and CSASP-100) to the total ion sink due to the particles (taking into account also the hygroscopic growth correction of the DMPS data) was less than 8% (the mean was about 0.7%) if the RH<96%, but it increased exponentially up to an order of magnitude (up to 84%) above the RH=96%. The corresponding numbers for the percentage of the total ion loss (due to particles and the extra ion loss in the forest canopy) are, in average, about 2 times smaller if the RH <96%; the maximum in the case of RH>96% was 76%. Thus, the coarse particles (above 0.52 μm) can be ignored in calculations of the ion sink at the Hyytiälä SMEAR II station only if to exclude the situations of high relative humidity, which are often accompanied with fog formation. Such situations were eliminated from the database to determine the ionization rate (given below), and set aside for future analysis due to the lack of complete information about the size distribution of fog droplets above 32 μm necessary for the ion sink calculation.

Considering the statistical estimate of the extra ion loss of $5.2 \times 10^{-3} \text{ s}^{-1}$ due to small ion deposition inside the forest canopy, in addition to the ion sink on aerosol particles

found in Sect. 4.3, results in the ionization rate of about $4.8 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$. The recalculated values of the parameters using the same method as described in Sect. 4.3 and taking into account also the effect of hygroscopic growth of particles on the ion sink below the $\text{RH}=96\%$ are about 15% and 20% higher – $5.5 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$ and $6.3 \times 10^{-3} \text{ s}^{-1}$ respectively. The estimates of the standard error (standard deviation of the sample mean) of the parameters are $0.11 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$ and 0.21 s^{-1} , respectively. The statistics of the ionization rate (q_{θ}), calculated from the balance equation corrected by the extra ion loss (S_{θ}) of $6.3 \times 10^{-3} \text{ s}^{-1}$, are given in Table 4. The mean values of the ionization rate (q_{θ}) in Table 4 are somewhat higher, compared to the value of $5.5 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$, given above due to the contribution of the data with relative humidity above 95% and wind speed less than 1 m s^{-1} . The extra ion loss further decreases the percentage of small ion loss due to ion-ion recombination down to 6.2%, in average, and the maximum down to 15.7% (see Table 4).

Correction of the size distributions of particles measured with the DMPS by hygroscopic growth factor, as well as taking into account the particle measurements above 500 nm at ambient humidity by the optical particle counters, changes the overall poor correlation between the small ion concentration and the ion sink depicted in Fig. 3a. The nonlinear correlation coefficient (R) increases from 0.59 (the determination coefficient $R^2=0.35$) up to 0.74 (0.55). The both above-mentioned factors have nearly the same contribution (0.09 and 0.06, respectively) to the increase in the correlation coefficient by decreasing the dispersion of data-points in the case of high humidity. In Fig. 3a, the effect is the most essential in the central region of the correlation field below the regression curve, where the highly deviated data-points are moved towards higher ion sinks up to about $6 \times 10^{-2} \text{ s}^{-1}$. The increase in the median ion sink is about 33%. In the case of Fig. 3b and Fig. 4, the contribution of large aerosol particles above 500 nm to the ion sink is insignificant and the hygroscopic growth correction of the DMPS data increases the ion sink by a factor of 1.18 and 1.14, respectively. The increase in the correlation coefficient is still less than 2%.

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Regarding the examples selected for case studies in Fig. 9, the general regularities between the small ion concentration and ion sink stayed nearly the same in all four days, if to compare the ion sinks from dry and wet particle size distributions. On 5, 10 and 14 April, the maxima of ion sinks by the DMPS data considering the hygroscopic growth correction of particle sizes were about 7%, 20% and 15% higher, respectively, but the changes in the correlation (determination) coefficients were insignificant. The ratios of wet to dry ion sinks were changing in the range of 1.06–1.27, 1.05–1.2 and 1.1–1.2 during 5, 10 and 14 April, respectively, depending on the diurnal variation in relative humidity. The minima were found to take place at local noon. On 2 April, during the nucleation burst in the afternoon, the changes in the ratio of wet to dry ion sinks were similar to that found for 10 April. However, the ratio was changing drastically in the early morning during the fog (03:30–06:30), when the maximum ratio of about 9 was recorded at 5 o'clock. Then the ion sink from the wet large particles (0.52–32 μm) was about 80% of the total ion sink due to aerosol particles, and the hygroscopic growth correction of the DMPS measurements (3–500 nm) increased the corresponding ion sink by a factor of 1.7. During the other days, the contribution of large particles was less than 6%, commonly 0.5–2%. Thus, taking into account the above-mentioned corrections in the ion sink, the data-points corresponding to the high relative humidity and fog event in Fig. 9a (brown squares) are spread towards considerably higher ion sinks up to $4.9 \times 10^{-2} \text{ s}^{-1}$. Therefore, considering all the data on 2 April, the determination coefficient between the concentration of small ions and ion sink increases from 87.1% to 89.5%.

5 Conclusions

On the basis of the measurements carried out during the BIOFOR III campaign, we can point out the following characteristics of the behaviour of positive air ions at boreal forest in early spring.

The concentration of small (or cluster) ions at the Hyytiälä station varied from 110

to 1183 cm^{-3} , the average was about 480 cm^{-3} . The highest concentrations were recorded during the nighttime calms and after the inflow of clean maritime air masses. The smallest concentrations below about 300 cm^{-3} belonged to fog events and polluted continental air masses with a high aerosol load. No clear indication of ion-induced nucleation (scavenging of small ions) was found considering the behavior of positive small ions during nucleation events.

The variation in the concentration of small ions can well be explained by the changes in ion loss due to aerosols (ion sink). Exceptions are the nighttime calms and weak wind, when the increase in the concentration of small ions (up to about 1000 cm^{-3}) is probably due to the effect of radon on the ionization rate of air. The concentration of small ions decreases independently of fine aerosol particles (3–500 nm) measured by the DMPS at high humidity (above 96%), presumably due to the increasing ion sink caused by hygroscopic growth of particles and scavenging by fog droplets. The nonlinear correlation between the small ion concentration and the ion sink for the selected hours (wind speed $>0.6\text{ m s}^{-1}$, relative humidity $<95\%$) was -87% . Study of the effect of the hygroscopic growth of aerosol particles on the ion sink and ionization rate demonstrates the importance of the hygroscopic growth correction of the aerosol particle size distributions measured by the DMPS, as well as the importance of the contribution of wet coarse particles above 500 nm in the case of high relative humidity ($>96\%$) and fog events.

In addition to small ion loss due to recombination and attachment to aerosol particles, the term of an “extra ion loss”, presumably due to small ion deposition on coniferous forest, is needed to explain the observations. This extra ion loss, which is nearly equal to the mean ion sink induced by aerosol particles, is significant and should be taken into account, when estimating the balance of small ions inside the forest. The loss of small ions by small ion recombination cannot be ignored; its contribution to the total ion loss was about 6% in average; the maximum of 16% was recorded in the case of Arctic clean air advection from the North Atlantic.

The ion sink on aerosols varied in the range of 4.5×10^{-4} – $6.6\times 10^{-2}\text{ s}^{-1}$ (commonly

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1×10^{-3} – $1.2 \times 10^{-2} \text{ s}^{-1}$) at the Hyytiälä station. It was mainly determined by the concentration of particles in the size range of 3–500 nm; the contribution of larger particles of the sizes of 0.52–32 μm was about 1–2% in average and always less than 8% if the relative air humidity was less than 96%. However, sometimes the contribution of coarse particles can be as high as 70%–80% in the case of high relative humidities and fog. In general, the main contribution was made by the particles in the Aitken and accumulation mode size ranges. The nucleation mode particles (3–20 nm) had an effect on the concentration of small ions only sometimes during an intensive generation of nanometer particles followed by a rapid growth of particles towards large sizes.

In the nucleation event days, the minimum in the ion sink occurred before the onset of the generation of nanometer particles or at about the burst peak. Before the onset, the fine aerosols (3–500 nm) were responsible for 37–88% (commonly 65–83%) of the total ion loss due to small ion recombination and attachment to the aerosol particles. The mean ion sink on aerosols was about 1.8 times smaller during the nucleation event days compared to the non-event days, and the mean concentration of small ions was correspondingly higher: 530 cm^{-3} versus 424 cm^{-3} .

The ion sink is affected by the total concentration of particles and by the changes in the effective attachment coefficient of small ions onto aerosol particles. The effective attachment coefficient (which is nearly proportional to the mean diameter of aerosol particles) varied in the range of 2×10^{-7} – $6.2 \times 10^{-6} \text{ cm}^3 \text{ s}^{-1}$. It showed a considerable diurnal variation during the nucleation burst days due to transformations in the aerosol particle size distribution initiated by the mixing of boundary layer and new particle formation. During the peak time of nanometer particle generation around noon, the effective attachment coefficient was almost always less than $6 \times 10^{-7} \text{ cm}^3 \text{ s}^{-1}$.

The average ionization rate of about $3.3 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$, calculated from the small ion balance equation applying experimental data of aerosol particle size distribution (with hygroscopic growth correction of particle sizes) and small ion concentration measurements, is too small for continental stations. The relationship between the concentration of small ions and the ion sink on aerosol particles is far from the dependence

**Positive air ions in
boreal forest air**

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

predicted by the balance equation of small ions. The best fit to experimental data was found assuming an extra ion loss of $6.3 \times 10^{-3} \text{ s}^{-1}$ (presumably due to small ion deposition on the coniferous forest) in addition to the ion sink due to particles. Therefore, the estimated average total ionization rate at the Hyytiälä station at a height of about 2 m above the ground during early spring, when the ground was wet and partly covered with snow, was found to be about $6.2 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$ with the standard deviation of sample of $1.9 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$.

The analysis of the correlation between the concentrations of nanometer aerosol particles and their charged fraction (positive air ions) showed a good qualitative consistency. Considering different nucleation event days when the concentration of positively charged nanometer particles (2.5–8 nm) exceeded 120 cm^{-3} , the linear correlation coefficients varied in the range of 84–98%. The correlation is also good for coarse nanometer particles and their charged fraction (8–ca. 20 nm); the correlation coefficients are in the range of 87–98%. These particles are almost quasi-steady state charged. The estimates of the charged fraction of nanometer aerosol particles (2.5–8 nm) are in the range of 3–6%, which are close to the steady state bipolar charging probability of 2.7% for the mean size of the fraction of 5.3 nm. Considering the measuring uncertainties, we cannot state that the excess of the charged nanometer particles of positive polarity is an indication of the overcharge of atmospheric aerosols due to an ion-induced nucleation process. Further study is necessary to specify the nucleation mechanism.

Acknowledgements. This research has in part been supported by the Nessling Foundation, the Nordic Center of Excellence (BACCI), the Estonian Science Foundation through grants 4622 and 6223 and the University of Tartu research project of PP1FY07913.

References

Aalto, P., Hämeri, K., Becker, E., Weber, R., Salm, J., Mäkelä, J. M., Hoell, C., O'Dowd, C. D., Karlsson, H., Hansson, H.-C., Väkevä, M., Koponen, I., Buzorius G., and Kulmala M.:

Physical characterization of aerosol particles during nucleation events, *Tellus*, 53B, 344–258, 2001.

Beig, G. and Brasseur, G. P.: Model of tropospheric ion composition: A first attempt, *J. Geophys. Res.* 105, 22 671–22 684, 2000.

5 Birmili, W., Berresheim, H., Plass-Dülmer, C., Elste, T., Gilge, S., Wiedensohler, A., and Uhrner, U.: The Hohenpeissenberg aerosol formation experiment (HAFEX): A long-term study including size-resolved aerosol, H₂SO₄, OH, and monoterpenes measurements, *Atmos. Chem. Phys.*, 3, 361–376, 2003,
<http://www.atmos-chem-phys.net/3/361/2003/>.

10 Birmili, W., Schwirn, K., Nowak, A., Petäjä, T., Rose, D., Hämeri, K., Aalto, P., Wiedensohler, A., Kulmala, M., and Boy, M. Hygroscopic growth of particle number size distributions at the atmospheric research station Hytytiälä, Finland, Report Series in Aerosol Science (Helsinki), 81A, 62–67, 2006.

Chalmers, J. A.: *Atmospheric Electricity*, Pergamon Press, Oxford, London, 1967.

15 Dolezalek, H.: The atmospheric electric fog effect, *Rev. Geophys.*, 1(2), 231–282, 1963.

Eisele, F. L. and Tanner, D. J.: Identification of ions in continental air, *J. Geophys. Res.*, 95, 20 539–20 550, 1990.

Froyd, K. D. and Lovejoy, E. R.: Experimental thermodynamics of cluster ions composed of H₂SO₄ and H₂O. 1. Positive ions, *J. Phys. Chem. A*, 107, 9800–9811, 2003a.

20 Froyd, K. D. and Lovejoy, E. R.: Experimental thermodynamics of cluster ions composed of H₂SO₄ and H₂O. 2. Measurements and ab initio structures of negative ions, *J. Phys. Chem. A*, 107, 9812–9824, 2003b.

Fuchs, N. A. and Sutugin, A. G.: High-dispersed aerosols, in: *Topics in Current Aerosol Research*, edited by: Hidy, G. M. and Brock, J. R., vol. 2, pp. 1–60, Pergamon, Oxford, 1971.

25 Hanson, D. R. and Eisele, F. L.: Measurement of prenucleation molecular clusters in the NH₃, H₂SO₄, H₂O system, *J. Geophys. Res.*, 107(D12), 4158, doi:10.1029/2001JD001100, 2002.

Hari, P. and Kulmala, M.: Station for Measuring Ecosystem–Atmosphere Relations (SMEAR II), *Boreal Environ. Res.*, 10, 315–322, 2005.

30 Hatakka, J., Paatero, J., Viisanen, Y., and Mattsson, R.: Variations of external radiation due to meteorological and hydrological factors in central Finland, *Radiochemistry*, 40(6), 515–519, MAIK Nauka/ Interperiodica Publishing, 1998.

Hirsikko, A., Laakso, L., Hörrak, U., Aalto, P. P., Kerminen, V.-M., and Kulmala, M.: Annual and size dependent variation of growth rates and ion concentrations in boreal forest, *Boreal Env.*

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Res., 10, 357–369, 2005
- Hoppel, W. A.: Ion-aerosol attachment coefficients, ion depletion, and the charge distribution on aerosols, *J. Geophys. Res.*, 90, 5917–5923, 1985.
- Hoppel, W. A. and Frick, G. M.: Ion-aerosol attachment coefficients and the steady-state charge distribution on aerosols in a bipolar ion environment, *Aerosol Sci. Technol.*, 5, 1–21, 1986.
- Hoppel, W. A. and Frick, G. M.: The nonequilibrium character of the aerosol charge distributions produced by neutralizers, *Aerosol Sci. Technol.*, 12, 471–496, 1990.
- Hoppel, W. A., Anderson, R. V., and Willett, J. C.: Atmospheric electricity in the planetary boundary layer, In: *The Earth's Electrical Environment*, National Academy Press, Washington, D.C., 149–165, 1986.
- Hoppel, W. A., Frick, G. M., Fitzgerald, J. W., and Larson, R. E.: Marine boundary layer measurements of new particle formation and the effects nonprecipitating clouds have on aerosol size distribution, *J. Geophys. Res.*, 99, 14 443–14 459, 1994.
- Hörrak, U., Salm, J., and Tammet, H.: Bursts of intermediate ions in atmospheric air, *J. Geophys. Res.*, 103, 13 909–13 915, 1998a.
- Hörrak, U., Mirme, A., Salm, J., Tamm, E., and Tammet, H.: Air ion measurements as a source of information about atmospheric aerosols, *Atmospheric Research*, 46, 233–242, 1998b.
- Hörrak, U., Salm, J., and Tammet, H.: Statistical characterization of air ion mobility spectra at Tahkuse Observatory: Classification of air ions, *J. Geophys. Res.*, 105, 9291–9302, 2000.
- Hörrak, U., Aalto, P., Salm, J., and Kulmala, M.: Characterization of air ions during nucleation events in boreal forest air, *Report Series in Aerosol Science (Helsinki)*, 59, 196–201, 2003a.
- Hörrak U., Salm J., and Tammet H.: Diurnal variation in the concentration of air ions of different mobility classes at a rural area, *J. Geophys. Res.*, 108(D20), 4653, doi:10.1029/2002JD003240, 2003b.
- Hörrak, U., Tammet, H., Aalto, P. P., and Kulmala, M: Charge distribution on nanometer aerosol particles in the atmosphere during nucleation burst events, *J. Aerosol Sci.*, 35, S299–S300, 2004a.
- Hörrak, U., Tammet, H., Aalto, P. P., and Kulmala, M.: Study of air ions and nanometer particles in the atmosphere during nucleation burst events, *Report Series in Aerosol Science (Helsinki)*, 71A, 184–189, 2004b.
- Hämeri, K., Väkevää, M., Aalto, P. P., Kulmala, M., Swietlicki, E., Zhou, J., Seidl, W., Becker, E., and O'Dowd, C. D.: Hygroscopic and CCN properties of aerosol particles in boreal forests, *Tellus*, 53B, 359–379, 2001.

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Positive air ions in
boreal forest air**

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- lida, K., Stolzenburg, M., McMurry, P., Dunn, M. J., Smith, J. N., Eisele, F., and Keady, P.: Contribution of ion-induced nucleation to new particle formation: Methodology and its application to atmospheric observations in Boulder, Colorado, *J. Geophys. Res.*, 111, D23201, doi:10.1029/2006JD007167, 2006.
- 5 Israëi, H.: Atmospheric Electricity, Vol. I, Israel Program for Scientific Translations, Jerusalem, 1970.
- Israëi, H.: Atmospheric Electricity, Vol. II, Israel Program for Scientific Translations, Jerusalem, 1973.
- Kim, T. O., Adachi, M., Okuyama, K., and Seinfeld, J. H.: Experimental measurement of competitive ion-induced and binary homogeneous nucleation in SO₂/H₂O/N₂ mixtures, *Aerosol Sci. Technol.*, 26, 527–543, 1997.
- 10 Kim, T. O., Ishida, T., Adachi, M., Okuyama, K., and Seinfeld, J. H.: Nanometer-sized particle formation from NH₃/SO₂/H₂O/air mixtures by ionizing irradiation, *Aerosol Sci. Technol.*, 29, 111–125, 1998.
- 15 Korhonen, P., Kulmala, M., Laaksonen, A., Viisanen, Y., McGraw, R., and Seinfeld, J.: Ternary nucleation of H₂SO₄, NH₃ and H₂O in the atmosphere, *J. Geophys. Res.*, 104, 26 349–26 353, 1999.
- Kulmala, M., Pirjola, L., and Mäkelä, J. M.: Stable sulphate clusters as a source of new atmospheric particles, *Nature*, 404, 66–69, 2000.
- 20 Kulmala, M., Hämeri, K., Aalto, P. P., Mäkelä, J. M., Pirjola, L., Nilsson, E. D., Buzorius, G., Rannik, Ü., Dal Maso, M., Seidl, W., Hoffman, T., Janson, R., Hansson, H.-C., Viisanen, Y., Laaksonen, A., and O’Dowd, C. D.: Overview of the international project on biogenic aerosol formation in the boreal forest (BIOFOR), *Tellus*, 53B, 324–343, 2001a.
- Kulmala, M., Dal Maso, M., Mäkelä, J. M., Pirjola, L., Väkevä, M., Aalto, P. P., Miiikkulainen, P., Hämeri, K., and O’Dowd, C. D.: On the formation, growth and composition of nucleation mode particles, *Tellus*, 53B, 479–490, 2001b.
- 25 Kulmala, M., Vehkamäki, H., Petäjä, T., Dal Maso M., Lauri, A., Kerminen, V.-M., Birmili, W., and McMurry, P. H.: Formation and growth rates of ultrafine atmospheric particles: a review of observations, *J. Aerosol Sci.*, 35, 143–176, 2004a.
- 30 Kulmala, M., Laakso, L., Lehtinen, K. E. J., Riipinen, I., Dal Maso, M., Anttila, T., Kerminen, V.-M., Hörrak, U., Vana, M., and Tammet, H.: Initial steps of aerosol growth, *Atmos. Chem. Phys.*, 4, 2553–2560, 2004b.
- Laakso L., Mäkelä, J. M., Pirjola, L., and Kulmala, M.: Model studies on ion-induced nucleation

**Positive air ions in
boreal forest air**

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- in the atmosphere, *J. Geophys. Res.*, 107(D20), 4427, doi:10.1029/2002JD002140, 2002.
- Laakso, L., Petäjä, T., Lehtinen, K. E. J., Kulmala, M., Paatero, J., Hörrak, U., Tammet, H., and Joutsensaari, J.: Ion production rate in a boreal forest based on ion, particle and radiation measurements, *Atmos. Chem. Phys.*, 4, 1933–1943, 2004a.
- 5 Laakso, L., Anttila, T., Lehtinen, K. E. J., Aalto, P. P., Kulmala, M., Hörrak, U., Paatero, J., Hanke, M., and Arnold, F.: Kinetic nucleation and ions in boreal forest particle formation events, *Atmos. Chem. Phys.*, 4, 2353–2366, 2004b.
- Lovejoy, E., Curtius, J., and Froyd, K.: Atmospheric ion-induced nucleation of sulfuric acid and water, *J. Geophys. Res.*, 109, doi:10.1029/2003JD004 460, 2004.
- 10 Luts, A. and Parts, T.-E.: Evolution of negative small air ions at two different temperatures, *J. Atmos. Solar-Terr. Phys.*, 64, 763–774, 2002.
- Mäkelä, J. M., Dal Maso, M., Pirjola, L., Keronen, P., Laakso, L., Kulmala, M., and Laaksonen, A.: Characteristics of the aerosol particle formation events observed at a boreal forest site in southern Finland, *Boreal Environment Research*, 4, 299–313, 2000.
- 15 Mäkelä, J. M., Salm, J., Smirnov, V. V., Koponen, I., Paatero, J., and Pronin, A. A.: Electrical charging state of fine and ultrafine particles in boreal forest air, *J. Aerosol Sci.*, 32, S149–S150, 2001a.
- Mäkelä, J. M., Yli-Koivisto, S., Hiltunen, V., Seidl, W., Swietlicki, E., Teinilä, K., Sillanpää, M., Koponen, I. K., Paatero, J., Rosman, K., and Hämeri, K.: Chemical composition of aerosol during particle formation events in Boreal forest, *Tellus*, 53B, 380–393, 2001b.
- 20 Mäkelä, J. M., Salm, J., Smirnov, V. V., Koponen, I., Paatero, J., and Pronin, A. A.: Measurements of the mobility distribution of air ions as a source of information for the study of aerosol generation, In: *Proceedings of 12th International Conference on Atmospheric Electricity*, 2003, Versailles, 793–796, 2003.
- 25 Nadykto, A. and Yu, F.: Uptake of neutral polar vapour molecules by charged particles: Enhancement due to dipole-charge interaction, *J. Geophys. Res.*, 108(D23), 4717, doi:10.1029/2003JD003664, 2003.
- Nagato, K. and Ogawa, T.: Evolution of tropospheric ions observed by an ion mobility spectrometer with a drift tube, *J. Geophys. Res.* 103, 13 917–13 925, 1998.
- 30 Napari, I., Kulmala, M., and Vehkamäki, H.: Ternary nucleation of inorganic acids, ammonia, and water, *J. Chem. Phys.*, 117, 8418–8425, 2002.
- Nilsson, E.D., Paatero, J., and Boy, M.: Effects of air masses and synoptic weather on aerosol formation in the continental boundary layer, *Tellus*, 53B, 462–478, 2001a.

Nilsson, E. D., Rannik, Ü., Buzorius, G., O'Dowd, C., Boy, M., Paatero J., and Laakso, L.: Effects of the continental boundary layer evolution, convection, turbulence and entrainment on aerosol formation, *Tellus*, 53B, 441–461, 2001b.

Parts, T.-E. and Luts, A.: Observed and simulated effects of certain pollutants on small air ion spectra: I. Positive ions, *Atmos. Environ.*, 38, 1283–1289, 2004.

Penttinen, L., Paatero, J., Viisanen, Y., and Kulmala, M.: Seasonal and diurnal variation of airborne radon-222 at Hyytiälä, Finland, *Report Series in Aerosol Science (Helsinki)*, 59, 79–82, 2003.

Porstendörfer, J.: Properties and behaviour of radon and thoron and their decay products in the air, *J. Aerosol Sci.*, 25, 219–263, 1994.

Raes, F. and Janssens, A.: Ion-induced aerosol formation in a $\text{H}_2\text{O}-\text{H}_2\text{SO}_4$ system. I. Extension of the classical theory and search for experimental evidence, *J. Aerosol Sci.*, 16, 217–227, 1985.

Raes, F. and Van Dingenen, R.: Simulation of condensation and cloud condensation nuclei from biogenic SO_2 in the remote marine boundary layer, *J. Geophys. Res.*, 97, 12 901–12 912, 1992.

Ramamurthi, M., Strydom, R., Hopke, P. K., and Holub, R. F.: Nanometer and ultrafine aerosols from radon radiolysis, *J. Aerosol Sci.*, 24, 393–407, 1993.

Reischl, G. P., Mäkelä, J. M., Karch, R., and Necid, J.: Bipolar charging of ultrafine particles in the size range below 10 nm, *J. Aerosol Sci.*, 27, 931–949, 1996.

Svenningsson, B., Rissler, J., Swietlicki, E., Mircea, M., Bilde, M., Facchini, M. C., Decesari, S., Fuzzi, S., Zhou, J., Mønster, J., and Rosenørn, T.: Hygroscopic growth and critical supersaturations for mixed aerosol particles of inorganic and organic compounds of atmospheric relevance, *Atmos. Chem. Phys. Discuss.*, 5, 2833–2877, 2005, <http://www.atmos-chem-phys-discuss.net/5/2833/2005/>.

Tamm, E., Hörrak, U., Mirme, A., and Vana, M.: On the charge distribution on atmospheric nanoparticles, *J. Aerosol Sci.*, 32, S347–S348, 2001.

Tammet, H.: The aspiration method for the determination of atmospheric ion-spectra, *Israel Program for Scientific Translations*, Jerusalem, 1970.

Tammet, H.: Aerosol electrical density: Interpretation and principles of measurement, *Report Series in Aerosol Science (Helsinki)*, 19, 128–133, 1991.

Tammet, H. and Kimmel, V.: Electrostatic deposition of radon daughter clusters on the trees, *J. Aerosol Sci.*, 29, S473–S474, 1998.

Positive air ions in boreal forest airU. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Tammet, H., Kimmel, V., and Israelsson, S.: Effect of atmospheric electricity on dry deposition of airborne particles from atmosphere, *Atmos. Environ.*, 35, 3413–3419, 2001.
- Tammet, H., Hörrak, U., Laakso, L., and Kulmala, M.: Factors of air ion balance in a coniferous forest according to measurements in Hyytiälä, Finland, *Atmos. Chem. Phys.*, 6, 3377–3390, 2006,
5 <http://www.atmos-chem-phys.net/6/3377/2006/>.
- Tunved, P., Hansson, H.-C., Kulmala, M., Aalto, P., Viisanen, Y., Karlsson, H., Kristensson, A., Swietlicki, E., Dal Maso, M., Strom, J., and Komppula, M.: One year boundary layer aerosol size distribution data from five nordic background stations, *Atmos. Chem. Phys.*, 3, 2183–2205, 2003,
10 <http://www.atmos-chem-phys.net/3/2183/2003/>.
- Vesala, T., Haataja, J., Aalto, P., Altimir, N., Buzorius, G., Garam, E., Hämeri, K., Ilvesniemi, H., Jokinen, V., Keronen, P., Lahti, T., Markkanen, T., Mäkelä, J.M., Nikinmaa, E., Palmroth, S., Palva, L., Pohja, T., Pumpanen, J., Rannik, Ü., Siivola, E., Ylitalo, H., Hari, P., and Kulmala, M.: Long-term field measurements of atmosphere-surface interactions in boreal forest combining forest ecology, micrometeorology, aerosol physics and atmospheric chemistry, *Trends in Heat, Mass and Momentum Transfer*, 4, 17–35, 1998.
15
- Wex, H., Kiselev, A., Stratmann, F., Zoboki, J., and Brechtel, F.: Measured and modeled equilibrium sizes of NaCl and $(\text{NH}_4)_2\text{SO}_4$ particles at relative humidities up to 99.1%, *J. Geophys. Res.*, 110, D21212, doi:10.1029/2004JD005507, 2005.
20
- Wilhelm, S., Eichkorn, S., Wiedner, D., Pirjola, L., and Arnold, F.: Ion-induced aerosol formation: new insights from laboratory measurements of mixed cluster ions $\text{HSO}_4^-(\text{H}_2\text{SO}_4)_a(\text{H}_2\text{O})_w$ and $\text{H}^+(\text{H}_2\text{SO}_4)_a(\text{H}_2\text{O})_w$, *Atmos. Environ.*, 38, 1735–1744, 2004.
- Yu, F. and Turco, R. P.: Ultrafine aerosol formation via ion-mediated nucleation, *Geophys. Res. Lett.*, 27, 883–886, 2000.
25
- Yu, F. and Turco, R. P.: From molecular clusters to nanoparticles: Role of ambient ionization in tropospheric aerosol formation, *J. Geophys. Res.*, 106, 4797–4814, 2001.
- Zhou, J., Swietlicki, E., Berg, O. H., Aalto, P. P., Hämeri, K., Nilsson, E. D., and Leck, C.: Hygroscopic properties of aerosol particles over the central Arctic Ocean during summer, *J. Geophys. Res.* 106, 32 111–32 123, 2001.
30

Positive air ions in boreal forest air

U. Hörrak et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Positive air ions in boreal forest air

U. Hörrak et al.

Table 1. Descriptive statistics of the positive small ion concentration (n), effective attachment coefficient (β_{eff}), the sink of small ions on aerosol particles of 3–500 nm (S_a), ion loss due to the recombination of small ions ($\alpha \cdot n$), the ionisation rate (q), the total concentration of aerosol particles (N_{tot}) and the ratio of ion loss due to the recombination to total ion loss due to particles (last row). 31 March–29 April 1999, Hyytiälä.

a) Nucleation event days (number of measurements 1813).

	Mean	Median	Min	Max	Lower Quartile	Upper Quartile	Percentile 10	Percentile 90	Std.Dev.
n , cm^{-3}	530	513	110	1183	430	613	364	731	145
β_{eff} , $10^{-6} \text{cm}^3 \text{s}^{-1}$	1.27	1.17	0.20	3.30	0.63	1.83	0.41	2.35	0.73
S_a , 10^{-3}s^{-1}	4.15	3.92	0.39	11.90	2.72	5.42	1.67	6.70	1.98
αn , 10^{-3}s^{-1}	0.795	0.769	0.165	1.775	0.645	0.92	0.545	1.096	0.218
q , $\text{cm}^{-3} \text{s}^{-1}$	2.49	2.30	0.77	7.57	1.84	2.93	1.47	3.74	0.94
N_{tot}	4072	3474	588	18980	2278	5343	1422	7358	2579
$\alpha n / (\alpha \cdot n + S_a)$	0.199	0.164	0.024	0.746	0.113	0.263	0.076	0.352	0.126

b) Non-event days (number of measurements 1754).

	Mean	Median	Min	Max	Lower Quartile	Upper Quartile	Percentile 10	Percentile 90	Std.Dev.
n , cm^{-3}	424	410	120	1175	282	539	219	659	180
β_{eff} , $10^{-6} \text{cm}^3 \text{s}^{-1}$	2.23	2.24	0.63	3.85	1.64	2.85	1.13	3.17	0.74
S_a , 10^{-3}s^{-1}	7.59	5.37	1.18	58.70	4.05	8.36	3.41	16.00	6.39
$\alpha \cdot n$, 10^{-3}s^{-1}	0.636	0.614	0.181	1.762	0.422	0.809	0.328	0.989	0.27
q , $\text{cm}^{-3} \text{s}^{-1}$	3.03	2.77	0.55	14.56	1.89	3.77	1.39	4.89	1.59
N_{tot}	3518	2910	536	25449	2050	4338	1388	5827	2623
$\alpha \cdot n / (\alpha \cdot n + S_a)$	0.110	0.104	0.005	0.483	0.050	0.152	0.022	0.201	0.073

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Positive air ions in
boreal forest air

U. Hörrak et al.

Table 2. Descriptive statistics of the concentration of positive aerosol ions: light intermediate ions (m), heavy intermediate ions (N), and the corresponding classes of nanometer aerosol particles (A) of the sizes of 2.5–8 nm and 8–20 nm during the nucleation event days/non-event days. The number of measurements during the nucleation event days is 1506 and on the non-event days 1503. 31 March–29 April 1999, Hyytiälä.

	Mean	Median	Min*	Max	Lower Quartile	Upper Quartile	Percentile 10	Percentile 90	Std.Dev.
m , cm^{-3}	67/36	57/33	0/0	428/130	38/21	81/48	27/11	125/63	45/21
N , cm^{-3}	96/50	77/47	0/0	414/295	41/26	140/66	17/13	196/89	72/35
$A_{2.5-8}$, cm^{-3}	579/78	123/19	0/0	12030/2290	22/4	671/66	4/0.2	1777/217	1067/164
A_{8-20} , cm^{-3}	1462/371	667/186	23/6	11040/7477	283/73	2186/389	144/38	3941/769	1667/657

* Negative values are corrected to be equal to zero.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Positive air ions in
boreal forest air

U. Hörrak et al.

Table 3. The classification of nucleation events and the growth rates of newly formed particles during the BIOFOR III campaign (Kulmala et al., 2001a), positively charged fractions of aerosol particles, the correlation coefficients between the concentrations of positive air ions and aerosol particles in coinciding size ranges, and the maximum concentrations of aerosol particles during various nucleation event days. 2–29 April 1999, Hyytiälä.

Date	Class	Growth rate (nm/h)	Size class of particles 2.5–8 nm			Size class of particles 8–20 nm		
			Charged fraction (%)	Correlation coefficient (%)	Maximum of aerosols (cm ⁻³)	Charged fraction (%)	Correlation coefficient (%)	Maximum of aerosols (cm ⁻³)
990 402	1	6.7	4.9	83.6	2050	3.8	98.1	6810
990 403	2	2.2	3.0	43.5	840	4.2	95.4	3590
990 404	1	3.5	3.2	94.5	6470	3.9	94.5	9040
990 405	1	6.7	3.5	98.3	12030	3.5	94.1	11040
990 406	1	3.5	5.0	91.3	3290	4.0	87.7	5640
990 408	2	5.4	4.7	85.5	1790	4.3	95.7	4240
990 410	3	2.2	6.0	89.6	3500	3.9	90.2	3650
990 412	1	4.4	3.2	88.2	4390	3.7	95.6	4630
990 413	2	2.2	3.8	52.4	1400	3.4	92.3	5320
990 414	1	5.4	4.4	94.3	3480	4.2	95.8	4780
990 419	1	4.4	4.0	94.4	4450	5.1	89.8	2430
990 421	3	6.7	1.2*	4.4	270	2.0*	35.4	1710
990 427	2	4.4	0.4*	5.3	1080	0.4*	2.9	800
990 429	3	3.5	5.6	91.4	2770	4.4	97.5	5370

* Estimates are of low significance.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 4. The statistics of positive small ion balance in the atmosphere corrected by ambient relative humidity: effective attachment coefficient (β_{eff}), the sink of small ions on aerosol particles of 3–500 nm (S_a), the ionization rate (q), the ratio of ion loss due to the recombination to total ion loss due to particles ($\alpha \cdot n / (\alpha \cdot n + S_a)$), and corrected by the extra ion loss due to the forest ($S_e = 6.28 \times 10^{-3} \text{ s}^{-1}$) the values of the ionization rate (q_e) and the ratio of $\alpha \cdot n / (\alpha \cdot n + S_a + S_e)$. The air temperature (T) and relative humidity (RH) are presented in the last rows of the tables as well. 31 March–29 April 1999, Hyytiälä.

a) Nucleation event days (number of measurements 1813).

	Mean	Median	Min	Max	Lower Quartile	Upper Quartile	Percentile 10	Percentile 90	Std.Dev.
$\beta_{\text{eff}}, 10^{-6} \text{ cm}^3 \text{ s}^{-1}$	1.54	1.33	0.20	4.43	0.68	2.21	0.44	3.07	1.00
$S_a, 10^{-3} \text{ s}^{-1}$	4.88	4.58	0.45	13.0	2.99	6.38	1.80	8.52	2.48
$q, \text{ cm}^{-3} \text{ s}^{-1}$	2.82	2.63	1.04	8.71	2.11	3.32	1.60	4.20	1.06
$\alpha \cdot n / (\alpha \cdot n + S_a)$	0.182	0.141	0.018	0.721	0.096	0.243	0.063	0.334	0.124
$q_e, \text{ cm}^{-3} \text{ s}^{-1}$	6.15	5.63	1.74	13.4	5.20	6.77	4.84	8.51	1.51
$\alpha \cdot n / (\alpha \cdot n + S_a + S_e)$	0.071	0.066	0.011	0.157	0.051	0.090	0.037	0.111	0.071
RH, %	60.7	58.7	25.4	98.3	44.7	75.6	36.4	89.5	19.0
$T, ^\circ\text{C}$	3.4	3.5	-6.2	15.6	0.8	5.9	-2.1	8.2	4.2

b) Non-event days (number of measurements 1754).

	Mean	Median	Min	Max	Lower Quartile	Upper Quartile	Percentile 10	Percentile 90	Std.Dev.
$\beta_{\text{eff}}, 10^{-6} \text{ cm}^3 \text{ s}^{-1}$	2.98	2.95	0.75	6.20	2.09	3.70	1.39	4.67	1.16
$S_a, 10^{-3} \text{ s}^{-1}$	9.74	6.94	1.71	66.4*	5.37	11.19	4.26	18.55	7.46
$q, \text{ cm}^{-3} \text{ s}^{-1}$	3.71	3.38	0.99	16.3*	2.56	4.36	2.14	5.45	1.70
$\alpha \cdot n / (\alpha \cdot n + S_a)$	0.090	0.081	0.005	0.393	0.036	0.125	0.018	0.177	0.065
$q_e, \text{ cm}^{-3} \text{ s}^{-1}$	6.38	6.07	1.81	17.8*	4.91	7.34	4.16	8.81	2.23
$\alpha \cdot n / (\alpha \cdot n + S_a + S_e)$	0.046	0.044	0.004	0.147	0.024	0.061	0.014	0.083	0.027
RH, %	77.5	79.5	35.9	100	63.8	94.8	50.9	98.0	18.0
$T, ^\circ\text{C}$	5.9	4.7	-1.7	18.8	2.8	8.2	1.4	12.1	4.3

* Elimination of the probable local pollution episodes (two events with 47 measurements), reduces the maximum of the ion sink (S_a) down to $3.47 \times 10^{-2} \text{ s}^{-1}$ in Table 4b and the corresponding maximum values of the ionization rates (q and q_e) down to about 10 ion pairs $\text{cm}^{-3} \text{ s}^{-1}$ and 16 ion pairs $\text{cm}^{-3} \text{ s}^{-1}$, respectively.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Positive air ions in
boreal forest air

U. Hörrak et al.




Fig. 1. Equivalent or single size ion-aerosol attachment coefficient as a function of particle size (2 nm–10 μm). Solid line: model calculations (Eqs. 4–6); dashed line: approximation function (Eq. 7); plus sign (+) and open circles (O): calculations based on Eq. (6) and the tabulated results of attachment coefficients for ions with the mobility of $1.3\text{ cm}^2\text{ V}^{-1}\text{ s}^{-1}$ and the mass of 200 amu; and of $1.8\text{ cm}^2\text{ V}^{-1}\text{ s}^{-1}$ and the mass of 100 amu, respectively (Hoppel and Frick, 1990).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

EGU

Positive air ions in
boreal forest air

U. Hörrak et al.




Fig. 2. Ion sink as a function of particle size (factor $dN/d\log(d)\cdot w(d)$) for the nucleation event days, non-event days and the average of the period of 31 March–29 April 1999. Hyytiälä station.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

EGU

Positive air ions in
boreal forest air

U. Hörrak et al.




Fig. 3. Scatterplot of the concentration of positive small air ions (n) versus ion sink (S_a) due to aerosol particles (3–500 nm) and a logarithmic regression line. **(a)** all the data, **(b)** selected data when no fog was recorded, relative humidity is less than 95%, and the wind speed is higher than 0.6 m s^{-1} . 31 March–29 April 1999, Hyttiälä station.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Positive air ions in
boreal forest air

U. Hörrak et al.




Fig. 4. Scatterplot of the concentration of positive small air ions (n) versus ion sink (S_a) due to aerosol particles (3–500 nm). Comparison of experimental data and the results derived from the balance equation assuming the ionization rate of $4.8 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$, an extra ion loss of 0.0052 s^{-1} and the recombination coefficient of $1.5 \times 10^{-6} \text{ cm}^3 \text{ s}^{-1}$. Selected data: relative humidity < 95%, wind speed $> 1 \text{ m s}^{-1}$. 31 March–29 April 1999, Hyttiälä station.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Positive air ions in
boreal forest air

U. Hörrak et al.




Fig. 5. Scatterplot between the concentrations of positive aerosol ions and aerosol particles: **(a)** aerosol ions (m ; 2.5–8 nm), or intermediate ions, versus nanometer aerosol particles (2.5–8 nm). The data points corresponding to 5 April are marked with red squares (\square). **(b)** Aerosol ions (N ; 8–ca. 20 nm), or large air ions, versus aerosol particles (8–20 nm). Nucleation event days, 31 March–29 April 1999, Hyytiälä station.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

EGU

Positive air ions in
boreal forest air

U. Hörrak et al.




Fig. 6. Diurnal variation in the concentration of positive air ion classes: small ions (n ; diameters less than 2.5 nm), charged fine nanometer particles (m ; 2.5–8 nm) and coarse nanometer particles (N ; 8–ca. 20 nm), as well as the total concentration of particles (P ; 2.5–8 nm) measured by the DMPS on 2 April (a), 5 April (b), 10 April (c) and 14 April 1999 (d) at Hyytiälä.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

EGU

Positive air ions in
boreal forest air

U. Hörrak et al.




Fig. 7. Diurnal variation in the concentration of positive small ions (n), ion sink (S_a) and the size distribution of ion sink due to aerosol particles (3–500 nm) on 2 April (**a**; left) and 10 April 1999 (**b**; right) at Hyytiälä.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Positive air ions in
boreal forest air

U. Hörrak et al.




Fig. 8. Diurnal variation in the concentration of positive small ions (n), ion sink (S_a) and the size distribution of ion sink due to aerosol particles (3–500 nm) on 5 April (a; left) and 14 April 1999 (b; right) at Hyytiälä.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

Positive air ions in
boreal forest air

U. Hörrak et al.




Fig. 9. Scatterplot between the positive small ion concentration and ion sink on 2 April **(a)**, 5 April **(b)**, 10 April **(c)** and 14 April 1999 **(d)** at Hyytiälä station. Specific data points: brown squares (\square) correspond to the fog event; green triangles (Δ) correspond to the period of weak wind less than 0.2 m s^{-1} . Specific data points have been excluded from the regression analysis.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU