



HAL
open science

Influence of modelled soil biogenic NO emissions on related trace gases and the atmospheric oxidizing efficiency

J. Steinkamp, L. N. Ganzeveld, W. Wilcke, M. G. Lawrence

► **To cite this version:**

J. Steinkamp, L. N. Ganzeveld, W. Wilcke, M. G. Lawrence. Influence of modelled soil biogenic NO emissions on related trace gases and the atmospheric oxidizing efficiency. *Atmospheric Chemistry and Physics Discussions*, 2008, 8 (3), pp.10227-10255. hal-00304213

HAL Id: hal-00304213

<https://hal.science/hal-00304213>

Submitted on 18 Jun 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Modelled NO soil
emissions, related
trace gases and
oxidizing efficiency**

J. Steinkamp et al.

Influence of modelled soil biogenic NO emissions on related trace gases and the atmospheric oxidizing efficiency

J. Steinkamp¹, L. N. Ganzeveld², W. Wilcke³, and M. G. Lawrence¹

¹Department of Atmospheric Chemistry, Max-Planck-Institute for Chemistry, Mainz, Germany

²Department of Environmental Sciences, Chairgroup Earth System Sciences, Wageningen University and Research Centre, Wageningen, Netherlands

³Geographic Institute, Johannes Gutenberg University, Mainz, Germany

Received: 22 April 2008 – Accepted: 2 May 2008 – Published: 30 May 2008

Correspondence to: J. Steinkamp (steinkam@mpch-mainz.mpg.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

The emission of nitric oxide (NO) by soils (SNOx) is an important source of oxides of nitrogen ($\text{NO}_x = \text{NO} + \text{NO}_2$) in the troposphere, with estimates ranging from 4 to 21 Tg of nitrogen per year. Previous studies have examined the influence of SNOx on ozone (O_3) chemistry. We employ the ECHAM5/MESSy earth system model to go further in the reaction chain and investigate the influence of SNOx on lower tropospheric NO_x , O_3 , peroxyacetyl nitrate (PAN), nitric acid (HNO_3), the hydroxyl radical (OH) and the lifetime of methane (τ_{CH_4}). We show that SNOx is responsible for a significant contribution to the NO_x mixing ratio in many regions, especially in the tropics. On the other hand in some regions SNOx has a negative feedback on the lifetime of NO_x through O_3 and OH, which results in regional increases in the mixing ratio of NO_x despite lower total emissions in a simulation without SNOx. Furthermore, the concentration of OH is substantially increased due to SNOx, resulting in an enhanced oxidizing efficiency of the global troposphere, reflected in a $\sim 10\%$ decrease in τ_{CH_4} due to soil NO emissions.

1 Introduction

Nitric oxide (NO) in the soil is produced by the microbial processes of nitrification and denitrification (Firestone and Davidson, 1989). The NO emission originates from a natural pool of nitrogen and a fraction from fertilizer application (Yienger and Levy II, 1995; Stehfest and Bouwman, 2006). The estimates of NO emitted yearly by soils (hereafter called SNOx) ranges from 4 to 21 Tg(N) (Yienger and Levy II, 1995; Davidson and Kingerlee, 1997, and references therein). NO reacts rapidly with other atmospheric compounds, establishing an equilibrium between NO and nitric dioxide (NO_2). These two species are frequently referred to the oxides of nitrogen (NO_x). SNOx is topped by the anthropogenic combustion of fossil fuels ($20\text{--}24 \frac{\text{Tg(N)}}{\text{yr}}$) (Holland et al., 1999) and is comparable to the production of NO_x from lightning and biomass burning, but especially in remote continental regions of the mid- and low-latitudes SNOx is the dom-

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

inant source of NO_x . Through reactions, deposition and stomatal uptake directly within the vegetation layer not all NO emitted by the soil escapes the canopy layer as NO_x (Yienger and Levy II, 1995; Ganzeveld et al., 2002b). In this work SNO_x refers to the flux from the vegetation to the atmosphere. The fraction of NO_x that reaches the atmosphere reacts as a catalyst for production of ozone (O_3), one of the greenhouse gases. This O_3 production is driven by the oxidation of carbon monoxide (CO) and volatile organic compounds (VOC), if the concentration of NO is higher than about 5–30 $\frac{\text{pmol}}{\text{mol}}$ (Brasseur et al., 1999). Atmospheric NO_x is also involved in the production of the hydroxyl radical (OH), which is responsible for the oxidation and depletion of methane (CH_4), another greenhouse gas. Beyond these climate related issues, high NO_x mixing ratios also have a direct impact on human health. NO_x is removed from the atmosphere by reaction with hydroxyl radicals (OH) or oxidation to dinitrogen pentoxide (N_2O_5) and subsequent deposition as nitric acid (HNO_3). It can also react with organic tracers to form peroxy nitrates, mainly peroxyacetyl nitrate (PAN), which, once it is lifted to higher altitudes, can be transported over large distances releasing NO_x when it is transported back downward again.

Previous model studies of the influence of SNO_x on atmospheric chemistry mainly focused either on the NO_x source itself, or on O_3 , mostly on a regional scale. Jaeglé et al. (2005) examined the global partitioning of NO_x sources using inverse modelling and the space-based NO_2 column derived by GOME (Global Ozone Monitoring Experiment). Their a posteriori SNO_x ($8.9 \frac{\text{Tg(N)}}{\text{yr}}$) is 68% greater than their a priori SNO_x ($5.3 \frac{\text{Tg(N)}}{\text{yr}}$). Based on this, Jaeglé et al. (2005) suggest that the influence of SNO_x on background O_3 could be underestimated in current chemistry transport models (CTMs). Bertram et al. (2005) come to a similar conclusion by inverse modelling using another satellite sensor (SCIAMACHY) above the western United States, computing an underestimation of 60%. Delon et al. (2007) modelled higher O_3 concentrations with higher SNO_x above Western Africa. For Europe, Simpson (1995) found that SNO_x hardly has any influence on controlling the O_3 mixing ratio. Isaksen and Hov (1987) already investigated the influence of changes in the emission intensity of different relevant trace gases

on the oxidizing efficiency through an increase in OH concentration with increased NO_x emissions, but they neglected SNOx.

In this study, we take these analyses a step further and follow the reaction chain from SNOx through O_3 and OH to its global influence on the oxidizing efficiency of the atmosphere. To do so, we compare two model runs with a state-of-the-art global 3-D global chemistry climate model. One is a simulation with all relevant emissions and reactions (BASE), and the second simulation is without SNOx (NOBIONO = “No biogenic NO”). We expect a considerable influence of SNOx on the mixing ratios and distribution of related global tropospheric trace gases (NO_x , PAN, HNO_3 , O_3 and OH). Furthermore the global oxidizing efficiency, indicated by the lifetime of CH_4 (τ_{CH_4}), is expected to decrease (τ_{CH_4} increases) if we exclude NO_x emission from soils.

In the following section we briefly describe the model setup. We then compare the relevant tracer mixing ratios from the BASE simulation versus the NOBIONO simulation. In the final section we present our conclusions and outlook.

2 Model description and setup

For this study the Modular Earth Submodel System version 1.2 (MESSy) coupled to the general circulation model ECHAM5 is employed. MESSy connects, through a standardized interface, submodels for different processes with bidirectional feedbacks (Jöckel et al., 2005, 2006). The meteorology for these simulations is driven by sea surface temperature (SST) from the AMIP11b dataset (Taylor et al., 2000). The calculation of SNOx in the BASE simulation is based on the algorithm of Yienger and Levy II (1995), which is the most widely used SNOx algorithm in CTMs (Ganzeveld et al., 2002a; Jaeglé et al., 2005; Delon et al., 2007). This calculation is performed in the submodel EMDEP (Ganzeveld et al., 2006). NO_x produced by lightning is calculated in the submodel LNOX ($2.2 \frac{\text{Tg(N)}}{\text{yr}}$). The remaining sources of NO_x ($49 \frac{\text{Tg(N)}}{\text{yr}}$) are read in from the offline EDGAR database (Olivier et al., 1994) by the submodel OFFLEM (Kerkweg et al., 2006). Other relevant emissions are calculated either by the EMDEP

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

or OFFLEM submodel.

A model spinup time of eleven months (January–November 1994) was chosen and the data of the period December 1994–December 1995 is analysed here. To achieve an identical meteorology of both simulations feedback through trace gases and water vapor is switched off. Table 1 recapitulates the setup of the two simulations.

3 Results and discussion

The emission of NO from soils in the BASE simulation accounts for 15% of the total annual global NO emissions (Table 2). The interannual variability of SNO_x is low in the model (Steinkamp, 2007). The largest emissions are calculated for tropical regions. During JJA there are some exceptions further north in northern America, Europe and north-eastern China. These are fertilizer induced emissions in agricultural regions (Fig. 1 and Table 2).

The data is analysed by season with a focus on the winter and summer season. There is a notable seasonal variation with larger SNO_x in the summer period of each hemisphere and with a larger contribution of SNO_x to the total NO emissions during the northern hemispheric spring and summer (Table 2). The first point can be explained by the temperature dependence and the second one by the greater landmasses in the Northern Hemisphere. In the northern mid-latitudes SNO_x plays a less important role relative to other NO_x emissions, except during the JJA period.

3.1 Influence of SNO_x on related trace gases

The column mean mixing ratios of NO_x, PAN, HNO₃ and O₃ and the column mean concentration of OH in the gridcells (weighted by the air mass in the gridcells) in the lower troposphere (below 500 hPa; hereafter “LT”) from the BASE simulation are compared with the values from the NOBIONO simulation.

As expected, in the LT the difference between the NO_x column mean mixing ratio in the NOBIONO simulation versus the BASE simulation is well-correlated with SNO_x in

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

all regions (Table 3). A low correlation is computed for the Northern Hemisphere during DJF, as could be expected due to the small SNO_x compared to the anthropogenic emissions.

In the low-latitudes and in the northern mid-latitudes the correlation between SNO_x and the difference in the PAN column mean mixing ratio of the two simulations is low (Table 3). This is likely due to the stronger dependence of PAN formation on VOC mixing ratios than on NO_x mixing ratios (Singh et al., 1986). In contrast, there is a much better correlation in the southern mid-latitudes between the difference in the LT PAN column mixing ratio and SNO_x. This suggests a dominating role of NO_x in the formation of PAN in the mid-latitudes of the Southern Hemisphere. This hypothesis is partially confirmed by the calculation of the correlation coefficient of the difference in the PAN column mixing ratio with SNO_x up to a lower altitude (750 hPa). R² increases slightly for the northern mid- and low-latitudes and stays nearly the same for the southern mid-latitudes (Table 3). In the low-latitudes, convective downdrafts and subsiding airmasses, combined with the strong temperature dependence of the decomposition of PAN decreases the correlation.

The correlation between SNO_x and the difference in the LT O₃ column mean mixing ratio is lower than for NO_x. This is partly due to the longer lifetime of O₃, which is better mixed in the LT. Furthermore the production of O₃ is not only determined by the NO_x mixing ratio, but also by the concentration of VOC. The correlation of the OH column mean concentration difference in the LT with SNO_x is similar to O₃. OH is a very short lived tracer, whose production depends on the one hand on the photolysis of O₃ and the water vapor concentration and on the other hand on the reaction of NO with HO₂ in the troposphere. This results, depending on the dominating reaction, in a higher or lower correlation of the OH column concentration difference versus SNO_x than the O₃ column mixing ratio difference versus SNO_x.

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3.1.1 NO_x

The global mean mixing ratio of NO_x in the LT during DJF decreases by 6% in the NOBIONO simulation compared to the BASE simulation. During JJA it decreases by 12%. In both cases the decrease in the mixing ratio is less than the contribution of SNO_x (12% and 17% respectively). The maximum decrease is 78% in DJF and 68% in JJA, while the maximum absolute decreases in the DJF and JJA periods are 139 and 209 $\frac{\text{pmol}}{\text{mol}}$, respectively (figures with absolute differences can be found in the supplement: <http://www.atmos-chem-phys-discuss.net/8/10227/2008/acpd-8-10227-2008-supplement.pdf>).

Interestingly, during DJF the mixing ratio above large parts of the Northern Hemisphere even increases by up to 3.5% (Fig. 2a), with the largest absolute increase of 15.2 $\frac{\text{pmol}}{\text{mol}}$ above Europe. In the JJA period the maximum relative increase of 4.8% is larger than in the DJF period, but the maximum absolute difference is only 3.0 $\frac{\text{pmol}}{\text{mol}}$ (Fig. 2b).

A similar result has been noted for model sensitivity simulations with and without NO_x from lightning (Stockwell et al., 1999; Labrador et al., 2005), in which a decrease in near surface NO_x mixing ratios was computed for similar regions with increasing production of NO_x by lightning. Although NO_x produced by lightning is formed in the free troposphere and SNO_x originates from the surface, we achieve comparable results with SNO_x. To explain why the NO_x mixing ratio decreases less than the relative decrease in the emission of the NOBIONO simulation compared to the BASE simulation, and why it increases during the DJF period in large areas in the Northern Hemisphere, the feedback through O₃ and OH has to be taken into account. Stockwell et al. (1999) assumed that the general increase in O₃ with lightning NO_x causes an increase in OH. This OH reduces the lifetime of NO_x (τ_{NO_x}) through Eq. (1) above regions with high non-lightning NO_x sources. Labrador et al. (2005) showed that the conversion to HNO₃ via N₂O₅ also contributes to the shorter τ_{NO_x} (Eq. 2) with higher NO_x emissions.



Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Similarly we find that without SNO_x, O₃ and OH levels decrease, resulting in enhanced τ_{NO_x} , and due to Eqs. (1) and (2) the NO_x mixing ratio increases in regions with low SNO_x. The changes in HNO₃, O₃ and OH related to this are discussed in the following sections.

In the vertical direction the strongest effects of SNO_x are simulated near the surface (DJF: 53%, JJA: 42%), and a decrease of up to 10 to 25% for higher altitudes in the zonal mean is calculated if SNO_x was switched off (Fig. 3). The effect of convective transport to higher altitudes has a stronger influence on the difference in the total burden between 500 and 250 hPa during DJF (relative: 11.3%, absolute: 1.85 Gg) than during JJA (relative: 6.5%, absolute: 1.22 Gg). This is because the main regions, where the convective transport is most effective, are in the Southern Hemisphere especially the Amazon Basin and the southern tropics of Africa (not shown).

3.1.2 PAN

The LT PAN mixing ratio decreases globally by 4% during DJF and 8% during JJA without SNO_x. In both periods the PAN mixing ratio decreases nearly everywhere above the continents (Fig. 4). Above the tropical oceans, especially during JJA, there is a high relative but a negligible absolute increase in the PAN mixing ratio associated with a decrease in SNO_x. As mentioned above, the formation of PAN in the northern mid- and low latitudes relies more on VOC than on NO_x, but more on NO_x in the southern mid-latitudes. This explains the larger decrease during DJF than during JJA. There is

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

also no increase of PAN in the Northern Hemisphere during DJF despite higher NO_x mixing ratios, which confirms a dominating role of VOC in PAN formation.

Interestingly, in the upper troposphere between 500 hPa and 250 hPa the largest decrease in the PAN mixing ratio is during DJF (7%), whereas it is 5% during JJA. In the zonal mean of the relative difference in PAN mixing ratio with and without SNOx (Fig. 5), the effect of convective transport in the lower latitudes is more effective during DJF than during JJA. At the higher altitudes PAN does not increase anymore, due to its longer lifetime resulting in better mixing.

The differences in the PAN mixing ratio should be interpreted with caution, because the model generally overestimates its levels compared to observations (Jöckel et al., 2006), though this may improve with a new isoprene oxidation scheme (Butler, personal communication).

3.1.3 HNO_3

The global LT mixing ratio of HNO_3 decreases by 16% (DJF) and 15% (JJA) without SNOx. The greatest decrease occurs above continental regions of the low-latitudes and in the summer months in the Northern Hemisphere (Fig. 6). The larger decrease in the mixing ratio of HNO_3 compared to the decrease of NO_x mixing ratio is because the formation of HNO_3 is not only determined by the NO_x mixing ratio, but also relies on the mixing ratios of O_3 and OH, which also decrease, as discussed in the following sections.

Nitric acid is mainly deposited on aerosol particles, taken up by cloud water or directly deposited on the earth's surface. The deposition of HNO_3 is decreased by 14% throughout the year without SNOx. During DJF the decrease is 12% and during JJA it is 15%.

3.1.4 O_3

The mixing ratio of O_3 in the NOBIONO simulation compared to the BASE simulation in the LT decreases by 6% in both seasons, with the greatest decline above the conti-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

nents (Fig. 7). The maximum relative decrease during DJF is 37% and during JJA it is 28%. The maximum absolute decrease ($17.6 \frac{\text{nmol}}{\text{mol}}$) occurs during DJF above Australia (Fig. 7a). In contrast to what was found for NO_x , there is no region with increasing O_3 mixing ratios. The decrease of the NO_x mixing ratio in the LT during JJA (12%) is less effective in reducing the O_3 mixing ratio than the decrease in the NO_x mixing ratio during DJF (6%). This is because the formation of O_3 through SNO_x competes with other strong sources of NO_x during JJA in the Northern Hemisphere, whereas SNO_x is relatively much more important for the formation of O_3 during DJF in the Southern Hemisphere. Furthermore, as was noted above for the PAN formation in the Northern Hemisphere the simulated O_3 production may be more dependent on VOC than NO_x .

In the zonal mean distribution (not shown) a similar pattern of the influence of convection can be seen as already discussed for NO_x and PAN. But due to the longer lifetime of O_3 the relative change has only a maximum of 17% in DJF and 14% in JJA. These changes are smaller than for NO_x and are more evenly distributed above all latitudes and also in the vertical direction. In the vertical direction there is, as with the horizontal, no increase in the O_3 mixing ratio throughout the troposphere.

3.1.5 OH

Excluding the contribution of SNO_x , the mean OH concentration declines by 10% during DJF and 8% during JJA in the LT. The largest relative decrease is 58% during DJF and 42% during JJA above the tropical land regions. During DJF the decrease is shifted to the southern tropics and to the northern tropics during JJA (Fig. 8). Note that during JJA an absolute increase above the Antarctic region is calculated, but the OH concentration here is less than $1 \times 10^4 \frac{\text{molec}}{\text{cm}^3}$.

The decrease is on the one hand induced directly by NO_x through Eq. (3), and on the other hand indirectly by the lower O_3 mixing ratio, leading to less primary OH production, and therefore in a decrease of the OH concentration in the LT.



Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The largest relative decrease in the zonal mean concentration of OH is 18% during DJF and 14% during JJA. This maximum of the relative decrease in the OH concentration without SNO_x is nearly detached from the surface, despite the surface source of SNO_x (Fig. 9). At the surface OH production is mainly related to the reaction of O(¹D) with water, while at higher altitudes it depends more on the reaction of NO with HO₂ (Eq. 3). In the zonal mean the shift to the Southern Hemisphere during DJF is stronger than the shift during JJA to the Northern Hemisphere.

3.2 Influence of SNO_x on the oxidizing efficiency

The oxidation of CO and VOC in the atmosphere is mainly driven by OH. As a measure for the oxidizing efficiency of the atmosphere, τ_{CH_4} is calculated for both simulations according to Lawrence et al. (2001). The trend of monthly mean values is depicted in Fig. 10. The mean τ_{CH_4} averaged for one year (December 1994 to November 1995) for the BASE simulation is 7.26 years and 7.93 years for the NOBIONO simulation, a 9.2% increase without SNO_x. The maximum prolongation of 0.89 years occurs in December 1994 (10.3%) and January 1995 (10.4%).

The changes in τ_{CH_4} are not equally distributed over the globe. In the southern and low-latitudes the relative influence is noticeably greater than in the northern latitudes (Fig. 11). This agrees with the smaller relative change in the OH concentration in the northern latitudes (Fig. 8). In the vertical direction, the relative changes are slightly larger above 500 hPa, even though the origin of SNO_x is at the surface. Beginning from the surface source of SNO_x and following the reaction chain from NO_x over O₃ and OH in each step, the relative difference of our two simulations becomes smaller near the surface and larger at higher altitudes. This trend corroborates the larger relative change of the oxidizing efficiency at higher altitudes. However, only ~15% of the absolute amount of CH₄ in the troposphere is oxidized above 500 hPa (Lawrence et al., 2001).

Labrador et al. (2004) modelled a decrease of 15% in τ_{CH_4} in a simulation with 5 Tg(N) NO_x produced by lightning relative to one with no lightning NO_x. Compared to this, SNO_x is somewhat less effective in altering the oxidizing efficiency of the atmo-

sphere, which is interesting, given that CH_4 oxidation is more effective near the surface where SNO_x is emitted, due to the strong temperature dependence of the reaction of OH with CH_4 . The change in the oxidizing efficiency due to lightning NO_x is larger than due to SNO_x , even though the total emission rate is lower. This is because at higher altitudes the $\text{NO}:\text{NO}_2$ ratio is greater, so that with more NO the NO_x lifetime is not diminished as strongly as near the surface. Furthermore at higher altitudes more NO results in higher OH yields by reaction with HO_2 .

4 Conclusions and outlook

The emission of NO from soils plays an important role for chemical reactions in the atmosphere. Lower global mean NO_x mixing ratios without SNO_x lead to lower global O_3 mixing ratios in the LT. The lower O_3 mixing ratios result in lower OH concentrations. This results in an enhanced lifetime of NO_x in regions with other dominating sources of NO_x . Hence the NO_x mixing ratios increases in some regions, despite lower emissions when SNO_x is neglected in our NOBIONO simulation. From this it follows, that although NO_x is a short-lived tracer it indirectly influences chemical processes in regions with low SNO_x through feedback with O_3 and OH . By following the reaction chain up to PAN and HNO_3 , we detected a dominating role of SNO_x compared to VOC in the mid-latitudes of the Southern Hemisphere. Also by following the reaction chain ($\text{SNO}_x \rightarrow \text{NO}_x \rightarrow \text{O}_3 \rightarrow \text{OH}$), the magnitude of relative effects are shifted step by step to higher altitudes in the troposphere.

Through reaction of NO with HO_2 SNO_x is directly involved in the production of OH . SNO_x also has, through O_3 , an indirect influence on OH production. With OH formed by SNO_x through these pathways τ_{CH_4} is decreased considerably, and the influence of SNO_x on the tropospheric oxidizing efficiency is considerable, in the range of 10%.

The notable modelled influence of SNO_x on directly and indirectly related trace gases shown in this work supports further efforts to improve the parameterization of SNO_x in CTMs, as also proposed by Jaeglé et al. (2005).

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Acknowledgements. We appreciate the help of the the modellers and the MESSy team, especially T. Butler, P. Jöckel, R. Sander, M. Tanarhte and H. Tost. We acknowledge the International Max Planck Research School on Atmospheric Chemistry and Physics for financial support.

References

- 5 Bertram, T. H., Heckel, A., Richter, A., Burrows, J. P., and Cohen, R. C.: Satellite measurements of daily variations in soil NO_x emissions, *Geophys. Res. Lett.*, 32, L24 812, doi:10.1029/2005GL024640, 2005. [10229](#)
- Brasseur, G. P., Orlando, J. J., and Tyndall, G. S. (Eds.): *Atmospheric chemistry and global change*, Oxford University Press, New York, 1999. [10229](#)
- 10 Davidson, E. A. and Kingerlee, W.: A global inventory of nitric oxide emissions from soils, *Nutr. Cycl. Agroecosys.*, 48, 37–50, 1997. [10228](#)
- Delon, C., Reeves, C. E., Stewart, D. J., Serça, D., Dupont, R., Mari, C., Chaboureau, J.-P., and Tulet, P.: Biogenic nitrogen oxide emissions from soils – impact on NO_x and ozone over West Africa during AMMA (African Monsoon Multidisciplinary Experiment): modelling study, *Atmos. Chem. Phys.*, 8, 2351–2363, 2008, <http://www.atmos-chem-phys.net/8/2351/2008/>. [10229](#), [10230](#)
- 15 Firestone, M. K. and Davidson, E. A.: Mikrobiological basis of NO and N₂O production and consumption in soil, In: *Exchange of trace gases between terrestrial ecosystems and the atmosphere*, Edited by: Andreae, M. O. and Schimel, D. S., 7–21, 1989. [10228](#)
- 20 Ganzeveld, L. N., Lelieveld, J., Dentener, F. J., Krol, M. C., Bouwman, A. F., and Roelofs, G.-J.: Global soil-biogenic NO_x emissions and the role of canopy processes, *J. Geophys. Res.*, 107, 4321, doi:10.1029/2001JD001289, 2002a. [10230](#)
- Ganzeveld, L. N., Lelieveld, J., Dentener, F. J., Krol, M. C., and Toelofs, G.-J.: Atmosphere-biosphere trace gas exchanges simulated with a single-column model, *J. Geophys. Res.*, 25 107, 4320, doi:10.1029/2001JD000684, 2002b. [10229](#)
- Ganzeveld, L. N., van Aardenne, J. A., Butler, T. M., Lawrence, M. G., Metzger, S. M., Stier, P., Zimmermann, P., and Lelieveld, J.: Technical Note: Anthropogenic and natural offline emissions and the online Emissions and dry DEPosition submodel EMDEP of the Modular Earth Submodel system (MESSy), *Atmos. Chem. Phys. Discuss.*, 6, 5457–5483, 2006, <http://www.atmos-chem-phys-discuss.net/6/5457/2006/>. [10230](#), [10242](#)
- 30

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Holland, E. A., Dentener, F. J., Brasswell, B. H., and Sulzman, J. M.: Contemporary and pre-industrial global reactive nitrogen budgets, *Biogeochemistry*, 46, 7–43, 1999. [10228](#)
- Isaksen, I. S. A. and Hov, Ø.: Calculation of trends in the tropospheric concentration of O₃, OH, CO, CH₄ and NO_x, *Tellus*, 39B, 271–285, 1987. [10229](#)
- 5 Jaeglé, L., Steinberger, L., Martin, R. V., and Chance, K.: Global partitioning of NO_x sources using satellite observations: relative roles of fossil fuel combustion, biomass burning and soil emissions, *Faraday Discuss.*, 130, 407–423, 2005. [10229](#), [10230](#), [10238](#)
- Jöckel, P., Sander, R., Kerkweg, A., Tost, H., and Lelieveld, J.: Technical Note: The Modular Earth Submodel System (MESSy) – a new approach towards Earth System Modeling, *Atmos. Chem. Phys.*, 5, 433–444, 2005, <http://www.atmos-chem-phys.net/5/433/2005/>. [10230](#)
- 10 Jöckel, P., Tost, H., Pozzer, A., Brühl, C., Buchholz, J., Ganzeveld, L., Hoor, P., Kerkweg, A., Lawrence, M. G., Sander, R., Steil, B., Stiller, G., Tanarhte, M., Taraborrelli, D., van Aardenne, J., and Lelieveld, J.: The atmospheric chemistry general circulation model ECHAM5/MESSy1: consistent simulation of ozone from the surface to the mesosphere, *Atmos. Chem. Phys.*, 6, 5067–5104, 2006, <http://www.atmos-chem-phys.net/6/5067/2006/>. [10230](#), [10235](#), [10242](#)
- 15 Kerkweg, A., Sander, R., Tost, H., and Jöckel, P.: Technical Note: Implementation of prescribed (OFFLEM), calculated (ONLEM) and pseudo-emissions (TNUDGE) of chemical species in the modular earth submodel system (MESSy), *Atmos. Chem. Phys.*, 6, 3603–3609, 2006, <http://www.atmos-chem-phys.net/6/3603/2006/>. [10230](#), [10242](#)
- 20 Labrador, L., von Kuhlmann, R., and Lawrence, M. G.: The effects of lightning-produced NO_x and its vertical distribution on atmospheric chemistry: sensitivity simulations with MATCH-MPIC, *Atmos. Chem. Phys.*, 5, 1815–1834, 2005, <http://www.atmos-chem-phys.net/5/1815/2005/>. [10233](#)
- 25 Labrador, L. J., von Kuhlmann, R., and Lawrence, M. G.: Strong sensitivity of the global mean OH concentration and the tropospheric oxidizing efficiency to the source of NO_x from lightning, *Geophys. Res. Lett.*, 31, L06 102, doi:10.1029/2003GL019229, 2004. [10237](#)
- Lawrence, M. G., Jöckel, P., and von Kuhlmann, R.: What does the global mean OH concentration tell us?, *Atmos. Chem. Phys.*, 1, 37–49, 2001, <http://www.atmos-chem-phys.net/1/37/2001/>. [10237](#), [10254](#), [10255](#)
- 30 Olivier, J. G. J., Bouwman, A. F., Vandermaas, C. W. M., and M, B. J. J.: Emission database for global atmospheric research (EDGAR), *Environ. Monit. Assess.*, 31, 93–106, 1994. [10230](#)
- Sander, R., Kerkweg, A., Jöckel, P., and Lelieveld, J.: Technical note: The new comprehensive

atmospheric chemistry module MECCA, Atmos. Chem. Phys., 5, 445–450, 2005,

<http://www.atmos-chem-phys.net/5/445/2005/>. [10242](#)

Simpson, D.: Biogenic emissions in Europe: 2. Implications for ozone control strategies, J. Geophys. Res., 100, 22 891–22 906, 1995. [10229](#)

5 Singh, H. B., Salas, L. J., and Viezee, W.: Global distribution of peroxyacetyl nitrate, Nature, 321, 588–591, 1986. [10232](#)

Stehfest, E. and Bouwman, L.: N₂O and NO emission from agricultural fields and soils under natural vegetation: summarizing available measurement data and modeling of global annual emissions, Nutr. Cycl. Agroecosys., 24, 207–228, 2006. [10228](#)

10 Steinkamp, J.: Globale Stickstoffmonoxid-Emissionen aus dem Boden: Literatúrauswertung und Evaluierung des Modells ECHAM5/MESSy, Diploma thesis, Johannes-Gutenberg University, Mainz/Germany, 2007. [10231](#)

Stockwell, D. Z., Giannakopoulos, C., Plantevin, P.-H., Carver, G. D., Chipperfield, M. P., Law, K. S., Pyle, J. A., Shallcross, D. E., and Wang, K.-Y.: Modelling NO_x from lightning and its impact on global chemical fields, Atmos. Environ., 33, 4477–4493, 1999. [10233](#)

15 Taylor, K., Williamson, D., and Zwiers, F.: The sea surface temperature and sea ice concentration boundary conditions for AMIP II simulations; PCMDI Report, Tech. Rep. 60, Program for Climate Model Diagnosis and Intercomparison, 2000. [10230](#)

Tost, H.: Global Modelling of Cloud, Convection and Precipitation Influences on Trace Gases and Aerosols, Ph.D. thesis, Rheinische Friedrich-Wilhelms-University, Bonn/Germany, 2006. [10242](#)

Tost, H., Jöckel, P., Kerkweg, A., Sander, R., and Lelieveld, J.: Technical note: A new comprehensive SCAVenging submodel for global atmospheric chemistry modelling, Atmos. Chem. Phys., 6, 565–574, 2006a. [10242](#)

25 Tost, H., Jöckel, P., and Lelieveld, J.: Influence of different convection parameterisations in a GCM, Atmos. Chem. Phys., 5475–5493, 2006b. [10242](#)

Tost, H., Jöckel, P., and Lelieveld, J.: Lightning and convection parameterization – uncertainties in global modelling, Atmos. Chem. Phys., 4553–4568, 2007. [10242](#)

30 Yienger, J. J. and Levy II, H.: Empirical model of global soil-biogenic NO_x emissions, J. Geophys. Res., 100, 11 447–11 464, 1995. [10228](#), [10229](#), [10230](#)

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. Setup of the ECHAM5/MESSy model and used submodels.

Horizontal resolution	T42 ($\sim 2.8^\circ \times 2.8^\circ$)
Vertical resolution	L31 (up to 10 hPa)
Internal timestep	20 min
Timestep of output	5 h
Period of simulation	1994–1995

Used submodels	Calculation of	Literature ref.
CLOUD	Clouds and precipitation	Jöckel et al. (2006)
CONVECT	Convection	Tost et al. (2006b)
CVTRANS	Convective tracer transport	Tost (2006)
EMDEP ^a	Emission and deposition	Ganzeveld et al. (2006)
JVAL	Rates of photolysis	Jöckel et al. (2006)
LNOX	Lightning NO _x	Tost et al. (2007)
MECCA	Chemical atmospheric reactions ^b	Sander et al. (2005)
OFFLEM	Offline emissions	Kerkweg et al. (2006)
RAD4ALL	Radiation	Jöckel et al. (2006)
SCAV	Wet deposition	Tost et al. (2006a)
TNUDGE	Tracer nudging	Kerkweg et al. (2006)
TROPOP	Calculation of the tropopause	Jöckel et al. (2006)

^a Extended version; soil NO emissions switched off in NOBIONO simulation

^b Tropospheric reaction with NMHC and without halogens

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.

Table 2. Simulated total NO_x emissions, SNO_x in Tg(N) of the BASE simulation and in brackets relative contribution of SNO_x to the total NO emissions for different regions and periods.

Season ^a	Global		Low-latitudes (30° N–30° S)		Mid-latitudes (30° N–60° N) (30° S–60° S)			
	total	soil	total	soil	total	soil	total	soil
DJF	12.53	1.53 (12%)	7.09	1.34 (19%)	4.93	0.06 (1%)	0.46	0.12 (26%)
MAM	11.93	2.03 (17%)	6.07	1.60 (26%)	5.40	0.35 (6%)	0.40	0.08 (20%)
JJA	13.67	2.34 (17%)	7.04	1.43 (20%)	6.15	0.88 (14%)	0.33	0.04 (12%)
SON	13.03	1.81 (14%)	7.33	1.45 (20%)	5.27	0.29 (6%)	0.38	0.07 (18%)
All	51.17	7.72 (15%)	27.52	5.82 (21%)	21.76	1.58(7%)	1.57	0.31 (20%)

^a DJF = December 1994, January, February 1995; MAM = March, April, May 1995; JJA = June, July, August 1995; SON = September, October, November 1995

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 3. Correlation coefficient (R^2) between surface SNO_x flux values and the difference (NOBIONO–BASE) of the tracer burden in the overlying lower troposphere (> 500 hPa for all gases and > 750 hPa for PAN values in brackets) by gridcell, both averaged over the corresponding period; only gridcells with a land surface fraction of at least 75% were included.

Season ^a	NO _x	PAN	HNO ₃	O ₃	OH	NO _x	PAN	HNO ₃	O ₃	OH
Global (N=2462)						Low-latitudes 30° N–30° S (N=646)				
DJF	0.86	0.53 (0.60)	0.43	0.62	0.58	0.68	0.17 (0.25)	0.18	0.29	0.22
MAM	0.90	0.47 (0.53)	0.60	0.58	0.64	0.79	0.14 (0.24)	0.42	0.29	0.33
JJA	0.82	0.27 (0.35)	0.37	0.37	0.43	0.69	0.07 (0.07)	0.15	0.21	0.22
SON	0.86	0.44 (0.52)	0.43	0.55	0.58	0.66	0.14 (0.24)	0.07	0.14	0.14
Year	0.88	0.46 (0.54)	0.51	0.56	0.60	0.72	0.11 (0.23)	0.16	0.19	0.21
Northern 30° N–60° N (N=637)			Mid-latitudes			Southern 30° S–60° S (N=46)				
DJF	0.55	0.02 (0.09)	0.40	0.07	0.31	0.80	0.45 (0.45)	0.62	0.67	0.39
MAM	0.82	0.19 (0.30)	0.43	0.20	0.27	0.76	0.71 (0.60)	0.45	0.70	0.59
JJA	0.75	0.10 (0.22)	0.26	0.13	0.18	0.83	0.46 (0.51)	0.54	0.46	0.69
AON	0.80	0.09 (0.19)	0.40	0.19	0.32	0.87	0.74 (0.77)	0.65	0.77	0.47
Year	0.77	0.12 (0.23)	0.33	0.16	0.16	0.79	0.66 (0.65)	0.64	0.78	0.56

^a See Table 2 for abbreviations

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.




Fig. 1. Simulated SNOx flux for **(a)** December 1994 to February 1995 and **(b)** June to August 1995 in $\frac{\text{ng}}{\text{m}^2 \text{s}}$.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.




Fig. 2. Relative difference ($\frac{\text{NO}_{\text{BIONO}} - \text{BASE}}{\text{BASE}} \times 100\%$) of the lower tropospheric mixing ratio of NO_x in % – values below 30 $\frac{\text{pmol}}{\text{mol}}$ in the BASE simulation excluded from calculation – averaged for **(a)** December, January, February and **(b)** June, July and August.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.




Fig. 3. Zonal mean relative difference ($\frac{\text{NOBIONO}-\text{BASE}}{\text{BASE}} \times 100\%$) of NO_x mixing ratio in % averaged for **(a)** December, January, February and **(b)** June, July and August. Note that the y-axis is linearly scaled, since the focus of this work lies in the lower troposphere.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 4. Relative difference ($\frac{\text{NO}_{\text{BIONO}} - \text{BASE}}{\text{BASE}} \times 100\%$) of the lower tropospheric mixing ratio of PAN in % – values below $50 \frac{\text{pmol}}{\text{mol}}$ in the BASE run excluded from calculation – averaged for (a) December, January, February and (b) June, July and August.




Fig. 5. Zonal mean relative difference ($\frac{\text{NOBIONO}-\text{BASE}}{\text{BASE}} \times 100\%$) of PAN mixing ratio in % averaged for **(a)** December, January, February and **(b)** June, July and August.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 6. Relative difference ($\frac{\text{NOBIONO}-\text{BASE}}{\text{BASE}} \times 100\%$) of the lower tropospheric mixing ratio of HNO_3 in % – values below $30 \frac{\text{pmol}}{\text{mol}}$ in the BASE simulation excluded from calculation – averaged for **(a)** December, January, February and **(b)** June, July and August.

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.




Fig. 7. Relative difference ($\frac{\text{NOBIONO}-\text{BASE}}{\text{BASE}} \times 100\%$) of the lower tropospheric mixing ratio of O₃ in % – values below $25 \frac{\text{nmol}}{\text{mol}}$ in the BASE simulation excluded from calculation – averaged for (a) December, January, February and (b) June, July and August.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.




Fig. 8. Relative difference ($\frac{\text{NOBIONO}-\text{BASE}}{\text{BASE}} \times 100\%$) of the lower tropospheric concentration of OH in % – values below $10^4 \frac{\text{molec}}{\text{cm}^3}$ in the BASE simulation excluded from calculation – averaged for **(a)** December, January, February and **(b)** June, July and August.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.




Fig. 9. Zonal mean relative difference ($\frac{\text{NOBIONO-BASE}}{\text{BASE}} \times 100\%$) of OH mixing ratio in % averaged for **(a)** December, January, February and **(b)** June, July and August.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.




Fig. 10. Seasonal cycle of monthly mean lifetime of CH₄ from December 1994 to December 1995 in years (calculated according to [Lawrence et al., 2001](#)).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelled NO soil emissions, related trace gases and oxidizing efficiency

J. Steinkamp et al.




Fig. 11. Relative increase of τ_{CH_4} ($\frac{\tau_{\text{CH}_4, \text{NOBIONO}} - \tau_{\text{CH}_4, \text{BASE}}}{\tau_{\text{CH}_4, \text{BASE}}} \times 100\%$) in various zonal subdomains of the atmosphere (calculated after Lawrence et al., 2001).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion