



HAL
open science

Estimation of erosion and sediment export from an agricultural catchment

O. Evrard, Guillaume Nord, Olivier Cerdan, Véronique Souchère, Yves Le Bissonnais, Philippe Bonté

► **To cite this version:**

O. Evrard, Guillaume Nord, Olivier Cerdan, Véronique Souchère, Yves Le Bissonnais, et al.. Estimation of erosion and sediment export from an agricultural catchment. EGU General Assembly 2010, May 2010, Vienne, Austria. pp.4697. hal-00533834

HAL Id: hal-00533834

<https://brgm.hal.science/hal-00533834>

Submitted on 8 Nov 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Estimation of erosion and sediment export from an agricultural catchment (1960 – 2000) confronting the outputs of an expert-based model and Cs-137 inventories

Olivier Evrard (1), Guillaume Nord (2), Olivier Cerdan (3), Véronique Souchère (4), Yves Le Bissonnais (5), and Philippe Bonté (1)

(1) Laboratoire des Sciences du Climat et de l'Environnement, Gif-sur-Yvette, France, (2) Institute of Environmental Assessment and Water Research (IDÆA), CSIC, Barcelona, Spain, (3) BRGM – Aménagement et risques naturels, Orléans, France, (4) INRA, UMR 1048, SAD-APT, Thiverval Grignon, France, (5) INRA, Laboratoire d'étude des Interactions Sol-Agrosystème-Hydrosystème, UMR LISAH, Montpellier, France

Soil erosion leads to important environmental problems (e.g. muddy floods, reservoir sedimentation) in cultivated areas of the European loess belt. This study aimed to quantify erosion and to determine the impact of rainfall seasonality and land use change on soil erosion over the last 40 years in a 94-ha cultivated catchment of Normandy (France). To this end, scenarios representative of the different land use conditions were simulated using the STREAM expert-based erosion model. A 13-yr long sequence of rainfall events was run with this model. Results showed that erosion increased dramatically after land consolidation (+168% on average). Interannual variability of erosion is important. After land consolidation, 79% of erosion was observed in summer and autumn, even though these seasons only accounted for 58% of annual rainfall kinetic energy. The bulk of erosion was hence produced by a few intense thunderstorms during this period. Thunderstorms correspond to 5% of rainfall events, but they generate 51% of total annual erosion after land consolidation (and up to 57% of erosion before land consolidation). Confrontation of the model outputs with the erosion rates derived from Cs-137 measurements suggested that soil redistribution within the catchment was very high but that sediment exports from the catchment remained limited (sediment delivery ratio between 1 – 10%). Erosion rates derived from Cs-137 measurements showed an important and organised spatial variability, but erosion rates integrated over larger areas remained in the same order of magnitude than those simulated by the model or were slightly higher. Water erosion would hence not be the only process generating erosion within this catchment. In this context, tillage erosion cannot be neglected to calculate the sediment budget over several decades. These findings show the necessity to simulate sequences of rainfall events to obtain reliable erosion predictions. They also demonstrate the interest of developing expert-based models simulating both water and tillage erosion at the catchment scale to conduct erosion studies over several decades.