

**Concentration and mobility of lithogenic trace metals in
soils : significance of anthropogenic lateral
redistributions.**

Ary Bruand

► **To cite this version:**

Ary Bruand. Concentration and mobility of lithogenic trace metals in soils : significance of anthropogenic lateral redistributions.. Comptes Rendus Géoscience, Elsevier Masson, 2002, 334, pp.581-582. 10.1016/S1631-0713(02)01789-3 . hal-00072495

HAL Id: hal-00072495

<https://hal-insu.archives-ouvertes.fr/hal-00072495>

Submitted on 6 Jun 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Concentration and mobility of lithogenic trace metals in soils: significance of anthropogenic lateral redistributions

Ary Bruand

Institut des sciences de la Terre d'Orléans (ISTO), université d'Orléans, Géosciences, BP 6759, 45067 Orléans cedex 2, France

1. Introduction

One of the main current concerns regarding soil quality is the presence of trace metals that constitutes a risk of toxicity for plants and animals [1, 5, 7, 11]. Trace metals in soils can have either a lithogenic origin when they inherit from the lithosphere or an anthropogenic origin when they result from man's activities (mining, metallurgy of non-ferrous metals, combustion of fossil energy sources, solid waste or sewage sludge) [5]. Recent studies showed that concentration of lithogenic trace metals in soils could not be explained without taking man's activities into account.

2. Redistribution of lithogenic trace metals in cultivated lands

Concentration of lithogenic trace metals in soils is usually explained by the weathering of primary minerals that are inherited from the bedrock. Weathering leads to accumulation or impoverishment of mineral phases with their consequences on trace metal concentration in saprolite and soil [2, 5]. Lateral redistribution of soil materials along slopes by erosion is also a major process explaining trace metal concentration in soils in cultivated lands [9]. Detailed spatial analysis of Cr concentration in soils developed in weathered metamorphic rocks showed that it varied according to material redistribution due to the effect of tillage: homogenisation of Cr concentration in the topsoil and accumulation of Cr-rich materials in areas located above former edges because of tillage erosion [10]. Thus, in cultivated lands, presence of lithogenic trace metals in soils results from both vertical (rock

weathering and pedogenesis) and lateral (tillage homogenisation and erosion) redistribution processes.

3. Consequences for mobility of anthropogenic trace elements in soils

Mobility of trace metals in soils depends on the chemical and physical properties of the soil and especially on their speciation [5, 7, 11]. Because of lateral redistribution processes, soil materials can be moved downslope and exposed to new conditions leading an increase in the mobility of trace metals [3, 11]. Quantin et al. [8] showed that under reducing conditions similar to those existing in waterlogged soils, chemical and bacterial processes can reduce Mn oxide and release Co and Ni that were located within the Mn-oxide lattice with an increase in their bioavailability as a consequence. The possible toxicity of Cr in waterlogged soils was also shown by Cooper [4] because of the presence of Cr(VI), which is soluble and toxic. A seasonal release of trace metals was recorded in wetlands by Olivie-Lauquet et al. [6]. Their results showed that redox processes in the presence of organic ligands cause the transfer of trace metals from the solid phase to the soil solution.

4. Conclusion

These new results show clearly that in cultivated lands and in land that have been cultivated for years in the past, the concentration of lithogenic trace metals in soils cannot be only explained by natural processes. They are affected by redistribution processes along hillslopes that are related to agricultural usage of soils. As a consequence, soil materials can be moved

towards areas where the mobility and toxicity of trace metals are much greater. Thus, a trace metal can be endogenous, but with a concentration and mobility strongly man-affected. Finally, such results

point out the necessity to study soil distribution as a result of both pedogenesis and modifications by man, and to analyse the consequences of soil material redistribution on metal trace mobility.

References

- [1] D. Baize, Teneurs totales en éléments traces métalliques dans les sols (France), Inra Éditions, Paris, 1997, 408 p.
- [2] D. Baize, T. Sterckeman, Of the necessity of knowledge of the natural pedo-geochemical background content in the evaluation of the contamination of soils by trace elements, *Sci. Total Environ.* 264 (2001) 127–139.
- [3] T. Becquer, E. Bourdon, J. Pétard, Disponibilité du nickel le long d'une toposéquence de sols développés sur roches ultramafiques de Nouvelle-Calédonie, *C. R. Acad. Sci. Paris, Ser. IIA* 321 (1995).
- [4] G.R.C. Cooper, Oxidation and toxicity of chromium in ultramafic soils in Zimbabwe, *Appl. Geochem.* 17 (2002) 585–592.
- [5] A. Kabata-Pendias, H. Pendias, Trace elements in soils and plants, CRC Press, Boca Raton, FL, 2001, 331 p.
- [6] G. Olivie-Lauquet, G. Gruau, A. Dia, C. Riou, A. Jaffrézic, O. Hénin, Release of trace elements in wetlands: role of seasonal variability, *Water Resour. Res.* 35 (2001) 943–952.
- [7] E. Podlesáková, J. Nemecek, R. Vácha, Mobility and bioavailability of trace elements in soils, in: I.K. Iskandar, M.B. Kirkham (Eds.), *Trace Elements in Soils. Bioavailability, Flux, and Transfer*, CRC Press, Boca Raton, FL, 2001, pp. 21–41.
- [8] C. Quantin, T. Becquer, J. Berthelin, Mn-oxide: a major source of easily mobilisable Co and Ni under reducing conditions in New Caledonia Ferralsols, *C. R. Geoscience* 334 (2002) 273–278.
- [9] S. Salvador-Blanes, S. Cornu, M. Hardy, I. Gay-Ovejero, V. Deschatrettes, D. Baize, D. King, Influence des substrats et des formations de versants sur la variabilité des teneurs naturelles en chrome de sols issus de roches métamorphiques, *C. R. Acad. Sci. Paris, Ser. IIA* 332 (2001) 681–687.
- [10] S. Salvador-Blanes, S. Cornu, D. King, Distribution d'un élément trace (Cr) dans un sol développé sur roches métamorphiques : variabilité à l'échelle d'un versant, *C. R. Geoscience* 334 (2002) 51–58.
- [11] F. Trolard, G. Bourrié, A. Jaffrézic, Distribution spatiale et mobilité ETM, in: D. Baize, M. Tercé (Eds.), *Les éléments traces métalliques dans les sols. Approches fonctionnelles et spatiales*, Inra Éditions, Paris, 2002, in press.