

Response of littoral chironomid community and organic matter to late glacial lake level and environmental changes at Lago dell'Accesa (Tuscany, Italy).

Laurent Millet, Boris Vannière, Michel Magny, Jacques-Louis de Beaulieu, Jean-Robert Disnar, Fatima Laggoun-Défarge, V. Verneaux, Anne-Véronique Walter-Simonnet, E. Ortu, Gilles Bossuet

► **To cite this version:**

Laurent Millet, Boris Vannière, Michel Magny, Jacques-Louis de Beaulieu, Jean-Robert Disnar, et al.. Response of littoral chironomid community and organic matter to late glacial lake level and environmental changes at Lago dell'Accesa (Tuscany, Italy).. *Journal of Paleolimnology*, Springer Verlag, 2007, 38, pp.525-539. 10.1007/s10933-007-9088-z . hal-00085520

HAL Id: hal-00085520

<https://hal-insu.archives-ouvertes.fr/hal-00085520>

Submitted on 11 May 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Response of littoral chironomid communities and organic matter to late glacial lake—level, vegetation and climate changes at Lago dell'Accesa (Tuscany, Italy)

Laurent Millet^{1, 2}, Boris Vanni re¹, Val rie Verneaux³, Michel Magny¹, Jean Robert Disnar⁴, Fatima Laggoun-D farge⁴, Anne V ronique Walter-Simonnet⁵, Gilles Bossuet¹, Elena Ortu¹ and Jacques-Louis de Beaulieu⁶

(1) Laboratoire de Chrono-Ecologie, UMR 6565 CNRS, UFR Sciences et Techniques, Universit  de Franche-Comt , 25030 Besan on cedex, France

(2) Laboratoire EDYTEM, UMR 5204, CISM, Universit  de Savoie, 73376 Le Bourget du Lac cedex, France

(3) Laboratoire de Biologie Environnementale, EA 3184 USC INRA, UFR Sciences et Techniques, Universit  de Franche-Comt , 25030 Besan on cedex, France

(4) Institut des Sciences de la Terre d'Orl ans, UMR 6113 CNRS, Universit  d'Orl ans, G osciences, BP 6759, 45067 Orl ans Cedex 2, France

(5) Laboratoire de G osciences, UFR Sciences et Techniques, Universit  de Franche-Comt , 25030 Besan on cedex, France

(6) Laboratoire IMEP, UMR 6116, Europ le M diterran en de l'Arbois, Bat. Villemin, BP 80, 13545 Aix-en-Provence cedex 4, France

Abstract

This study focuses on the response of lacustrine littoral chironomid communities to late glacial changes in limnological, environmental and climate conditions in the Mediterranean context. Late glacial chironomid (Diptera: Chironomidae) assemblages, organic petrography and geochemistry were analysed in a sediment core from the littoral zone of Lago dell'Accesa (Tuscany, Italy), where the lake-level fluctuations and the vegetation history have been previously reconstructed. Comparison of the chironomid stratigraphy to other proxies (pollen assemblages, organic petrography and geochemistry, lake-level) and regional climate reconstruction suggested the predominant influence of lake-level changes on the littoral chironomid fauna. The main lowering events that occurred during the Oldest and the Younger Dryas were followed by higher proportions of taxa typical of littoral habitats. A complementary study of organic matter suggested the indirect impact of lake-level on the chironomids through changes in humic status and habitat characteristics, such as the type of substrate and aquatic macrophyte development. Several chironomid taxa, such as *Glyptotendipes*, *Microtendipes* and *Cricotopus* type *patens*, were identified as possible indicators of low lake-level in the late glacial records. Nevertheless, this study suggested that parallel analyses of organic matter and chironomid assemblages may be needed to circumvent misinterpretation of littoral chironomid assemblage stratigraphy. There was a weak response of the chironomid assemblages to small lake-level lowerings that corresponded to the Older Dryas and Preboreal oscillations. A higher level of determination, e.g. to the species group level, may be necessary to increase the sensibility of the indicators to lake-level changes.

Keywords Chironomids - Organic matter - Late glacial - Paleolimnology - Lake-level - Italy

Introduction

Subfossil chironomids are one of the most promising proxies of paleoenvironmental and paleoclimatic changes (Walker 2001). The long persistence of larval remains (head capsules) in the sediment allows reconstructing past assemblages after extraction and determination. Changes in the composition of the chironomid fauna are interpreted in reference to ecological preferences of the taxa involved. A great variety of environmental factors are known to potentially affect chironomid communities such as oxygen conditions (Verneaux and Aleya 1998), trophic level (Saether 1979; Woodward and Schulmeister 2006), humic status (Brinkhurst 1974), type of substrate and habitat (Maitland 1979; Gandouin et al. 2006) and temperature (Walker et al. 1991; Barley et al. 2006). In mid-latitude and northern Europe, the late glacial period was marked by numerous and strong temperature changes (Magny et al. 2006b). Here, chironomid studies of late glacial records have focused on quantitative reconstruction of temperature considered as the main driving factor of chironomid successions (e.g. Heiri and Millet 2005). These reconstructions provided information on late glacial climate variability and its forcing mechanisms (Magny et al. 2006b).

In the Mediterranean zone and in particular in Italy, while many late glacial lacustrine records are available, chironomid remains have rarely been used to reconstruct late glacial climate and environmental variability (e.g. Manca et al. 1996). Recent high-resolution studies of marine and lacustrine sequences suggest close similarities in the timing of late glacial climate variability between the Mediterranean and the North Atlantic regions (Huntley et al. 1999, Asioli et al. 2001). Nevertheless, the magnitude of summer temperature changes during the major transitions of the last deglaciation was probably lower in southern Europe than in northern Europe (Renssen and Isarin 2001 and references therein). At Lago dell'Accesa (Tuscany, Italy), multiproxy reconstruction involving vegetation and lake-level changes suggested that hydrological changes were probably strong, especially for low-elevation Mediterranean sites (Magny et al. 2006a). Velle et al. (2005) suggested that in times of low amplitude temperature changes, other environmental factors may control the chironomid response. In Mediterranean lake records, the predominant role of temperature changes on late glacial littoral chironomid fauna should then be carefully studied.

In this study, besides the chironomid assemblages, petrographic and geochemical analyses of the sediment organic matter (OM) were performed on a late glacial sediment core from Lago dell'Accesa (Tuscany, Italy), where lake-level variations and vegetation history already have been reconstructed (Magny et al. 2006a; Drescher-Schneider et al. 2007). The aim of this work was to assess the response of the littoral chironomid communities to late glacial limnological, environmental and climate changes in a Mediterranean lake. We paid special attention to the lake-level impact on the littoral chironomid fauna through changes in humic status and habitat characteristics. The final goal was to provide some backgrounds for the interpretation of future late glacial chironomid works in Mediterranean lakes, and more specifically to define some valuable indicators of lake-level changes from chironomid and OM records.

Site description

Lago dell'Accesa is located in the Grosseto province (Tuscany, Italy, Fig. 1). The small catchment (5 km²) lies at the southern margin of the Colline Metallifere (alt. max.: 1060 m

a.s.l.). The catchment is delimited by several low elevation hills (max alt. 352 m) and lies between three main geological formations: Permian schist to the north-east, highly karstified Rhaetian dolomite limestone to the north and west, and Eocene schist to the south and west (Merciai 1933). The lake is of karstic origin and is fed by run-off from the catchment and by a small karstic inlet (Inferno spring, Fig. 1). The eastern outlet forms the source of the Bruna River. The maximum lake depth is 42 m and the mean depth is 21.3 m (Fig. 1). In 1912, the outlet floor was deepened by about 50 cm in order to extend the cultivated area around the lake (Merciai 1933). The subsequent lake lowering led to the emergence of a large littoral platform (ca. 150 m wide).

Material and methods

Coring

The coring point is located on land 50 m from the present-day western lakeshore within the former littoral platform (Fig. 1). This site was chosen following several observations: (i) preliminary pollen analysis indicated that the sedimentary accumulation corresponding to the late glacial period is there around 5 m thick and can then provide a good temporal resolution for paleoenvironmental studies, (ii) the zone is fully accessible to a corer fixed on a caterpillar APAGEO[®]. Two overlapping cores series, 17.50 m long each, were retrieved using a Russian corer. Magnetic susceptibility was used to correlate stratigraphically the two sediment columns and build a synthetic record.

Chironomid analysis

Sampling intervals were 10 cm in the main part of the core and 5 cm in the 13.30–11.90 m section. In total, 84 samples, 2 cm thick, were used for subfossil chironomid analysis. Chironomid head capsules were extracted from sediment following the procedure described by Walker (2001). Identification of chironomid head capsules was performed according to Wiederholm (1983) and Klink and Moller-Pillot (2003), in most cases, to the genus or species group level. For the taxon names, the designation ‘gr.’ indicates that the taxon could be any species within the species group and the use of the designation ‘type’ indicates that the taxon is morphologically similar to a specific species. *Cricotopus* type *patens* was identified according to the description of the species given by Hirvenoja (1973). *Parakiefferiella* type *bathophila* was recognised from Wiederholm (1983). Specimens referred to as *Parakiefferiella* sp. A were described in Brooks et al. (1997). *Chironomus* type *plumosus* specimens were differentiated from other *Chironomus* from the combination of a long fourth lateral tooth of the mentum and three inner teeth on the mandibule (Hofmann 1971).

Relative abundances of the taxa were calculated for each sample. Only chironomid taxa occurring in at least two samples, with a relative abundance of more than 2% in at least one sample, were included for further analysis. The Hill’N2 (Hill 1973) diversity index was calculated for each sample. A stratigraphic diagram of taxon relative abundances was produced using the TILIA and TILIA-Graph software package (Grimm 1991). The chironomid assemblage zones were defined from a stratigraphically constrained cluster analysis performed by Coniss (Grimm 1991). The number of statistical zones was assessed by comparison with the broken stick model (Bennett 1996).

To help the diagram interpretation, we cumulated the relative abundances of taxa usually considered in the literature as shallow-water living larvae: *Dicrotendipes*, *Glyptotendipes*,

Labrundinia, *Limnophyes*, *Phaenopsectra*, *Polypedilum* type *sordens*, *Stenochironomus* and *Xenochironomus* (Littoral biotope taxa in Fig. 3). The only Palaeartic species of *Labrundinia* is found in bogs (Fittkau and Roback 1983). *Labrundinia* is one of the dominant chironomid taxa in the submerged vegetation of the littoral zone of Lago di Vico (Mastrantuono 1991). *Polypedilum sordens*, *Glyptotendipes* and *Dicrotendipes* are widely recognised as being associated with aquatic macrophytes (Thut 1968; Pinder and Reiss 1983; Klink and Moller-Pillot 2003). *Phaenopsectra* lives in very shallow zones in permanent lakes and ponds (Palomäki 1989). *Limnophyes* is a terrestrial genus scattered in aquatic biotopes, *Stenochironomus* mine in submerged wood remains and *Xenochironomus* are obligate miners in freshwater sponges (Pinder and Reiss 1983; Klink and Moller-Pillot 2003).

Organic geochemistry and petrography

Total Organic Carbon (TOC) contents and Hydrogen Index (HI) values were determined by Rock-Eval[®] pyrolysis (Espitalié et al. 1985a, b) with a model 6 device (Vinci Technologies). The analyses were carried out on 50–100 mg of crushed samples under standard conditions. TOC (%) accounts for the quantity of organic carbon present in the sediment. HI (mg HC g⁻¹ TOC) is the amount of hydrocarbonaceous products (HC) released during pyrolysis (S2 signal intensity, in mg HC g⁻¹ dry sediment) normalized to TOC

$$HI = [S2 \cdot TOC^{-1}] \times 100$$

TOC and HI indicate OM abundance and hydrogen richness, respectively. TOC contents depend not only on autochthonous OM productivity and allochthonous OM inputs, but also on oxygen conditions, which determine OM degradation in the water column and sediments. In stratigraphic records, TOC measurements can be affected by the sedimentation rate. For example, a high influx of minerogenic particles entails a dilution of OM contents.

The HI index is controlled by the type (i.e. the source) and the degree of degradation of the OM. It may be used to assess main variations in lacustrine-deposit OM quality (Talbot and Livingstone 1989; Meyers and Lallier-Vergès 1999; Jacob et al. 2004). An HI decrease is mainly due to a larger contribution of terrestrial and littoral (e.g. aquatic macrophytes) organic material in the total sediment OM (Disnar et al. 2003). However, OM oxidation also entails a decrease of the HI (Espitalié et al. 1985a, b; Jacob et al. 2004).

The petrographic study (palynofacies) of the Accessa record involved the microscopic examination under transmitted light of 26 samples after acid hydrolysis of organic remains. Organic residues were mounted as palynofacies slides and examined at 20 and 50× magnification. Chromatic and textural characteristics of organic remains were used to identify and quantify OM compounds. Organic remains were classified into two groups: material of phytoplanktonic origin and material from vascular plants. The first group is the authigenic fraction. It mainly consists of greyish amorphous OM and translucent membrane fragments (Lallier-Vergès et al. 1993). The second group is composed of well-preserved ligno-cellulosic tissues, reddish amorphous OM, yellow-orange gelified particles and dark organic components. They belong to higher plants and likely originate from emergent plants or local vegetation growing around the lake. This material may represent the post-burial state of autochthonous higher plants when the system is functioning as a shallow lake or a peat swamp (Sifeddine et al. 1996). It may also derive from the surrounding soil (Noël et al. 2001). The surfaces of each OM particle type were counted with a grid reticle in the eye-piece of the

microscope. About 3000 items per sample were counted to calculate relative frequencies with an estimated error of about 10%.

Age–depth model

The age–depth model of the sequence (Fig. 2) was developed from a linear interpolation of five tephra layers and two radiocarbon dates (Table 1). More precise information about the age–depth model, including analysis and dating of tephra layers, can be found in Magny et al. (2006a). Age of tephra layers was determined with reference to the work of Wulf et al. (2004) in the laminated sediment of Lago di Monticchio. The pollen stratigraphy allowed the identification of the main late glacial climatic events: Oldest Dryas, late glacial Interstadial and Younger Dryas (Drescher-Schneider et al. 2007). The age of these events was obtained using our age–depth model (Fig. 2) and appeared to be in agreement with the age found in other Mediterranean terrestrial and marine records (Magny et al. 2006a).

Results and interpretations

General comments

Twenty-seven chironomid taxa were identified from the entire core (Fig. 3). Despite the processing of large quantities of sediment (up to 50 g), less than 10 head capsules (h.c.) were recovered in 5 samples (16.20 m, 15.95 m, 14.51 m, 14.30 m, 14.09 m). These samples were removed from the data set for further analysis. Seven chironomid assemblage biozones (CAZ-Ac1 to 7) were identified throughout the diagram (Fig. 3).

TOC values fluctuate between 0.13 and 2.39% throughout the record (Fig. 4). However, most values are gathered in the 0.42–0.85% range (Fig. 4) that is relatively low in comparison to other late glacial lacustrine records (e.g. Ramrath et al. 1999). This difference can tentatively be explained by the dilution of the OM by carbonate minerals that were produced in rather great abundance on the littoral platform, where were deposited the studied sediments. Minimum and maximum HI values reach 280 and 834 mg HC g⁻¹ TOC, respectively. However, a S2 versus TOC diagram (Fig. 5) reveals that all the data points are gathered between two lines passing through the origin of the axes, the slope of these two lines corresponding to HI values of 300 and 750 mg HC g⁻¹, respectively. The fact that these two lines do not show any tendency to intersect the abscissa axis rules out any mineral matrix effect that would have lowered HI values (Espitalié et al. 1985b; Langford and Blanc-Valleron 1990). In addition, the total absence of points between the lower line and the abscissa axis reveals that no sample contains notably weathered OM that would be characterised by a rather low HI. The HI values 300 and 750 mg HC g⁻¹ delineated by the two lines defined here above, are typical for well-preserved ligno-cellulosic debris and phytoplankton, respectively. At least in a first approach, in the absence of any notable mineral matrix effect and OM alteration, the gathering of all the data points between these two lines delineating HI values typical for the two major allochthonous/autochthonous OM end-members allow us to consider the HI of any sample as resulting from the mixing of variable proportions of these two materials (and thus is indicative of these proportions). However, in such an approach it must also be born in mind that in lacustrine environments and especially in sediments deposited near a palaeoshore, remains of aquatic macrophytes can add to the debris of woody land plants constituting the true allochthonous OM contribution (Meyers and Lallier-Vergès 1999; Jacob et al. 2004). The results suggest an overall dominance of

lacustrine remains in total sediment OM. This assumption is supported by the palynofacies analysis. The proportion of terrestrial or paludal OM (TOM, Fig. 4) represents between 4.9 and 47.4% throughout the whole record.

Changes in chironomid and OM can be compared with the vegetation history reconstructed from pollen analysis (Drescher-Schneider et al. 2007) and lake-level variations inferred from a specific sedimentological method (Magny et al. 2006a).

Oldest Dryas

The Oldest Dryas pollen zone is represented by the CAZ-Ac1 within the chironomid record (Fig. 3). The chironomid community is dominated by *Parakiefferiella* type *bathophila*, *Paratanytarsus*, *Glyptotendipes* and *Microtendipes* (Fig. 3). The low lake-level reconstructed during this period (Magny et al. 2006a) led to the development of a littoral habitat fauna composed of taxa like *Glyptotendipes*, *Dicrotendipes*, *Xenochironomus* and *Phaenopsectra* (Fig. 3). Development of the aquatic macrophyte flora near the coring site is suggested by the high relative abundance of *Glyptotendipes*, a vegetation mining taxon (Thut 1968; Pinder and Reiss 1983; Klink and Moller-Pillot 2003). The high percentage of TOM and the low HI (Fig. 4) are most probably caused by a greater contribution of aquatic macrophytes remains in the sediment. A terrestrial origin of the OM remains would be in contradiction with the poor arboreal cover indicated by the high percentages of non-arboreal pollen (NAP curve in Fig. 6). High relative abundance of *Microtendipes* may be explained by the high proportion of carbonated concretions (Oncolites and Cawlfower morphotypes, Magny et al. 2006a) and coarse organic particles in the sediment. A substrate preference of *Microtendipes* for similar littoral habitat was found by McGarrigle (1980).

The first part of the Oldest Dryas is marked by the only occurrence of *Stictochironomus* for the whole record (Fig. 3). *Stictochironomus* can be found in the littoral zone of cold lakes with low organic accumulation and well-oxygenated waters (Hofmann 1986). Given the overall low water depth during the Oldest Dryas, cool conditions seemed to prevail. In the lower part of the Oldest Dryas, the lower abundance of *Glyptotendipes* (Fig. 3) and the higher HI (Fig. 4) points to a less developed aquatic macrophyte belt, probably limited by cold climate.

Late glacial Interstadial

The late glacial Interstadial is represented by CAZ-Ac2 and CAZ-Ac3 in the chironomid stratigraphy (Fig. 3). The transition from CAZ-Ac1 to CAZ-Ac2 is abrupt (Fig. 3). *Microtendipes*, *Glyptotendipes* and other littoral habitat taxa sharply decline in abundance. New dominant taxa (*Chironomus*, *Cricotopus* type *patens*, *Psectrocladius* type *sordidellus* and *Ablabesmyia*) characterise both CAZ-Ac2 and CAZ-Ac3.

The late glacial Interstadial is marked by a decrease in TOC contents (Fig. 4). This decline cannot be interpreted as a lower organic input to the sediment since the sedimentation rate is much higher than in the previous zone (Fig. 2). HI increases up to its maximum for the whole record, whereas TOM percentages decline. Several successive episodes of high lake-levels (Fig. 6) were reconstructed during the late glacial Interstadial (Magny et al. 2006a).

Consequently to these higher water levels, the distance of the lake paleoshore from the coring site increases. The typical littoral habitats including wood and aquatic plant remains had then little influence on the chironomid fauna. The sediment OM was dominated by phytoplankton-like particles of pelagic origin. The higher proportion of fine OM in the sediment led to the dominance of taxa, such as *Ablabesmyia* and *Psectrocladius* type *sordidellus*.

Several episodes of lake-level lowering of similar magnitude as during the Oldest Dryas have been reconstructed during the late glacial Interstadial (Fig. 6, Magny et al. 2006a). Nevertheless, responses of both the chironomid community and the OM to these low lake-level episodes were different than during the Oldest Dryas. The low lake-level event between 15.50 m and 15.85 m corresponds to a peak of *Chironomus* (Fig. 3). This period is also marked by higher TOM percentages as well as lower HI (Fig. 4). The low lake-level probably favoured the development of aquatic macrophytes in the vicinity of the coring site. Some members of *Chironomus* are known to survive under anoxia and are typical of the deep zone of eutrophic lakes, whereas others live in littoral habitats, such as aquatic macrophyte belts (Butler et al. 1999). The high relative abundance of *Chironomus* observed between 15.50 and 15.85 m (Fig. 3) may then be linked to the development of these littoral-living *Chironomus*. The lake-level lowering identified at 13.65–14.05 m seemed to have no visible effect on OM, but corresponds to a high relative abundance of *Chironomus*, most probably littoral species (Fig. 3). Between 14.85 m and 15.10 m, a low lake-level event correlated with the Older Dryas pollen zone (Magny et al. 2006a). It was followed by the dominance of *Cricotopus* type *patens* and the occurrence of *Xenochironomus* in the chironomid community (Fig. 3). In Fennoscandia, *Cricotopus patens* is found in submerged mosses in lakes and in smaller bodies of standing water (Hirvenoja 1973).

Younger Dryas

The first part of the Younger Dryas (13.00–13.65 m) was marked by high lake-levels (Magny et al. 2006a). Nevertheless, a short lowering event could have occurred between 13.30 and 13.45 m (lake-level curve in Fig. 6). In the OM record, this episode is characterised by notable TOM percentages and a drop in HI (Fig. 4). The low lake-level led to growth of aquatic macrophytes near the coring site as suggested by the increase in the remains of littoral aquatic macrophytes in the sediment OM. In the chironomid assemblages, this short event was followed by an increase in the assemblage diversity (Hill's N2 in Fig. 3) probably due to the diversification of the benthic habitat. A slight rise in relative abundance of littoral habitat taxa, such as *Labrundinia*, *Glyptotendipes* and *Polypedilum* type *sordens*, can also be recognised (Fig. 3).

The second part of the Younger Dryas (13.00–12.20 m) was characterised by a lacustrine regressive episode (Fig. 6, Magny et al. 2006a). Impacts of this lowering were found in both the OM content and the chironomid stratigraphy. In the OM record, HI and TOM percentages reached similar values as during the Oldest Dryas (Fig. 4). An increase in allochthonous inputs of terrestrial plants and soil particles is inconsistent with the strongly reduced vegetation cover as indicated by the high NAP percentages in the pollen stratigraphy (Fig. 6). Thus, the observed trend was due to the accumulation of organic remains from the aquatic macrophytes belt that grew in the near vicinity of the coring site. The chironomid community (CAZ-Ac5 and CAZ-Ac6) was dominated by typical littoral taxa like *Polypedilum* type *sordens* and *Labrundinia*. The littoral environment favoured the development of *Microtendipes* and sustained a well diversified fauna as indicated by the higher assemblage diversity (Hill's N2 in Fig. 3).

Early Holocene

At the Younger Dryas/Holocene transition, a short hiatus was followed by a brief lake lowering (Fig. 6, Magny et al. 2006a). Imprints of this event were found in both chironomid and organic records. The high relative abundance of littoral habitat taxa was maintained briefly posterior to the beginning of the Holocene (Fig. 6). This period was also marked by low HI and high TOM percentages as during the second phase of the Younger Dryas (Fig. 4).

The higher lake-level reconstructed after 12.10 m coincides with the transition CAZ-Ac6/CAZ-Ac7, which was characterised by the decline of littoral taxa combined to the dominance of *Chironomus*, *Cryptochironomus*, *Procladius* and *Tanytarsus* (Fig. 3). *Chironomus* specimens found during CAZ-Ac7 belong to the *plumosus*-group (Fig. 3), which is more common of deep zones in eutrophic polyhumic lakes with high phytoplanktonic production and high nutrient concentrations (Saether 1979). This group is also known to live in anoxic conditions due to physiological and behavioural adaptations (Heinis and Crommentuijn 1992). The OM stratigraphy indicates the dominance of autochthonous organic remains in the total OM in the sediment, as shown by high HI (Fig. 6). The age–depth model (Fig. 2) indicates a similar sedimentation rate as during the Younger Dryas. A higher organic influx might then be deduced from the increase in TOC contents. The higher lake-level, combined with a warmer climate as indicated by the pollen record (Drescher-Schneider et al. 2007), led to a large increase in lacustrine production. The excessive organic production led to OM accumulation in the sediment (Fig. 6). The following oxygen depletion, also possibly favoured by the increase in the lake-level, most likely induced the strong dominance of taxa resistant to hypoxia, such as *Chironomus* gr. *plumosus*, and a drop in the assemblage diversity (Hill's N2 in Fig. 3).

The high lake-level phase was interrupted at 11.60 m by a water lowering that possibly was correlated to the Preboreal Oscillation (Magny et al. 2006a). The sharp peak of *Cricotopus* type *patens* and the presence of a few specimens of *Labrundinia* found at the time might be the result of the development of a littoral habitat.

Discussion

Controlling factors for the chironomid community and the sediment organic matter

Main shifts in chironomid and OM were concomitant with major lake-level changes. The low lake-level during the Oldest and Younger Dryas corresponded to the dominance of littoral habitat taxa in the chironomid community and higher proportions of aquatic macrophyte remains in the OM composition. Changes in chironomid and OM records were also synchronous with some major modifications in the pollen assemblages. Reduced arboreal cover (e.g. during Oldest Dryas and Younger Dryas) were concomitant with a high proportion of littoral remains in the OM (Fig. 6) so that there is probably no cause–effect relationship between vegetation on the watershed and OM in the sediment. Both respond in concert to changes in the water balance. During the Oldest and Younger Dryas, the decline of arboreal cover was driven by dry and probably cold conditions (Drescher-Schneider et al. 2007). Our work showed that dry phases (Fig. 6) followed by the reduction in arboreal cover and concurrently by the drop in lake-level (Magny et al. 2006a) led to the consecutive development of aquatic macrophytes in the vicinity of the coring point. On the contrary, wet episodes led to a concurrent increase in vegetation cover around the lake (NAP decrease in

Fig. 6) and to a higher lake-level (Fig. 6) determining a higher contribution of algal organic remains in the sediment (HI increase in Fig. 6). Lake-level fluctuations have then a major impact on the chironomid assemblages. At the coring point, the influence of water depth oscillations on the chironomid assemblages was most probably indirect through multiple impacts on lake properties (Velle et al. 2005). The proxies analysed in this study enlightened the influence of some benthic habitat characteristics (the humic status and the substrate type).

In mid-latitude and northern European lakes, a cold-adapted chironomid fauna dominated the Oldest and Younger Dryas (Brooks and Birks 2001). At the late glacial Interstadial and the early Holocene, it was replaced by typical warm lake assemblages (e.g. Millet et al. 2003). At Lago dell'Accesa, analogue changes in the chironomid community were not recorded. Chironomids living in temperate to warm environments maintained a high relative abundance throughout the late glacial period. A short event, coinciding with the first part of the Oldest Dryas could correspond to colder conditions, as indicated by the presence of *Stictochironomus* in the chironomid assemblage. This implies that temperature changes played only a minor role on the late glacial littoral chironomid succession at Lago dell'Accesa. In their climate quantifications for the two major warming phases of the last deglaciation at 14.7 and 11.5 ka cal. BP, Renssen and Isarin (2001) showed that in southern Europe, the increase in July temperature was less intense than in north-western Europe. Moreover, they suggested July air temperature around 17–20°C at sea-level during the Oldest and Younger Dryas in the area of Lago dell'Accesa. At this low elevation Mediterranean site (157 m a.s.l) summer temperature variations were probably too low to produce significant modifications in littoral fauna composition, whereas marked changes in benthic habitat characteristics were found as one of the main controlling factor of the chironomid assemblages.

Littoral chironomids and organic matter as lake-level indicators

Our study suggests that the chironomid assemblage and the OM were sensitive indicators to the major changes in lake-level. The results show that taxa such as *Microtendipes*, *Dicrotendipes*, *Glyptotendipes*, *Labrundinia*, *Limnophyes*, *Phaenopsectra*, *Polypedilum* type *sordens*, *Stenochironomus*, *Xenochironomus* and *Cricotopus* type *patens* should be considered as low lake-level indicators in littoral records. Korhola et al. (2000) already showed that *Microtendipes* and *Dicrotendipes* seemed to prefer shallower lakes in northern Fennoscandia. Barley et al. (2006) found also low water-depth optima for *Glyptotendipes*, *Dicrotendipes* and *Microtendipes* in their northwest North American training set. From our study, other taxa such as *Chironomus* type *plumosus* should be recognised as deeper water indicators. Taken alone, the indicator value of these chironomid taxa could be questioned. For example, *Glyptotendipes* may be associated with macrophytes, but this taxon also includes several species of collector-feeders that filter food materials from the water column (Armitage et al. 1995). Parallel analyses of chironomids with organic geochemistry and petrology studies may circumvent this uncertainty. For example, a combination of high relative abundances of *Glyptotendipes* with a high proportion of OM remains from aquatic macrophytes provides less questionable indication of a low lake-level during the Oldest Dryas at Lago dell'Accesa. A dominance of *Chironomus* type *plumosus* concurring with a high accumulation of planktonic OM implies a higher water level, as was found during the early Holocene at Lago dell'Accesa.

Century-scale oscillation like the Older Dryas and the Preboreal Oscillation only had a minor impact on our chironomid and OM stratigraphy. There is probably a need for a higher taxonomic resolution of the chironomid head capsules. The indicator value of a taxon is better

at the species or species group level than at the genus level. For example, *Chironomus* includes many species with very different ecological preferences (Butler et al. 1999). In the Lago dell'Accesa littoral record, high relative abundances of this genus was found both during high and low lake-level periods. Nevertheless, a high proportion of *Chironomus* type *plumosus* seemed to be linked to high lake-level events unlike the *Chironomus* type *anthracinus*, which may include typical species from the littoral environment.

Chironomid and OM analysis also provided original information on lake-level variations. In the first part of the Younger Dryas, brief changes in the relative frequency of carbonated concretion morphotypes were not interpreted as significant evidence of a low lake-level event (Magny et al. 2006a). However, synchronous shifts in chironomid assemblages and OM, which occurred during this period, strengthened the hypothesis of a short but well marked low lake-level event.

Conclusion

Late glacial chironomid (Diptera: Chironomidae) assemblages, organic petrography and geochemistry were analysed in a sediment core from the littoral zone of a Mediterranean lake, Lago dell'Accesa (Tuscany, Italy), where lake-level and vegetation history previously have been reconstructed. We tried to assess the response of chironomid community and sediment OM to lake-level, vegetation and climate changes. The proxies analysed suggested that chironomid succession was influenced indirectly by lake-level variations through their impact on benthic habitat characteristics and more specifically on the humic status and substrate type. At this low elevation Mediterranean site, small temperature changes played only a minor role on the late glacial littoral chironomid fauna. In this geographical area, interpretation of future littoral chironomid stratigraphy should take into consideration the predominant influence of lake-level variations. Our results showed that littoral chironomid fauna can be used in combination with OM analysis as a useful indicator of past lake-level changes in Mediterranean lakes. These proxies give original and complementary information on lake-level variations independent of other approaches. Nevertheless, an improvement of the identification level of chironomid remains is needed to refine lake-level reconstruction.

Acknowledgements

Financial support for this study was provided by the French CNRS (Program ECLIPSE). The authors express their sincere thanks to Mouna el Mekki and Didier Kéravis for technical assistance and to James Dat for his help with the English language. Many thanks also to Gaute Velle and the anonymous reviewer for helpful comments.

References

Armitage P, Cranston PS, Pinder LCV (1995) The Chironomidae. The biology and ecology of non-biting midges. Chapman & Hall, London

- Asioli A, Trincardi F, Lowe JJ, Ariztegui D, Langone L, Oldfield F (2001) Sub-millennial scale climatic oscillations in the central Adriatic during the Lateglacial: palaeoceanographic implications. *Quat Sci Rev* 20:1201–1221
- Barley EM, Walker IR, Kurek J, Cwynar LC, Mathewes RW, Gajewski K, Finney BP (2006) A northwest North American training set: distribution of freshwater midges in relation to air temperature and lake depth. *J Paleolimnol* 36:295–314
- Bennett KD (1996) Determination of the number of zones in a biostratigraphical sequence. *New Phytol* 132:155–170
- Brinkhurst RO (1974) *The benthos of lakes*. Macmillan Press Ltd., London
- Brooks SJ, Birks HJB (2001) Chironomid-inferred air temperatures from Lateglacial and Holocene sites in north-west Europe: progress and problems. *Quat Sci Rev* 20:1723–1741
- Brooks SJ, Mayle FE, Lowe JJ (1997) Chironomid-based Late-glacial climatic reconstruction for southeast Scotland. *J Quat Sci* 12:161–167
- Butler MG, Reznicka RL, Giovannielli KM (1999) Utility of species-level identification of *Chironomus* larvae in biological assessment of prairie lakes. *Int Ver Theor Angew Limnol Verh* 27:273–277
- Disnar JR, Guillet B, Keravis D, Di-Giovanni C, Sebag D (2003) Soil OM (SOM) characterisation by Rock-Eval pyrolysis: scope and limitations. *Org Geochem* 34:327–343
- Di Vito MA, Isaia R, Orsi G, Southon J, de Vita S, D'Antonio M, Pappalardo L, Piochi M (1999) Volcanism and deformation since 12,000 years at the Campi Flegrei caldera (Italy). *J Volcanol Geoth Res* 91:221–246
- Drescher-Schneider R, Beaulieu JL, Magny M, Walter-Simonnet AV, Bossuet G, Millet L, Bruggiapaglia E, Drescher A (2007) Vegetation history, climate and human impact over the last 15 000 years at Lago dell'Accesa (Tuscany, Central Italy). *Veget Hist Archeobot*. doi:10.1007/s00334-006-0089-z
- Espitalié J, Deroo G, Marquis F (1985a) La pyrolyse Rock Eval et ses applications. 2de partie. *Rev Inst Fr Pet* 40:755–784
- Espitalié J, Deroo G, Marquis F (1985b) La pyrolyse Rock Eval et ses applications. 3ème partie. *Rev Inst Fr Pet* 41:73–89
- Fittkau EJ, Roback SS (1983) The larvae of Tanyptodinae (Diptera: Chironomidae) of the Holarctic region—Keys and diagnoses. *Entomol Scan* 19(Suppl):33–110
- Gandouin E, Maasri A, Van Viet-Lanoë B, Franquet E (2006) Chironomid (Insecta: Diptera) assemblages from a gradient of lotic and lentic waterbodies in river floodplains of France: a methodological tool for paleoecological applications. *J Paleolimnol* 35:149–166
- Grimm EC (1991) *TILIA and TILIA GRAPH Software*. Illinois State Museum

- Heinis FH, Crommentuijn T (1992) Behavioural responses to changing oxygen concentrations of deposit feeding chironomid larvae (Diptera) of littoral and profundal habitats. *Arch Hydrobiol* 124:173–185
- Heiri O, Millet L (2005) Reconstruction of Late Glacial summer temperatures from chironomid assemblages in Lac Lautrey (Jura, France). *J Quat Sci* 20:31–44
- Hill MO (1973) Diversity and evenness: a unifying notation and its consequences. *Ecology* 54:427–432
- Hirvenoja M (1973) Revision der gattung *Cicotopus* van der Wulp und ihrer Verwandten (Diptera, Chironomidae). *Ann Zool Fenn* 10:1–363
- Hofmann W (1971) Zur Taxonomie und Palökologie subfossiler Chironomiden (Dipt.) in Seesedimenten. *Arch Hydrobiol Beih* 6:1–50
- Hofmann W (1986) Chironomid analysis. In: Berglund BE (ed) *Handbook on holocene palaeoecology and palaeohydrology*. Wiley, Chichester, pp 715–727
- Huntley B, Watts WA, Allen JRM, Zolitschka B (1999) Palaeoclimate, chronology and vegetation history of the Weichselian Lateglacial: comparative analysis of data from three cores at Lago Grande di Monticchio, southern Italy. *Quat Sci Rev* 18:945–960
- Jacob J, Disnar JR, Boussafir M, Sifeddine A, Turcq B, Albuquerque ALS (2004) Major environmental changes recorded by lacustrine sedimentary organic matter since the Last Glacial Maximum under the tropics (Lagoa do Caço, NE Brazil). *Palaeogeogr Palaeoclim Palaeoecol* 205:183–197
- Klink AG, Moller Pillot HKM (2003) Chironomidae larvae. Key to the higher taxa and species of the lowlands of Northwestern Europe. World Biodiversity Database CD-ROM series. Expert Center for Taxonomic Identification, University of Amsterdam, Amsterdam
- Korhola A, Olander H, Blom T (2000) Cladoceran and chironomid assemblages as quantitative indicators of water depth in subarctic Fennoscandian lake. *J Paleolimnol* 24:43–54
- Lallier-Vergès E, Sifeddine A, Beaulieu JL, Reille M, Tribouvillard N, Bertrand P, Mongenot T, Thouveny N, Disnar JR, Guillet B (1993) Sensibilité de la sédimentation organique aux variations climatiques du Tardi-Würm et de l'Holocène - le lac du Bouchet (Haute-Loire, France). *Bull Soc Geol France* 164:661–673
- Langford FF, Blanc-Valleron MM (1990) Interpreting Rock-Eval pyrolysis data using graphs of pyrolysable hydrocarbons vs. total organic carbon. *Am Assoc Petrol Geol Bull* 74:799–804
- Magny M, Beaulieu JL, Drescher-Schneider R, Vannièrè B, Walter-Simonnet A-V, Millet L, Bossuet G, Peyron O (2006a) Climatic oscillations in central Italy during the Last Glacial–Holocene transition: the record from Lake Accesa. *J Quat Sci* 21(4):311–320
- Magny M, Aalbersberg G, Bégeot C, Benoit-Ruffaldi P, Bossuet G, Disnar JR, Heiri O, Laggoun-Defarge F, Millet L, Peyron O, Vannièrè B, Walter-Simonnet AV (2006b)

Environmental and climatic changes in the Jura mountains (eastern France) during the Last Glacial–Interglacial transition: a multi-proxy record from Lake Lautrey. *Quat Sci Rev* 25(5/6):414–445

Maitland PS (1979) The distribution of zoobenthos and sediments in Loch Leven, Kinross, Scotland. *Arch Hydrobiol* 85:98–125

Manca M, Nocentini AM, Belis CA, Comoli P, Corbella L (1996) Invertebrate fossil remains as indicators of late Quaternary environmental changes in Latium crater lakes (L Albano and L Nemi). *Mem Ist Ital Idrobiol* 55:149–176

Mastrantuono L (1991) Zoobenthos associated with submerged macrophytes in littoral areas of Lake Vico (Italy) some relations between fauna structure and water quality. *Limnetica* 7:153–162

McGarrigle ML (1980) The distribution of chironomid communities and controlling sediment parameters. In: Murray DA (ed) *Chironomidae. Ecology, systematics, cytology and physiology*. Pergamon Press, Oxford, pp 275–282

Merciai G (1933) Il Lago dell'Accesa presso Massa Maritima. *Soc Toscanese Sci Nat* 43:29–50

Meyers P, Lallier-Vergès E (1999) Lacustrine sedimentary organic matter records of Late Quaternary paleoclimates. *J Paleolimnol* 21:345–372

Millet L, Verneaux V, Magny M (2003) Lateglacial paleoenvironmental reconstruction using subfossil chironomid assemblages from Lake Lautrey (Jura, France). *Arch Hydrobiol* 156:405–429

Noël H, Garbolino E, Brauer A, Lallier-Vergès E, Beaulieu JL, Disnar JR (2001) Human impact and soil erosion during the last 5000 years as recorded on lacustrine sedimentary organic matter – Lac d'Annecy, the French Alps. *J Paleolimnol* 25:229–244

Palomäki R (1989) The chironomid larvae in the different depth zones of the littoral in some Finnish lakes. *Acta Biol Debr Oecol Hung* 3:257–266

Pinder LCV, Reiss F (1983) The larvae of Chironominae (Diptera: Chironomidae) of the Holarctic region-Key and diagnoses. *Entomol Scand Suppl* 19:293–435

Ramrath A, Nowaczyk NR, Negendank JFW (1999) Sedimentological evidence for environmental changes since 34,000 years BP from Lago di Mezzano, central Italy. *J Paleolimnol* 21:423–435

Renssen H, Isarin RFB (2001) The two major warming phases of the last deglaciation at 14.7 and 11.5 ka cal BP in Europe: climate reconstructions and AGCM experiments. *Glob Planet Change* 30:117–153

Saether OA (1979) Chironomid communities as water quality indicators. *Holarct Ecol* 2:65–74

Schmidt R, Van Den Bogaard C, Merkt J, Müller J (2002) A new Lateglacial chronostratigraphic tephra marker for the south-eastern Alps: The Neapolitan Yellow Tuff (NYT) in Längsee (Austria) in the context of a regional biostratigraphy and palaeoclimate. *Quat Int* 88:45–56

Sifeddine A, Bertrand P, Lallier-Vergès E, Patience AJ (1996) Lacustrine organic fluxes and paleoclimatic variations during the last 15 ka: Lac du Bouchet (Massif Central, France). *Quat Sci Rev* 15:203–211

Stuiver M, Reimer PJ, Bard E, Beck W, Burr GS, Hughen KA, Kromer B, McCoarman G, Van Der Plicht J, Spurk M (1998) INTCAL 98 Radiocarbon age calibration, 24 000-0 cal. BP. *Radiocarbon* 40:1041–1083

Talbot MR, Livingstone DA (1989) Hydrogen index and carbon isotopes of lacustrine organic matter as lake-level indicators. *Palaeoclim Palaeogeogr Palaeoecol* 70:121–137

Thut RN (1968) A study of the profundal bottom fauna of lake Washington. *Ecol Monogr* 39:79–100

Velle G, Brooks SJ, Birks HJB, Willassen E (2005) Chironomids as a tool for inferring Holocene climate: an assessment based on six sites in southern Scandinavia. *Quat Sci Rev* 24:1429–1462

Verneaux V, Aleya L (1998) Bathymetric distributions of chironomid communities in ten French lakes: implications on lake classification. *Arch Hydrobiol* 142:209–228

Walker IR (2001) Midges: Chironomidae and related Diptera. In: Last WM, Smol JP (eds) *Tracking Environmental change using lake sediments. Volume 4: Zoological Indicators*. Kluwer Academic Publishers, Dordrecht, pp 43–66

Walker IR, Smol JP, Engstrom DR, Birks HJB (1991) An assessment of Chironomidae as quantitative indicators of past climatic change. *Can J Fish Aquat Sci* 48:975–987

Wiederholm T (1983) Chironomidae of the Holarctic region. Keys and diagnoses. Part 1. Larvae. *Entomol Scand Suppl* 19:1–457

Woodward CA, Shulmeister J (2006) New Zealand chironomids as proxies for human-induced and natural environmental change: transfer functions for temperature and lake production (chlorophyll a). *J Paleolimnol* 36:407–429

Wulf S, Kraml M, Brauer A, Keller J, Negendank JFW (2004) Tephrochronology of the 100 ka lacustrine sediment record of Lago Grande di Monticchio (southern Italy). *Quat Int* 122:7–30

Figures




Fig. 1 Study site location and Lago dell'Accesa main characteristics




Fig. 2 Age–depth model built from tephra layers and radiocarbon dates. The age of the main late glacial events identified from the pollen stratigraphy are also indicated




Fig. 3 Chironomid assemblage diagram from Lago dell'Accesa showing relative abundances and assemblage diversity (N2, Hill 1973) relative to depth in the core. Percentages of littoral biotope taxa were cumulated (dark curve: three point running mean). The taxa are distributed from left to right according to the depth along the core of their maximum relative abundance. Grey bars indicate samples in which less than 10 head capsules were recovered. Stratigraphic

boundaries of the statistical significant biozones (CAZ-Ac: Chironomid assemblage zone of Accesa) are indicated. I.A.C.P = Intra Alleröd Cold Phase; P.B.O. = Preboreal Oscillation




Fig. 4 Organic geochemistry and petrography diagram from Lago Dell'Accesa. TOC (Total Organic Carbon) in grams of carbon per gram of dried sediment (%) (dark curve: three-point running mean); HI: Hydrogen Index in milligrams of hydrocarbon per gram TOC (dark curve: three-point running mean); TOM (Terrestrial Organic Matter) in percentages of the palynofacies. I.A.C.P = Intra Alleröd Cold Phase; P.B.O. = Preboreal Oscillation




Fig. 5 S2 rock-Eval peak versus Total Organic Carbon. S2 in milligrams of Hydrocarbons released during pyrolysis per gram of dried sediment. $HI = [S2 \cdot TOC^{-1}] \times 100$




Fig. 6 Main results derived from pollen (NAP: Non-Arboreal Pollen, Drescher-Schneider et al. 2007), chironomid and OM analysis, and the lake-level events derived from Magny et al. (2006a). The age of the major late glacial events was determined by using the age–depth model (Fig. 2). See Figs. 3 and 4 for details

Tables

Table 1 Tephra layers and radiocarbon dates used to build the age–depth model

Depth (cm)		Age	Calibrated age (1sigma)	Laboratory number	References
1040.5	¹⁴ C from peat	8,830 ± 50 BP	9,775–10,147 cal. yrs BP	POZ-9970	
1158.5	¹⁴ C from wood fragments	9,890 ± 40 BP	11,296–11,226 cal. yrs BP	VERA-2807	
1200	Tephra: Pomici Principali	9,760 ± 300 BP			De vito et al. (1999)
1305	Tephra: Pomici Principali (1, main)	10,320 ± 50 BP 12,180 varve yrs BP			Wulf et al. (2004)
1542	Tephra: Neapolitan Yellow Tuff	12,300 ± 300 BP 14,120 varve yrs BP			Schmidt et al. (2002); Wulf et al. (2004)
1605	Tephra: Tufi Biancastri	14,557 varve yrs BP			Wulf et al. (2004)
1660	Tephra: Amendolare	15,550 varve yrs BP			Wulf et al. (2004)

Radiocarbon dates were calibrated using CALIB 4.3 (Stuiver et al. 1998)

