

# What do we know about Fossil Organic Carbon stock and fluxes in supergene environments?

Yoann Copard, Christian Di Giovanni

► **To cite this version:**

Yoann Copard, Christian Di Giovanni. What do we know about Fossil Organic Carbon stock and fluxes in supergene environments?. 20th Colloquium of African Geology, 2004, Orléans, France. hal-00089065

**HAL Id: hal-00089065**

**<https://hal-insu.archives-ouvertes.fr/hal-00089065>**

Submitted on 9 Aug 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# What do we know about Fossil Organic Carbon stock and fluxes in supergene environments?

Yoann Copard, Christian Di-Giovanni  
ISTO – UMR 6113 du CNRS, rue de St Amand, bâtiment Géosciences, BP 47,  
45067 Orléans cedex 2  
email: yoann.copard@univ-orleans.fr; christian.di-giovanni@univ-orleans.fr

CO<sub>2</sub> concentration is regulated by processes controlling the carbon fluxes occurring between different carbon pools such as atmosphere, ocean, hydrosphere, soil, biomass, and lithosphere. However, some controversies persist on the nature and quantification of carbon stocks and fluxes acting in supergene environment such as soils and rivers pools. For example, Meybeck (1993) considers that FOC released by weathering of sedimentary and metamorphic rocks can reflect up to 40-45% of POC annually carried by major rivers, instead of others numerous authors (Holland, 1978; Adams and Faure, 1996) that do not take into account this FOC yield. Up to now, at a global scale, we only know an initial estimation of the annual organic yield induced by carbonates and shales chemical weathering that reach 0.1 Gt C y<sup>-1</sup> (Di-Giovanni et al., 2002). No FOC fluxes have been reported for chemical weathering of sands and sandstones, as well as those inherited from mechanical weathering of all sedimentary rocks, although mechanical weathering is 4 to 11 higher than chemical one (Probst, 1992; Gaillardet, pers. com.). In addition to these gaps concerning FOC fluxes, the role of erosional processes on those of FOC in global carbon cycle (source or sink of CO<sub>2</sub>) and the FOC stock available in surficial layer of sedimentary rocks and rapidly reload in supergene carbon pools, remain unresolved.

The influence of weathering processes on the release and fate of FOC in supergene carbon cycle was approached through our recent results from a study carried out on 2 experimental watersheds (Draix; Copard et al. submitted). Input of FOM inherited from Jurassic marls in soils through alterites, where occur chemical weathering and bioerosion, experiences a significant, but not complete, mineralisation of its carbon. Input of FOM in streams, controlled by mechanical weathering, experiences no FOM mineralisation. Hence, for the global carbon cycle, FOM reload in supergene carbon cycle can act either as a maintaining of an ancient sink, when gully erosion is dominant, or as a source of carbon, when (bio)chemical erosion is preponderant.

At a global scale, we estimated FOC stock in the uppermost bedrock to a depth of 1m. That first estimate is of the same order of magnitude than soil OC (generally also calculated to a depth of 1 m), but obviously do not exhibit the same turnover of its OC.

We focus now to the preliminary estimate of FOC fluxes in world major watersheds that store about one half of the global FOC stock. For each of these watersheds, we attempt to calculate these fluxes by taking into account: their bedrock lithological composition, their mechanical and chemical rock denudation, their land cover type overlaps.

## References

- Adams J., Faure H., 1996. *In* Global continental change: The context of palaeohydrology (Brandson J. et al., Eds). Geological Society Special Publication 115, 27-42.
- Copard Y., Di-Giovanni Ch., Martaud T., Albéric P., Olivier J.E. submitted to Earth Surface processes and landforms.
- Di-Giovanni Ch., Disnar J.R., Macaire J.J., 2002. *Global and Planetary Change* 32, 195-210.
- Holland H.D., 1978. *The chemistry of atmosphere and oceans*. Wiley Interscience Publishers: Chichester, New York
- Meybeck M., 1993. *In* Interaction of C, N, P, and S, Biogeochemical Cycles on Global Change (Wollast R. et al., Eds), Springer Verlag, 163-193.
- Probst J.L., 1992. *Mémoire des Sciences Géologiques* 94, 161p.

**keywords:** Fossil Organic Carbon, carbon cycle, major watersheds, chemical weathering, mechanical weathering, carbon sink, carbon source

This work is a contribution of the GIS "Bassins de Draix, étude de l'érosion en montagne" and was financially supported by the CNRS program INSU PNSE-ACI 2001-2002: Influence de l'érosion sur les flux de matière organique fossile dans les géosystèmes continentaux actuels (sols, cours d'eau) : bassins expérimentaux de Draix, Alpes de Haute Provence, France.