



# Consequences of transformation of native Cerrado in pasture land on the porosity of a clayey-sand red latosol

Luíz Carlos Balbino, Ary Bruand, Michel Brossard, M. F. Guimarães, P. Quétin

## ► To cite this version:

Luíz Carlos Balbino, Ary Bruand, Michel Brossard, M. F. Guimarães, P. Quétin. Consequences of transformation of native Cerrado in pasture land on the porosity of a clayey-sand red latosol. inconnu, 2000, 3p. hal-00089690

**HAL Id: hal-00089690**

**<https://insu.hal.science/hal-00089690>**

Submitted on 22 Aug 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# CONSEQUENCES OF TRANSFORMATION OF NATIVE CERRADO IN PASTURE LAND ON THE POROSITY OF A CLAYEY-SAND RED LATOSOL.

L.C. Balbino<sup>1,2</sup>, A. Bruand<sup>1,3</sup>, M. Brossard<sup>4</sup>, M. F. Guimarães<sup>5</sup> and P. Quétin<sup>1</sup>

1: Science du Sol, INRA, BP 20619, 45166 Olivet cedex, France (Luiz.Balbino@orleans.inra.fr)

2: Embrapa CNPAF, BP 179, 74001-970 Goiânia, GO Brazil

3: ISTO, Université d'Orléans, Géosciences, BP 6759, 45067 Orléans cedex 2, France

4: IRD/Embrapa CPAC, BP 7091, 71619-970 Brasília, DF Brazil

5: Universidade Estadual de Londrina, Dept. de Agrônoma, BP 6001, 86051-990 Londrina, PR Brazil

Key-words: *Brachiaria decumbens*, structure, bulk density, mercury porosimetry, scanning electron microscopy

In the Brazilian savannah (*Cerrados* area), about 60 % of the surface area is used as pasture land and 30 % of the latter is cultivated pasture. Many studies have reported that pastures degrade quickly and degradation would be more intense when the soil was tilled prior pasture sowing. Pasture degradation (decrease in the pasture quality and yield, weed invasion) would be partly a consequence of the soil structure evolution (Balbino & Di Stefano, 1999). Origin of pasture degradation is still under discussion and analysis of the literature showed that there is a lack of quantitative data. The aim of this study is to perform a detailed analysis of the porosity in a low productivity cultivated pasture and to compare the results with the porosity of a similar soil still under native vegetation (*Cerradão*, i.e. Scierophylous forest).

## Material and methods

The study site is located in the *Mato Grosso do Sul* state, *Brasilândia* county (21° 22,53' S 52° 11,73' E). Mean annual rainfall is 1600 mm. The soil is a sandy-clay latosol (*Latossolo vermelho*) developed on deeply weathered cretaceous materials (*Bauru* sandstones). Clay fraction is kaolinite mainly with iron oxy-hydroxides in small proportion. Two sites with similar soils were selected: one in native forest (*cerrado*) and another one in a low productivity pasture of brachiara (*Brachiaria decumbens*) 20 years old. The soil was described down to 1 m depth with particular attention to structure. Bulk density measurements were performed using cylinder 98 cm<sup>3</sup> in volume. In the laboratory we investigated the pore size distribution from 360 down to 0.006 µm using mercury porosimetry (Fig. 1). Thin sections of undisturbed samples were examined in scanning electron microscopy using back-scattered electron emission (Bruand et al., 1996).

## Results and discussion

Under pasture, the upper horizon (0–3 cm) was massive with horizontal roots. Then from 3 to 15 cm depth, well developed and small to medium polyhedral aggregates associated with a few micro-aggregates were observed. Roots are mainly located between the polyhedral aggregates. Under forest, small to medium polyhedral aggregates with lots of micro-aggregates were encountered from 0 to 15 cm depth. Roots were located between and within the polyhedral aggregates. From 15 to 100 cm depth, aggregates were micro-aggregates mainly in a closer packing under pasture than under forest. A few polyhedral aggregates were present under pasture but their number and size decreased with depth. Biological activity (termites, ants and earthworms) was smaller under pasture than under forest and located close to the surface mainly.

Under forest, the bulk density ranged from 0.80 in the first top centimetres to 1.03 at 15 cm depth when under pasture it ranged from 1.08 to 1.22 from the surface to 13 cm depth (Table). These bulk density values showed a much smaller pore volume in the topsoil under pasture. From 18 to 100 cm depth, bulk density showed also smaller porosity under pasture.

Analysis of the structure using microscopy showed the presence of polyhedral aggregates a few millimetres to one centimetre in size and micro-aggregates 50–100 µm in size (Fig. 2). These two types of aggregate were present in different proportions according to the site and horizon. Analysis of the porosity showed that pore were (Fig. 1): (i) pores larger than several hundreds of micrometers in size which resulted from biological activity (root, ants and termites channels) and polyhedral aggregate packing, (ii) pores a few micrometers to one hundred micrometers in size which resulted from micro-aggregate packing and (iii) pores which resulted from the packing of clay particles within polyhedral aggregates and micro-aggregates and which were not visible in scanning electron

microscopy because they are too small. Comparison of porosity within every profile showed a decreased in the large pores with depth and an increase in the pores corresponding to micro-aggregates packing because they are more numerous with depth.

The mercury intrusion curve enabled the identification of several classes of pores (Bruand et al., 1993; Curmi et al., 1994). Results showed the presence of two classes of pores A and B which corresponded to pores with equivalent pore diameters ranging from 0.022 to 0.025  $\mu\text{m}$  and from 18 to 58  $\mu\text{m}$  in equivalent diameter, respectively (Fig. 1). Class B corresponded to pores resulting from the packing of micro-aggregates. Large pores resulting from polyhedral aggregates packing and biological activity were not measured or just partially because they were too large to be studied using mercury porosimetry. Results showed that the volume of pores B was greater in the 0–3 cm layer and much smaller (from 20 to 40 % smaller) in the 3–17 cm layer under pasture than under forest (Table). Pores B were also smaller under pasture than under forest as shown by the smaller  $D_{m,B}$ . Under pasture, the greater  $V_{p,B}$  in the 0–3 cm layer would result from a great under-estimation of  $V_{p,B}$  in the 0–3 cm layer under forest because the micro-aggregates were in a very loose packing leading to too large pores to be measured using mercury porosimetry. From 17 to 100 cm depth,  $V_{p,B}$  was similar or smaller (from 5 to 25 % smaller) under pasture. This is consistent with field observation which showed a less developed structure under pasture. This smaller  $V_{p,B}$  might result also from a smaller amount of micro-aggregates because of a smaller clay content (Table). The volume of pores belonging to class A ranged from 0.046 to 0.112  $\text{cm}^3 \text{g}^{-1}$  (Table). These pores resulted from the packing of clay particles within the polyhedral and micro-aggregates. The no or very little variation of  $D_{m,A}$  showed no or very little variation of both the size and packing of the elementary kaolinite particles. Such a result is consistent with the close relationship between  $V_{p,A}$  and the clay content ( $r^2 = 0.95$ ).

## Conclusion

Our results showed a strong evolution of the structure which led to a decrease in the porosity, particularly in the topsoil layers. Except the very small pores which result from the packing of the kaolinite particles within aggregates, all the pores were less developed under pasture. Field observations showed also a structure less developed from 15 to 100 cm depth under pasture and mercury porosimetry showed that micro-aggregates would be in a closer packing under pasture than under forest. However, this will need further analysis to be confirmed.

- Balbino, L.C., Di Stefano J.G., 1999. Projet PROCITROPICOS: Intégration de l'agriculture et de l'élevage par les systèmes de semis direct, pp. 409-417, In: Rasolo, F. and M. Raunet (eds), Gestion agrobiologique des sols et des systèmes de culture. Actes de l'atelier international, Antsirabe, Madagascar, 23-28 mars 1998, Cirad, Collection Colloques, 658p.
- Bruand, A., D'Acqui, L.P., Nyamugafata, A.P., Darthout, R., Ristori, G.G., 1993. Analysis of porosity in a tilled "crusting soil" in Zimbabwe. *Geoderma*, 59, 235-248.
- Bruand, A., Cousin, I., Nicoulaud, B., Duval, O., Bégon J.C., 1996. Backscattered electron scanning images of soil porosity for analysing soil compaction around roots. *Soil Science Society of America Journal*, 60, 895-901.
- Curmi, P., Kertzman, F.F., Quieroz Neto, J.P., 1994. Degradation of structure and hydraulic properties in an Oxisol under cultivation (Brazil). In: *Soil Micromorphology: Studies in Management and Genesis*. Developments in Soil Science 22, Elsevier, Amsterdam, 569-579.




Figure 1: Pore volume distribution for the 18–30 cm layer under forest.

Table: Physical and chemical data for the soils studied.<sup>a</sup>

	Horizon	PSD				OC	pH <sub>w</sub>	CEC	D <sub>b</sub>	Mercury porosimetry				$\mu\text{m}$
		<2	2–50	50–200	200–2000					Pore class A		Pore class B		
										V <sub>p,A</sub>	D <sub>m,A</sub>	V <sub>p,B</sub>	D <sub>m,B</sub>	
cm	g kg <sup>-1</sup>				cmol <sub>c</sub> kg <sup>-1</sup>		cm <sup>3</sup> g <sup>-1</sup>		$\mu\text{m}$		cm <sup>3</sup> g <sup>-1</sup>			
Forest	0–3	32.1	8.6	14.2	45.1	20.1	3.9	3.14	0.80	0.090	0.023	0.189	58	
	3–6	35.3	7.2	30.2	27.4	14.2	3.9	2.63	1.03	0.101	0.023	0.196	51	
	6–15	29.6	6.8	32.6	31.0	-	-	-	1.03	0.083	0.023	0.256	29	
	18–30	30.3	8.1	31.1	27.5	7.5	4.0	2.42	1.02	0.092	0.023	0.252	50	
	30–40	31.0	4.7	35.2	29.1	-	-	-	-	0.089	0.023	0.165	48	
	60–80	36.5	4.4	32.7	26.4	4.1	4.2	2.49	0.94	0.105	0.023	0.290	50	
	90–100	44.4	3.7	30.4	21.5	-	-	-	0.92	0.112	0.023	0.255	48	
Pasture	0–3	18.6	4.0	36.2	41.3	14.2	5.8	3.33	1.08	0.046	0.025	0.209	27	
	3–6	21.2	4.5	30.0	44.2	11.7	5.3	2.62	1.17	0.055	0.022	0.162	18	
	10–13	21.5	4.2	30.5	43.8	9.2	5.2	2.22	1.22	0.058	0.023	0.154	18	
	14–17	22.1	6.4	30.6	40.9	7.9	5.3	2.22	-	0.065	0.023	0.182	35	
	18–30	23.0	9.2	29.7	38.1	6.2	5.2	2.01	1.21	0.068	0.023	0.183	55	
	30–40	26.4	4.7	30.2	38.7	-	-	-	1.22	0.075	0.025	0.164	42	
	60–80	30.6	6.0	32.1	31.4	3.6	5.0	1.83	1.09	0.084	0.023	0.243	56	
	90–100	32.8	4.4	29.8	33.0	-	-	-	1.13	0.088	0.023	0.194	27	

<sup>a</sup>PSD is particle size distribution; OC is organic carbon content; CEC is cation-exchange capacity; D<sub>b</sub> is bulk density; V<sub>p,A</sub> and D<sub>m,A</sub> are the volume of pores A and their modal equivalent diameter, respectively; V<sub>p,B</sub> and D<sub>m,B</sub> are the volume of pores B and their modal equivalent diameter, respectively.




Figure 2: Back-scattered electron scanning image (x60, WD = 18 mm, 15 kV) of the 90–100 cm layer under forest (left) and of the 0–3 cm layer under pasture (right). Voids occupied by resin are black, homogeneous grey areas are quartz and heterogeneous dark grey areas are porous clayey phase.