

## **Induration of ferrallitic microaggregated soils (Northeastern Brazil).**

Mathieu Lamotte, Fernando B.R. Silva, René Boulet, Ary Bruand, Coelho Araujo Filho J., J.C. Leprun, Emmanuel Fritsch, Heloisa Filizola, Arnaud Vannier, Bruno Boulange

► **To cite this version:**

Mathieu Lamotte, Fernando B.R. Silva, René Boulet, Ary Bruand, Coelho Araujo Filho J., et al.. Induration of ferrallitic microaggregated soils (Northeastern Brazil).. World Congress of Soil Science, 1998, Montpellier, France. 7p. hal-00090054

**HAL Id: hal-00090054**

**<https://hal-insu.archives-ouvertes.fr/hal-00090054>**

Submitted on 28 Aug 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Enregistrement scientifique N° : 136  
Symposium N° : 4  
Présentation : Oral

## **Induration of ferrallitic microaggregated soils (Northeastern Brazil)**

### **Induration de sols microagrégés ferrallitiques (Nord-est du Brésil)**

**LAMOTTE Mathieu<sup>1</sup>, SILVA Fernando B. R.<sup>2</sup>, BOULET René<sup>1</sup>, BRUAND Ary<sup>3</sup>,  
ARAUJO FILHO J. Coelho de<sup>2</sup>, LEPRUN Jean-Claude<sup>4</sup>, FRITSCH Emmanuel<sup>1</sup>,  
FILIZOLA Heloisa F.<sup>5</sup>, VANNIER Arnaud<sup>3</sup>, BOULANGE Bruno<sup>1</sup>**

- <sup>1</sup> ORSTOM-NUPEGEL, Universidade de São Paulo, Instituto de Geociências / DGG, CP 11348, 05422-970 São Paulo - SP, Brazil (*mlamotte@usp.br*)  
<sup>2</sup> EMBRAPA-CNPS-UEP Recife, rua Antônio Falcão 402, Boa Viagem, 51020-240 Recife - PE, Brazil (*fbjc@elogica.com.br*)  
<sup>3</sup> INRA Orléans, Science du Sol, 45160 Ardon, France (*ary.bruand@orleans.inra.fr*)  
<sup>4</sup> ORSTOM, 911 avenue Agropolis, BP 5045, 34032 Montpellier cedex 1, France  
<sup>5</sup> EMBRAPA-CNPMA, CP 69, 13820-000 Jaguariúna - SP, Brazil

Studies of soils in the northeastern region of Brazil showed the widespread occurrence of hard subsurface horizons that strongly restrict the penetration of plant roots, water and ploughing tools, and that consequently reduce the agricultural yields. The objective was to characterise the groundmass of these hard horizons and to identify the cementing agent responsible for the hardness. Samples were collected from friable subsoil horizons which are latosolic horizons, and from hard subsoil horizons which are either *fragipã* or *duripã* according to the Brazilian soil classification. The mechanical stability of the groundmass was measured. Thin-sections were prepared and examined in scanning electron microscopy using backscattered electrons. The thin sections were also used for microprobe analysis.

The results showed a great stability of *fragipã* and *duripã* when they are shacked in water. This stability clearly indicated that they are indurated horizons and related to duripans. The results showed also that the groundmass of the indurated horizons differed from friable latosolic horizons by the continuity of the solid phase. The clayey material responsible for the continuity consisted mainly of kaolinite with small variation of the aluminium and iron contents. The difference of consistence between the non-indurated and indurated horizons would be related to an increase in the aluminium content and a decrease in the iron content. Thus, aluminium compounds would play a major role as cementing agent within the clayey material.

Key words: ferrallitic soils, duripan, microanalysis, iron, microaggregate, Brazil  
Mots clés: sol ferrallitique, duripan, microanalyse, fer, microagrégation, Brésil

Enregistrement scientifique N° : 136

Symposium N° : 4

Présentation : Oral

## **Induration of ferrallitic microaggregated soils (Northeastern Brazil)**

## **Induration de sols microagrégés ferrallitiques (Nordeste du Brésil)**

**LAMOTTE Mathieu<sup>1</sup>, SILVA Fernando B. R. <sup>2</sup>, BOULET René<sup>1</sup>, BRUAND Ary<sup>3</sup>,  
ARAÚJO FILHO J. Coelho de<sup>2</sup>, LEPRUN Jean-Claude<sup>4</sup>, FRITSCH Emmanuel<sup>1</sup>,  
FILIZOLA Heloisa F.<sup>5</sup>, VANNIER Arnaud<sup>3</sup>, BOULANGE Bruno<sup>1</sup>**

<sup>1</sup> ORSTOM-NUPEGEL, Universidade de São Paulo, Instituto de Geociências / DGG, CP 11348, 05422-970 São Paulo - SP, Brazil (*mlamotte@usp.br*)

<sup>2</sup> EMBRAPA-CNPS-UEP Recife, rua Antônio Falcão 402, Boa Viagem, 51020-240 Recife - PE, Brazil (*fbjc@elogica.com.br*)

<sup>3</sup> INRA Orléans, Science du Sol, 45160 Ardon, France (*ary.bruand@orleans.inra.fr*)

<sup>4</sup> ORSTOM, 911 avenue Agropolis, BP 5045, 34032 Montpellier cedex 1, France

<sup>5</sup> EMBRAPA-CNPMA, CP 69, 13820-000 Jaguariúna - SP, Brazil

### **Introduction**

Studies of soils in the north-eastern region of Brazil showed the widespread occurrence of hard subsurface horizons that strongly restrict the penetration of plant roots, water and ploughing tools, and that consequently reduce the agricultural yields (Embrapa, 1975). Hard horizons are present in topographic depressions of coastal plateaux (Silva & Araujo Filho, 1989; Silva *et al.*, 1993). The objective of the present study was to characterise the groundmass of the hard horizons and friable latosolic horizons, the latter being closely associated with the hard horizons within the toposequences. The results enabled discussion of cementing agent responsible for induration within the studied soils and were compared with those earlier published by Silva *et al.* (1997).

### **Material and methods**

#### *Site*

The site is near Boca da Mata, 50 km west of Maceió (Alagoas State). The climate is warm and humid with a mean annual air temperature of 25° C, a mean annual precipitation of 1640 mm and a mean annual evapotranspiration of 1450 mm. For 3-4

months, between October and January, rainfall can be smaller than 50 mm, i.e. much less than evapotranspiration. The landscape consists of plateaux (altitude of 120 m) which formed a discontinuous coastal band 15-75 km wide. The plateaux show numerous depressions variable in shape (depressions with or without outlet) and extension (from 5 m wide and 0.3 m deep to 3 km wide and 1 m deep). Originally, the plateaux were covered by dense forest; the deforestation process started about 300 years ago and the increasing of the sugar cane production aggravated the process in this century. Soils differentiated on the Barreira deposit (Kistler, 1954; Mabesoone & Alheiros, 1988). A soil survey of the site (Silva & Araujo Filho, 1989) showed that the soil cover consists mainly of *Podzólicos* and *Latosolos amarelos* (Embrapa, 1988; Oliveira *et al.*, 1992) which are Ultisols and Oxisols, respectively (Soil Survey Staff, 1975). Soils exhibited friable subsoil horizons (latosolic horizons) excepted in the depressions where they exhibited hard subsoil horizons overlaying latosolic horizons. *Podzois*, which are Spodosols, occurred also locally in the centre of the largest depressions.

#### *Field sampling*

A trench, 50 m long and 3 m deep was excavated from the edge to the centre of a depression for a detailed field study (Boulet *et al.*, 1998). The mineralogical composition consists invariably of kaolinite (about 85 %), quartz (about 10 %), iron oxi-hydroxides (about 4 %), anatase and zircon (about 1 %). The trench showed friable subsoil horizons, which are latosolic horizons (*L*), and hard subsoil horizons. The latter can be easily distinguished in the field by the lack of macroporosity and high strength. They can be sampled and broken only with difficulties under both wet and dry conditions. According to the Brazilian soil classification (Embrapa, 1988; Oliveira *et al.*, 1992), two types of hard horizons were identified: (i) *fragipã* (*F*) which become softer after immersion and shaking in water, (ii) *duripã* (*D*) which are always very hard even when immersed and shacked in water. The trench showed lateral transition between *L*, *F* and *D* subsoil horizons. Undisturbed samples were collected from these subsoil horizons at about 1 m depth. Collection of undisturbed samples from the *F* and *D* horizons required a bell-shaped saw with carbide teeth.

#### *Laboratory study*

Aggregate stability tests were carried out in triplicate with air-dried samples 3-5 mm in diameter. The procedure (Le Bissonnais, 1996) consisted of submitting the samples to three different treatments of disaggregation: (i) fast wetting by immersion in distilled water, (ii) slow wetting by capillary rise, (iii) immersion in ethanol before immersion in distilled water and mechanical shaking. After sieving in ethanol, the amount of aggregates remaining on the 50, 100, 200, 500, 1000, 2000 µm sieves and the amount of < 50 µm material were determined for each treatment.

After impregnation of oriented and undisturbed samples, thin-sections were prepared following the method of Fitzpatrick (1984). The surfaces were polished and coated with carbon for examination in scanning electron microscopy using backscattered electrons (Bruand *et al.*, 1996) and for electron microprobe analysis. Observations were carried out in scanning electron microscopy with a Cambridge 90B instrument (magnification from x 300 to 1500) using the backscattered electron emission.

Si, Al, Fe, Ti, Mn, K, Mg, Ca, Na, P, S and Cl contents were determined on the thin sections using a Camebax Datanim electron microprobe equipped with four wavelength-dispersive spectrometers (WDS) and linked with the ZAF-MBXCOR quantitative analysis software. The accelerating voltage was 15 kV, the probe current was 10  $\mu\text{A}$  and the count time was 90-120 s. Chemical determinations concerned small areas (1-2  $\mu\text{m}^2$ ) or larger surfaces (25-50  $\mu\text{m}^2$ ).

## Results and discussion

### *Mechanical stability of the groundmass fabric*

Stability tests showed that for the *L* horizon, 15 % of the sample remains > 2 mm after the fast wetting treatment and 30 % after slow wetting or mechanical shaking treatment. The *F* and *D* horizons showed very high stability. For the *F* horizon, 74 % of the sample remains > 2 mm after the fast wetting treatment, 77 % after the slow wetting treatment and 87 % after the mechanical treatment. The stability was higher for the *D* horizon (87 to 95 % of the sample remain > 2 mm whatever the treatment). The < 2 mm material resulting from stability tests was mainly constituted of > 100  $\mu\text{m}$  aggregates: only 10 % of the < 2 mm material was < 100  $\mu\text{m}$  for the *L* horizon and about 2 % for the *F* and *D* horizons.

Field observations suggested that the *D* horizon differed from the *F* horizon by cementation because the *D* horizon were always very hard even when immersed and slacked in water. Laboratory results showed high variation of stability between friable and hard horizons and low variation of stability between the *F* and *D* horizons. This low variation would be related more to the degree of induration than to any difference of origin for the consistence: cemented for the *duripã* and uncemented for the *fragipã*.

### *Groundmass fabric*

Thin section showed high proportion of clay in the groundmass of *L*, *F* and *D* horizons. The surface area occupied by clay and by coarse grains (both silt and sand) was about 35-45 % and 20-25 %, respectively. The coarse grains were mainly quartz, ranging from fine silt to coarse sand. They were well sorted and subangular to rounded grains.

Observations also revealed microaggregation of clay in the *L*, *F* and *D* horizons but it was more developed in the *L* horizon. The groundmass fabric of the latter was heterogeneous, mainly agglomeroplasmic, and secondarily porphyrosquelic (Brewer, 1964). The agglomeroplasmic fabric consisted of dense microaggregates (5-100  $\mu\text{m}$  in diameter) which were composed of closely packed clay (with or without incorporated fine silt). Most voids were either packing voids (2-100  $\mu\text{m}$ ) between microaggregates or vughs (100-250  $\mu\text{m}$ ) although rare channels were also present. These voids were polyconcave and were well connected. Some of the microaggregates were coated and linked one together by clay material. The coatings and bridges (1-2  $\mu\text{m}$  thick) differed from aggregates by their grey level on the backscattered electron scanning images (BESI). The aggregates were lighter than the coatings and bridges. The porphyrosquelic

fabric showed microaggregates similar to those which were described for the agglomeroplastic fabric, but they were very closely packed and embedded in a clay material which exhibited on the BESI a grey level darker than for the microaggregates. The porosity was lower than for the agglomeroplastic fabric. The voids consisted of packing voids (2-20  $\mu\text{m}$ ) between microaggregates, vughs or channels (100-250  $\mu\text{m}$ ) and planar voids (10-20  $\mu\text{m}$  in width). These voids varied in shape and were weakly connected. Clay-coatings and clay-bridges occurred also in some places.

The groundmass fabric of *F* and *D* horizons were also heterogeneous but mainly porphyroquelic, and secondarily agglomeroplastic. Thus, it was denser than for the *L* horizon. The surface area occupied by porphyroquelic fabric was largely greater than in the *L* horizon. The amount of aggregates covered by clay coating and linked by clay-bridges was also greater for the agglomeroplastic fabric. Consequently, the porosity was lower and the groundmass was more continuous in the *F* and *D* horizons than in the *L* horizon.

Finally, SEM observations showed that the *L*, *F* and *D* horizons have a microaggregated fabric but there is a gradient of continuity for the solid phase which increase from the *L* to the *F* horizons and from the latter to the *D* horizon. The continuity of the solid phase appeared in relation with the closed packing arrangement of the microaggregates and with the presence of clayey material covering and linking the microaggregates one together. This continuity and, consequently the low porosity (small voids weakly connected) could partly explain the hardness of the *F* and *D* horizons.

#### *Chemical composition of the fine material*

Results of microanalysis showed close composition of the clayey material between *L*, *F* and *D* horizons. The  $\text{SiO}_2$  content (about 50.2 %) was similar for all the studied horizons. The small difference of composition between *L*, *F* and *D* horizons concerned the  $\text{Al}_2\text{O}_3$  and  $\text{Fe}_2\text{O}_3$  contents. The  $\text{Al}_2\text{O}_3$  content increased from 42.6 % in the *L* horizon to 44.6 % in the *F* and *D* horizons. The  $\text{Fe}_2\text{O}_3$  content decreased progressively from 4.8 % in the *L* horizon to 4.3 % in the *F* horizon and then 2.7 % in the *D* horizon. The Si/Al atomic ratio of the clayey material was about 1.01 in the *L* horizon and about 0.95 in the *F* and *D* horizons. The results were consistent with the homogeneous and dominantly kaolinic nature of the clayey material. These results confirmed that both a decrease in the  $\text{Fe}_2\text{O}_3$  content and an increase in  $\text{Al}_2\text{O}_3$  content play a major role in the induration process as earlier discussed by Silva *et al.* (1997).

#### **Conclusion**

Even if the hard *F* and *D* horizons presented differences of consistence in the field, the laboratory study with air-dried samples showed low variation of stability. Their great stability even when they are immersed and shacked in water show that they are indurated horizons and related to duripans.

The fabric appeared mainly porphyroquelic in the *F* and *D* horizons and dominantly agglomeroplastic in the *L* horizon. Thus the groundmass of the indurated horizons differed from those of the non-indurated horizon by the continuity of the solid phase.

The clayey material responsible for the continuity consisted mainly of kaolinite with small variation of the aluminium and iron contents. The difference of consistence between the non-indurated and indurated horizons would be related to an increase in the aluminium content and a decrease in the iron content. Thus, aluminium compounds would play a major role as cementing agent within the clayey material. The cementation would be closely associated with a decrease in the iron content, probably because of the dissolution of iron oxi-hydroxide including a high proportion of Fe-Al substitutions. This process would explain both the decreases in the Si/Al ratio and Fe content, which were associated to the induration process for the studied soil.

### **Acknowledgements**

The authors thank the FAPESP (São Paulo) for the funding (Project 94/2471-8) and the Usina Triunfo (Boca da Mata) for technical assistance in the field. We thank also H. Gaillard and C. Le Lay (INRA-Orléans), G. Drouet and O. Rouer (CNRS-Orléans) for technical assistance in the laboratory.

### **References**

- Boulet, R., Fritsch, E., Filizola, H.F., Araujo Filho, J.C. (de), Leprun, J.C., Silva, F.B.R. (e), Balan, E. & Tessier, D. (1998). Iron bands, fragipan and duripan in the northeastern plateaux of Brazil. Structure, mineralogy and genesis. In: 11th International Clay conference, Ottawa, 1997, submitted.
- Brewer, R. (1964). Fabric and mineral analysis of soils. John Wiley & Sons, Inc., New York, 470 p.
- Bruand, A., Cousin, I., Nicoullaud, B., Duval, O. & Begon, J.C. (1996). Backscattered electron-scanning images of soil porosity for analysing soil compaction around roots. *Soil Sci. Soc. Am. J.*, 60, 895-901.
- Embrapa (1975). Levamento exploratório-reconhecimento de solos do Estado de Alagoas. Embrapa/ CPP, Boletim Técnico, 36 - SUDENE/DRN, Série Recursos de Solos, 5 (Recife), 35, 531 p.
- Embrapa (1988). Critérios para distinção de classes de solos e de fases de unidades de mapeamento. Embrapa/SNLCS (Rio de Janeiro), 11, 67 p.
- Fitzpatrick, E.A. (1984). Micromorphology of soils. Chapman & Hall Ltd (London), 433 p.
- Klister, P. (1954). Historical resume of the Amazon basin. Petrobras (104 -A), Bélem
- Le Bissonnais, Y. (1996) Aggregate stability and assessment of soil crustability and erodibility: I. Theory and methodology. *Eur. J. Soil Sci.*, 47, 425-437
- Mabesoone, J.M. & Alheiros, M.M. (1988). Origem da bacia sedimentar costeira de Pernambuco. *Rev. Bras. Geo.*, 18, 476-482
- Oliveira, J.B. (de), Jacomine, P.K.T. & Camargo, M.N. (1992). Classes gerais de solos do Brasil. Guia auxiliar para seu reconhecimento. FUNEP (Jaboticabal), 201 p.
- Silva, F.B.R. (e) & Araujo Filho, J.C. (de) (1989). Levatamento detalhado (1/10 000) dos solos da Usina Triunfo (Boca da Mata, Estado de Alagoas). Embrapa/CPATSA-CNPS (Recife)

- Silva, F.B.R. (e), Riche, G.R., Tonneau, J.P., Souza Neto, N.C., Brito, L.T., Correia, R.C., Cavalcanti, A.C., Silva, F.H.B.B., Silva, A.B. & Araujo Filho, J.C. (de) (1993). Zoneamento agroecológico do Nordeste: diagnostico do quadro natural e agro-socio-econômico. Embrapa/CPATSA-CNPS (Petrolina), 387 p.
- Silva, F.B.R. (e), Leprun J.C., Boulet R., 1997 - Duripãs em solos dos tabuleiros costeiros do Nordeste do Brasil: Mineralogia, micromorfologia e gênese. Cong. Bras. de Ciencia do Solo, 26, Rio de Janeiro, CD-ROM
- Soil Survey Staff (1975). Soil taxonomy. A basic system of soil classification for making and interpreting soil surveys. U.S. Gov. Printing Office, U.S.D.A. Handbook (Washington, D.C.), vol. 436, 754 p.

Key words: ferrallitic soils, duripan, microanalysis, iron, microaggregate, Brazil  
Mots clés: sol ferrallitique, duripan, microanalyse, fer, microagrégation, Brésil