

**Variation of the kaolinite and gibbsite content at regional and local scale in Latosols of the Brazilian Central Plateau Variabilité à l'échelle régionale et locale de la teneur en kaolinite et gibbsite des latosols du plateau central brésilien.**

Adriana Reatto-Braga, Ary Bruand, Éder de Souza Martins, Fabrice Muller, Euzebio Medrado da Silva, Osmar Abilio de Carvalho Jr, Michel Brossard

► **To cite this version:**

Adriana Reatto-Braga, Ary Bruand, Éder de Souza Martins, Fabrice Muller, Euzebio Medrado da Silva, et al.. Variation of the kaolinite and gibbsite content at regional and local scale in Latosols of the Brazilian Central Plateau Variabilité à l'échelle régionale et locale de la teneur en kaolinite et gibbsite des latosols du plateau central brésilien.. Comptes Rendus Géoscience, Elsevier Masson, 2008, 340 (11), pp.741-748. 10.1016/j.crte.2008.07.006 . insu-00326369

**HAL Id: insu-00326369**

**<https://hal-insu.archives-ouvertes.fr/insu-00326369>**

Submitted on 2 Oct 2008

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

C.R Geoscience

Surface Geosciences (Pedology)

**Variation of the kaolinite and gibbsite content at regional and local scale  
in Latosols of the Brazilian Central Plateau**

Adriana Reatto <sup>a, b</sup>, Ary Bruand <sup>b, \*</sup>, Eder S. Martins <sup>a</sup>, Fabrice Muller <sup>b</sup>, Euzebio M. Silva <sup>a</sup>,  
Osmar Abílio de Carvalho Jr <sup>c</sup>, Michel Brossard <sup>d</sup>

<sup>a</sup> Empresa Brasileira de Pesquisa Agropecuária (Embrapa Cerrados), BR 020, km 18, 73310-970, Planaltina, Distrito Federal, Brazil.

<sup>b</sup> Université d'Orléans, CNRS/INSU, Université de Tours, Institut des Sciences de la Terre d'Orléans (ISTO), 1A rue de la Férellerie, 45071 Orléans Cedex 2, France.

<sup>c</sup> Universidade de Brasília (UnB), Departamento de Geografia, 70910-000, Brasília, Brazil

<sup>d</sup> Institut de Recherche pour le Développement (IRD), Unité Valpédo, BP 64501, 34394 Montpellier Cedex 5, France;

\*Corresponding author [Ary.Bruand@univ-orleans.fr](mailto:Ary.Bruand@univ-orleans.fr) (A. Bruand)

Presented by Georges Pédro

**Abstract**

Mineralogy of the Latosols of the Brazilian Central Plateau remains under discussion in the absence of clear relationship with their age according to their geomorphic location. The aim of this study was thus to clarify the origin the kaolinite and gibbsite content variation by studying a regional toposequence and using data from the literature. Chemical composition and soil color were used to discuss mineralogy. The mineralogy of the clay fraction was also

discussed by using X-ray diagrams. Our results showed, that the large variation of kaolinite and gibbsite content can be explained by taken into account both their local and regional location, the variation of the hematite and goethite content remaining limited. The model that is proposed to explain such a variation combines a regional component which is mainly associated to the age of the geomorphic surface and a local component which is mainly associated to the hydraulic conditions along the toposequence.

**Keywords: Oxisol, Ferralsol, mineralogy, iron oxyhydroxide, soil color, biome Cerrado**

### **Résumé**

#### **Variabilité à l'échelle régionale et locale de la teneur en kaolinite et gibbsite des Latosols du Plateau Central Brésilien**

La minéralogie des Latosols du Plateau Central Brésilien reste discutée en l'absence de relation clairement établie avec leur âge qui est fonction de leur position géomorphologique. L'objectif de cette étude est par conséquent de clarifier l'origine de la variation de teneur en kaolinite et gibbsite en étudiant une toposéquence régionale et les données de la littérature. La minéralogie a été discutée à partir de la composition chimique et de la couleur du sol. Elle a aussi été discutée à l'aide des données de la diffraction des rayons X. Les résultats montrent que la variation élevée de la proportion de kaolinite et de gibbsite des Latosols peut être expliquée en prenant en compte à la fois leur localisation régionale et locale. Le modèle proposé combine en effet une composante régionale qui est principalement liée à l'âge de la surface géomorphologique et une composante locale qui est elle principalement liée aux conditions hydriques le long de la toposéquence.

**Mots-clés : Oxisol, Ferralsol, oxy-hydroxyde de fer, couleur du sol, biome Cerrado**

## 1. Introduction

The Latosols of the Brazilian Soil Taxonomy [8], which are Oxisols in the Soil Taxonomy [33] and Ferralsols in the World Reference Base [12] cover approximately 40% of the Brazilian Central Plateau [24]. This region that corresponds to 24% of Brazilian territory is composed of two main geomorphic surfaces developed during the Cretaceous Superior and Tertiary: (i) the South American Surface (SAS) which is the oldest and mainly made up of tablelands called chapadas with smoothly convex plane portions with an elevation ranging from 900 to 1,200 m, (ii) and the Velhas Surface (VS) characterized by moderate and convex slopes at an elevation below 900 m [23].

In the Central Plateau, the Latosols are Red Latosols (~28%) where the hematite is the main iron oxyhydroxide, Yellow Red Latosols (~10%) where hematite and goethite are present in similar proportion and Yellow Latosols (~2%) where goethite is the main iron oxyhydroxide. Besides iron oxyhydroxides, gibbsite and kaolinite were shown to be the main associated minerals in Latosols of the SAS and VS, respectively [39]. However, several studies showed high proportions of kaolinite in Latosols of the SAS and high proportions of gibbsite in Latosols of the VS. Indeed, Resende [27] studied a topolithosequence 67 km-long across the SAS and VS and showed high proportion of kaolinite in Red Latosols and Yellow Red Latosols developed in clay sediments on the SAS. Curi and Franzmeier [6] studied a toposequence 200 m-long on the VS with Latosols developed in weathered basalts and found Red Latosols upslope with a high proportion of gibbsite. Macedo and Bryant [14] studied a hydrosequence 3 km-long on the SAS and found Yellow Red Latosols downslope with similar proportion of kaolinite and gibbsite. Several authors [10, 11, 15, 16, 20] studied Latosols located on the two geomorphic surfaces and recorded a variable proportion of gibbsite and kaolinite for Latosols developed on the same surface. Thus, the mineralogy of the Latosols of the Brazilian Central Plateau remains under discussion because it appears weakly related to their age according to their location on the two main geomorphic surfaces. In this

context, the aim of this study is (i) to analyze the mineralogy of these Latosols by studying Latosols along a regional toposequence and using data from the literature, (ii) and to show that a model consistent with our data and those from the literature can be proposed.

## 2. Material and Methods

Ten Latosols (L) developed in different parent materials were selected for study along a 350 km-long toposequence across the SAS (L1 to L4) and VS (L5 to L10). Location and basic properties of these Latosols can be found in [26] and Table 1. The Latosols L5 and L6 were located on the upper VS, L7 and L8 on the intermediate VS, and L9 and L10 on the lower VS. The Latosols L7 and L8 are those also studied by Volland *et al.* [37, 38] and similar to those studied by Balbino *et al.* [1, 2, 3]. A set of 25 samples was collected in the diagnostic horizons Bw<sub>1</sub>, Bw<sub>2</sub> and when possible Bw<sub>3</sub> of the Latosols selected. The SiO<sub>2</sub>, Al<sub>2</sub>O<sub>3</sub>, and Fe<sub>2</sub>O<sub>3</sub> contents were determined on the <2-mm material after dissolution in 1:1 H<sub>2</sub>SO<sub>4</sub> [5, 7, 15, 30, 35]. This acid attack enables dissolution of the clays, Fe oxyhydroxides and Al hydroxides [22, 28, 31].

The SiO<sub>2</sub> and Al<sub>2</sub>O<sub>3</sub> extracted with sulfuric acid were used to compute the kaolinite (*K*) and gibbsite (*Gb*) content as following [4, 28]:

$$K = S_{SiO_2} / K_{SiO_2} \quad (1)$$

where *K* is the kaolinite content (%) of the sample, *S*<sub>SiO<sub>2</sub></sub> the SiO<sub>2</sub> content of the sample recorded with sulphuric acid extraction (%), *K*<sub>SiO<sub>2</sub></sub> the specific proportion of SiO<sub>2</sub> of the kaolinite and equaled to 0.465.

The goethite (*Gt*) and hematite (*Hm*) contents were computed by combining two equations relating *Gt* and *Hm* as following:

$$S_{Fe_2O_3} = Gt_{Fe_2O_3} \times Gt + Hm_{Fe_2O_3} \times Hm \quad (2)$$

$$Hm / (Hm + Gt) = (RI - 3.50) / 8.33 \quad (3)$$

where  $S_{Fe_2O_3}$  is the  $Fe_2O_3$  content (%) of the sample recorded with sulphuric acid extraction,  $Gt_{Fe_2O_3}$  is the specific proportion of  $Fe_2O_3$  in the goethite and equaled to 0.899 for a non Al-substituted goethite and to 0.675 for a 33% Al-substituted goethite [32],  $Hm_{Fe_2O_3}$  is the specific proportion of  $Fe_2O_3$  in the hematite and equaled to 1 for a non Al-substituted hematite and to 0.890 for a 16% Al-substituted hematite [32],  $RI$  is the red index [9, 21, 30] and equaled to:

$$RI = (M + C/V) \quad (4)$$

with  $M$  a parameter related to the hue ( $M$  was 10 for 10R, 7.5 for 2.5YR, 5 for 5YR, 2.5 for 7.5YR and 0 for 10YR),  $C$  the chroma and  $V$  the value of the Munsell notation [9, 21, 30].

The gibbsite content of the sample was computed as following:

$$Gb = \left\{ \left[ S_{Al_2O_3} - (Gt \times Gt_{Al_2O_3}) - (Hm \times Hm_{Al_2O_3}) - (K \times K_{Al_2O_3}) \right] / Gb_{Al_2O_3} \right\} \quad (5)$$

where  $Gb$  is the gibbsite content (%) of the sample,  $S_{Al_2O_3}$  the  $Al_2O_3$  content of the sample recorded with sulphuric acid extraction (%),  $K_{Al_2O_3}$  the specific proportion of  $Al_2O_3$  of the kaolinite and equaled to 0.395,  $Gb_{Al_2O_3}$  the specific proportion of  $Al_2O_3$  of the gibbsite and equaled to 0.654. Equation (1) and (5) assumed kaolinite and gibbsite without any substitution.

The mineralogy of the  $< 2 \mu m$  fraction of the  $Bw_2$  horizons was determined by using X-ray diffraction on powder samples by using a Thermo Electron ARL XTRA diffractometer [29]. The  $SiO_2$ ,  $Al_2O_3$ , and  $Fe_2O_3$  contents of the  $< 2 mm$  material of 162  $Bw$  horizons collected in Latosols of the Central Plateau and earlier published [25] were also used to discuss the mineralogy of Latosols.

### 3. Results and Discussion

#### 3.1. Composition and mineralogy of the Latosols along the regional sequence studied

In the Bw horizons studied, the Fe<sub>2</sub>O<sub>3</sub> content ranged from 15 to 33%, the Al<sub>2</sub>O<sub>3</sub> content from 43 to 68% and the SiO<sub>2</sub> content from 11 to 36% (Fig. 1a). For those belonging to Latosols developed on the SAS, the Fe<sub>2</sub>O<sub>3</sub> content ranged from 15 to 33%, the Al<sub>2</sub>O<sub>3</sub> content from 54 to 68% and the SiO<sub>2</sub> content from 11 to 24%. On the other hand, for those belonging to Latosols developed on the VS, the Fe<sub>2</sub>O<sub>3</sub> content ranged from 18 to 24%, the Al<sub>2</sub>O<sub>3</sub> content from 43 to 52% and the SiO<sub>2</sub> content from 22 to 36%, (Fig. 1a). The range of Fe<sub>2</sub>O<sub>3</sub> content is consistent with the one recorded by Melfi and Pédro [19] for Latosols of the Central Plateau.

The results showed a relatively small variation of the iron oxyhydroxide content between the Latosols studied whatever the Al-substitution rate since *Gt+Hm* ranged from 13 to 27% in the absence of Al-substitution and from 15 to 29% when the goethite and hematite were 33% and 16% Al-substituted, respectively (Fig. 2 a and b). On the other hand, there was a large variation of the kaolinite and gibbsite content with *K* ranging from 17 to 67% and *Gb* from 15 to 65% with non Al-substituted goethite and hematite and *K* ranging from 18 to 69% and *Gb* from 13 to 62% when the goethite and hematite were 33% and 16% Al-substituted, respectively (Fig. 2 a and b). Thus, the Latosols sampled along the regional toposequence studied were gibbsitic Latosols on the SAS (L1 to L4) and kaolinitic Latosols on the VS (L5 to L10) (Fig. 2a). The mineralogy obtained with data from sulfuric acid extraction was consistent with the X-ray diagrams recorded for the < 2 μm fraction of the Bw<sub>2</sub> horizons studied (Fig. 3). X-ray diagrams showed also a greater kaolinite content in the L3 than in L10 and a close gibbsite content between the latter, thus indicating again no sharp variation of mineralogy between the Latosols developed on the SAS and VS (Fig. 3).

### *3.2. Mineralogy of Latosols located in the Brazilian Central Plateau*

Results from sulphuric extractions earlier published [25] were used to describe the mineralogy of <2-mm material of Latosols as done above for the Latosols of the regional

toposequence studied. The  $\text{Fe}_2\text{O}_3$  contents ranged from 9 to 34%, the  $\text{Al}_2\text{O}_3$  content from 36 to 78% and the  $\text{SiO}_2$  content 9 to 42% (Fig. 1b). For the Bw horizons of Latosols developed on the SAS, the  $\text{Fe}_2\text{O}_3$  content ranged from 9 to 34%, the  $\text{Al}_2\text{O}_3$  content from 39 to 78% and the  $\text{SiO}_2$  content from 9 to 39%. On the other hand, for the Bw horizons of Latosols developed on the VS, the  $\text{Fe}_2\text{O}_3$  content ranged from 18 to 33%, the  $\text{Al}_2\text{O}_3$  content from 36 to 60% and the  $\text{SiO}_2$  content from 13 to 42%, (Fig. 1b).

The  $\text{Fe}_2\text{O}_3$ ,  $\text{Al}_2\text{O}_3$  and  $\text{SiO}_2$  contents were used to compute  $K$ ,  $Gb$  and  $Gt+Hm$  as done for the Bw horizons of the regional toposequence studied. In the absence of Al-substitution in goethite and hematite, results showed that  $K$  and  $Gb$  ranged from 11 to 78% and from 1 to 77%, respectively (Fig. 2c). On the other hand, with 33% Al-substituted goethite and 16% Al-substituted hematite, results showed that  $K$  and  $Gb$  ranged from 12 to 79% and from 0 to 75%, respectively (Fig. 2c). Results showed also a large overlapping of the mineralogical composition range between Latosols developed on the SAS and those developed on the VS (Fig. 2c). Indeed, for the Bw horizons of Latosols developed on the SAS,  $K$  ranged from 11 to 75% and  $Gb$  from 3 to 77% with non Al-substituted goethite and hematite, and  $K$  ranged from 12 to 78% and  $Gb$  ranged from 0 to 75% with 33% Al-substituted goethite and 16% Al-substituted hematite. On the other hand, for the Bw horizons of Latosols developed on the VS,  $K$  ranged from 21 to 78% and  $Gb$  from 1 to 57% with non Al-substituted goethite and hematite, and  $K$  ranged from 22 to 79% and  $Gb$  ranged from 0 to 55% with 33% Al-substituted goethite and 16% Al-substituted hematite. Results showed also that  $Gt+Hm$  from 9 to 31% in the absence of Al-substitution and from 9 to 35% when the goethite and hematite were 33% and 16% Al-substituted respectively (Fig. 2 c and d) without any relationship with the location of Latosols on the two main geomorphic surfaces.

### 3.3. Variation of the kaolinite and gibbsite content at the regional and local scale



Macedo and Bryant [14] and Motta *et al.* [20] showed that the Latosols distribution on the SAS was closely related to the soil hydraulic regime thus explaining the Red Latosol, Yellow Red Latosols and Yellow Latosol sequence according to local variation of the topography. As a consequence, the Latosols distribution appeared roughly independent of the underlying geological material [20]. Motta *et al.* [20] suggested that more attention should be devoted to geomorphology to explain variation of Latosols characteristics and particularly their mineralogy. Melfi and Pédro [17, 18] showed that Latosols mineralogy should be related to their geochemical functioning that is characterized by an hydrolytic environment according to landscape history at both regional and geological scale. Tardy [34] discussed the kaolinite/gibbsite ratio in tropical soils and showed that the kaolinite–gibbsite equilibrium would be preferentially controlled by variation of the hydraulic conditions along of the toposequences. Finally, Lucas [13] showed that the spatial distribution in equatorial areas of the secondary minerals such as kaolinite, gibbsite and goethite can be related to their stability in aqueous solutions and then to the amount of the water percolating the soils. Thus, as discussed by Lucas [13], the higher the volume of water percolating the profile is, the lower the soil-solution concentrations are.

On the basis of these results, we plotted the altitude at which every Latosols was located on the SAS and VS according to the  $Gb/(Gb+K)$  ratio. Figure 4 shows that  $Gb/(Gb+K)$  varies according to the local topographic location of every Latosol (Axe 1) and to the regional topographic location of every Latosols (Axe 2). Locally, Latosols located on the slope showed higher  $Gb/(Gb+K)$  ratio than those located on the plateau of the same portion of landscape (Fig. 4). At the regional scale, our results showed the  $Gb/(Gb+K)$  ratio increased with the altitude thus explaining the trend to an increase in the  $Gb/(Gb+K)$  ratio value with the altitude, the age of the surface increasing itself with the altitude. Thus, the Axe 2 shows a regional variability that is mainly related to time. More the topographic surface is old, more the

Latosols are old, higher is the weathering and consequently the hydrolysis processes intensity resulting in a higher gibbsite content in the Bw studied as discussed by Vitte [36] and Melfi and Pédro [17, 18]. On the other hand, the Axe 1 shows a local variability that would be mainly related to the volume of water percolating the soil. Indeed, because of local topographic characteristics, water can percolate more or less easily, maintaining the Fe, Si and Al concentrations that result from mineral hydrolysis at values that are more or less favorable to hydrolysis process continuation. Thus according to the local topographic location, the higher the volume of water percolating the Latosol is, the higher hydrolysis process is, and the higher resulting gibbsite content is.

#### **4. Conclusion**

Our results showed that the kaolinite and gibbsite content in the Latosols developed on the South American Surface and Velhas Surface of the Brazilian Central Plateau can be explained by taken into account both their local and regional location. The model proposed combines (i) a regional variation which would be mainly associated to the age of the surface, the more the surface being old, the more SiO<sub>2</sub> removal from the soil being developed and thus the gibbsite content being high compared to the kaolinite content, (ii) and a local variation which would be mainly associated the hydraulic conditions along the toposequence at the scale of several hectometers or a few kilometers, the gibbsite content being the highest where SiO<sub>2</sub> removal is the easiest as upslope and on the plateau border.

#### **Acknowledgements**

We thank the Empresa Brasileira de Pesquisa Agropecuária (EMBRAPA) for its financial support of A. Reatto's work in France. This research is part of the project Embrapa Cerrados -

IRD, n°0203205 (Mapping of the Biome Cerrado Landscape and Functioning of Representative Soils).

## References

- [1] L. C. Balbino, A. Bruand, M. Brossard, M. F. Guimarães, Comportement de la phase argileuse lors de la dessiccation dans des Ferralsols microagrégés du Brésil : rôle de la microstructure et de la matière organique, *Comptes Rendus de l'Académie des Sciences* 332 (2001) 673-680.
- [2] L. C. Balbino, A. Bruand, M. Brossard, M. Grimaldi, M. Hajnos, M. F. Guimarães, Changes in porosity and microaggregation in clayey Ferralsols of the Brazilian Cerrado on clearing for pasture, *European Journal of Soil Science* 53 (2002) 219–230.
- [3] L. C. Balbino, A. Bruand, I. Cousin, M. Brossard, P. Quéting, M. Grimaldi, Change in the hydraulic properties of a Brazilian clay Ferralsol on clearing for pasture, *Geoderma* 120 (2004) 297–307.
- [4] A. Bruand, R. Prost, Analyse minéralogique quantitative d'un échantillon de sol: utilisation des données concernant la composition chimique de l'échantillon, *Agronomie* 8(1) (1988), 15-22.
- [5] M. N. Camargo, J. M. Kimble, F. H. Beinroth, Classification, Characterization and Utilization of Oxisols, Part 2: Field Trip Background, Site and Pedon Descriptions, Analytical Data, in: *Proceedings of the Eight International Soil Classification Workshop, Brazil*, SNLCS-Embrapa, Soil Management Support Services and Soil Conservation Service–United States Department of Agriculture, University of Puerto Rico, 1986.
- [6] N. Curi, D. Franzmeier, Toposequence of Oxisols from Central Plateau of Brazil, *Soil Science Society of America Journal* 48 (1984) 341–346.

- [7] Embrapa, Manual de métodos de análise de solo. 2ed. rev. atual. Empresa Brasileira de Pesquisa Agropecuária, Rio de Janeiro, RJ, 1997.
- [8] Embrapa, Sistema Brasileiro de Classificação de Solos, Empresa Brasileira de Pesquisa Agropecuária, Rio de Janeiro, RJ, 1999.
- [9] M. A. Figueiredo, J. D. Fabris, A. F. D. C. Varajão, P.R. C. Couceiro, I. S. Loutfli, I. S. Azevedo, V. K. Garg, Óxidos de ferro de solos formados sobre gnaiss do Complexo Bação, Quadrilátero Ferrífero, Minas Gerais, Pesquisa Agropecuária Brasileira 41(2) (2006) 313-321.
- [10] J.B.V. Gomes, N. Curi, D.G. Schulze, J.J.G.S.M. Marques, J.C. Ker, P.E.F. Motta, Mineralogia, morfologia e análise microscópica de solos do Bioma Cerrado, Revista Brasileira de Ciência do Solo 28 (2004a) 679–694.
- [11] J.B.V. Gomes, N. Curi, P.E.F. Motta, J.C. Ker, J.J.G.S.M. Marques, D.G. Schulze, Análise de componentes principais de atributos físicos, químicos e mineralógicos de solos do Bioma Cerrado, Revista Brasileira de Ciência do Solo 28 (2004b) 137–153.
- [12] IUSS Working Group WRB, World reference base for soil resources 2006, World Soil Resources Reports N° 103. FAO, Rome, 2006.
- [13] Y. Lucas, D. Nahon, S. Cornu, F. Eyrolle, Genèse et fonctionnement des sols en milieu équatorial, in: F. Fabre (Ed.), Géochimie de la surface, pédologie, hydrologie, Le point sur..., Sciences de la Terre VII, Comptes Rendus de l'Académie des Sciences, Elsevier, Paris, 2001, pp. 1-16.
- [14] J. Macedo, R. B. Bryant, Morphology, mineralogy, and genesis of a hydrosequence of Oxisols in Brazil, Soil Science Society of America Journal 51 (1987) 690-698.
- [15] J.J.G.S.M. Marques, Trace element distributions in Brazilian Cerrado soils at the landscape and micrometer scales, Ph. D. Thesis, Purdue University, USA, 2000 (available at <http://www.dcs.ufla.br/marques>).

- [16] J.J. Marques, D. G. Schulze, N. Curi, S. A. Mertzman, Major element geochemistry and geomorphic relationships in Brazilian Cerrado soils, *Geoderma* 119 (2004) 179 – 195.
- [17] A. J. Melfi, G. Pédro, Estudo geoquímico dos solos e formações superficiais do Brasil. Parte 1 – Caracterização e repartição dos principais tipos de evolução pedogeoquímica, *Revista Brasileira de Geociências* 7 (1977) 271 – 286.
- [18] A. J. Melfi, G. Pédro, Estudo geoquímico dos solos e formações superficiais do Brasil. Parte 2 – Considerações sobre os mecanismos geoquímicos envolvidos na alteração superficial e sua repartição no Brasil, *Revista Brasileira de Geociências* 8 (1978) 11 – 22.
- [19] A. J. Melfi, G. Pédro, B. Volkoff, Natureza e distribuição dos compostos ferríferos nos solos do Brasil, *R. Bras. Ci. Solo* 3 (1979) 47 – 54.
- [20] P. E. F. Motta, A. Carvalho Filho, J. C. Ker, N. R. Pereira, W. Carvalho Junior, P. Blancaneaux, Relações solo-superfície geomórfica e evolução da paisagem em uma área do Planalto Central Brasileiro. *Pesquisa Agropecuária Brasileira* 37 (2002) 869-878.
- [21] N. Kampf, E. Kamt, P. Schneider, Óxidos de ferro em Latossolos do Brasil Sudeste e Sul, in: *Anais da III Reunião de Classificação, Correlação de Solos e Interpretação de Aptidão Agrícola*, Embrapa-SNLCS, Rio de Janeiro-RJ, Série Documentos, 12 (1988) 153-183.
- [22] J. C. Ker, Latossolos do Brasil: uma revisão, *Geonomos* 5 (1998) 17-40.
- [23] Radambrasil, Levantamentos de recursos naturais, Folha SD. 23. Brasília, Ministério de Minas e Energia - Secretaria Geral 19, 1984.
- [24] A. Reatto, J.R. Correia, S.T. Spera, Solos do Bioma Cerrado: aspectos pedológicos, in: S.M. Sano, S.P. Almeida, (Eds), *Cerrado: ambiente e flora*, Empresa Brasileira de Pesquisa Agropecuária, Planaltina-DF, 1998, pp. 47 - 88.
- [25] A. Reatto, E.S. Martins, E.M. Guimarães, S.T. Spera, J.R. Correia, Variabilidade mineralógica de latossolos da área da Embrapa Cerrados em relação aos do Bioma Cerrado, *Boletim de Pesquisa Embrapa Cerrados* 20 (2000) 1-29.

- [26] A. Reatto, A. Bruand, E. M. Silva, E. S. Martins, M. Brossard, Hydraulic properties of the diagnostic horizon of Latosols of a regional toposequence across the Brazilian Central Plateau, *Geoderma* 139 (2007) 51-59.
- [27] M. Resende, Mineralogy, chemistry, morphology and geomorphology of some soils of the Central Plateau of Brazil, Ph. D. Thesis, Purdue University, USA, 1976.
- [28] M. Resende, A. F. C. Bahia Filho, J. M. Braga, Mineralogia da argila de latossolos estimada por alocação a partir do teor total de óxidos do ataque sulfúrico, *Revista Brasileira de Ciência do Solo* 11 (1987) 17–23.
- [29] M. Robert, D. Tessier, Méthode de préparation des argiles des sols pour des études minéralogiques, *Ann. Agron.* 25 (6) (1974) 859–882.
- [30] D. P. Santana, Soil formation in a toposequence of Oxisols from Patos de Minas region, Minas Gerais, Brazil. Ph. D. Thesis, Purdue University, USA, 1984.
- [31] C. E. G. R. Schaefer, J. D. Fabris, J. C. Ker, Minerals in the clay fraction of Brazilian (Oxisols): a review, *Clay Minerals* 43 (2008) 1-18.
- [32] U. Schwertmann, R. M. Taylor, Iron Oxides, in: J.B. Dixon, S. B. Weed (Eds.), *Minerals in Soil Environments*, Soil Science Society of America, Madison, Wisconsin, USA, 1989, pp. 379-438.
- [33] Soil Survey Staff, *Keys to Soil Taxonomy*, 10th ed. USDA-Natural Resources Conservation Service, Washington, DC, 2006.
- [34] Y. Tardy, *Pétrologie des Latérites et des Sols Tropicaux*, Masson, Paris, 1993
- [35] L. Vettori, As relações Ki e Kr na fração argila e na terra fina, in: *Anais do VII Congresso Brasileiro de Ciência do Solo*, Piracicaba-SP, SBCS, 1959.
- [36] A. C. Vitte, Considerações sobre a teoria da etchplanação e sua aplicação nos estudos das formas de relevo nas regiões tropicais quentes e úmidas, *Terra Livre* 16 (2001) 11 – 24.

- [37] N. Volland-Tuduri, M. Brossard, A. Bruand, H. Garreau, Direct analysis of microaggregates shrinkage for drying: Application to microaggregates from a Brazilian clayed Ferralsol, *Comptes Rendus de l'Académie des Sciences* 336 (2004) 1017–1024.
- [38] N. Volland-Tuduri, A. Bruand, M. Brossard, L. C. Balbino, M. I. L. Oliveira, E. S. Martins, Mass proportion of microaggregates and bulky density in a Brazilian clayed Oxisol, *Soil Science Society American Journal* 69 (2005) 1559–1564.
- [39] R. M. Weaver, *Soils of the Central Plateau of Brazil: chemical and mineralogical properties*, Agronomy Mimeo 74-8, Department of Agronomy, Cornell University, Ithaca, New York, 1974.

## Figures

Fig. 1. SiO<sub>2</sub>, Al<sub>2</sub>O<sub>3</sub> and Fe<sub>2</sub>O<sub>3</sub> relative contents in the Bw horizons of the Latosols of the regional toposéquence studied (a) and those of Bw horizons from the literature (b): Latosols located on the South American Surface (+) and Velhas Surface (▲).

Fig. 1. Contenu relatifs en SiO<sub>2</sub>, Al<sub>2</sub>O<sub>3</sub> et Fe<sub>2</sub>O<sub>3</sub> dans les horizons Bw des Latosols de la toposéquence régionale étudiée (a) et ceux des horizons Bw issus de la littérature (b) : Latosols situés sur la Surface Sud Américaine (+) et sur la Surface Velhas (▲).

Fig. 2. Kaolinite, gibbsite, and (goethite + hematite) relative contents in the Bw horizons of the Latosols of the regional toposéquence studied (a, with non Al-substituted goethite and hematite and b, with 33% Al-substituted-goethite and 16% Al-substituted hematite) and in Bw horizons from the literature (c, with non Al-substituted goethite and hematite and d, with 33% Al-substituted-goethite and 16% Al-substituted hematite): Latosols located on the South American Surface (+) and Velhas Surface (▲).

Fig. 2. Contenus relatifs en kaolinite, gibbsite, and (goethite + hematite) dans les horizons Bw des Latosols de la toposéquence régionale étudiée (a, calculé avec une goethite et une hématite sans substitution par Al et b, calculé avec une goethite substituée par Al à 33% et une hématite substituée par Al à 16%) et dans les horizons Bw issus de la littérature (c, calculé avec une goethite et une hématite sans substitution par Al et d, calcul avec une goethite substituée par Al à 33% et une hématite substituée par Al à 16%) : Latosols situés sur la Surface Sud Américaine (+) et ceux situés sur la Surface Velhas (▲).

Fig. 3. X-ray diagrams of the oriented <2 μm fraction (powder) of horizons Bw of the Latosols studied.

Fig. 3. Diagrammes de rayons-X de la fraction <2 μm (poudre) des horizons Bw des Latosols étudiés.



Fig. 4. Altitude of every Latosol (L) according to the gibbsite/(gibbsite + kaolinite) ratio computed with non Al-substituted goethite and hematite (◆) and both 33% Al-substituted goethite and 16% Al-substituted hematite (▲). Every Latosols was also located on its portion of landscape according to the local topography (Axe 1: local variation associated to the hydraulic condition along the toposequence and Axe 2: regional variability according to the age of the surface). SAS: South American Surface, VS: Velhas Surface (VS – I: Upper level, VS – II: Intermediate level, VS – III: Lower level).

Fig. 4. Altitude de chaque Latosol (L) en fonction du rapport gibbsite/(gibbsite + kaolinite) calculé avec une goethite et une hémate non substituée par Al (◆) et avec à la fois une goethite substituée par Al à 33% et une hémate substitué par Al à 16% (▲). Chaque Latosol est localisé sur sa portion de paysage (Axe 1: variabilité locale liée aux conditions hydriques le long de la toposéquence et Axe 2: variabilité régionale en fonction de l'âge de la surfaces. SAS: Surface Sud Américaine, VS: Surface Velhas (VS - I: Niveau Supérieur, VS - II: Niveau Intermédiaire, VS – III: Niveau Inférieur).

## Tables

Table 1 – General characteristics of the Latosols studied.

Tableau 1 - Caractéristiques générales des Latosols étudiés.






Fig. 1. SiO<sub>2</sub>, Al<sub>2</sub>O<sub>3</sub> and Fe<sub>2</sub>O<sub>3</sub> relative contents in the Bw horizons of the Latosols of the regional toposéquence studied (a) and those of Bw horizons from the literature (b): Latosols located on the South American Surface (+) and Velhas Surface (▲).

Fig. 1. Contenus relatifs en SiO<sub>2</sub>, Al<sub>2</sub>O<sub>3</sub> et Fe<sub>2</sub>O<sub>3</sub> dans les horizons Bw des Latosols de la toposéquence régionale étudiée (a) et ceux des horizons Bw issus de la littérature (b) : Latosols situés sur la Surface Sud Américaine (+) et sur la Surface Velhas (▲).




Fig. 2. Kaolinite, gibbsite, and (goethite + hematite) relative contents in the Bw horizons of the Latosols of the regional toposequence studied (a, with non Al-substituted goethite and hematite and b, with 33% Al-substituted-goethite and 16% Al-substituted hematite) and in Bw horizons from the literature (c, with non Al-substituted goethite and hematite and d, with 33% Al-substituted-goethite and 16% Al-substituted hematite): Latosols located on the South American Surface (+) and Velhas Surface (▲).

Fig. 2. Contenus relatifs en kaolinite, gibbsite, and (goethite + hematite) dans les horizons Bw des Latosols de la toposequence regionale etudiee (a, calcule avec une goethite et une hematite sans

substitution par Al et b, calculé avec une goethite substituée par Al à 33% et une hématite substituée par Al à 16%) et dans les horizons Bw issus de la littérature (c, calculé avec une goethite et une hématite sans substitution par Al et d, calcul avec une goethite substituée par Al à 33% et une hématite substituée par Al à 16%) : Latosols situés sur la Surface Sud Américaine (+) et ceux situés sur la Surface Velhas (▲).




Fig. 3. X-ray diagrams of the oriented  $<2 \mu\text{m}$  fraction (powder) of horizons Bw of the Latosols studied.

Fig. 3. Diagrammes de rayons-X de la fraction  $<2 \mu\text{m}$  (poudre) des horizons Bw des Latosols étudiés.




Fig. 4. Altitude of every Latosol (L) according to the gibbsite/(gibbsite + kaolinite) ratio computed with non Al-substituted goethite and hematite (◆) and both 33% Al-substituted goethite and 16% Al-substituted hematite (▲). Every Latosols was also located on its portion of landscape according to the local topography (Axe 1: local variation associated to the hydraulic condition along the toposequence and Axe 2: regional variability according to the age of the surface). SAS: South American Surface, VS: Velhas Surface (VS – I: Upper level, VS – II: Intermediate level, VS – III: Lower level).

Fig. 4. Altitude de chaque Latosol (L) en fonction du rapport gibbsite/(gibbsite + kaolinite) calculé avec une goethite et une hématite non substituée par Al (◆) et avec à la fois une goethite substituée par Al à 33% et une hématite substitué par Al à 16% (▲). Chaque Latosol est localisé sur sa portion de paysage (Axe 1: variabilité locale liée aux conditions hydriques le long de la toposéquence et Axe 2: variabilité régionale en fonction de l'âge de la surfaces. SAS: Surface Sud Américaine, VS: Surface Velhas (VS - I: Niveau Supérieur, VS - II: Niveau Intermédiaire, VS – III: Niveau Inférieur).

Table 1 – General characteristics of the Latosols studied.

Tableau 1 - Caractéristiques générales des Latosols étudiés.

Latosols	Geomorphic Surface	Altitude (m)	Position along the toposequence	Slope length (km)	Declivity (%)
L1	South American	1050	median	3	< 1
L2	South American	1200	median	5	2
L3	South American	1190	median	5	2
L4	South American	1180	down	12	3
L5	Velhas, Superior Level	920	median-up	12	< 1
L6	Velhas, Superior Level	880	down	20	6
L7	Velhas, Intermediate Level	820	median-up	20	2
L8	Velhas, Intermediate Level	805	median-down	7	2
L9	Velhas, Inferior Level	785	median-up	15	< 1
L10	Velhas, Inferior Level	760	down	15	7