



## Observing and modelling Mars' airglow

Cyril Simon, Olivier Witasse, François Leblanc, Guillaume Gronoff, Jean Lilensten, Jean-Loup Bertaux

### ► To cite this version:

Cyril Simon, Olivier Witasse, François Leblanc, Guillaume Gronoff, Jean Lilensten, et al.. Observing and modelling Mars' airglow. European Planetary Science Congress 2008, Sep 2008, Münster, Germany. pp.EPSC2008-A-00111. insu-00366192

HAL Id: insu-00366192

<https://insu.hal.science/insu-00366192>

Submitted on 14 Jan 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Observing and modelling Mars' airglow

C. Simon (1), O. Witasse (1), F. Leblanc (2), G. Gronoff (3), J. Lilensten (3), J.-L. Bertaux (4)


(1) Research Scientific Support Department, ESA-ESTEC, Noordwijk, The Netherlands  
[cyril.simon@rssd.esa.int](mailto:cyril.simon@rssd.esa.int), (2) Service d'Aéronomie, Université Paris 6/CNRS, Institut Pierre Simon Laplace, Verrières-le-Buisson, France, (3) Laboratoire de Planétologie de Grenoble, Université Joseph Fourier/CNRS, Grenoble, France, (4) Service d'Aéronomie, Université de Versailles-Saint-Quentin/CNRS, Institut Pierre Simon Laplace, Verrières-le-Buisson, France

### Abstract

SPICAM is the UV spectrometer onboard ESA's Mars Express, which has been recording for more than four years upper atmospheric emissions ([1], [2]). 70000 dayglow limb spectra were analysed using an updated and variable point spread function, yielding altitude profiles of the Cameron bands ( $a^3\Pi - X^1\Sigma$ ) of CO, the  $\text{CO}_2^+$  ultraviolet doublet ( $B^2\Sigma_u^+ - X^2\Pi_g$ ) at 289 nm and the oxygen emission at 297.2 nm. The seasonal evolution of these emissions (Fig. 1) exhibits an unexpected increase around  $L_s = 140$  degrees, which is correlated with a sudden increase of neutral densities due to a dust storm event. A comparison with a newly-developed kinetic electron and airglow model ([3], [4]) yields promising results on emission altitude profiles. Improvements on the neutral atmosphere and cross sections are discussed. The new data analysis performed enables the search for faint emissions underlying the main Cameron emission bands. This comparison and analysis work is being extended to nightside auroral emissions using multi-instrument data on Mars Express.

### References

- [1] Bertaux, J.-L., Koralev, O., Perrier, S., Quémerais, E. and the SPICAM team (2006) *JGR*, **111**, E10S90, doi:10.1029/2006JE002690.
- [2] Leblanc, F., Chaufray, J.-Y., Lilensten, J., Witasse, O., Bertaux, J.-L., (2006) *JGR*, **111**, E09S11, doi:10.1029/2005JE002664.
- [3] Simon, C., Witasse, O., Leblanc, F., Gronoff, G., Bertaux, J.-L., (2008) *submitted to PSS*
- [4] Gronoff, G., Lilensten, J., Simon, C., Barthélémy, M., Leblanc, F., Dutuit, O., (2008) *A&A*, **482**, 1015-1029.



**Fig. 1** Seasonal evolution of the total emission intensities (in kR) of the Cameron bands, the  $\text{CO}_2^+$  ultraviolet doublet and OI(2972Å).  $L_s$  is the Solar Longitude expressed in degrees. Two peaks are present, one near perihelion ( $L_s \sim 290^\circ$ ), the other during northern spring ( $L_s \sim 140^\circ$ ) which is correlated with an unexpected dust storm event.