



**HAL**  
open science

## Molecular signature of Late Bronze Age archaeological layers preserved in littoral settings in Lake le Bourget (French Alps).

Jérémy Jacob, Jean-Robert Disnar, Yves Billaud, Fabien Arnaud, Claude Le Milbeau

► **To cite this version:**

Jérémy Jacob, Jean-Robert Disnar, Yves Billaud, Fabien Arnaud, Claude Le Milbeau. Molecular signature of Late Bronze Age archaeological layers preserved in littoral settings in Lake le Bourget (French Alps).. International Meeting of Organic Geochemistry - IMOG, Sep 2009, France. 1 p. insu-00442920

**HAL Id: insu-00442920**

**<https://insu.hal.science/insu-00442920>**

Submitted on 23 Dec 2009

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Molecular signature of Late Bronze Age archaeological layers preserved in littoral settings in Lake le Bourget (French Alps).

Jérémy Jacob<sup>a,\*</sup>, Jean-Robert Disnar<sup>a</sup>, Yves Billaud<sup>b</sup>, Fabien Arnaud<sup>c</sup>, Claude LeMilbeau<sup>a</sup>

<sup>a</sup> Institut des Sciences de la Terre d'Orléans. Université d'Orléans/Université François Rabelais de Tours, UMR 6113 du CNRS/INSU. 1A rue de la Férollerie, 45071 Orléans, France. (\* corresponding author: [Jeremy.jacob@univ-orleans.fr](mailto:Jeremy.jacob@univ-orleans.fr))

<sup>b</sup> Ministère de la Culture, DRASSM 58 bis rue des Marquisats 74000 Annecy, France.

<sup>c</sup> EDYTEM, UMR 5204 CNRS/Université de Savoie, Campus Savoie Technolac, F-73376, le Bourget du Lac, France.

The Final Bronze Age was a key period for Human history in the Alps since numerous human occupations developed on lake shores attest to an increased anthropication of alpine ecosystems at that time [1]. Successions of abandonment and occupations were most probably climatically-driven through lacustrine level variations [2]. Our recent results on a sediment core drilled in Lake le Bourget provide a detailed record of the evolution of past human activities for this period by using a molecular biomarker specific of millet cultivation [3]. These original results motivated the research of new molecular biomarkers of human activities.

This study focuses on biomarker imprints of archaeological levels that correspond to former settlement layers preserved in sub-aqueous conditions (3 m depth) at the Chatillon station of Lake le Bourget (Chindrieux, France). A 70 cm long core has been retrieved from Final Bronze Age IIIb archaeological layers that cover the Neolithic lacustrine chalks. The stratigraphy has been established by subaquatic archaeological investigations, the chronological frame being based on artefacts, dendrochronology and radiocarbon dates.

Archaeological levels are peculiarly well preserved with numerous vegetal debris (wood and bark shaves and twigs, seeds, charcoals...), ceramics, shells (bivalves and gasteropods) and gravels interspaced with more argillaceous levels.

Lipids from 20 samples collected in the different stratigraphic units were extracted and separated into neutrals and acidic compounds. The neutral compounds were further separated into aliphatics, aromatics, ethers and ketones before identification and quantitation by gas chromatography-mass spectrometry.

Triterpenes were abundant in archaeological layers whereas few were detected in lacustrine chalks. The aliphatic and aromatic fractions are mainly composed of des-A-triterpenes and aromatic derivatives of triterpenes (tetranor-oleana(ursa, lupa)heptaenes, dinor-oleana(ursa, lupa)trienes and tetraenes, lanosta(eupha)pentaenes and hexaenes) that largely exceed n-alkane abundances. Miliacin is the dominant compound detected in the ether fraction with concentrations ranging from 0 (in lacustrine chalks) to 6 µg/g sediment in archaeological levels. The ketone

fraction contains abundant and original pentacyclic triterpene ketones such as friedelin (up to 3 µg/g) and a compound identified as adian-5-en-3-one [4] (up to 2 µg/g). The variability in triterpene imprints probably reflects variable sources of organic material in archaeological levels and differential physico-chemical conditions in the medium.



Fig. 1. Subaquatic photo of the Chatillon station during an archaeological survey revealing the stratigraphic succession of archaeological levels and the disposition of wooden piles. Credits : E. Champelovier / Drassm.

## References

- [1] Billaud Y., Marguet A. (2007). In : "Environnements et cultures à l'âge du Bronze en Europe occidentale", Mordant C., Richard H. and Magny M. (Eds.). 129th National Congress of Historical and Scientific Societies, Besançon, 2004, p. 211-225.
- [2] Magny, M. (2004). *Quat. Int.* 113, 65-79.
- [3] Jacob, J. et al. (2008). *Journal of Archaeological Science* 35, 814-820.
- [4] [http://www.blog-orleans-valde Loire.fr/isto\\_frog/2008/12/le-spectre-du-nouvel-an.html](http://www.blog-orleans-valde Loire.fr/isto_frog/2008/12/le-spectre-du-nouvel-an.html)