



HAL
open science

Annual fossil organic carbon delivery due to mechanical and chemical weathering of marly badlands areas

Y. Graz, Christian Di Giovanni, Yoann Copard, N. Mathys, A. Cras, V. Marc

► To cite this version:

Y. Graz, Christian Di Giovanni, Yoann Copard, N. Mathys, A. Cras, et al.. Annual fossil organic carbon delivery due to mechanical and chemical weathering of marly badlands areas. *Earth Surface Processes and Landforms*, 2012, 37, pp.1263-1271. 10.1002/esp.3232 . insu-00782723

HAL Id: insu-00782723

<https://insu.hal.science/insu-00782723>

Submitted on 19 Mar 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Annual fossil organic carbon delivery due to mechanical and chemical weathering of marly badlands areas

Yann Graz,^a Christian Di-Giovanni,^{a*} Yoann Copard,^b Nicolle Mathys,^c Alexandre Cras^d and Vincent Marc^d

^a *Université d'Orléans, Université François Rabelais - Tours, CNRS/INSU Institut des Sciences de la Terre d'Orléans - UMR 6113, Campus Géosciences 1A rue de la Férellerie 45071 Orléans cedex 2.*

^b *UMR CNRS 6143 M2C, University of Rouen, department of Earth Sciences, place E. Blondel, Bat. Irese A, Mont Saint Aignan Cedex, France.*

^c *Cemagref Grenoble, Research Unit Erosion Torrentielle, Neige et Avalanches, BP 76, 38402 Saint Martin d'Hères, France.*

^d *Laboratoire d'Hydrogéologie, Université d'Avignon, Faculté des Sciences, 33 rue Louis Pasteur, 84000 Avignon.*

** Corresponding author: C. Di-Giovanni, Institut des Sciences de la terre d'Orléans, Université d'Orléans/Université François Rabelais de Tours, CNRS/INSU UMR 6113, 1A rue de la Férellerie, 45072 Orléans, France. E-mail : Christian.Di-Giovanni@univ-orleans.fr; fax : (33)02-38-41-72-12*

Abstract

A key issue in the study of the carbon cycle is to well constrain stocks and fluxes in and between C-reservoirs. Amongst these, the role and importance of Fossil Organic Carbon (FOC) release by weathering of outcropping sedimentary rocks on continental surfaces is still debated and lacks measured constraint. Our work focuses on FOC fluxes due to chemical and mechanical weathering of marls in two experimental watersheds showing a typical badlands geomorphology (Draix watersheds, Laval and Moulin, Alpes de Haute Provence, France). Organic matter from bedrocks, soil litters and riverine

particles are characterized by Rock-Eval 6 pyrolysis. FOC fluxes due to mechanical weathering are then estimated by monitoring the annual particulate solid exports at the outlets of the watersheds (1985-2005 period). FOC fluxes from chemical weathering were calculated using Ca^{2+} concentrations in dissolved loads (year 2002) to assess the amount of FOC released by the dissolution of carbonate matrix. Results show that FOC delivery is mainly driven by mechanical weathering, with a yield ranging from 30 to 59 $\text{t km}^{-2} \text{ yr}^{-1}$ in Moulin (0.08 km^2) and Laval (0.86 km^2) catchments respectively (1985-2005 average). The release of FOC attributed to chemical weathering reach 2.2 to 4.2 t km^{-2} for the year 2002. These high FOC fluxes from badlands are similar to those observed in tectonically active mountain catchments. At a regional scale, badlands outcropping within the Durance watershed do not exceed 0.25 % in area of the Rhône catchment, but could annually deliver 12 000 t yr^{-1} of FOC. This flux could correspond to 27 % of the total Particulate Organic Carbon (POC) load exported by the Rhône River to the Mediterranean Sea. At a global scale, our findings suggest that erosion of badlands may contribute significantly to the transfer of FOC from continental surfaces to depositional environments.

Keywords: fossil organic matter, experimental watersheds, organic carbon fluxes, mechanical weathering, chemical weathering.

1. Introduction

Continental surfaces are composed of a combination of various sedimentary, metamorphic and plutonic rocks. Sedimentary rocks as shales, carbonates and

sandstones represent 65 % of the total continental surface (Amiotte-Suchet *et al.*, 2003). All of these rocks contain a variable Fossil Organic Carbon (FOC) content, generally comprised of between 0.01 to near 50.00 wt. % - with an average of 0.25 to 0.97 wt. % for carbonates and shales (Ronov and Yaroshevsky, 1976). At outcrop, FOC is generally considered fully mineralized and acts as a CO₂ source for the atmosphere (Berner, 1989; Berner and Canfield, 1989; Berner and Maasch, 1996). However, this assumption is in disagreement with numerous observations which suggest a natural resistance and FOC occurrence in various modern carbon reservoirs, including soils (Lichtfouse *et al.*, 1997a, 1997b; Di-Giovanni *et al.*, 1998a, 1999a, 1999b, 2000; Copard *et al.*, 2006; Graz *et al.*, 2010, 2011), rivers (Kao and Liu, 1996; Di-Giovanni *et al.*, 2000; Massiello *et al.*, 2001; Blair *et al.*, 2003, 2004; Raymond and Bauer, 2001, 2004; Copard *et al.*, 2006; Graz *et al.*, 2010, 2011; Hilton *et al.*, 2011) and recent sediments (Combaz, 1980; Tyson, 1995; Di-Giovanni *et al.*, 1997, 1998b, 1999b, 1999c, 2000; Eglinton *et al.*, 1997; Leithold *et al.*, 2001; Blair *et al.*, 2003; Dickens *et al.*, 2004a, 2004b; Wakeham *et al.*, 2004; Galy *et al.*, 2008a, 2008b), which indicates that the FOC is not completely oxidized. However, while the occurrence of FOC in various C-reservoirs is undisputable, the global flux of weathered FOC to the hydrosphere still remains unknown.

At a global scale, it is assumed that rivers carry between 0.04 to 0.08 Gt yr⁻¹ of particulate FOC to the world's oceans (Meybeck, 1993, 2005; Blair *et al.*, 2003). This input mainly originates from mechanical weathering of outcropping rocks or soils. Nevertheless, FOC release from the continents can not be directly related to the estimated FOC input to world's ocean, as an unknown fraction of FOC can be partially (Fredericks *et al.*, 1983; Lo and Cardott, 1995; Petsch *et al.*, 2000; Copard *et al.*, 2002)

or fully mineralised (Berner, 1989; Petsch *et al.*, 2000) during its storage and / or transfer within continental surfaces. Until now, we only know that the modelled FOC delivery due to chemical weathering (chemFOC) of shales and carbonates can reach between 0.04 to 0.09 Gt yr⁻¹ (Di-Giovanni *et al.*, 2002; Copard *et al.*, 2007). FOC flux originating from chemical weathering of sandstones is supposed negligible (Amiotte-Suchet *et al.*, 2003). For these latter fluxes, the lack of global estimate is related to the complex dependence on climatic and geological conditions. These chemFOC and FOC loads to oceans are estimated to be the same order of magnitude of the particulate Organic Carbon (OC) load delivered to world's oceans (between 0.09 and 0.43 Gt yr⁻¹, Ittekkot, 1988; Degens *et al.*, 1991; Berner, 1992; Meybeck, 1993; Ludwig *et al.*, 1996; Stallard, 1998; Schlunz and Schneider, 2000). Yet, the impact of FOC fluxes remains uncertain.

Here we quantify the flux of FOC due to mechanical and chemical weathering in two small experimental watersheds over annual to decadal timescales. The FOC is attributable to Jurassic marls weathering that contribute to 80 % of the suspended load of the “moyenne Durance”; one of the main tributaries of the Rhône River (Fournier, 2004). A classical geochemical analysis (Rock-Eval 6 pyrolysis) was used to characterize organic matter in studied samples (bedrocks, soil litters and riverine particles). FOC fluxes were estimated by monitoring the particulate (1985-2005 period) and dissolved (year 2002) exports at the outlet of the watersheds.

2. Material and methods

2.1. Study area

The study area corresponds to two experimental watersheds Laval (0.86 km²) and Moulin (0.08 km²) located near Digne (44° 08' 33''N, 06° 22' 05''E ; Alpes-de-Haute-Provence, France; Fig. 1) and overlying Callovo-Oxfordian marly limestones (Mathys *et al.*, 2003; Mathys, 2006). Such rocks are very sensitive to weathering and develop typical badlands morphology with V-shapes gullies, producing very high sediment loads (>10 g/L) during floods (Mathys *et al.*, 2003). These two experimental watersheds deliver a sediment yield of 5 700 t km⁻² yr⁻¹ in Moulin and 11 200 t km⁻² yr⁻¹ in Laval (Mathys *et al.*, 2003) comparable to other badlands located around the Mediterranean Sea (Nadal Romero *et al.*, 2011). These watersheds present different vegetal cover densities: Laval (32 %), and Moulin (46 %) (Mathys *et al.*, 2003).

2.2. Previous results

72 watershed samples (bedrock: 5 samples; soils: 15 samples; regolith profiles (bedrocks to soils): 44 samples; riverine particles: 8 samples) were previously examined in order to distinguish fossil and recent organic matter (Copard *et al.*, 2006; Graz *et al.*, 2011). We used complementary techniques: optical (palynofacies methods), geochemical (Rock-Eval 6 pyrolysis, C/N ratio), molecular (gas chromatography/mass spectrometry) and isotopic (stable C isotopic composition) analyses. These investigations were completed by quantitative palynofacies analysis performed on 82 samples from Laval watershed (bedrock, regolith profiles, soils: 72 samples; riverine particles: 10 samples, Graz *et al.*, 2010). Fossil and recent organic markers were identified and tracked through the different organic carbon reservoirs. The results permit us to (i) quantify bedrock organic content varying between 0.10 (limestones) and 0.50 – 0.70 wt. % (marls) - such values are of the same order as the average value (0.50

wt. %) obtained by Artru (1972) on 6 000 callovo-oxfordian marls samples; (ii) distinguish recent and fossil organic matter and highlight the contribution of Fossil Organic Matter (FOM containing FOC), which can be found in all the studied organic carbon reservoirs. FOM contribution to total organic matter varies from 5.5 to 70.0 % in soils, 20.0 to 95.0 % in regolith profiles and 47.0 to 91.0 % in riverine particles (Graz *et al.*, 2010, 2011).

2.3. Strategy, organic matter characterization and samples design

Compilation of the data (Copard *et al.*, 2006; Graz *et al.*, 2010, 2011, see section 2.2.) indicates that FOM contribution was only quantified for 18 published riverine samples between years 2002 and 2008. However, this small number of samples is not sufficient to estimate annual FOM (and thus FOC) fluxes at the watersheds outlets. In order to address this issue, a calibration of Rock-Eval 6 parameters using previous quantitative palynofacies results was performed (bedrock, regolith profiles, soils and riverine particles, Graz *et al.*, 2010). The obtained HI / FOM relation was applied to 60 other riverine samples in order to directly distinguish fossil and recent organic carbon (see section 3.1.). These samples were collected by using automatic sampling between 1999 and 2008 at Laval (30 suspended load / 6 bed load) and Moulin (24 suspended load) outlets.

Rock-Eval 6 pyrolysis (RE6, Vinci Technologies®), a method previously developed for petroleum purposes and the analysis of sedimentary rocks, is now successfully applied for recent material (soils and recent sediments, Di-Giovanni *et al.*, 1997, 1998b; Noël *et al.*, 2001; Disnar *et al.*, 2003; Copard *et al.*, 2006; Sebag *et al.*, 2006). The protocol consists of two successive stages performed under a temperature program of 30 °C min⁻¹

¹. The former consists of pyrolysis of 100 mg of crushed sample in an oven. Hydrocarbon and oxygenated products released during a temperature increase from 200 to 650 °C are removed via a N₂ flow and quantified with flame ionization and infrared detectors respectively. The second stage consists of oxidation in an oven of the carbonaceous residue subjected to a temperature increase from 400 to 750°C (Lafargue *et al.*, 1998). Analysis of the pyrolysis signals provides some classical parameters such as Hydrogen Index (HI, in mg HC g⁻¹ TOC), which can be defined as an indicator of the hydrogen richness of the organic matter. The Organic Carbon content (OC) is given by the Total Organic Carbon (TOC, expressed in wt. %), equal to the sum of pyrolysed OC and residual OC provided by the oxidation stage.

A first study dealing with the development of quantitative palynofacies was already published (Graz *et al.*, 2010). This method allowed us to determine organic particles mass concentrations and to discriminate Recent Organic Matter (ROM) from Fossil Organic Matter (FOM containing FOC) in 82 samples from Laval watershed (bedrock, regolith profiles, soils: 72 samples; riverine particles: 10 samples). Rock-Eval pyrolysis was also performed for all these samples. HI values (11 to 180 mg HC g⁻¹ TOC) and the relative proportion of ROM (0.0 to 88.0 %) were used to calibrate Rock Eval parameters in order to obtain fossil and recent carbon proportions in all studied samples (see 3.1.). Results are listed online in supplementary material.

2.4. Calculation of FOC fluxes released by bedrocks mechanical (mecFOC) and chemical weathering (chemFOC)

Drain riverine particles are the product of mechanical weathering of the bedrock (Di-

Giovanni *et al.*, 2000; Copard *et al.*, 2006). mecFOC can be calculated by considering FOC content of bedrocks and their annual fluxes. Such calculations were possible since the studied watersheds were previously equipped by Cemagref in order to estimate both the water discharge and fine and coarse solid transports (see Richard and Mathys, 1999; Mathys *et al.*, 2003; Klotz *et al.*, 2005; Mathys, 2006; for further details). At the outlet of Laval and Moulin catchments, a gauging station, with a calibrated flume is equipped with a series of recorders (e.g. floating device, ultrasonic sensor, radar sensor). Upstream of this flume a sediment trap retains the coarse material. This method gives a total amount of bed load conveyed by the streams. In the gauging flume, an automatic sampler takes samples during floods with a program referring to both the level of water and the time lag between two samples, giving a discontinuous estimation of fine solid transport during a flood. Since 1995, prototypes of optical fiber sensors were installed and provide measures sediment concentration of the flow using back scattering properties of the mixture. As a result, combined sediment concentration and discharge measurements allow one to calculate the suspended sediment yield for flood events and on annual timescales.

The dissolution of rocks that contain carbonates produces dissolved carbonates and an insoluble residue that can contain organic carbon if the parent rocks contained organic matter. If the yield of dissolved carbonates and the initial carbonates content of the parent rocks are known, then the yield of the insoluble residue can be calculated (Di-Giovanni *et al.*, 2002). Furthermore, if the initial FOC content in the parent rock is known, then the amount of FOC yielded from the dissolution of the carbonate matrix can be estimated. Calculations were only possible for the year 2002. Indeed, Cras (2005) carried out a multi-scale hydrochemical investigation over one year in Laval and

Moulin catchments, including both high flow and low flow periods. Automatic water samplers were used to collect water during the high flows periods whereas low flow waters were hand sampled. According to the flood events, the sampling frequency could reach 5 min. During low flow periods, a weekly sampling was carried out. Mean concentration estimates were calculated from 163 and 208 samples in Laval (including 6 flood events) and in Moulin (including 6 flood events) respectively. Cras *et al* (2007) showed that rainfall did not contain significant amount of Ca²⁺. Accordingly, we can consider that Ca²⁺ measured at the outlet of the watersheds mainly originated from bedrock chemical weathering.

3. Results

3.1. Rock-Eval 6 pyrolysis parameters calibration

There is a significant correlation between previously determined proportions of ROM (Recent Organic Matter - by opposition to FOM - Graz *et al.*, 2010) and HI values from Rock-Eval pyrolysis ($R^2 = 0.93$, Fig. 2) for 82 samples from Laval watershed (bedrock, regolith, soil and riverine particles, see online supplementary material for more details). Accordingly, a relationship between HI values and ROM % can be defined and lead to the quantification of FOM % (FOM % = 100 - ROM %) in riverine suspended sediments following the equation (Eq.1):

$$\text{(Eq 1) HI (mg HC.g}^{-1}\text{ TOC)} = 1.7927 \times \text{ROM (\%)} + 10.292$$

Uncertainties on HI values (5%) are widely lower than those on palynofacies results (15 %, Graz *et al.*, 2010). Uncertainties on calculated FOM % values are then close to 15 %.

3.2. FOC content of riverine particles

Riverine particles present low TOC and HI values (0.46 to 0.65 % and 19 to 72 mg HC g⁻¹ TOC respectively, Tables 1 and 2) that are similar, but slightly higher, to those observed in bedrocks samples (Graz *et al.*, 2010).

The observed relationship between HI values and ROM % (Fig. 2) permits to estimate FOC and Recent Organic Carbon (ROC) contents (wt. %) (FOC = TOC x FOM %; and ROC = TOC x ROM %) in all studied riverine samples (Laval, Table 1 and Moulin, Table 2). FOC content appears constant in all studied samples and close to 0.5 wt. % (+/- 0.07, 60 samples). This is in agreement with previous results (18 samples) showing that riverine particles of these watersheds are the direct product of bedrocks mechanical weathering (Di-Giovanni *et al.*, 2000; Copard *et al.*, 2006; Graz *et al.*, 2010).

The higher ROC values (from 0.1 to 0.2 wt. %) are obtained for Moulin suspended load samples, which also exhibits a large area of vegetation cover (46 %, Mathys *et al.*, 2003). The lower ones (< 0.1 %) are obtained for Laval suspended load samples, where the vegetal cover area is somewhat lower (32 %, Mathys *et al.*, 2003).

3.3. FOC fluxes released by bedrocks mechanical weathering

Measurements of solid export values at the outlets (Klotz *et al.*, 2005; Mathys, 2006) are highly variable from one year to another but also from one basin to another (Table 3). The export loads at Laval outlet range from 4352 - 21148 t km⁻² yr⁻¹ compared to


1350 - 9990 t km⁻² yr⁻¹ at Moulin outlet. With respect to the spatial scale (area) and badlands context, these magnitudes are comparable to those observed for numerous badlands area, located around the Mediterranean Sea (Nadal-Romeo *et al.*, 2011).

Considering that the FOC content of these loads is close to 0.5 wt. % (Tables 1 and 2, see also section 3.2.), mecFOC can be calculated. Obtained values range between 22 and 106 t km⁻² yr⁻¹ for Laval, and between 7 and 50 t km⁻² yr⁻¹ for Moulin (Table 3). The inter-annual FOC fluxes average 59 and 30 t km⁻² yr⁻¹ respectively.

3.4. FOC fluxes release by bedrocks chemical weathering

Exported total dissolved load were 340 and 570 t km⁻² for the year 2002 respectively for Laval and Moulin (Table 4; Cras, 2005). Total Dissolved Solid (TDS) concentrations reached an average of 13.4 and 11.7 mmol l⁻¹ respectively in watershed waters and Ca²⁺ concentrations reached 2.5 and 2.4 mmol l⁻¹ (Table 4).

Ca²⁺ concentrations measured at the outlets of watersheds originate from the bedrocks carbonate dissolution as rainwater does not contain Ca²⁺ (Cras *et al.*, 2007). This dissolution is illustrated by the following equation (Eq.2):



On this basis, the estimation of the chemical weathered bedrock mass (t km⁻² yr⁻¹) can be obtained following three steps (Eqs.3, 4 and 5).

$$\text{(Eq.3) Annual Ca}^{2+} \text{ fluxes (t km}^{-2} \text{ yr}^{-1}) = \mathbf{M} \times \text{Ca}^{2+} \text{ \%}$$

Where M is dissolved load exported from the watersheds in 2002 (Table 4) and Ca% is the Ca^{2+} content in TDS from riverine waters (18.7 and 20.5 for Laval and Moulin respectively).

$$\text{(Eq.4) Annual mass of dissolved CaCO}_3 \text{ (t km}^{-2} \text{ yr}^{-1}) = \text{annual Ca}^{2+} \text{ fluxes} \times 2.5$$

Where 2.5 correspond to the molar mass of Ca reported to the molar mass of CaCO_3 (100 / 40).

$$\text{(Eq.5)}$$

$$\text{Mass of altered bedrock (t km}^{-2} \text{ yr}^{-1}) = \text{Annual mass of dissolved CaCO}_3 \times (100 / \text{Tc})$$

Where Tc is the CaCO_3 content of watersheds bedrocks (i.e. 35 %, Simmonet *et al.*, 1995).

Considering the bedrocks average TOC (0.50 wt. %, Artru, 1972; Graz *et al.*, 2010) and the annual mass of altered bedrocks, chemFOC (t $\text{km}^{-2} \text{ yr}^{-1}$) can be calculated as follows:

$$\text{(Eq.6) chemFOC (t km}^{-2} \text{ yr}^{-1}) = \text{Mass of altered bedrock (t km}^{-2} \text{ yr}^{-1}) \times \text{bedrock TOC (0.50 wt. \%)}$$

FOC fluxes inherited from chemical weathering of bedrocks (chemFOC) can be further mineralized, dissolved or stored as a insoluble residue. They reach 2.2 and 4.2 t $\text{km}^{-2} \text{ yr}^{-1}$ for Laval and Moulin watersheds, respectively.

4. Discussion

4.1. FOC fluxes

For these watersheds, FOC input in continental surfaces is mainly controlled by mechanical weathering (average mecFOC from 30 to 59 t km⁻² yr⁻¹, Table 3) while chemFOC fluxes, inherited from chemical weathering of bedrocks and considered as an insoluble residue, are much lower (from 2 to 4 t km⁻² yr⁻¹, year 2002). This strong difference between mechanical and chemical FOC estimates is not a surprise with respect to the geomorphological context (badlands) of these watersheds, marked by predominant gully erosion.

However in our study, the estimated chemFOC fluxes in Moulin catchment (4.2 t km⁻² yr⁻¹) are twice that of Laval catchment (2.2 t km⁻² yr⁻¹). This suggests that chemical weathering is more efficient within the former catchment. One of the possible causes to explain this variation would be the higher extent of forest area in Moulin (46 %) than in Laval (32 %, Mathys *et al.*, 2003). Indeed, it is well known that a large area of forest cover leads to a decrease in the runoff intensity (e.g. Campy and Macaire, 2003). Such context would favor the storage of FOC as an insoluble residue; which can be stored and subsequently mineralized in regoliths or soils or even exported if a future mechanical weathering occurs prior to its mineralization. mecFOC fluxes are globally twice higher for Laval (59 t km⁻² yr⁻¹) than for Moulin (30 t km⁻² yr⁻¹) catchments. Again, a possible cause for this variation is the lower forest cover in Laval catchment (32%). This could also be exacerbated by the higher mean slope values observed for Laval (58%, Mathys *et al.*, 2003), which should inhibit the development of forest cover and promote the mechanical weathering via an efficient runoff and erosion processes (Dietrich *et al.*, 2003).

mecFOC yields from the studied catchments are of the same order of magnitude as FOC yields previously reported in suspended sediments from tectonically active mountain catchments in USA/California (9 t km⁻² yr⁻¹ Eel river, Blair *et al.*, 2003; Leithold *et al.*, 2006; 11 t km⁻² yr⁻¹ Santa Clara river, Blair *et al.*, 2003; Komada *et al.*, 2004), New Zealand (14 t km⁻² yr⁻¹ Southern Alpines rivers, Hilton *et al.*, 2008; 29 t km⁻² yr⁻¹ Waipaoa river, Gomez *et al.*, 2003; Leithold *et al.*, 2006) and Taiwan (19 t km⁻² yr⁻¹, Lanyang Hsi, Kao and Liu, 1996, from 12 to 246 t km⁻² yr⁻¹, Hilton *et al.*, 2011). Our results suggest that high mecFOC transfers in particulate load are not restricted to tectonically active mountains belts. Steep terrain and weak sedimentary rocks give rise to the high calculated yields herein. As such, the delivery of FOC to depositional setting may be more widespread than previously thought (e.g. Blair *et al.*, 2003; Hilton *et al.*, 2011).

4.2. Fate of FOC in continental surfaces

The riverine particles at outlets of these watersheds reach the Rhône delta in less than five days (IRS, 2001). This suggests a very short transit time and little storage of FOC within the Durance and the Rhone catchments, which can be considered as a conduit for transporting FOC to the Mediterranean Sea. Consequently FOC delivered by mechanical weathering could significantly contribute to Rhône rivers organic content. Considering that, (i) at a regional scale (Durance catchment), watershed badlands outcrops area, where erosion of FOC is likely, reach 233 km² (Brochot, 1999; Rey *et al.*, 2007), (ii) all of these badlands present a similar erosion rate (Saignon and Savournon watersheds, Olivry and Hoorelbeck, 1990; Rey, 2002; Ray *et al.*, 2002) which are also comparable to other badlands area located around the Mediterranean sea

(Nadal Romero *et al.*, 2011) and (iii) the average mecFOC we obtained is $52 \text{ t km}^{-2} \text{ yr}^{-1}$, such formations covering at best 0.25 % of the Rhône watershed (97800 km^2), could likely deliver $12\,000 \text{ t yr}^{-1}$ of FOC to the Mediterranean Sea. This flux could correspond to 27 % of the total particulate organic carbon annually delivered by the Rhône river ($44\,080 \text{ t yr}^{-1}$, Ludwig *et al.*, 1996). This C-kerogen (FOC) has been recorded in marine sediments of Gulf of Lion (Gadel and Ragot, 1973) that are likely fed by Rhone sediments (Cauwet *et al.*, 1990). Such FOC reburial in marine sediments suggests the balance between OC burial and FOC oxidation maintaining atmospheric O_2 constant (Bernier, 1989; Bernier and Canfield, 1989) is not completed (Hilton *et al.*, 2011). This also means that O and C cycles should be revisited in the light of this FOC route for which a significant part, strongly controlled by the chemical and physical properties of kerogen (Copard *et al.*, 2007; Galy *et al.*, 2008b), may be reburied (e.g. Hilton *et al.*, 2011).

FOC delivered by chemical dissolution of bedrocks with carbonate matrix (e.g. marls) can be mineralized, dissolved or stored in regoliths and soils. Its resident time in soils could be close to 10^4 years (Keller and Bacon, 1998; Petsch *et al.*, 2000). However this duration is purely theoretic and should be considered with care as the mineralization rate firstly depends on chemical and physical properties of FOC. On the whole, the recalcitrant character of FOC increases with the richness of C content associated to H depleted content of kerogen (Joseph and Oberlin, 1983). This implies that higher burial temperature of kerogen produces material more recalcitrant to chemical weathering as oxidation, hydrolysis or biodegradation. In other terms, a mature kerogen reaching the gas window (close to the methagenesis, Durand, 1980) as metha-anthracites or the ultimate C-member, graphite, could be persistent over a long period in continental

surfaces and can be exported to oceans (Galy *et al.*, 2008b; Bouchez *et al.*, 2010). For some lower degree of thermal evolution, kerogen resistance toward weathering is governed by its chemical composition inherited from the biological precursors and previous processes experienced before burial. Accordingly, a kerogen showing a ligno-cellulosic origin (type III in a Van Krevelen diagram) with a high proportion of aromatic sheets, exhibits a higher resistance toward weathering than a kerogen consisting in bacterial and/or algale origin (type I and II in a Van Krevelen diagram) with rather a higher proportion of aliphatic compounds likely to be preferentially attacked by oxidation (Copard, 2002 and references therein). However analysis of soil OM modulates this simplistic postulate where some rich-aliphatic compounds appear to be highly resistant towards soil processes (e.g. Lichtfouse *et al.*, 1998). As some field examples, mineralization rates of FOC of outcropping humic coals ranged from different ranks (from low to high volatile bituminous coals) are comprised between 20 to 45% (Fredericks *et al.*, 1983; Lo and Cardott, 1995; Copard *et al.*, 2002) while C-kerogen of outcropping black shales (phytoplanktonic-derived origin) present a complete mineralization (Petsch *et al.*, 2000). For the FOM disseminated in outcropping marls of the Draix experimental catchments, FOM is dominantly a type III kerogen (Graz *et al.*, 2010, 2011). During Jurassic, these marls were deposited in marine environments where a mixture between marine and continental organic matter classically occurs (Artru, 1972). Without any further alterations, and regarding the maturity level reached by the kerogen disseminated in these rocks during burial (attested by a low Tmax values, <410°C, from the RE6 pyrolysis, Copard *et al.*, 2006; Graz *et al.*, 2011), this kerogen would have exhibited some HI values higher than 400 mg HC g⁻¹ TOC (Espitalié *et al.*, 1985). However, for the examined rocks, HI values

never exceed 180 mg HC g⁻¹ TOC and highlight that this kerogen is already weathered. This means that the kerogen has already lost a significant (but unknown) amount of labile FOC prior analyses, during a chemical weathering as the weathering front can dip several meters from the soil surface (e.g. Petsch *et al.*, 2000). This kerogen, or more precisely this recalcitrant “residual kerogen”, is thus resistant against further weathering. This could explain why the observed mineralization rates of FOC along some previously studied regolith profiles in these catchments never exceed 30 % (Graz *et al.*, 2011). As a consequence, a significant part of FOC released by chemical weathering of bedrocks can be stored in regoliths.

Unfortunately, constraint on the chemical composition of kerogen is not sufficient to predict its persistence in continental surfaces. External factors of climate, geological context and more recently of Human disturbances (e.g. Land Use Change on forest cover) in continental surfaces advance or mitigate the mineralization rate. As an example the aggressiveness of climate, both temperatures than precipitations amount as seen in tropical climate, would improve the mineralization rate of any OM including kerogen with a higher biodegradation or oxidation processes. If this FOC escapes from such processes, soil can be further eroded and deliver its FOC to rivers as part of the particulate load.

5. Conclusions

This study proposes a first assessment of FOC fluxes release by mechanical and chemical weathering of marls in badlands. With values ranging from 30 to 59 t km⁻² yr⁻¹, fluxes due to mechanical weathering are predominant and are the same order of

magnitude than those previously estimated from tectonically active mountain catchments (9 to 246 t km⁻² yr⁻¹). These significant values are not surprising in badlands due to weak substrate and high local slope in gully valleys which drive high physical erosion rates. At the scale of the lower course of the Rhone river, we assume that such geological formations could annually export 12 000 t of FOC into the Mediterranean Sea. At global scale, rivers draining such badlands, whose lithology is mainly composed of marls and clayed rocks (Nadal Romero *et al.*, 2011) rich in organic carbon, could be also a significant source of FOC in continental surfaces. This FOC can be conveyed downstream of the river with the clastic load to ocean sediments with implications for the global C and O biogeochemical cycles (Bernier 1989; Blair *et al.*, 2003).

FOC fluxes caused by the chemical dissolution of the bedrock during weathering are lower and reach respectively 2.2 and 4.2 t km⁻² yr⁻¹ for Laval and Moulin watersheds. Despite a high resistance of this FOC against weathering, this could promote storage as insoluble residues (soils, regoliths), and a part of this carbon can either be mineralized or dissolved.

At global scale, the fate of FOC depends on several parameters as the physical and chemical properties of FOC (origin, temperature recorded during OM burial), the duration of exposure to weathering, but also on the nature and hence the aggressiveness of the weathering processes (e.g. hydrolysis, bacterial activity).

Our results confirm that the release of FOC by mechanical and chemical weathering of sedimentary rocks does not simply act as a source of C to the atmosphere, but can also supply other carbon reservoirs (soils, riverine particles, sediments). Fossil contribution should be taken into account by other studies based on organic matter data and the role of FOC in the carbon cycle has to be revisited.

Acknowledgements

This work is a contribution of the GIS "Bassins de Draix, étude de l'érosion en montagne" piloted by the CEMAGREF and was financially supported by the CNRS program INSU PNSE-ACI 2001: "*Influence de l'érosion sur les flux de matière organique fossile dans les géosystèmes continentaux actuels (sols, cours d'eau) : bassins expérimentaux de Draix, Alpes de Haute Provence, France*". Authors thank Marielle Hatton, Emily Moore and Jean-Emmanuel Olivier for their technical / analytical supports and corrections. They also warmly thank Prof. Lane and Robert Hilton for their constructive comments.

References

- Amiotte-Suchet P, Probst JL, Ludwig W. 2003. Worldwide distribution of continental rock lithology: implications for the atmospheric/soil CO₂ uptake by continental weathering and alkalinity river transport to the oceans *Global Biogeochemical Cycles* 17: DOI: 10.1029/2002GB001891.
- Artru P, 1972. Les terres noires du basin rhodanien (Bajocien supérieur à oxfordien moyen) : stratigraphie, sédimentologie, géochimie. Doctoral thesis, Lyon University Grenoble, 182 pp.
- Berner RA. 1989. Biogeochemical cycles of carbon and sulphur and their effect on atmospheric oxygen over Phanerozoic time. *Paleogeography Paleoclimatology Paleoecology* 75: 97-122.
- Berner RA, Canfield DE. 1989. A new model for atmospheric oxygen over phanerozoic time. *American Journal of Science* 289: 333-361.

- Berner RA. 1992. Comments on the role of marine of marine sediment burial as a repository for anthropogenic CO₂. *Global Biogeochemical Cycles* 6: 1-2.
- Berner RA, Maasch KA. 1996. Chemical weathering and controls on atmospheric O₂ and CO₂: Fundamental principles were enunciated by J.J. Ebelmen in 1845. *Geochimica and Cosmochimica Acta* 60: 1633-1637.
- Blair NE, Leithold EL, Ford ST, Peeler KA, Holmes JC, Perkey DW. 2003. The persistence of memory: The fate of ancient sedimentary organic carbon in a modern sedimentary system. *Geochimica and Cosmochimica Acta* 67: 63–73.
- Blair NE, Leithold RL, Aller RC. 2004. From bedrock to burial: the evolution of particulate organic carbon across coupled watershed-continental margin systems. *Marine Chemistry* 92: 141-156.
- Bouchez J, Beyssac O, Galy V, Gaillardet J, France-Lanord C, Maurice L, Moreira-Turcq P. 2010. Oxidation of petrogenic organic carbon in the Amazon floodplain as a source of atmospheric CO₂, *Geology* 38: doi:10.1130/G30608.1.
- Brochot S. 1999. Les bassins versants de Draix, expérimentaux... et représentatifs. Les bassins versants expérimentaux de Draix laboratoire d'étude de l'érosion en montagne – actes du séminaire, Draix, Le Brusquet, Digne. Cemagref Editions, Antony : 223-234.
- Campy M, Macaire JJ. 2003. *Géologie de la surface. Erosion, transfert et stockage dans les environnements continentaux*, second Ed. Dunod, Paris, 440 p.

- Cauwet G, Gadel F, Sierra MMD, Donard O, Ewald M. 1990. Contribution of the Rhone river to organic-carbon inputs to the Northwestern Mediterranean sea. *Continental Shelf Research* 10 : 1025-1037.
- Combaz A. 1980. Les kérogènes vus au microscope. In *Kerogen. Insoluble Matter from Sedimentary Rocks*, Durand B (ed). Technip: Paris; 56–112.
- Copard Y. 2002. Altération diagénétique et post-diagénétique (thermicité et oxydation) des charbons carbonifères du Massif Central français (Saint-Etienne, Graissessac et autres lieux). Doctoral thesis, University of Orléans, 245pp.
- Copard Y, Disnar JR, Becq-Giraudon JF. 2002. Erroneous maturity assessment given by Tmax and HI Rock-Eval parameters on highly mature weathered coals. *International Journal of Coal Geology* 49: 57-65.
- Copard Y, Di-Giovanni C, Martaud T, Albéric P, Olivier JE. 2006. Using Rock-Eval 6 pyrolysis for tracking fossil organic carbon in modern environments: implications for the roles of erosion and weathering. *Earth Surface Processes and Landforms* 31: 135-153.
- Copard Y, Amiotte-Suchet P, Di-Giovanni C. 2007. Storage and release of fossil organic carbon related to weathering of sedimentary rocks. *Earth and Planetary Science Letters* 258: 345-357.
- Cras A. 2005. Etude de la dynamique de fonctionnement hydrologique des bassins versants torrentiels marneux. Apport du traçage naturel. Doctoral thesis, Université d'Avignon et des pays de Vaucluse (France), 161pp.
- Cras A, Marc V, Travi Y. 2007. Hydrological behaviour of sub-Mediterranean alpine headwater streams in a badlands environment. *Journal of Hydrology* 339: 130-144. doi: 10.1016/j.jhydrol.2007.03.004

Commentaire [yc1]: Tu as vraiment cité ma thèse qq part...?

- Degens ET, Kempe S, Richey JE. 1991. Biogeochemistry of major world rivers. SCOPE 42: 323-344.
- Dickens AF, Gélinas Y, Hedges JI. 2004a. Physical separation of combustion and rock sources of graphitic black carbon in sediments. *Marine Chemistry* 92: 215-223.
- Dickens AF, Gélinas Y, Hedges JI. 2004b. Reburial of fossil organic carbon in marine sediments. *Nature* 427 : 336-339.
- Dietrich WE, Bellugi DG, Sklar LS, Stock JD, Heimsath AM, Roering JJ. 2003. Geomorphic transport laws for predicting landscape form and dynamics prediction in geomorphology. *Prediction in geomorphology, Geophysical Monograph* 135: doi: 10.1029/135GM09.
- Di-Giovanni C, Bertrand Ph, Campy M, Disnar JR. 1997. Contribution de matière organique méso-cénozoïque dans un flux organique terrigène tardi et post-glaciaire (bassin de Chaillexon, Doubs, France). *Bulletin de la Société géologique de France* 168 : 553-559.
- Di-Giovanni C, Disnar JR, Bichet V, Campy M. 1998a. Sur la présence de matière organique méso-cénozoïque dans les humus actuels (bassin de Chaillexon, Doubs, France). *Comptes rendus de l'Académie des Sciences Paris* 326 : 553-559.
- Di-Giovanni C, Disnar JR, Bichet V, Campy M, Guillet B. 1998b. Geochemical characterization of soil organic matter and variability of a past glacial detrital organic supply (Chaillexon lake, France). *Earth Surface Processes and Landforms* 23: 1057-1069.
- Di-Giovanni C, Disnar JR, Campy M, Macaire JJ. 1999a. Variability of the ancient organic supply in modern humus. *Analisis* 27: 398-402.

- Di-Giovanni C, Disnar JR, Turpin S, Bréheret JG. 1999b. Estimation de la contribution des matières organiques remaniées au stock organique des sols et sédiments lacustres (bassin des Peyssières, hautes Alpes, France). *Bulletin de la Société géologique de France* 170 : 121-129.
- Di-Giovanni C, Disnar JR, Campy M. 1999c. Des particules charbonneuses, témoins des variations de l'érosion chimique d'un bassin versant calcaire durant l'Holocène (bassin de Chaillexon, Doubs, France). *Comptes rendus de l'Académie des Sciences Paris* 328 : 167-172.
- Di-Giovanni C, Disnar JR, Bakyono JP, Keravis D, Millet F, Olivier JE. 2000. Application de l'étude de la matière organique à l'analyse de l'érosion : exemple du bassin versant du Moulin, dans les terres noires des Alpes de Haute Provence, France. *Comptes rendus de l'Académie des Sciences Paris* 331: 7-14.
- Di-Giovanni C, Disnar JR, Macaire JJ. 2002. Estimation of the annual yield of organic carbon released from carbonates and shales by chemical weathering. *Global and Planetary Change* 32: 327-343.
- Disnar JR, Guillet B, Keravis D, Di-Giovanni C, Sebag D. 2003. Soil organic matter (SOM) characterization by Rock-Eval pyrolysis : scope and limitations. *Organic Geochemistry* 34: 327-343.
- Durand, B. 1980. *Kerogen: insoluble organic matter from sedimentary rocks*. Technip, Paris, 525 pp.
- Eglinton TI, Benitez-Nelson BC, Pearson A, McNichol AP, Bauer JE, Druffel ERM. 1997. Variability in Radiocarbon Ages of Individual Organic Compounds from Marine Sediments. *Science* 277, DOI: 10.1126/science.277.5327.796.

- Espitalié J, Deroo G, Marquis F. 1985. La pyrolyse Rock-Eval et ses applications, Partie 1-2. *Revue de l'Institut Français du Pétrole* 40: 563-579.
- Fournier L. 2004. Modélisation de la production des apports sédimentaires dans le bassin versant de la Durance. Master thesis, Université Paris XI, 50 pp.
- Fredericks PM, Warbrooke P, Wilson MA. 1983. Chemical changes during natural oxidation of high volatile bituminous coal. *Organic Geochemistry* 5: 89-97.
- Gadel F, Ragot JP. 1973. Sur l'allochtonie de la fraction organique particulaire des dépôts quaternaires récents du Golfe du Lion. In Tissot B, Bienner F (Eds). *Adv. Org. Geochem.* 21. Technips, Paris; 619-628.
- Galy V, France-Lanord C, Lartiges B. 2008a. Loading and fate of particulate organic carbon from the Himalaya to the Ganga–Brahmaputra delta. *Geochimica et Cosmochimica Acta*, 72: 1767-1787.
- Galy V, Beyssac O, France-Lanord C, Eglinton T. 2008b. Recycling of Graphite During Himalayan Erosion: A Geological Stabilization of Carbon in the Crust. *Science* 7 322: 943-945. DOI: 10.1126/science.1161408.
- Gomez B, Trustrum NA, Hicks DM, Rogers KM, Page MJ, Tate KR. 2003. Production, storage, and output of particulate organic carbon: Waipaoa River basin, New Zealand, *Water Resource Research* 39: doi:10.1029/2002WR001619.
- Graz Y, Di-Giovanni C, Copard Y, Laggoun-Defarge F, Boussafir M, Lallier-Verges E, Baillif P, Perdereau L, Simonneau A. 2010. Quantitative palynofacies analysis as a new tool to study transfers of fossil organic matter in recent terrestrial environments. *International Journal of Coal Geology* 84: 49-62.
- Graz Y, Di-Giovanni C, Copard Y, Elie M, Faure P, Laggoun-Défarge F, Lévêque J, Michels R, Olivier J.E. 2011. Occurrence of fossil organic matter in modern

- environments: optical, geochemical and isotopic evidence. *Applied Geochemistry* 26: 1302-1314.
- Hilton RG, Galy A, Hovius N. 2008. Riverine particulate organic carbon from an active mountain belt: Importance of landslides. *Global Biogeochemical Cycles* 22: GB1017, doi:10.1029/2006GB002905.
- Hilton RG, Galy A, Hovius N, Homg M J, Chen H. 2011. Efficient transport of fossil organic carbon to the ocean by steep mountain rivers: An orogenic carbon sequestration mechanism. *Geology* 39: 71-74.
- IRS 2001. Etude globale pour une stratégie de réduction des risques dus aux crues du Rhône. Institution interdépartementale des bassins Rhône-Saône (Ed)., Valence ; 40pp.
- Ittekkot V. 1988. Global trends in the nature of organic matter in river suspensions. *Nature* 332: 436-438.
- Joseph D, Oberlin A. 1983. Oxidation of carbonaceous matter – I Elemental analysis (C, H, O) and IR spectrometry. *Carbon* 21, 6: 559-564.
- Kao SJ, Liu KK. 1996. Particulate organic carbon export from a subtropical mountainous river (Lanyang Hsi), in Taiwan. *Limnology and Oceanography* 41: 1749-1757.
- Keller C K. and Bacon D H., 1998. Soil respiration and georespiration distinguished by transport analyses of vadose CO₂, ¹³CO₂, and ¹⁴CO₂. *Global biogeochemical cycles* 12: 361-372.
- Klotz S, Mathys N, Olivier JE. 2005. *Annuaire de l'ORE Draix 2005*. Document Cemagref (Ed)., France; 34pp.

- Komada T, Druffel ERM, Trumbore SE. 2004. Oceanic export of relict carbon by small mountainous rivers. *Geophysical Research Letters* 31: L07504, doi:10.1029/2004GL019512
- Lafargue, E, Marquis F, Pillot D. 1998. Rock-Eval 6 applications in hydrocarbon exploration, production and soil contamination studies. *Revue de l'Institut Français du Pétrole* 53: 421-437.
- Leithold EL, Blair NE. 2001. Watershed control on the carbon loading of marine sedimentary particles. *Geochimica and Cosmochimica Acta* 65: 2231-2240.
- Leithold EL, Blair NE, Perkey DW. 2006. Geomorphologic controls on the age of particulate organic carbon from small mountainous and upland rivers. *Global Biogeochemical Cycles* 20: GB3022, doi:10.1029/2005GB002677.
- Lichtfouse E, Bardoux G, Mariotti A, Balesdent J, Ballentine DC, Mack SA. 1997a. Molecular ^{13}C and ^{14}C evidence for the allochthonous and ancient origin of C16-C18 n-alkanes in modern soils. *Geochimica and Cosmochimica Acta* 61: 1891-1898.
- Lichtfouse E, Budzinski E, Garrigues P, Eglinton TI. 1997b. Ancient polycyclic aromatic hydrocarbons in modern soils: ^{13}C , ^{14}C and biomarkers evidence. *Organic Geochemistry* 26: 353-359.
- Lichtfouse E, Chenu C, Baudin F, Leblond C, Da Silva M, Behar F, Derenne S, Largeau C, Wehrung, P, Albrecht P. 1998. A novel pathway of soil organic matter formation by selective preservation of resistant straight-chain biopolymers: chemical and isotope evidence. *Organic Geochemistry* 28, 411-415.
- Lo HB, Cardott BJ. 1995. Detection of natural weathering of Upper Mc Alester and Woodford shale, Oklahoma. *Organic Geochemistry* 22: 73-83.

Commentaire [yc2]: Tu continues la vérification...

- Lopez Saez J, Corona C, Stoffel M, Rovéra G, Astrade L, Berger F. 2011. Mapping of erosion rates in marly badlands based on a coupling of anatomical changes in exposed roots with slope maps derived from LiDAR data. *Earth Surface Processes and Landforms* 36, 1162-1171.
- Ludwig W, Probst JL, Kempe S. 1996. Predicting the oceanic input of organic carbon by continental erosion. *Global Biogeochemical Cycles* 10: 23-41.
- Mathys N, Brochot S, Meunier M, Richard D. 2003. Erosion quantification in the small marly experimental catchments of Draix (Alpes de Haute Provence, France). Calibration of the ETC rainfall-runoff-erosion model. *CATENA* 50: 527-548.
- Mathys N. 2006. Analyse et modélisation à différentes échelles des mécanismes d'érosion et de transport de matériaux solides. Cas des petits bassins versants de montagne sur marnes (Draix, Alpes de Haute Provence). Doctoral thesis, Institut national polytechnique de Grenoble, 346 pp.
- Masiello CA, Druffel ERM. 2001. Carbon Isotope Geochemistry of the Santa Clara River. *Global Biogeochemical Cycles* 15: 407-416.
- Meybeck M. 1993. Riverine transport of atmospheric carbon: sources, global typology and budget. *Water, Air, and Soil Pollution* 70: 443-463.
- Meybeck M. 2005. Origins and behaviours of carbon species in world rivers. In: Roose and Lal (Eds). *Erosion and Carbon Dynamics, Advances in Soil Sciences Series*. CRC Boca Raton; 209-238.
- Nadal-Romero E, Martínez-Murillo J-F., Vanmaercke M., Poesen J, 2011. Scale-dependency of sediment yield from badland areas in Mediterranean environments *Progress in Physical Geography* ; 1–36.

- Noël H, Garbolino E, Brauer A, Lallier-Vergès E, de Beaulieu JL, Disnar JR. 2001. Human impact and soil erosion during the last 5000 yrs as recorded in lacustrine sedimentary organic matter at Lac d'Annecy, the French Alps. *Journal of Paleolimnology* 25: 229-244.
- Olivry JC, Hoorelbeck J. 1990. Erodabilité des terres noires de la vallée du Buëch (France, Alpes du Sud). *Cah. ORSTOM, ser. Pédot.* XXV; 95-110
- Petsch ST, Berner RA, Eglinton TI. 2000. A field study of the chemical weathering of ancient sedimentary organic matter. *Organic Geochemistry* 31: 475–487.
- Raymond PA, Bauer JE. 2001. Riverine export of aged terrestrial organic matter to the North Atlantic Ocean. *Nature* 409: 497-500.
- Raymond PA, Bauer JE. 2004. Controls on the variability of organic matter and dissolved inorganic carbon ages in northeast US rivers. *Marine Chemistry* 92: 353-366.
- Rey F. 2002. Influence de la distribution spatiale de la végétation sur la production sédimentaire de ravines marneuses dans les Alpes du sud. Doctoral thesis, Université de Grenoble I, 194 pp.
- Rey F, Robert Y, Vento O, 2002. Influence de la végétation forestière sur la formation de dépôts sédimentaires en terrain marneux. *Géomorphologie : relief, processus, environnement*, 1 : 85-92.
- Rey F, Abraham A, Gallicher-Lavanne C, Burylo M, Eugène RE, Mermin E, Puech C. 2007. Etude de la faisabilité d'une opération de réhabilitation écologique de ravines marneuses érodées pour la rétention durable des sédiments à l'échelle du bassin versant de la Durance. Cemagref (Ed). Rapport d'étude de l'Action 8 « Expérimenter le contrôle de l'érosion des versants de marnes noires » du Plan Durance Multi-Usages, 51 pp.

- Richard D, Mathys N. 1999. History, technical and scientific context of the experimental basins of Draix. Characteristics, available data and main results after ten years of monitoring. Document Cemagref (Ed), France; 20 pp. <http://excerpts.numilog.com/books/9782853625142.pdf>
- Ronov AB, Yaroshevsky AA. 1976. A new model for the chemical structure of the Earth crust. *Geochemistry International* 13: 1761-1795.
- Schlunz B, Schneider RR. 2000. Transport of terrestrial organic carbon to the oceans by rivers: re-estimating flux- and burial rates. *International Journal of Earth Sciences* 88: 599-606.
- Sebag D, Disnar JR, Guillet B, Di-Giovanni C, Verrecchia EP, Durand A. 2006. Monitoring organic matter dynamics in soil profiles by 'Rock-Eval pyrolysis': bulk characterization and quantification of degradation. *European Journal of Soil science* 57: 344-355.
- Simmonnet JP, Richey P, Pardon C. 1995. Contribution à l'étude des mécanismes et bilans de l'érosion chimique des Terres Noires du bassin de la Durance: exemple des bassins représentatifs expérimentaux de Draix (NE Digne). In *Compte Rendu de Recherche en érosion et hydraulique torrentielle*, 3, Cemagref (Ed). Antony : France; 189-199.
- Stallard RF. 1998. Terrestrial sedimentation and the carbon cycle: Coupling weathering and erosion to carbon burial. *Global Biogeochemical Cycles* 12: 231-257.
- Tyson RV. 1995. *Sedimentary Organic Matter: Organic Facies and Palynofacies*. Chapman and Hall, London, 615 pp.

Wakeham SG, Forrest J, Masiello C, Gaelinas Y, Alexander C, leavitt, P. 2004.
Hydrocarbons in Lake Washington Sediments. A 25-Year Retrospective in an Urban
Lake. *Environmental Science and Technology* 38: 431-439.

Figures and Tables captions

Fig. 1. Geographical location of Draix experimental watersheds (after Lopez Saez *et al.*, 2011).

Fig. 2. HI (mg HC g⁻¹ TOC) vs. Recent Organic Matter proportions (ROM %) for 82 samples (Laval watershed). Uncertainties on HI values: 5 %; uncertainties on ROM proportions: 15 %.

Table 1. Rock-Eval 6 parameters (TOC and HI), ROM / FOM proportions, and ROC / FOC contents for 36 Laval riverine particles.(Suspended load (SL) and bed load (BL)).

Table 2. Rock-Eval 6 parameters (TOC and HI), ROM / FOM proportions, and ROC / FOC contents for 24 Moulin riverine particles.(Suspended load (SL) and bed load (BL)).

Table 3. Sediment and FOC (mecFOC) yields for the Laval and Moulin watersheds.

Table 4. Dissolved load yield for Laval and Moulin catchments, Ca²⁺ and Total Dissolved Solid (TDS) concentrations in watershed waters (year 2002; after Cras, 2005).

table 1

Samples	TOC %	HI (mg HC g ⁻¹ TOC)	ROM % (% 15%)	FOM % (% 15%)	FOC wt. % (% 15%)	ROC wt. % (% 15%)
Laval SL						
03/05/2002a	0.55	25	8.2	91.8	0.5	<0.1
Laval SL						
03/05/2003b	0.51	33	12.7	87.3	0.4	0.1
Laval SL						
13/05/2002	0.55	36	14.3	85.7	0.5	0.1
Laval SL						
24/05/2002	0.51	27	9.3	90.7	0.5	<0.1
Laval SL						
26/08/2002	0.53	34	13.2	86.8	0.5	0.1
Laval SL						
02/09/2002	0.52	27	9.3	90.7	0.5	<0.1
Laval SL						
05/09/2002	0.50	30	11.0	89.0	0.4	0.1
Laval SL						
23/10/2002	0.55	24	7.6	92.4	0.5	<0.1
Laval SL						
06/07/2006	0.47	21	6.0	94.0	0.4	<0.1
Laval SL						
07/07/2006	0.56	25	8.2	91.8	0.5	<0.1
Laval SL						
12/07/2006b	0.65	23	7.1	92.9	0.6	<0.1
Laval SL						
12/07/2006a	0.56	20	5.4	94.6	0.5	<0.1
Laval SL						
18/07/2006b	0.46	19	4.9	95.1	0.4	<0.1
Laval SL						
18/07/2006a	0.51	20	5.4	94.6	0.5	<0.1
Laval SL						
08/09/2006	0.62	19	4.9	95.1	0.6	<0.1
Laval SL						
15/09/2006	0.52	25	8.2	91.8	0.5	<0.1
Laval SL						
25/09/2006	0.58	19	4.9	95.1	0.6	<0.1
Laval SL						
20/10/2006a	0.54	26	8.8	91.2	0.5	<0.1
Laval SL						
20/10/2006b	0.53	28	9.9	90.1	0.5	0.1
Laval SL						
20/10/2006c	0.52	29	10.4	89.6	0.5	0.1
Laval SL						
07/12/2006a	0.52	25	8.2	91.8	0.5	<0.1
Laval SL						
07/12/2006b	0.52	23	7.1	92.9	0.5	<0.1
Laval SL						
04/06/2007	0.54	33	12.7	87.3	0.5	0.1
Laval SL						
05/06/2007b	0.60	32	12.1	87.9	0.5	0.1
Laval SL						
05/06/2007a	0.62	24	7.6	92.4	0.6	<0.1
Laval SL						
16/09/2007	0.53	21	6.0	94.0	0.5	<0.1
Laval SL						
22/11/2007	0.48	19	4.9	95.1	0.5	<0.1
Laval SL						
11/01/2008a	0.61	23	7.1	92.9	0.6	<0.1
Laval SL						
11/01/2008b	0.55	24	7.6	92.4	0.5	<0.1
Laval SL						
11/01/2008c	0.53	28	9.9	90.1	0.5	0.1
Laval BL						
24/09/1999a	0.49	14	2.1	97.9	0.5	<0.1
Laval BL						
24/09/1999b	0.46	11	0.4	99.6	0.5	<0.1
Laval BL						
24/09/1999c	0.59	29	10.4	89.6	0.5	0.1
Laval BL						
Laval BL	0.42	14	2.1	97.9	0.4	<0.1

20/11/2006a						
Laval BL						
20/11/2006b	0.45	18	4.3	95.7	0.4	<0.1
Laval BL						
20/11/2006c	0.60	22	6.5	93.5	0.6	<0.1
Average	0.53	24	7.7	92.3	0.5	<0.1

table 2

Samples	TOC %	HI (mg HC g ⁻¹ TOC)	ROM % (*1.15 %)	FOC % (*1.15 %)	FOC wt. % (*1.15 %)	ROC wt. % (*1.15 %)
Moulin SL 13/05/2002	0.60	72	34.4	65.6	0.4	0.2
Moulin SL 03/05/2002	0.56	30	11.0	89.0	0.5	0.1
Moulin SL 05/08/2002	0.53	28	9.9	90.1	0.5	0.1
Moulin SL 27/08/2002a	0.54	43	18.2	81.8	0.4	0.1
Moulin SL 27/08/2002b	0.51	29	10.4	89.6	0.5	0.1
Moulin SL 02/09/2002	0.56	45	19.4	80.6	0.5	0.1
Moulin SL 05/09/2002	0.55	38	15.5	84.5	0.5	0.1
Moulin SL 12/09/2002	0.54	35	13.8	86.2	0.5	0.1
Moulin SL 06/07/2006a	0.53	21	6.0	94.0	0.5	<0.1
Moulin SL 06/07/2006b	0.62	45	19.4	80.6	0.5	0.1
Moulin SL 12/07/2006	0.54	19	4.9	95.1	0.5	<0.1
Moulin SL 08/09/2006a	0.54	24	7.6	92.4	0.5	<0.1
Moulin SL 08/09/2006b	0.54	20	5.4	94.6	0.5	<0.1
Moulin SL 15/09/2006a	0.53	23	7.1	92.9	0.5	<0.1
Moulin SL 15/09/2006b	0.53	21	6.0	94.0	0.5	<0.1
Moulin SL 24/09/2006	0.52	19	4.9	95.1	0.5	<0.1
Moulin SL 20/10/2006a	0.53	45	19.4	80.6	0.4	0.1
Moulin SL 20/10/2006b	0.58	52	23.3	76.7	0.4	0.1
Moulin SL 07/12/2006	0.51	22	6.5	93.5	0.5	<0.1
Moulin SL 05/06/2007a	0.58	17	3.7	96.3	0.6	<0.1
Moulin SL 05/06/2007b	0.52	40	16.6	83.4	0.4	0.1
Moulin SL 23/11/2007	0.53	21	6.0	94.0	0.5	<0.1
Moulin SL 12/01/2008a	0.58	33	12.7	87.3	0.5	0.1
Moulin SL 12/01/2008b	0.50	26	8.8	91.2	0.5	<0.1
Average	0.54	32	12.1	87.9	0.5	0.1

table 3

year	Laval		Moulin	
	sediment yield (t km ⁻² yr ⁻¹)	FOC yield (t km ⁻² yr ⁻¹)	sediment yield (t km ⁻² yr ⁻¹)	FOC yield (t km ⁻² yr ⁻¹)
1985	6188	31		
1986	10812	54		
1987	11288	56		
1988	7004	35	3564	18
1989	4352	22	1350	7
1990	12308	62	6534	33
1991	9180	46	5346	27
1992	21148	106	9990	50
1993	11968	60	6156	31
1994	18360	92	9396	47
1995	9452	47	4374	22
1996	17884	89	8424	42
1997	8704	44	4536	23
1998	7140	36	4590	23
1999	9996	50	4590	23
2000	13668	68	5562	28
2001	8772	44	6588	33
2002	12716	64	6642	33
2003	15300	77	5130	26
2004	17952	90	9180	46
2005	13328	67	5670	28
Average	11787	59	5979	30

table 4

catchment	dissolved load yield (Cras, 2005) (t km ⁻² yr ⁻¹)	[Ca ²⁺] (Cras, 2005) (mmol l ⁻¹)	average TDS (Cras, 2005) (mmol l ⁻¹)
Laval	340	2.5	13.4
Moulin	570	2.4	11.7

fig1




Fig2

