



Local time variations of the hydrogen corona observed by SPICAV-UV/VEX

Jean-Yves Chaufray, Jean-Loup Bertaux, Eric Quémerais, François Leblanc

► **To cite this version:**

Jean-Yves Chaufray, Jean-Loup Bertaux, Eric Quémerais, François Leblanc. Local time variations of the hydrogen corona observed by SPICAV-UV/VEX. International Venus Conference 2016, Apr 2016, Oxford, United Kingdom. insu-01298370

HAL Id: insu-01298370

<https://hal-insu.archives-ouvertes.fr/insu-01298370>

Submitted on 6 Apr 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LOCAL TIME VARIATIONS OF THE HYDROGEN CORONA OBSERVED BY SPICAV-UV/VEX.

J-Y. Chaufray, J-L. Bertaux, E. Quémerais, and F. Leblanc, *LATMOS-IPSL, CNRS, UVSQ, UPMC, Guyancourt, France (chaufray@latmos.ipsl.fr)*

Introduction:

As all the terrestrial planets, Venus is surrounded by an hydrogen corona coming from the photodissociation of the water vapor followed by its vertical transport at very high altitudes. From the UV observations of the resonant scattering Lyman-alpha emission, two components have been observed in this corona. A cold component corresponding to hydrogen population at equilibrium with the global atmosphere dominant below ~ 2000 km. A suprathermal component, produced by collisions between energetic protons with the cold component in the upper atmosphere. Due to its large energy, this hot population is able to reach very high altitudes, and escape and becomes dominant above ~ 4000 km (e.g. Bertaux et al. 1982).

Observations:

The hydrogen corona of Venus has been observed systematically from the UV spectrometer SPICAV aboard Venus Express (Bertaux et al. 2007). It has been observed at dayside, confirming most of the results obtained by past studies, and for the first time at the nightside. As shown in Fig. 1, the two components corona is seen through the two scale heights profile obtained at limb above the dayside (LT = 12H). The nightside profiles are very different due to the low illumination of the shadow by multiple scattering.




Fig. 1 Altitude profile of the Lyman- α brightness at different local times.

Results:

Chaufray et al. (2012) have analyzed the first profiles obtained at the dayside and derive a cold hydrogen density of $\sim 1-2 \times 10^5 \text{ cm}^{-3}$ and an exospheric temperature $\sim 250-300$ K at 250 km. A larger brightness was observed at the morning terminator (LT = 6h) compared to the evening terminator (LT=18h),

confirming an asymmetry observed from the mass spectrometer aboard Pioneer Venus Orbiter (PVO), but the hydrogen density needed to reproduce the profile at the morning terminator was lower than the hydrogen density derived by this instrument.

At higher altitudes, the brightness was dominated by the interplanetary emission making difficult an accurate retrieval of the hot hydrogen density. The hydrogen density derived was better constrained at the middle of the observed altitude range (~ 5000 km) and the hot hydrogen density estimated to $\sim 50 \text{ cm}^{-3}$.

More recently, Chaufray et al. (2015) reproduce the first brightness profiles at the nightside evening terminator (LT = 20H), using a radiative transfer model including both cold and hot hydrogen populations as well as the interplanetary emission partly scattered by the Venusian hydrogen corona.

The cold hydrogen density and temperature of the cold hydrogen population derived at the exobase was $4 \times 10^6 \text{ cm}^{-3}$ and temperature between 175-225 K, confirming a cooler and denser nightside corona compared to the dayside corona. The hot hydrogen density derived at 5000 km was $\sim 100 - 200 \text{ cm}^{-3}$ larger than the derived hot hydrogen density at the dayside and presenting a very short variability of a factor 2 in only few days. This variability could be driven by a variation of the dayside ionosphere.

In this presentation, we will summarize these recent results and present other observations obtained at the nightside morning side (LT = 4H) (Fig; 1), confirming the large variability of the hydrogen density with local time at the nightside.

Bibliography:

Bertaux, J-L., V.M. Lepine, V.G. Kurt, and A.S. Smirnov, 1982, *Icarus*, 52, 221-224

Bertaux, J-L. et al., 2007, *Planet. Space Sci.*, 55, 1673-1700

Chaufray, J-Y., J-L. Bertaux, E. Quémerais, E. Villard, and F. Leblanc, 2012, *Icarus*, 217, 767-778

Chaufray, J-Y., J-L. Bertaux, E. Quémerais, F. Leblanc, and S. Sulis, 2015, *Icarus*, 262, 1-8