



Reactions of acetone oxide stabilized Criegee intermediate with SO₂, NO₂, H₂O and O₃

Alexandre Kukui (1), Hui Chen (2), Shan Xiao (3), Wahid Mellouki (2), and Veronique Daële (2)

(1) Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E), CNRS / Université d'Orléans, Orléans, France, (2) Institut de Combustion Aérodynamique Réactivité et Environnement (ICARE), CNRS / Université d'Orléans, Orléans, France, (3) Department of Environmental Science & Engineering, Fudan University, Shanghai, China

Atmospheric aerosol particles represent a critical component of the atmosphere, impacting global climate, regional air pollution, and human health. The formation of new atmospheric particles and their subsequent growth to larger sizes are the key processes for understanding of the aerosol effects. Sulphuric acid, H₂SO₄, has been identified to play the major role in formation of new atmospheric particles and in subsequent particle growth. Until recently the reaction of OH with SO₂ has been considered as the only important source of H₂SO₄ in the atmosphere. However, recently it has been suggested that the oxidation of SO₂ by Criegee biradicals can be a significant additional atmospheric source of H₂SO₄ comparable with the reaction of SO₂ with OH.

Here we present some results about the reactions of the acetone oxide stabilized Criegee intermediate, (CH₃)₂=OO, produced in the reaction of 2,3-dimethyl-butene (TME) with O₃.

The formation of the H₂SO₄ in the reaction of acetone oxide with SO₂ was investigated in the specially constructed atmospheric pressure laminar flow reactor. The Criegee intermediate was generated by ozonolysis of TME. The H₂SO₄, generated by addition of SO₂, was directly monitored with Chemical Ionization Mass Spectrometer (SAMU, LPC2E). Relative rates of reactions of acetone oxide with SO₂, NO₂, H₂O and ozone were determined from the dependencies of the H₂SO₄ yield at different concentrations of the reactants.

Atmospheric applications of the obtained results are discussed in relation to the importance of this additional H₂SO₄ formation pathway compared to the reaction of OH with SO₂.