Resin exudation and resinicolous communities on *Araucaria humboldtensis* in New Caledonia

Christina Beimforde, Leyla J. Seyfullah, Vincent Perrichot, Kerstin Schmidt,
Jouko Rikkinen, Alexander-R. Schmidt

► To cite this version:

Christina Beimforde, Leyla J. Seyfullah, Vincent Perrichot, Kerstin Schmidt, Jouko Rikkinen, et al.. Resin exudation and resinicolous communities on *Araucaria humboldtensis* in New Caledonia. *Arthropod-Plant Interactions*, 2017, 11 (4), pp.495-505. 10.1007/s11829-016-9475-3 . insu-01408477

HAL Id: insu-01408477

<https://hal-insu.archives-ouvertes.fr/insu-01408477>

Submitted on 10 Jul 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

2 Resin exudation and resinicolous communities on *Araucaria* 3 *humboldtensis* in New Caledonia

4 Christina Beimforde¹ · Leyla J. Seyfullah¹ · Vincent Perrichot² ·
5 Kerstin Schmidt³ · Jouko Rikkinen^{4,5} · Alexander R. Schmidt¹

6 Received: 23 June 2016 / Accepted: 31 October 2016
7 © Springer Science+Business Media Dordrecht 2016

8 **Abstract** Conifers of the endemic species *Araucaria*
9 *humboldtensis* on Mont Humboldt in New Caledonia
10 exhibit extensive resin exudation. The resin flows of these
11 threatened trees are here shown to be induced by two beetle
12 species, which bore into branches and branchlets, leading
13 to abundant outpouring of resin, which gradually solidifies
14 into often drop-shaped resin bodies. The exudate is colo-
15 nized by a resinicolous and likely insect-vectored asco-
16 mycete, *Resinogalea humboldtensis*, which is only known
17 from Mont Humboldt. The fungus grows into fresh resin
18 and eventually develops ascomata on the surface of solid-
19 ifying resin. The solidified resin is also colonized by
20 another fungus, a dematiaceous hyphomycete. Based on
21 protein coding (CO1, CAD, ArgK) and ribosomal (LSU)
22 genes, the larger branch-boring beetle is a weevil of the
23 tribe Araucariini, which represents the sister group of all

other cossonine weevils. The smaller beetle species 24
belongs to the longhorn beetles (Cerambycidae). The 25
strong host specificity of the Araucariini, along with the 26
occurrence of two unique fungi, suggests that the resin- 27
associated community is native and has evolved on the 28
endemic conifer host. The formation of large amber 29
deposits indicates massive resin production in the past, but 30
the environmental triggers of exudation in Mesozoic and 31
Cenozoic ecosystems remain unclear. Our observations 32
from Mont Humboldt support the notion that the occur- 33
rences of small drop-shaped amber pieces in Triassic to 34
Miocene amber deposits were linked to ancient insect 35
infestations. 36

Keywords Amber · *Araucaria humboldtensis* · 38
Araucariini · Coleoptera · Resiniculous fungi · 39
Resinogalea · Weevils 40

A1 Handling Editor: Heikki Hokkanen.

A2 **Electronic supplementary material** The online version of this
A3 article (doi:10.1007/s11829-016-9475-3) contains supplementary
A4 material, which is available to authorized users.

A5 ✉ Alexander R. Schmidt
A6 alexander.schmidt@geo.uni-goettingen.de

A7 ¹ Department of Geobiology, University of Göttingen,
A8 Göttingen, Germany

A9 ² CNRS UMR 6118 Géosciences and OSUR, Université
A10 Rennes 1, Rennes, France

A11 ³ Institute of Ecology, Friedrich Schiller University Jena, Jena,
A12 Germany

A13 ⁴ Finnish Museum of Natural History, University of Helsinki,
A14 P.O. Box 7, 00014 Helsinki, Finland

A15 ⁵ Department of Biosciences, University of Helsinki,
A16 P.O. Box 65, 00014 Helsinki, Finland

Introduction 41

Some conifers and angiosperms produce large amounts of 42
resin from wounds in the wood in order to seal injuries and 43
to prevent microbial infections and infestations by arthro- 44
pods (Farrell et al. 1991; Gershenzon and Dudareva 2007; 45
Howe and Schaller 2008). Due to its compounds (ter- 46
penoids or phenolics), resin not only provides a mechanical 47
barrier but also protects plants through its toxic properties 48
(Bednarek and Osbourn 2009; Rautio et al. 2011; Sipponen 49
and Laitinen 2011). 50

Fossilized plant resins (ambers) date back to the Car- 51
boniferous 320 million years ago (Bray and Anderson 52
2009), but are found only in trace quantities until the Early 53
Cretaceous (Schmidt et al. 2012). In contrast, massive 54
amber deposits have been preserved in Cretaceous 55

(Barremian-Campanian) and in Eocene to Miocene sediments (Penney 2010). Worldwide, accumulation of this amber occurred in widespread Cretaceous and Cenozoic forests with resinous tree species. However, it remains uncertain why the trees produced so much resin at these particular times. Climate changes, the advent of wood-infesting insects and microorganisms, and/or even the appearance of certain conifer or angiosperm species might have triggered the massive resin outpourings (Schmidt et al. 2012; Labandeira 2014; Dal Corso et al. 2015).

Mesozoic resin-producing tree species typically include conifers of the Araucariaceae and the extinct family Cheirolepidiaceae (Langenheim 1995, 2003; Roghi et al. 2006a, b; Nohra et al. 2015). Later, probably since the Late Cretaceous and early Cenozoic, angiosperms such as Dipterocarpaceae (Rust et al. 2010) and Fabaceae (Langenheim 1995, 2003) started to contribute significantly to the amber fossil record. Among today's gymnosperms, many Pinaceae and Araucariaceae produce large quantities of resin (Langenheim 1995, 2003). While species of the former family are widespread in the Northern Hemisphere, extant Araucariaceae are predominantly found in mid- and southern latitudes (Borneo, Philippines, Chile, Argentina, southern Brazil, New Caledonia, New Zealand, Norfolk Island, Australia and New Guinea) with a major diversity centre in New Caledonia (Eckenwalder 2009). The disjunct extant distribution and the fossil record of the latter family suggest that araucarian conifers were more widespread in the past and probably represented a major component of Mesozoic forest ecosystems in both hemispheres (e.g. Lele 1956; Miller 1977; Stockey 1982; Hill 1995; Kunzmann 2007).

Presently the island of Grande Terre in New Caledonia has 19 Araucariaceae species of which 13 are endemic (Jaffré 1995; Gaudeul et al. 2012), and it has often been termed a Gondwanan refuge (Holloway 1979; Morat 1993a, b). However, recent studies suggest that the species richness of New Caledonia's araucarians is rather a result of adaptive radiation in the post-Eocene era, mainly forced by unusual edaphic conditions, i.e. the widespread occurrence of ultramafic soils (Setoguchi et al. 1998; Gaudeul et al. 2012; Escapa and Catalano 2013; Kranitz et al. 2014; Grandcolas et al. 2015). Nevertheless, with an evolutionary history of more than 200 million years, Araucariaceae are among the oldest extant conifers (Kunzmann 2007). In this respect, the conifer forests of New Caledonia offer an optimal site for the study of plant–environment interactions that are responsible for triggering resin production, past and present.

Here we show that the widespread and substantial resin exudation of *Araucaria humboldtensis* Buchholz 1949, an endangered (IUCN red list status) endemic New Caledonian mountain conifer, is induced by at least two species of beetles

and that the exuded resin provides the habitat and nutrient source for at least two unique types of resinicolous fungi. *Araucaria humboldtensis* occurs sporadically in the montane forests of Mont Mou, Mont Kouakoué and Montagne des Sources, but only close to the summit of Mont Humboldt is it the dominant timberline species (Fig. 1a). Our analysis of the plant–animal–fungal interactions at this unique location indicates that the wood-boring beetles are not recently introduced species, but native to New Caledonia and ecologically important in providing continuous substrate for two endemic, potentially ancient fungi. Our overall observations also provide insights into possible means of resin production and amber formation in earth history: for example, the similarity between the Mont Humboldt resin droplets and those preserved as Triassic amber is striking. We propose that not all ancient resin productions should necessarily be interpreted as anomalies caused by specific triggers such as high humidity (Dal Corso et al. 2015) or fire, but may have accumulated over time in humid forest environments. There is some indication also that the ancient resin surfaces may have supported specialized communities rather similar to that now described from Mont Humboldt (McKellar et al. 2011; Tuovila et al. 2013).

Materials and methods

Field work

Resin flows of *Araucaria humboldtensis* on Mont Humboldt were examined and photographed in October 2005 and in November 2011. Samples of resin with fungi and wood-boring insects were taken on 9 November 2011 from trees close to the refuge (elevation 1345 m, coordinates 21°52'57.52"S, 166°24'46.20"E), approximately 300 m north of the building (elevation 1320 m, coordinates 21°52'46.79"S, 166°24'49.17"E), and along the summit trail approximately 300 m east of the shelter (elevation 1380 m, coordinates 21°52'54.89"S, 166°24'55.85"E). Fungi were stored dry in sealed containers, and insects were preserved in 80% ethyl alcohol for storage and transport.

Repository

Specimens of *Resinogalea humboldtensis* are deposited in the herbaria at the MNHN Paris and in Helsinki (see Rikkinen et al. 2016). Lumps of resin with the dematiaceous hyphomycete are housed in the Geoscientific Collections of the Georg August University Göttingen (GZG.BST.21894a–e). Beetle specimens are housed in the collection of the laboratory Géosciences Rennes (University Rennes I), collection numbers VP-NC-221 (Araucariini) and VP-NC-202 (Cerambycidae).

Fig. 1 Resinous *Araucaria humboldtensis* on Mont Humboldt in New Caledonia. **a** Cloud forest of Mont Humboldt with *A. humboldtensis* as dominating tree species. **b** Tree with resin

outpourings on several branches. **c** Massive resin outpouring and death of distal branch end after infestation by weevil larvae of the Araucariini tribe

Microscopy and imaging

Samples were investigated under a Carl Zeiss Stereo Discovery V8 dissection microscope, and under a Carl Zeiss AxioScope A1 compound microscope, each equipped with a Canon EOS 5D digital camera. In some instances, incident and transmitted light were used simultaneously. Figure 4a, c–e shows digitally stacked photomicrographic composites of up to 150 individual focal planes, obtained by using the software package HeliconFocus 6.0 (HeliconSoft, <http://www.heliconsoft.com>) for an enhanced illustration of three-dimensional structures.

Cultivation of resinicolous fungi

For in vitro cultivation experiments, mycelia of the dematiaceous hyphomycete and spores from the mazaeium of *Resinogalea* were extracted and transferred to diverse sugar-based media: malt yeast extract agar (MYA, Ahmadjian 1967), malt extract agar (MEA, Blakeslee 1915) and potato dextrose agar (PDA, Roth). Additionally, Canada balsam and/or small pieces of *Araucaria humboldtensis* resin were provided as potential nutrient sources for the fungi.

176 DNA extraction and amplification

177 DNA was isolated using DNeasy Blood and Tissue Kit
178 (Qiagen) by following the manufacturer's protocol for
179 tissue samples. For the beetles, we amplified and
180 sequenced fragments of the mitochondrial gene cyto-
181 chrome oxidase 1 (COI), the D2–D3 segment of the nuclear
182 large ribosomal subunit (28S), the nuclear protein arginine
183 kinase (ArgK) and carbamoyl-phosphate synthase 2—as-
184 partate transcarbamylase—dihydroorotase (CAD). The ITS
185 region of the resinicolous fungi was amplified using pri-
186 mers ITS1F and ITS4. All primers used in this study are
187 provided in supplementary information (Table S1). PCRs
188 were performed in a 25- μ l volume containing final con-
189 centrations of 0.5 μ M of each primer, 0.5 μ M of each
190 dNTP, 1.25 units of GoTaq Hot Start DNA polymerase
191 (Promega), Green PCR buffer with a final concentration of
192 1.5 mM $MgCl_2$ and 1–2 μ l template DNA. A typical PCR
193 cycle consisted of 2 min initial heating to activate the DNA
194 polymerase and to ensure that the template DNA has
195 denatured, 40 cycles of 94 °C for 45–60 s of denaturation,
196 50–56 °C for 45–60 s of annealing, and 72 °C for 45–60 s
197 of elongation and a final elongation step at 72 °C for
198 10 min. PCR products were purified using Quick PCR
199 Purification Kit from Qiagen. PCR products were
200 sequenced in both directions with a MegaBACE 1000
201 automated sequencing machine and DYEnamic ET Primer
202 DNA Sequencing Reagent (Amersham Biosciences, Little
203 Chalfont, UK). All sequences were assembled and edited
204 using BioEdit version 5.0.9 (Hall 1999) and SeaView 4
205 (Gouy et al. 2010).

206 Phylogenetic analysis of the beetles

207 Since no adults of the two beetle species were available for
208 species identification, we substantiated morphological
209 diagnostics with comprehensive molecular phylogenetic
210 analysis. We combined our DNA sequence data obtained
211 from four genes (COI, LSU, ArgK and CAD) of the large
212 beetle species and from the COI gene for the small beetle
213 species with data from the National Center for Biotech-
214 nology Information (NCBI). All accession numbers are
215 provided in supplementary notes (Table S2). Data sets for
216 each gene were aligned separately using MAFFT version 6
217 (Kato and Toh 2008) with subsequent manual adjustment
218 to minimize the number of possible false homologies using
219 BioEdit 5.0.9. (Hall 1999) and SeaView 4 (Gouy et al.
220 2010). Unalignable regions and introns were excluded by
221 using the mask function in BioEdit 5.0.9. All genes were
222 subsequently combined in a super matrix using BioEdit
223 5.0.9. Bayesian analyses were performed using Markov
224 chain Monte Carlo (MCMC) in MrBayes 3.1.2 (Ronquist
225 and Huelsenbeck 2003). Evolutionary models with six

substitution rates, gamma distributed rate variation and a
proportion of invariable sites (GTR + I + G) were applied
to each gene separately by allowing MrBayes to estimate
specifications for the gamma shape parameter, proportion
of invariance and rate matrix for each partition. Phyloge-
netic analyses were performed using two parallel runs, each
with four chains, for 10 million generations and sampling
parameters every 1000 generations.

Most likely trees were sampled by using a burn-in of
25%, and a 50% majority rule consensus tree was gener-
ated. All analyses were performed on the freely available
computational resource CIPRES (Miller et al. 2010).
Assessing convergence and sufficient chain mixing (ef-
fective sample sizes >200) was observed using Tracer 1.5
(Rambaut and Drummond 2009). Resulting trees were
visualized using FigTree (Rambaut 2006–2009, [http://tree.
bio.ed.ac.uk/software/figtree/](http://tree.bio.ed.ac.uk/software/figtree/)).

Results and discussion

Field observations

A large proportion of *Araucaria humboldtensis* trees on
Mont Humboldt exhibited conspicuous resin outpourings
that were commonly associated with the death of branch-
lets or sometimes even entire branches (Fig. 1b, c). There
was no indication of damage caused by high winds or any
other external mechanical impact. The *Araucaria hum-
boldtensis* forest is located between approximately 1250
and 1500 m elevation, a region with daily fog and rainfall
ensuring constant high humidity year-round. Consequently,
while wildfires are common on the lower slopes of Mont
Humboldt, the timberline forest is not subjected to fire,
which is another possible trigger of enhanced resin pro-
duction (Scott 2000; Brasier et al. 2009; Najarro et al.
2010). Instead, we observed that the resin outpourings
consistently occurred on branches that had been infested by
wood-boring beetles (Figs. 1, 2, 3).

Two differently sized species of wood-boring beetles
were identified from the resin-exuding branches and
branchlets. Larvae of both species caused substantial
damage to the interior of the branch. Larvae of the smaller
beetle species were predominantly found in the distal thin
green branchlets (Figs. 2a–c, 3e), whereas larvae and
pupae of the larger species occurred in the wider woody
parts of mature branches (Figs. 1c, 3a–d). The boring
activity of the smaller species induced the production of
abundant resin drops of predominantly 3–4 mm size that
solidified on the leaves and branchlet tips. Borings of the
larger species often led to the death of distal branch ends or
of entire branches (Fig. 1c). This is likely because the
maximum width of the mature larvae is only slightly

Fig. 2 **Drop-shaped resin** pieces on *Araucaria humboldtensis* (a–c) and from Triassic and Eocene amber deposits (d, e). **a** Resin droplets on branchlets exuded after infestation by small beetles. **b**, **c** **Small resin** outpourings composed of several resin droplets.

d Amber droplets from the Triassic (Carnian) Heiligkreuz Formation in the Italian Dolomites. **e** **Drop-shaped** piece of Eocene Baltic amber. Scale bars 5 mm

smaller than mature branches of *Araucaria humboldtensis* so that both xylem and phloem are heavily damaged by the insect larvae (Fig. 3a, b).

Resin flows induced by the larger beetle species were drop-shaped to irregular, sometimes forming masses surrounding an entire branch (Fig. 1c). These larger resin outpourings commonly consist of several successive resin flows and obviously persist over many months, possibly even years (Figs. 1c, 4b).

Wood-boring beetles

The larger beetle species represents a weevil (family Curculionidae). According to our Bayesian analysis (Fig. 5), it belongs to the subfamily Cossoninae and likely represents a member of the Araucariini group, a tribe erected by Kuschel (1966) and presently accommodating the

neotropical genus *Araucarius* Kuschel 1966, and the four Oceanian genera *Coptocorynus* Marshall 1947, *Mastersiella* Lea 1896, *Xenocnema* Wollaston 1973 and *Inosomus* Broun 1882 (Alonso-Zarazaga and Lyal 1999). The phylogenetic analysis placed the smaller beetle species clearly outside the Curculionidae, and the morphological features of the larvae are characteristic for the family Cerambycidae. However, because of insufficient molecular data in public databases, we were not able to assign the smaller species to any group with any certainty.

The phylogenetic relationships of the Cossoninae (Fig. 5) revealed by our analysis are congruent with the results of Jordal et al. (2011). The monophyletic Araucariini tribe forms the sister group to the remaining Cossoninae, and the Mont Humboldt weevil constitutes the first-order sister clade to the remaining Araucariini. Although only ambiguously supported (0.79 pp, Fig. 5),

Fig. 3 Beetle infestation in *Araucaria humboldtensis*. **a** Fragment of a branch with branchlets died off after erosion of the branch's interior by weevil larvae. **b** Weevil larva feeding the tissue of a branch. **c**,

d Larva (**c**) and pupa (**d**) of a weevil of the Araucariini tribe. **e** Small beetle larvae, a representative of the Cerambycidae, from the interior of the branchlet. Scale bars 1 cm (**a**), 5 mm (**b–d**) and 1 mm (**e**)

the affiliation of the weevil from Mont Humboldt to the Araucariini group is conceivable because multiple independent Bayesian analysis with different taxon samplings for the Cossoninae all grouped the weevil from Mont Humboldt to the Araucariini tribe (data not shown).

Jordal et al. (2011) recently confirmed that the subfamily Cossoninae is monophyletic and includes the tribe Araucariini. Within the Cossoninae, only members of the monophyletic Araucariini are restricted to araucarian host plants, while the remaining Cossoninae comprise lineages feeding on angiosperms or Pinaceae. Both larvae and adults of Araucariini live inside the bark and phloem of their araucarian hosts (Mecke et al. 2005). Since mature branches of *Araucaria humboldtensis* were only slightly larger in diameter than the larvae and pupae of the weevil specimens within (Fig. 3a, b), it is difficult to deduce the preferred food of the larvae. However, phloem tissue was certainly included in their diet.

New Caledonia's Araucariaceae are believed to have diversified in post-Eocene times, mainly in response to the highly unusual edaphic conditions on the island (Setoguchi et al. 1998; Kranitz et al. 2014; Grandcolas et al. 2015). Not surprisingly, Mecke et al. (2005) also showed that New Caledonia harbours a wide variety of largely undescribed

weevil species that associate with different *Araucaria* species. Due to their specialization on araucarian hosts and basal position within the Cossoninae, the Araucariini are suspected to represent an archaic lineage within the Cossoninae (Kuschel 1966, 2000; Sequeira and Farrell 2001; Sequeira et al. 2000). The extreme host specificity may in turn be related to the stable morphology and anatomy of its araucarian hosts. Fossil data demonstrate that the phloem anatomy of araucarians has not changed significantly since Cretaceous times (Stockey 1994), and the trees are also known to have produced resin by the mid-Cretaceous (Nohra et al. 2015). Concurrently, resin flows of araucarian conifers in humid forest ecosystems have existed for tens of millions of years and potentially allowed the evolution of highly specialized resinicolous organisms and associations (see Mecke et al. 2005).

Resiniculous fungi

The semi-solid resin of *Araucaria humboldtensis* on Mont Humboldt provided suitable substrate for at least two unique species of resinicolous fungi (Fig. 4). The first fungus (Fig. 4a), a teleomorphic ascomycete, was recently described as new and named *Resinogalea humboldtensis*

Fig. 4 Resiniculous fungi from *Araucaria humboldtensis*. **a** Ascoma of *Resinogalea humboldtensis* exposing ascospores on **top**. Hyphae inside the resin are visible in the **lower left** of the image. **b** Solidified resin densely overgrown by a dematiaceous hyphomycete. **c** Close-up of (**b**) showing the **dark** hyphae. **d** Growing young colony of the

dematiaceous hyphomycete with formation of conidia. **e** Conidiophores of the dematiaceous hyphomycete arising forming aggregated superficial hyphae and bearing simple acropetal chains of conidia. Scale bars 200 μ m (**a**), 1 mm (**b**), 100 μ m (**c**, **d**) and 50 μ m (**e**)

Rikkinen et A. R. Schmidt 2016. The pale mycelium grows immersed in semi-solidified resin of *A. humboldtensis* (Fig. 4a, lower left), and ascomata are formed when the substrate solidifies. As the mycelium grows entirely submerged in the resin and because individual hyphae grow in random orientations, the fungus seems to utilize resin as a food source. For more details on the ecology, anatomy and taxonomy of *R. humboldtensis*, see Rikkinen et al. (2016).

While ascomata of *Resinogalea* were found in only a small minority of resin flows on Mont Humboldt, almost all hardening and solidified resin surfaces were colonized by a highly conspicuous dematiaceous hyphomycete (Fig. 4b–e). The dark hyphae of the fungus grew only on the hardened resin surface and did not penetrate into semi-solidified resin. Synnemata consisting of aggregated hyphae produce

simple acropetal chains of rounded to ovoid ornamented conidia 4–7 μ m long (Fig. 4e). Sometimes the conidiophores are reduced to mere conidiogenous cells with simple acropetal chains of more rounded conidia (Fig. 4d). On the basis of ITS sequences, the fungus belongs to the Mycosphaerellaceae, but its closer affinities remain unresolved.

Resiniculous fungi represent a polyphyletic ecological assemblage including, for example, many ascomycetes of the order Mycocaliciales (e.g. Rikkinen 1999, 2003a, b; Rikkinen et al. 2014, 2016; Tuovila et al. 2011a, b, 2012, 2013; Tuovila 2013). Many resinicolous mycocalicioids are highly substrate specific, a feature most likely related to the unique chemical compositions of many plant exudates (Lagenheim 2003). *Chaenothecopsis neocaledonica* from *Agathis ovata* (C.

Fig. 5 50% majority rule consensus phylogram from Bayesian analyses (MrBayes) showing phylogenetic relationships of the Cossoninae based on nuclear ribosomal (LSU) and protein coding (CO1, ArgK, CAD) sequence data of 18 Cossoninae species and two Cryptorhynchinae species used as outgroup. **Numbers** at nodes indicate posterior probabilities (pp) for node support. **Node supports** of 1.0 pp and less are shown

383 Moore ex Vieill.) Warb. 1900 is the only resinicolous myco- 403
 384 calicioid fungus so far known from araucarian exudates 404
 385 (Rikkinen et al. 2014). However, considering the high diver- 405
 386 sity of Araucariaceae in New Caledonia, many new resini- 406
 387 colous fungi may still await discovery on the island. 407

388 A resinous community 408

389 *Araucaria humboldtensis* is the dominant tree species in 409
 390 the timberline forest of Mont Humboldt. The trees produce 410
 391 a steady supply of fresh resin induced by at least two wood- 411
 392 boring beetle species. The phylogenetic placement of the 412
 393 weevil species within the Araucariini tribe suggests a long- 413
 394 lasting interaction between the narrow endemic conifer 414
 395 host and the beetle species that has specialized to bore into 415
 396 its branches and feed on their internal tissue. Interestingly, 416
 397 the occurrence of *Resinogalea* further indicates that the 417
 398 beetle-induced resin outpourings must have been common 418
 399 and continuous enough to allow the evolution and contin- 419
 400 ued existence of a unique ascomycete species specific to 420
 401 this unusual substrate. It is also likely that *Resinogalea* 421
 402 *humboldtensis* and possibly the dematiaceous hyphomycete 422
 423
 424

are dispersed by adult beetles. The ascomata of *Resino-* 403
galea have long and slender stalks and well-developed 404
 mazaedia (spore masses), which both represent typical 405
 calicioid features that are suspected to promote insect 406
 dispersal. Mature ascospores accumulate into the maza- 407
 edium and are then easily attached to roaming insects 408
 (Rikkinen 1995, 2003a; Tuovila et al. 2011a; Prieto and 409
 Wedin 2013). Also, the widespread production of syn- 410
 nemata by the anamorphic stages of many ascomycetes 411
 (e.g. Seifert 1985) is undoubtedly partly explained by 412
 adaptations for animal dispersal. Thus, the two unique 413
 fungi on *A. humboldtensis* resin may depend on the wood- 414
 boring beetles, not only for the production of suitable sub- 415
 strate, but also in their dispersal. 416

Despite repeated efforts, we could not induce the 417
 ascospores of *Resinogalea humboldtensis* to germinate and 418
 were thus unable to culture the fungus. The dematiaceous 419
 hyphomycete did grow, but very slowly and exclusively on 420
 the original substrate, i.e. small pieces of *Araucaria hum-* 421
boldtensis resin. No growth was observed on sugar-based 422
 media or Canada balsam (Pinaceae resin). When small 423
 pieces of the natural substrate and sugar were provided, 424

growth was restricted to the resin. This strongly suggests an inability to metabolize sugar-based carbohydrates and exemplifies the level of substrate specialization in resinicolous fungi. The inability to use Pinaceae resin together with the apparent narrow endemism in New Caledonia suggests that the dematiaceous hyphomycete may only occur on the resin of this one *Araucaria* species or that it is at least restricted to New Caledonia.

A concurrent example of extreme substrate specificity of a resinicolous fungus to a narrowly endemic conifer host was previously reported from California, where *Mycocalicium sequoiae* only lives on the resinous exudate that issues from the exposed heartwood of fire-scarred but still living trunks of *Sequoiadendron giganteum* (Bonar 1971). Two collections of the fungus have also been reported from *Sequoia sempervirens*, but their identity should be confirmed by molecular methods.

We conclude that the wood-boring beetles that cause highly conspicuous resin flows on *Araucaria humboldtensis* are not introduced species and probably do not pose a serious threat to the existence of the critically endangered tree species. Conversely, they induce and maintain a continuum of substantial resin flows that apparently provide the only substrate for two species of unique resinicolous fungi. There is reason to believe that the ‘triangle association’ between the endemic conifer host, the endemic weevil and the two endemic fungi evolved in the humid forests of New Caledonia and may be of considerable antiquity. Divergence time estimates by Kranitz et al. (2014) suggest that the New Caledonian *Araucaria* species diversified in the Miocene–Pliocene between 19 and 3 Ma and that *A. humboldtensis* appears to be approximately 5 Ma old, suggesting that the common history of the association may date back to the early Pliocene or even further.

Palaeoecological implications

We propose that the extant *Araucaria humboldtensis* forest offers a model for a type of enhanced resin production in ancient ‘amber forests’, i.e. past forest ecosystems from which amber deposits derive.

The reasons for massive Mesozoic and Cenozoic amber accumulations remain obscure. It has been suggested that the evolution of certain wood-boring insect species and subsequent large-scale insect outbreaks might have caused substantial tree damage followed by enhanced resin release (McKellar et al. 2011; Peris et al. 2015). Concurrently, it has been suggested that Coleoptera were unlikely to have been inducers for the resin outpourings that resulted in early–middle Cretaceous ambers, based on the sparse record of wood-boring beetle families such as Curculionidae, Cerambycidae or Buprestidae in these ambers (Peris

et al. 2016). However, this is based on the scarcity of adult specimens preserved and does not consider the possibility of larval stages boring into branchlets, which would have a low probability of being engulfed by resin flows. Our current observations from Mont Humboldt show that even a moderate population of specialized insects can trigger and maintain the continuous production of considerable amounts of resin. Ambers are known to vary in size from tiny droplets to massive pieces of up to 10 kg (e.g. Weitschat and Wichard 2002; Krumbiegel and Krumbiegel 2005), and their primary shape depends on resin viscosity and how and where the resin was produced on the ancient source trees. The shape and size of the resin droplets produced by *Araucaria humboldtensis* correspond closely to those of certain amber pieces, such droplets from the Triassic Heiligkreuz Formation in the Italian Dolomites (Fig. 2d, Roghi et al. 2006a, b; Schmidt et al. 2006, 2012), Late Cretaceous (Turonian–Santonian) of north-western and southern France (Saint Martin et al. 2013; Néraudeau et al. 2017) and from the Eocene of the Baltic area (Fig. 2e).

The occurrence of Triassic amber exclusively in a narrow Carnian-aged time horizon exemplifies a probable connection between enhanced resin production and climate change (Schmidt et al. 2012), and a connection to the Carnian Pluvial Event (Breda et al. 2009; Roghi et al. 2010), a global episode of atmospheric perturbation with dramatic changes in a monsoonal climate due to massive volcanism (Roghi et al. 2010; Preto et al. 2010; Dal Corso et al. 2012). Previously, Langenheim (1994) suggested that resin production can be enhanced in response to increased water availability, and so on Mont Humboldt the continuous high atmospheric humidity may well promote resin exudation responses during insect infestations. We presume, however, that massive resin productions during particular periods of Earth history were unlikely to be caused by any single event or trigger, but more likely reflect complex organismal interactions in humid forests, involving not only trees and wood-boring insects but potentially also many other resin-associated organisms, including resinicolous ascomycetes. This is supported by the fact that resinicolous *Chaenothecopsis* species have been found in Palaeogene ambers of Europe (Beimforde et al. 2014; Rikkinen and Poinar 2000; Tuovila et al. 2013), demonstrating that their special mode of nutrition was already developed at least 35 million years ago.

Acknowledgements We would like to thank Jérôme Munzinger (Montpellier) for help in the preparation of our field work in New Caledonia. Stefano Castelli and Guido Roghi (Padova) kindly provided the images of the Triassic amber droplets. Fieldwork and collection in southern New Caledonia were kindly permitted by the Direction de l’Environnement (Province Sud), permit number 17778 delivered in November 2011. Partial support for fieldwork was

provided to V.P. by grant OSUR (Univ. Rennes 1) from program AO1.P4 in 2011. We would also like to thank two anonymous reviewers for their helpful suggestions.

References

- Ahmadjian V (1967) A guide to the algae occurring as lichen symbionts: isolation, culture, cultural physiology, and identification. *Phycologia* 6:127–160
- Alonso-Zarazaga MA, Lyal CHC (1999) A world catalogue of families and genera of Curculionidea (Insecta: Coleoptera) (excepting Scolytidae and Platypodidae). *Entomopraxis*, Barcelona
- Bednarek P, Osbourn A (2009) Plant-microbe interactions: chemical diversity in plant defense. *Science* 324:746–747
- Beimforde C, Feldberg K, Nylander S, Rikkinen J, Tuovila H, Dörfelt H, Gube M, Jackson D, Reitner J, Seyfullah LJ, Schmidt AR (2014) Estimating the Phanerozoic history of the Ascomycota lineages: combining fossil and molecular data. *Mol Phylogenet Evol* 77:307–319
- Blakeslee AF (1915) Linder's roll tube method of separation culture. *Phytopathology* 5:68–69
- Bonar L (1971) A new *Mycocalicium* on scarred *Sequoia* in California. *Madrono* 21:62–69
- Brasier M, Cotton L, Yenney Y (2009) First report of amber with spider webs and microbial inclusions from the earliest Cretaceous (c.140 Ma) of Hastings, Sussex. *J Geol Soc* 166:989–997
- Bray PS, Anderson KB (2009) Identification of Carboniferous (320 million years old) class Ic amber. *Science* 326:132–134
- Breda A, Preto N, Roghi G, Furin S, Meneguolo R, Ragazzi E, Fedele P, Gianolla P (2009) The Carnian Pluvial Event in the Tofane area (Cortina d'Ampezzo, Dolomites, Italy). *Geo Alp* 6:80–115
- Dal Corso J, Mietto P, Newton RJ, Pancost RD, Preto N, Roghi G, Wignall PB (2012) Discovery of a major negative $\delta^{13}\text{C}$ spike in the Carnian (Late Triassic) linked to the eruption of Wrangellia flood basalts. *Geology* 40:79–82
- Dal Corso J, Gianolla P, Newton RJ, Franceschi M, Roghi G, Caggiati M, Raucsik B, Budai T, Haas J, Preto N (2015) Carbon isotope records reveal synchronicity between carbon cycle perturbation and the “Carnian Pluvial Event” in the Tethys realm (Late Triassic). *Glob Planet Change* 127:79–90
- Eckenwalder JE (2009) *Conifers of the world: the complete reference*. Timber Press, Portland
- Escapa IH, Catalano SA (2013) Phylogenetic analysis of Araucariaceae: integrating molecules, morphology, and fossils. *Int J Plant Sci* 174:1153–1170
- Farrell BD, Dussourd DE, Mitter C (1991) Escalation of plant defense: do latex/resin canals spur plant diversification? *Am Nat* 138:881–900
- Gaudeul M, Rouhan G, Gardner MF, Hollingsworth PM (2012) AFLP markers provide insights into the evolutionary relationships and diversification of New Caledonian *Araucaria* species (Araucariaceae). *Am J Bot* 99:68–81
- Gershenzon J, Dudareva N (2007) The function of terpene natural products in the natural world. *Nat Chem Biol* 3:408–414
- Gouy M, Guindon S, Gascuel O (2010) Seaview version 4: a multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Mol Biol Evol* 27:221–224
- Grandcolas P, Muriene J, Robillard T, Desutter-Grandcolas L, Jourdan H, Guilbert E, Deharveng L (2015) New Caledonia: a very old Darwinian island? *Philos Trans R Soc B* 363:3309–3317
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* 41:95–98
- Hill RS (1995) Conifer origin, evolution and diversification in the southern hemisphere. In: Enright NJ, Hill RS (eds) *Ecology of the southern conifers*. University Press, Cambridge, pp 10–29
- Holloway JD (1979) A survey of the Lepidoptera, biogeography and ecology of New Caledonia, Series entomologica, vol 15. The Hague, Dr W Junk, Netherlands
- Howe GA, Schaller A (2008) Direct defense in plants and their induction by wounding and insect herbivores. In: Schaller A (ed) *Induced plant resistance to herbivory*. Springer, Stuttgart, pp 7–29
- Jaffré T (1995) Distribution and ecology of the conifers of New Caledonia. In: Enright NJ, Hill RS (eds) *Ecology of the Southern Conifers*. Melbourne University Press, Melbourne, pp 171–196
- Jordal BH, Sequeira AS, Cognato AI (2011) The age and phylogeny of wood boring weevils and the origin of subsociality. *Mol Phylogenet Evol* 59:708–724
- Katoh K, Toh H (2008) Recent developments in the MAFFT multiple sequence alignment program. *Brief Bioinform* 9:286–298
- Kranitz ML, Biffin E, Clark A, Hollingsworth ML, Ruhsam M, Gardner MF, Thomas P, Mill RR, Ennos AR, Gaudeul M, Lowe AJ, Hollingsworth PM (2014) Evolutionary diversification of New Caledonian *Araucaria*. *PLoS ONE* 10:e110308. doi:10.1371/journal.pone.0110308
- Krumbiegel G, Krumbiegel B (2005) Bernstein—Fossile Harze aus aller Welt. Goldschneekverlag im Quelle & Meyer Verlag GmbH & Co, Wiebelsheim
- Kunzmann L (2007) Araucariaceae (Pinopsida): aspects in palaeobiogeography and palaeobiodiversity in the Mesozoic. *Zool Anz* 246:257–277
- Kuschel G (1966) A cossonine genus with bark-beetle habits, with remarks on relationships and biogeography (Coleoptera: Curculionidae). *N Z J Sci* 9:3–29
- Kuschel G (2000) La fauna curculiónica (Coleoptera: Curculionidea) de la *Araucaria araucana*. *Rev Chil Entomol* 27:41–51
- Labandeira CC (2014) Amber. *Paleontol Soc Pap* 20:163–216
- Langenheim JH (1994) Higher plant terpenoids: a phytocentric overview of their ecological roles. *J Chem Ecol* 20:1223–1280
- Langenheim JH (1995) Biology of amber-producing trees: focus on case studies of *Hymenaea* and *Agathis*. In: Anderson KB, Creeling JC (eds) *Amber, resin and fossil resin*. American chemical society symposium series, vol 617, pp 1–31
- Langenheim JH (2003) Plant resins. In: *Chemistry, evolution, ecology, ethnobotany*. Timber Press, Portland
- Lele KM (1956) Plant fossils from Parsora in the south Rewa Gondwana basin, India. *Palaeobotanist* 4:23–34
- McKellar RC, Wolfe AP, Muehlenbachs K, Tappert R, Engel MS, Cheng T, Sánchez-Azofeifa GA (2011) Insect outbreaks produce distinctive carbon isotope signatures in defensive resins and fossiliferous ambers. *Proc R Soc B* 278:3219–3224
- Mecke R, Mille C, Engels W (2005) *Araucaria* beetles worldwide: evolution and host adaptations of a multi-genus phytophagous guild of disjunct Gondwana derived biogeographic occurrence. *Pró Araucária Online* 1:1–18
- Miller CN (1977) Mesozoic conifers. *Bot Rev* 43:217–280
- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES science gateway for inference of large phylogenetic trees. In: *Proceedings of the gateway computing environments workshop (GCE)*, New Orleans
- Morat P (1993a) Our knowledge of the flora of New Caledonia: endemism and diversity in relation to vegetation types and substrates. *Biodivers Lett* 1:72–81
- Morat P (1993b) The terrestrial biota of New Caledonia. *Biodivers Lett* 1:69–71
- Najarro M, Peñalver E, Pérez-de la Fuente R, Ortega-Blanco J, Menor-Salván C, Barrón E, Soriano C, Rosales I, López del Valle R, Velasco F, Tornos F, Daviero-Gomez V, Gomez B,

- Delclòs X (2010) Review of the El Soplao amber outcrop, Early Cretaceous of Cantabria, Spain. *Acta Geol Sin* 84:959–976
- Néraudeau D, Perrichot V, Batten DJ, Boura A, Girard V, Jeanneau L, Nohra YA, Polette F, Saint Martin S, Saint Martin J-P, Thomas R (2017) Upper Cretaceous amber from Vendée, north-western France: age dating and geological, chemical, and palaeontological characteristics. *Cretac Res* 70:77–95
- Nohra YA, Perrichot V, Jeanneau L, Le Pollès L, Azar D (2015) Chemical characterization and botanical origin of French ambers. *J Nat Prod* 78:1284–1293
- Penney D (2010) Biodiversity of fossils in amber from the major world deposits. Siri Scientific Press, Manchester
- Peris D, Philips TK, Delclòs X (2015) Ptinid beetles from the Cretaceous gymnosperm-dominated forests. *Cretac Res* 52:440–452
- Peris D, Ruzzier E, Perrichot V, Delclòs X (2016) Evolutionary and paleobiological implications of Coleoptera (insecta) from Tethyan-influenced Cretaceous ambers. *Geosci Front* 7:695–706
- Preto N, Kustatscher E, Wignall PB (2010) Triassic climates—state of the art and perspectives. *Palaeogeogr Palaeoclimatol Palaeoecol* 290:1–10
- Prieto M, Wedin M (2013) Dating the diversification of the major lineages of Ascomycota (Fungi). *PLoS ONE* 8:e65576. doi:10.1371/journal.pone.0065576
- Rambaut A (2006–2009) FigTree. Tree figure drawing tool version 1.3.1, Institute of Evolutionary Biology, University of Edinburgh (<http://tree.bio.ed.ac.uk/software/figtree/>)
- Rambaut A, Drummond AJ (2009) Tracer. MCMC trace analysis tool version v1.5.0, University of Edinburgh (<http://tree.bio.ed.ac.uk/software/tracer/>)
- Rautio M, Sipponen A, Lohi J, Lounatmaa K, Koukila-Kähkölä P, Laitinen K (2011) In vitro fungistatic effects of natural coniferous resin from Norway spruce (*Picea abies*). *Eur J Clin Microbiol Infect Dis* 31:1783–1789
- Rikkinen J (1995) What's behind the pretty colours? A study on the photobiology of lichens. *Bryobrothera* 4:1–239
- Rikkinen J (1999) Two new species of resinicolous *Chaenothecopsis* (Mycocaliciaceae) from western North America. *Bryologist* 102:366–369
- Rikkinen J (2003a) New resinicolous ascomycetes from beaver scars in western North America. *Ann Bot Fenn* 40:443–450
- Rikkinen J (2003b) *Chaenothecopsis nigripunctata*, a remarkable new species of resinicolous Mycocaliciaceae from western North America. *Mycologia* 95:98–103
- Rikkinen J, Poinar GO Jr (2000) A new species of resinicolous *Chaenothecopsis* (Mycocaliciaceae, Ascomycota) from 20 million year old Bitterfeld amber, with remarks on the biology of resinicolous fungi. *Mycol Res* 104:7–15
- Rikkinen J, Tuovila H, Beimforde C, Seyfullah LJ, Perrichot V, Schmidt AR (2014) *Chaenothecopsis neocaledonica* sp. nov.: the first resinicolous mycocalicioid fungus from Araucariaceae. *Phytotaxa* 173:49–60
- Rikkinen J, Beimforde C, Seyfullah LJ, Perrichot V, Schmidt K, Schmidt AR (2016) *Resinogalea humboldtensis* gen. et sp. nov., a new resinicolous fungus from New Caledonia, placed in Bruceomycetaceae, a new family of Ascomycota. *Ann Bot Fenn* 53:205–215
- Roghi G, Ragazzi E, Gianolla P (2006a) Triassic amber of the Southern Alps. *Palaio* 21:143–154
- Roghi G, Kustatscher E, van Konijnenburg van Cittert JHA (2006b) Late Triassic Plants from the Julian Alps (Italy). *Boll Soc Paleontol Ital* 45:133–140
- Roghi G, Gianolla P, Minarelli L, Pilati C, Preto N (2010) Palynological correlation of Carnian humid sub-events throughout western Tethys. *Palaeogeogr Palaeoclimatol Palaeoecol* 290:89–106
- Ronquist F, Huelsenbeck JP (2003) Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19:1572–1574
- Rust J, Singh H, Rana RS, McCann T, Singh L, Anderson K et al (2010) Biogeographic and evolutionary implications of a diverse paleobiota in amber from the Early Eocene of India. *Proc Natl Acad Sci USA* 107:18360–18365
- Saint Martin S, Saint Martin J-P, Girard V, Néraudeau D (2013) Organismes filamenteux de l'ambre du Santonien de Belcodène (Bouches-du-Rhône, France). *Ann Paléontol* 99:339–360
- Schmidt AR, Ragazzi E, Coppelotti O, Roghi G (2006) A micro-world in Triassic amber. *Nature* 444:835
- Schmidt AR, Jancke S, Lindquist EE, Ragazzi E, Roghi G, Nascimbene P, Schmidt K, Wappler T, Grimaldi DA (2012) Arthropods in amber from the Triassic Period. *Proc Natl Acad Sci USA* 109:14796–14801
- Scott AC (2000) The pre-quaternary history of fire. *Palaeogeogr Palaeoclimatol Palaeoecol* 164:281–329
- Seifert KA (1985) A monograph of *Stilbella* and some allied hyphomycetes. *Stud Mycol* 27:1–235
- Sequeira AS, Farrell BD (2001) Evolutionary origin of Gondwanan interactions: how old are *Araucaria* beetle herbivores? *Biol J Linn Soc* 74:459–474
- Sequeira AS, Normark BB, Farrell BD (2000) Evolutionary assembly of the conifer fauna: distinguishing ancient from recent associations in bark beetles. *Proc R Soc B* 267:2359–2366
- Setoguchi H, Osawa TA, Pintaud JC, Jaffré T, Veillon JM (1998) Phylogenetic relationships within Araucariaceae based on rbcL gene sequences. *Am J Bot* 85:1507–1516
- Sipponen A, Laitinen K (2011) Antimicrobial properties of natural coniferous rosin in the European Pharmacopoeia challenge test. *APMIS* 119:720–724
- Stockey RA (1982) The Araucariaceae: an evolutionary perspective. *Rev Palaeobot Palynol* 37:133–154
- Stockey RA (1994) Mesozoic Araucariaceae: morphology and systematic relationships. *J Plant Res* 107:493–502
- Tuovila H (2013) Sticky business—diversity and evolution of *Mycocaliciales* (Ascomycota) on plant exudates. Publications from the Department of Botany, University of Helsinki, vol 44, pp 1–142
- Tuovila H, Cobbinah JR, Rikkinen J (2011a) *Chaenothecopsis khayensis*, a new resinicolous calicioid fungus on African mahogany. *Mycologia* 103:610–615
- Tuovila H, Larsson P, Rikkinen J (2011b) Three resinicolous North American species of Mycocaliciales in Europe with a re-evaluation of *Chaenothecopsis oregana* Rikkinen. *Karstenia* 51:37–49
- Tuovila H, Rikkinen J, Huhtinen S (2012) Nomenclatural corrections in calicioid fungi. *Karstenia* 52:73–74
- Tuovila H, Schmidt AR, Beimforde C, Dörfelt H, Grabenhorst H, Rikkinen J (2013) Stuck in time—a new *Chaenothecopsis* species with proliferating ascomata from *Cunninghamia* resin and its fossil ancestors in European amber. *Fungal Divers* 58:199–213
- Weitschat W, Wichard W (2002) Atlas of plants and animals in Baltic Amber. Pfeil-Verlag, Munich