Tracing the sources and cycling of phosphorus in river sediments using oxygen isotopes: Methodological adaptations and first results from a case study in France

Pistocchi Chiara, Federica Tamburini, Gérard Gruau, André Ferhi, Dominique Trevisan, Jean-Marcel Dorioz

► To cite this version:

Pistocchi Chiara, Federica Tamburini, Gérard Gruau, André Ferhi, Dominique Trevisan, et al.. Tracing the sources and cycling of phosphorus in river sediments using oxygen isotopes: Methodological adaptations and first results from a case study in France. Water Research, 2017, 111, pp.346-356. 10.1016/j.watres.2016.12.038 . insu-01424862

HAL Id: insu-01424862

<https://hal-insu.archives-ouvertes.fr/insu-01424862>

Submitted on 3 Jan 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Tracing the sources and cycling of phosphorus in river sediments using oxygen isotopes: Methodological adaptations and first results from a case study in France

Pistocchi Chiara, Federica Tamburini, Gerard Gruau, André Ferhi, Dominique Trevisan, Jean-Marcel Dorioz

PII: S0043-1354(16)30982-4

DOI: [10.1016/j.watres.2016.12.038](https://doi.org/10.1016/j.watres.2016.12.038)

Reference: WR 12588

To appear in: *Water Research*

Received Date: 10 June 2016

Revised Date: 22 December 2016

Accepted Date: 23 December 2016

Please cite this article as: Chiara, P., Tamburini, F., Gruau, G., Ferhi, A., Trevisan, D., Dorioz, J.-M., Tracing the sources and cycling of phosphorus in river sediments using oxygen isotopes: Methodological adaptations and first results from a case study in France, *Water Research* (2017), doi: 10.1016/j.watres.2016.12.038.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Tracing the sources and cycling of phosphorus in river sediments using oxygen isotopes: methodological adaptations and first results from a case study in France

Pistocchi Chiara^{1*}, Federica Tamburini¹, Gerard Gruau², André Ferhi³, Dominique Trevisan⁴, Jean-Marcel Dorioz⁴

¹Group of Plant Nutrition, Inst. Of Agricultural Sciences, Swiss Federal Institute of Technology (ETH Zürich)

²CNRS-Université de Rennes I, UMR Géosciences Rennes

³UPMC – Université Pierre et Marie Curie – Paris 6

⁴INRA, UMR Carrel

Abstract

An essential aspect of eutrophication studies is to trace the ultimate origin of phosphate ions (P-PO₄) associated with the solid phase of river sediments, as certain processes can make these ions available for algae. However, this is not a straightforward task because of the diversity of allochthonous and autochthonous sources that can supply P-PO₄ to river sediments as well as the existence of in-stream processes that can change the speciation of these inputs and obscure the original sources. Here, we present the results of a study designed to explore the potentials, limitations and conditions for the use of the oxygen isotope composition of phosphate ($\delta^{18}\text{O}_\text{p}$) extracted from river sediments for this type of tracing. We first tested if the method commonly applied to soils to purify P-PO₄ and to measure their $\delta^{18}\text{O}_\text{p}$ concentrations could be adapted to sediments. We then applied this method to a set of sediments collected in a river along a gradient of anthropogenic pressure and compared their isotopic signatures with those from

* Corresponding author.

E-mail address: Chiarapi@ethz.ch

samples that are representative of the potential P-PO₄ inputs to the river system (soils and riverbank material). The results showed that following some adaptations, the purification method could be successfully transposed to river sediments with a high level of P-PO₄ purification (> 97%) and high $\delta^{18}\text{O}_\text{p}$ measurement repeatability and accuracy (< 0.4‰). The values for the potential allochthonous sources varied from 11.8 to 18.3‰, while the $\delta^{18}\text{O}_\text{p}$ value for the river sediments ranged from 12.2 to 15.8‰. Moreover, a sharp increase (> 3‰) in the sediment $\delta^{18}\text{O}_\text{p}$ value immediately downstream from the discharge point revealed the strong impact of municipal wastewater. The calculation of the theoretical equilibrium $\delta^{18}\text{O}_\text{p}$ values using the river water temperature and $\delta^{18}\text{O}_\text{w}$ showed that the downstream sediments were in equilibrium, which was not the case for the upstream sediments. This difference could be related to the contrast between the short residence time of the transfer system in the catchment head, which can preserve the isotopic variability of the source materials, and the longer residence times and higher P bioavailability in the lower catchment, possibly fostering the recycling of P-PO₄ by the biota and the equilibration of the oxygen isotope signature in P-PO₄. These results demonstrate the potential of the isotopic approach to assess the sources and in-stream turnover of sedimentary P in river systems.

KEYWORDS: Phosphorus transfer, Agriculture, In-stream processes, Eutrophication, Phosphate oxygen isotopes

1. Introduction

Phosphorus (P) associated with the solid phase and designated as “particulate-P” often accounts for the major part of phosphorus transferred from terrestrial to aquatic ecosystems (Kleinman et

al., 2011; Kronvang et al., 2007; Sharpley et al., 1999). The impact of particulate-P on the trophic state and eutrophication of the receiving aquatic ecosystems, especially lentic ecosystems, depends on the size of its potentially bioavailable fraction. The amount of P that occurs as inorganic phosphate ions (P-PO_4 pools), either in solution (dissolved P-PO_4), sorbed onto, or precipitated within stream and river sediments, is thought to represent the most readily bioavailable P pool (Moorlegghem et al., 2013). Therefore, an essential aspect of studies dedicated to the restoration of eutrophicated aquatic ecosystems is to quantify P-PO_4 pools in river water and sediments and to trace the ultimate origin of these pools (Kleinman et al., 2011; Mellander et al., 2015).

However, this quantification is not a straightforward task because of the diversity of the allochthonous and autochthonous sources that may supply P-PO_4 to river sediments. Several allochthonous compartments in the catchment can supply particulate-P to river sediments, including pristine and weathered basement rocks, forested or agricultural soils, riverbank sediments, etc. For example, the autochthonous addition of P-PO_4 to the sediments can occur through sorption onto the sediments of dissolved P-PO_4 ions supplied by sewage effluents or the in-stream mineralization of sewage organic matter (Poulenard et al., 2008). The in-stream retention of dissolved P-PO_4 by the sediments primarily occurs during low-flow periods when the retention time of the sediment is long enough to allow the necessary physical-chemical reactions to develop (Dorioz et al., 1998a, 1989; Ensign et al., 2006; Pistocchi et al., 2012) and is particularly intense immediately downstream of the point source pollution inputs.

Autochthonous P-PO_4 pools can also come from the recycling of allochthonous particulate-P by aquatic flora or by microorganisms in the sediment. The biological uptake by biofilm is another important seasonal process that is involved in the retention of P-PO_4 ions by the sediment (Wang et al., 1999). These in-stream recycling and retention processes, which help the P to turn over within the river ecosystem, are expected to be triggered by the bioavailability of

allochthonous P and to regulate the internal productivity of the river system. Their quantification can be used to assess the efficiency of a river system to recycle and transform any particulate P into more bioavailable P-PO₄, but an efficient recycling can eventually obscure the ultimate origin of the P-PO₄ exported at the outlet.

It is often impossible to quantify and trace the different sources of particulate-P that are potentially involved in eutrophication using only the variability in P speciation as determined by classical chemical extraction methods (Burrus et al., 1989; Pacini and Gächter, 1999; Poulenard et al., 2009; Walling et al., 2008). As a result, there a need to develop tools or/and strategies that can be reliably used to trace this important P compartment, particularly its dominant inorganic P-PO₄ fraction, and to understand the extent and rate at which particulate-P is biologically recycled and modified within streams and rivers.

Oxygen isotopes could provide one such possible tool. The oxygen isotopic composition of phosphate ($\delta^{18}\text{Op}$) has already been successfully used to trace P sources and study the P-PO₄ cycle in soils, freshwater and marine sediments (Davies et al., 2014; Goldhammer et al., 2011; Gooddy et al., 2016; Gruau et al., 2005; Melby et al., 2013; Tamburini et al., 2014). This approach is based on: i) the existence of measurable differences in the abundance ratio for ¹⁸O and ¹⁶O atoms in the phosphate group of potential P-PO₄ sources (e.g. primary P-rich minerals, chemical fertilizers, chemical detergents, phosphate from wastewater, etc.); and ii) the fact that biologically mediated processes have the possibility to modify $\delta^{18}\text{Op}$ in a predictable way (Blake et al., 2005). The first property allows the oxygen isotope composition of P-PO₄ to be potentially used as a source tracer in environmental systems, whereas the second property allows it to be used to quantify the rate of biological recycling of P-PO₄. Two enzymatic processes are mainly responsible for changes in the $\delta^{18}\text{Op}$ value. One is the isotope equilibration between oxygen in P-PO₄ and oxygen in ambient water. This process, which is caused by reversible phosphorylation/dephosphorylation reactions that are catalyzed by the

inorganic intracellular pyrophosphatase enzyme (Blake et al., 2005), is temperature dependent and has been described by an empirical equation (Longinelli and Nuti, 1973) that was recently revised by Chang and Blake (2015). The second is the kinetic isotope fractionation produced by the enzymatic hydrolysis of phosphoesters (Liang and Blake, 2006; von Sperber et al., 2015, 2014), the magnitude of which depends on the type of enzyme involved (e.g. acid or alkaline phosphatases) and the final effect of which on the isotopic signature of the recycled P-PO₄ also depends on the $\delta^{18}\text{O}_\text{P}$ value for the original substrate, i.e. inheritance effect (Liang and Blake, 2006). Small kinetic isotope effects are associated with the initial stages of the abiotic reaction, such as the sorption of P-PO₄ onto the solid phases (Jaisi et al., 2010; Melby et al., 2013b) or biological uptake (Blake et al., 2005).

In freshwater ecosystem studies, the use of $\delta^{18}\text{O}_\text{P}$ is at an early stage (Davies et al., 2014). For river sediments, the lack of a standardized method to extract, purify and analyze $\delta^{18}\text{O}_\text{P}$ is a hinderance. In this study, we first adapted the method that is commonly used for soils to sediments and validated the capability of this method to provide precise and accurate results. We then applied this method to riverbed sediments from a river in the French Alps with the following objectives: i) to characterize the variability of the $\delta^{18}\text{O}_\text{P}$ signatures in the riverbed sediments and the potential external sediment sources (soils, riverbank materials) in an exemplar river system; ii) to explore the potential, limitations and conditions of using the isotopic tool to assess the respective role of external inputs and internal recycling processes to build up the P-PO₄ budget of the riverbed sediments in this river.

2. Materials and methods

2.1 Site description and sampling

The Redon River is a small alpine torrential tributary (i.e. a shallow high-gradient, high-energy stream with an annual average discharge of $0.5 \text{ m}^3/\text{s}$ at the outlet) of Lake Geneva (France).

The sediment dynamic of this river has been already intensively studied (Al-Charideh, 1988; Dorioz et al., 1998b; Feuillade and Dorioz, 1992; Hamid et al., 1989; Poulenard et al., 2008) making it a particularly appropriate site for the present exploratory study. The total catchment surface area is 3,300 ha with a land use that consists mainly of forests (1,100 ha) in its upland part and grasslands (1,200 ha) and cultivated farmlands (300 ha) in its lowland part. Several wetlands (20 ha) are present, while urbanized areas are concentrated near the outlet, close to the shore of Lake Geneva. The Redon River is impacted by both point and diffuse sources of P. The main point source of P is located in the lower part of the river and consists of inputs of predominantly dissolved P from a wastewater treatment plant (WWTP), whereas the main diffuse sources occur in the upper part of the catchment, consisting mainly of particulate-P from agricultural and forested lands (Fig. 1).

The soils are mainly Cambisols, formed from the weathering products of quaternary moraine material consisting of a large range of mainly calcareous sedimentary, metamorphic and igneous rocks from the Rhône glacier (Nicoud et al., 1993). The soil mineralogy is comprised of a mixture of minerals inherited from the weathering of this heterogeneous parent material and their secondary weathering products. Previous studies (Dorioz et al., 1998a, 1998b) have shown that the Redon River sediments are CaCO_3 -rich (10 to 20% CaCO_3 , $\text{pH} > 7$). Their fine ($< 200 \mu$) fraction is dominated by loamy and sandy materials, with an organic matter content between 2 to 4.5%. The clay fractions (7-11%) consist mostly of soil-derived illite, biotite and chlorite (Poulenard et al., 2008). The water in the Redon River is bicarbonated-calcic.

[Figure 1]

2.2 Sampling strategy and sample preparation

Details regarding the sampling sites and sampling conditions are given in Tables 1 and 2. Three types of samples were collected: i) eight riverbed sediments collected at six locations (A, B, C, D, E and F in Fig. 1) along a downstream gradient of land use that changes from forested to mixed agricultural/urban areas, thus representing a gradient of increasing anthropogenic pressure; this includes two samples collected in streamlets draining a small agricultural sub-catchment (locations A and B in Fig. 1) that is representative of the type of agriculture practiced in the Redon River catchment (Dorioz and Ferhi, 1994); ii) four samples representative of the potential sources of river sediments, including cultivated topsoil and deep horizon, riverbank material, and soil from a 2 ha wetland, hydraulically connected to the river; and iii) two samples from sediments from Lake Geneva representing an average reference for the local superficial lacustrine sediments (collected in 2012 and provided by the EDYTEM laboratory).

Given that it was difficult to access the river bed, it was not possible to visit all six riverbed sediment sampling locations at the same time. Three locations were first sampled at the end of March 2012 (B, C and F), and five (A, B, D, E and F) were sampled (again) at the beginning of September 2012. Both periods were low-flow periods, where the September period corresponds to the lowest weekly discharge period of the year. Samples were taken in areas of preferential accumulation of fine sediments. At each location, single samples of the entire sediment matrix were taken manually at 3-5 different points and then mixed to obtain an average sample (0.5-1 kg dry weight). Simultaneously, the electric conductivity (EC) was measured and a water sample was collected to determine the SRP (soluble reactive phosphorus) concentration and water oxygen isotope composition ($\delta^{18}\text{O}_w$). Hourly water temperatures were obtained from automatic recorders at the outlet. The riverbank sample was collected in the upstream, forested part of the catchment (see Fig. 1). The cultivated soil samples were collected in the same small

sub-catchment used to sample the streamlet sediments (Fig. 1), as this sub-catchment is known to represent zones of maximum soil erosion risk in the Redon catchment (Dorioz and Ferhi, 1994). In addition to taking a topsoil sample, a deeper soil sample was taken at a depth of approximately -70 cm to account for the possible erosion of deep soil materials along ditches or/and soil embankments. These two samples were obtained by pooling five subsamples (200-300 g) collected in different locations within a single field or its embankment. The pooled samples were then air-dried and sieved at 200 μm before analysis. No river water samples were collected for the analysis of the $\delta^{18}\text{O}_\text{p}$ values for the dissolved P-PO_4 , although these could have been useful in the interpretation of the riverbed sediment data. The decision not to sample the river water was due to the very low concentrations of dissolved P-PO_4 recorded in this river (Table 1), which makes it extremely difficult to analyze the $\delta^{18}\text{O}_\text{p}$ in the current state of the analytical protocols.

[Table 1]

[Table 2]

2.2 Sediment and soil P chemical analyses

The total phosphorus (Pt) and Olsen P (P_{Olsen}) concentrations were determined in all of the samples. P_{Olsen} (Olsen et al., 1954), which is considered as being representative of the bioavailable fraction of P in soils, has been shown to correlate linearly with P extracted from the sediment in bio-assays with algae (Jordan-Meille, 1998). The Pt concentrations were measured after a hot sodium hydroxide digestion. The total inorganic and organic P in the samples were determined according to Saunders and Williams (1955) as described by Fares et

al. (1974). The spectrophotometric determination of P-PO₄ in all of the extracts/digestates was carried out using the molybdenum blue method (Murphy and Riley, 1962) at 882.0 nm on a Cary 50 UV-Vis spectrometer (Agilent).

2.3 Isotopic analyses and data treatment

2.3.1 Extraction and purification protocol

We used the extraction and purification protocol initially developed by Tamburini et al. (2010) for soils (hereafter referred as to Tb10), which we adapted for sediments. The Tb10 protocol consists of the following six steps:

- i) extraction of phosphate from the soil/sediment with 1 M HCl, which in sequential extraction mainly targets the P-PO₄ pool (HCl-P) associated with Ca, but in our case (singular extraction) also P sorbed onto Fe and Al oxides;
- ii) dissolved organic matter (DOM) removal from the solution using an Amberlite resin,
- iii) ammonium phospho-molybdate (APM) precipitation and subsequent dissolution,
- iv) magnesium ammonium phosphate (MAP) precipitation and dissolution,
- v) cation removal by a cation exchange resin (BioRad, AG50 x8),
- vi) silver phosphate (Ag₃PO₄) precipitation through the addition of an Ag-ammine solution to the purified phosphate solution.

The goal of these steps, after HCl extraction, is to avoid sources of oxygen (i.e. DOM, nitrate, etc.) other than P-PO₄ and other contaminants, e.g. chloride, in the final silver phosphate precipitate.

We modified the Tb10 protocol in several ways as follows (Fig. 2 in the supplementary material). First, the samples were sieved at 200 µm instead of 2 mm, since this finer fraction contains the bulk of the bio-geochemically active P in river sediments (Poulenard et al., 2008). The liquid-to-solid ratio in the extraction step was modified to 100:1 instead of 50:1, since the

pre-tests showed that, with our highly carbonated sediments, the solution did not reach the targeted pH (1 ± 0.5). The quantity of sediment was adjusted to have at least 10 μ moles of P for the purification, corresponding to 10 to 20 g of sieved sediments. The DOM removal step using an Amberlite resin was not performed as the subsequent steps successfully eliminated the organic matter (see below). Finally, for samples with a high calcium carbonate content, an additional step had to be introduced to account for the formation of a calcium molybdate precipitate during the APM precipitation. To eliminate the precipitate, we repeated the subsequent MAP precipitation step twice.

The 1 M HCl extraction may hydrolyze some of the organic or condensed P (Turner et al., 2005) and the released Pi may bias the isotopic composition of the inorganic P pool targeted here. To assess the existence of such a possible artifact, extractions were run in duplicate using ^{18}O -labeled and unlabeled reagents. An ^{18}O -labelled 1M HCl solution was prepared by adding approximately 0.05 ml of ^{18}O -enriched water (97‰, Aldrich) to 1 liter of 1 M HCl solution. Labeled and unlabeled solutions were analyzed for $\delta^{18}\text{O}_w$. If hydrolysis occurred, it should result in different isotopic compositions depending on whether labeled or unlabeled reagents were used, i.e. a higher $\delta^{18}\text{O}_p$ for the labeled sample compared to the unlabeled one. In this case, we calculated the “real” isotopic value by a revised version of the mass balance equation used by McLaughlin et al. (2006b):

$$\delta^{18}\text{O}_p = \frac{(\delta^{18}\text{O}_{pl} \cdot \delta^{18}\text{O}_{wnl} - \delta^{18}\text{O}_{pnl} \cdot \delta^{18}\text{O}_{wl})}{(\delta^{18}\text{O}_{pl} - \delta^{18}\text{O}_{pnl} - \delta^{18}\text{O}_{wl} + \delta^{18}\text{O}_{wnl})} \quad (1)$$

where $\delta^{18}\text{O}_{pl}$ and $\delta^{18}\text{O}_{pnl}$ are the isotopic composition of P- PO_4 for the labeled and unlabeled samples, respectively, and $\delta^{18}\text{O}_{wl}$ and $\delta^{18}\text{O}_{wnl}$ are the isotopic composition of water for the labeled and unlabeled reagents, respectively.

The efficiency and reliability of the modified Tb10 protocol used here were evaluated by considering the repeatability of the extraction/purification process, the final Ag_3PO_4 yield and its purity. The purity was assessed by i) dissolving the obtained Ag_3PO_4 and comparing the

actual concentration of phosphate in the solution with the theoretical concentration, i.e. the amount expected if all of the solid mass was Ag_3PO_4 , and ii) determining the C and N concentration of the obtained Ag_3PO_4 on the elemental analyzer (Flash 2000, Termo Scientific); these two concentrations are used as indicators of possible contamination by organic matter and/or nitrate. The purity of the obtained Ag_3PO_4 crystals was also assessed through quantification of the oxygen yield during mass spectrometry analyses. This was done by graphically comparing the weight of the Ag_3PO_4 standards and samples introduced into the machine against the peak area of the produced CO (“oxygen yield”), which is proportional to the oxygen content in the sample. If other sources of oxygen or contaminants were present in the sample, the oxygen yield should be different from the standard Ag_3PO_4 . P- PO_4 losses at each step of the purification were also quantified (see Pistocchi et al. 2014 for more details).

2.3.2 Isotopic analyses and calculation of the theoretical equilibrium values

The oxygen isotope composition of the purified Ag_3PO_4 was measured using a thermal conversion elemental analyzer (Varioppyro Cube Elementar), coupled to an IsoPrime100 isotopic ratio mass spectrometer (IRMS) at the Plant Nutrition Group laboratory (ETH Zurich). Two to three replicates of each sample were weighed (300 to 600 μg) in silver capsules with a small added amount of glassy carbon powder. The temperature of the TC/EA furnace was kept at 1450°C for pyrolysis. During each run, repeated measurements of an internal standard (Acros Organics $\text{Ag}_3\text{PO}_4 > 97.5\%$, $\delta^{18}\text{O}_\text{p} = +14.2\text{‰}$) and two international benzoic acid standards (IAEA 601 and IAEA 602, $\delta^{18}\text{O} = +23.3\text{‰}$ and $+71.4\text{‰}$, respectively, IAEA, Vienna, Austria) were included and used to calibrate and correct the instrumental drift. The $\delta^{18}\text{O}_\text{p}$ values presented in this paper are reported in the standard δ notation with respect to VSMOW (Vienna Standard Mean Oceanic Water), i.e. as ‰ deviation values. Analytical uncertainties, as determined from the replicate analysis of the standards, were estimated to be below 0.3‰.

In order to assess the extent to which the sedimentary P-PO₄ pools investigated in this study could be secondary recycled P-PO₄ (i.e. P-PO₄ that has been cycled by the river biota), a theoretical equilibrium $\delta^{18}\text{O}_p$ value – i.e. the oxygen isotopic composition the phosphate should have if it is in equilibrium with the ambient water – was calculated for each sample using the following equation (Chang and Blake, 2015):

$$\delta^{18}\text{O}_p = (\delta^{18}\text{O}_w + 10^3) \cdot e^{\frac{\left[\left(14.43 \cdot \frac{10^3}{T}\right) - 26.54\right]}{10^3}} - 10^3 \quad (2)$$

where $\delta^{18}\text{O}_p$ is the calculated oxygen isotopic composition of the P-PO₄ sample at equilibrium in ‰, $\delta^{18}\text{O}_w$ is the measured oxygen isotopic composition of the water in ‰, and T is the temperature in degrees Kelvin.

Considering the stability of the $\delta^{18}\text{O}_w$ values during low-flow periods in the Redon River due to the dominance of the ground water contribution to the discharge during these periods (Al-Charideh, 1988), we used the $\delta^{18}\text{O}_w$ of the river water measured at each sampling date as the reference isotopic value for water in equation 2. We considered both the minimum and maximum temperatures measured over the period running between the sampling date and the closest big storm flow event (around 10-20 days; Table 1 in the Supplementary Material), since the sediments are (partially) renewed at each storm flow. The temperatures measured at the time of sampling fell within these ranges. For samples 9 to 12 (sources), we used the $\delta^{18}\text{O}_w$ values measured in the local water sources, including rain, groundwater and soil water data, and the extremes for the soil temperature measured within the previous 10 days at an agro-meteorological station close to the studied sites. This allowed us to have a conservative and time-integrated estimate of the equilibrium interval. For these samples, we also calculated the deviation from equilibrium using the average stream water temperature and isotopic composition, to verify whether or not their $\delta^{18}\text{O}_p$ values would fall within the equilibrium range once transferred into the river. For each sample, the deviation from equilibrium was calculated

as the interval (in ‰) between the measured $\delta^{18}\text{O}_\text{p}$ values and the closest values for the equilibrium range calculated from equation 2

3. Results

3.1 Method validations

Repeated extractions and purifications ($n = 3$ or 4) on the same sediment sample yielded $\delta^{18}\text{O}_\text{p}$ values with a standard deviation < 0.4 ‰. This repeatability can be considered as acceptable, since it is within the range of the precision obtained on the repeated standards measurements on the TC/EA-IRMS.

The purity of Ag_3PO_4 was 97%, as determined by analyzing the P- PO_4 in the redissolved Ag_3PO_4 (see section 2.3.1, Table 2 in the Supplementary Material). The carbon and nitrogen concentrations were generally lower than 1% and 0.1%, respectively (Table 2 in the Supplementary Material). The C content was comparable to that measured in the internal standard Ag_3PO_4 (0.5%) and much lower than the 23-29% or 0.2% reported by Weiner et al. (2011) for C and N, respectively. The efficiency of the purification procedure was confirmed by the oxygen yield given by the mass spectrometer (Fig. 1 in the Supplementary Material). All of the samples were close to the regression line defined by the Ag_3PO_4 standards, indicating that the amount of oxygen in the samples was close to the stoichiometry of the pure compound. Most of the labeled/unlabeled pairs did not show significantly different $\delta^{18}\text{O}_\text{p}$ values. Only four samples (2, 5, 8 and 9) showed different labeled and unlabeled values, meaning that the organic P or condensed phosphate was hydrolyzed during the HCl extraction. These values have been corrected with equation 1, as all of the other values are the average of the labeled and unlabeled results.

3.2 P speciation

Table 3. P speciation and isotopic data

The P content and speciation varied strongly from one sample to another, with the Pt value ranging from 459 to 1256 mgP/kg, Pi from 179 to 631 mgP/kg and Po from 36 to 450 mgP/kg. P_{Olsen} accounted for 0.7 to 10.6 % of Pt depending on the samples. The highest P_{Olsen} concentrations (65-67 mgP/kg, representing 8.4 to 10.6% of Pt) were found in the river sediments sampled immediately downstream from the WWTP discharge. These variations are the consequence of differences in the source, erosion and deposition conditions of the sediments as well as differences in the P inputs that accumulate along the river (Table 1). The downstream sediments (samples 5 to 8) and streamlet sediments (samples 1 and 2) were enriched in Pt, Pi and P_{Olsen} compared to the sediments coming from the head of the catchment (samples 3) or to the riverbank material (sample 12). The same relative enrichment is observed for the cultivated topsoil (sample 10) compared to its lower horizon (sample 11). Pt and P_{Olsen} were weakly and positively correlated ($r^2 = 0.49$, $p < 0.01$) as well as the P_{Olsen}/Pi ratio with Pt concentrations of $r^2 = 0.76$ ($p < 0.01$), providing evidence that the P enrichment process recorded by the downstream sediments is accompanied by an increase in the most reactive fraction of the total P- PO_4 pool. The highest Po concentrations were found in the topsoil samples (9 and 10).

3.3. Measured isotopic signatures and comparison with the equilibrium values

The measured $\delta^{18}O_p$ values for the HCl extracts varied between +11.8‰ (deep soil horizon, sample 11) and +18.3‰ (wetland soil, sample 9; Fig. 2a). No significant statistical relationships were found between the $\delta^{18}O_p$ values for the samples and their Pt, P_{Olsen} or Pi content or ratios, nor with the water characteristics, notably the SRP concentrations. The

riverbank sample collected in the head of the catchment (sample 12) corresponds to a weakly weathered moraine, as confirmed by field observations, i.e. no visible weathering and no vegetation or biofilm on the banks, since they are eroded at each storm flow, and by the P speciation data (very low P_{Olsen} and P_o fractions; see Table 3). This sample is representative of both the principal source of the upstream, unpolluted sediments and the parent material of most of the soils in this catchment (CIPEL, 1988). Its $\delta^{18}\text{O}_p$ value (+15.9‰) indicates that it has a mixed igneous/sedimentary origin, as this value is intermediate between the $\delta^{18}\text{O}_p$ values reported for igneous or metamorphic apatite (approximately +7‰; Tamburini et al., 2012) and for sedimentary apatite (between +17 and +23‰; Tamburini et al., 2014). Interestingly, the variability in the $\delta^{18}\text{O}_p$ values for the allochthonous sources that may possibly contribute toward the river particulate-P load was higher than the variability in the riverbed sediments. Whereas while the soil, riverbank and riverbed sediments collected in the head of the catchment (samples 1 to 3 and 9 to 12) are representatives of these sources (Dorioz and Ferhi, 1994) and exhibit $\delta^{18}\text{O}_p$ values varying by > 6 ‰ (from +11.8 to +18.3‰), this variation is reduced to approximately 2‰ for the sediments immediately upstream/downstream the WWTP and to 1‰ for the outlet sediments (+12.8 to +13.8‰; Fig. 2a), the $\delta^{18}\text{O}_p$ signatures of which were comparable to those recorded by the lake sediments (samples 13 and 14; Table 3).

[Figure 2]

We compared the measured $\delta^{18}\text{O}_p$ values with the theoretical equilibrium calculated with Equation 2 (see Table 2). The deviation between the calculated and measured $\delta^{18}\text{O}_p$ values is presented in Fig. 2b. The $\delta^{18}\text{O}_p$ value for the riverbank moraine sample (sample 12) was above the calculated equilibrium by ca. 2‰, as was the wetland soil sample (9). This was not the case for the cultivated soil samples that showed $\delta^{18}\text{O}_p$ values that were either very close to (-0.1‰,

topsoil, sample 10) or slightly lower than (-0.7‰, deep horizon, sample 11) the calculated equilibrium values.

The riverbed sediment samples collected in the streamlets of the agricultural sub-catchment (samples 1, 2) exhibited $\delta^{18}\text{O}_\text{p}$ values that were lower than the calculated equilibrium values by ca. 2‰. For the sediments located within the main hydrogeological network, the differences between the measured and calculated equilibrium values generally tended to be lower (see Table 3 and Fig. 2b): ca. $\pm 1.5\text{‰}$ in the middle part, upstream and downstream from the WWTP discharge point (samples 4 to 6) and $< 0.5\text{‰}$ for the sediments collected at the outlet (samples 7 and 8) and the lacustrine sediments (samples 13 and 14).

4. Discussion

4.1 Significance and variability of the measured and equilibrium $\delta^{18}\text{O}_\text{p}$ values

The present study shows that the method originally established and optimized by Tamburini et al. (2010) for soils is suitable for CaCO_3 -rich riverbed sediment samples, following some modifications to the methodology. However, as for soils, it should be kept in mind that the extraction is not entirely specific with regards to the nature of the extracted P-PO_4 . First, 1 M HCl can be used to extract the fraction that is primarily composed of Ca-bound P, which can contain a broad spectrum of P compounds (more or less crystallized apatite forms, CaCO_3 -bound P, etc.). This reagent can also extract some P-PO_4 sorbed onto Fe and Al oxides, if used as a single extractant. Thus, the $\delta^{18}\text{O}_\text{p}$ signatures obtained with this method correspond to a pooled P-PO_4 isotope signature of a mixture of different P fractions with a different degree of stability, exchangeability and bioavailability (Jarvie et al., 2002) rather than to the signature of a geochemically specific P-PO_4 compartment. Despite this limitation, the oxygen isotope composition of P-PO_4 extracted from sediment using HCl can provide information that is not

available with other methods, including the extent to which a P pool has been biologically processed or derived from primary inputs or secondary anthropogenic sources (Tamburini et al., 2014; Davis et al., 2014).

The calculation of the equilibrium $\delta^{18}\text{O}_\text{p}$ values is another critical point for the interpretation of the $\delta^{18}\text{O}_\text{p}$ isotope data. Information regarding the kinetics of the equilibrium is only derived from cell-free or soil incubation experiments (Blake et al., 2005; Melby et al., 2013), but not from conditions that are close to the natural state. Our strategy was to take samples during low-flow periods because of the stability of the $\delta^{18}\text{O}_\text{w}$ value and the temperature, but there is still the question of how to choose appropriate time points when calculating the theoretical equilibrium values with equation 2. This is also relevant for soils where sharp variations in the $\delta^{18}\text{O}_\text{w}$ value in the soil surface layers due to the progressive evaporation of the soil water (Hsieh et al., 1998) might be a major source of uncertainty. Amelung et al. (2015) suggest calculating the equilibrium $\delta^{18}\text{O}_\text{p}$ value with the annual average $\delta^{18}\text{O}_\text{w}$ for rainwater, arguing that the biological recycling of Ca-bound P in soil is a long-term process. Using an annual rainwater $\delta^{18}\text{O}_\text{w}$ value and temperature values, the calculated equilibrium value for the agricultural soil samples (10 and 11) would increase by approximately +2‰, resulting in measured signatures below the equilibrium.

Except for a small number of studies (Elsbury et al., 2009; Gooddy et al., 2016; Markel et al., 1994; McLaughlin et al., 2006a), freshwater ecosystems have been poorly explored in terms of $\delta^{18}\text{O}_\text{p}$. Our results expand the global $\delta^{18}\text{O}_\text{p}$ library for these systems, comparing, for the first time, the oxygen isotope signatures of HCl-P extracted from river sediment samples from upstream to downstream with those of the main potential allochthonous sources in the catchment. The isotope signatures of the analyzed river sediments, ranging from +12.2 to +15.8‰, lie in the interval reported in previous studies, i.e. +8.6 to 22.5 (Markel et al., 1994),

+12.4 to 23.5‰ (Davies et al., 2014) and +9.7 to +20.0‰ (Tamburini et al., 2014). The same is true for our soil samples (Amelung et al., 2015; Tamburini et al., 2014; Zohar et al., 2010).

4.2 Biological recycling versus tracing the sources using $\delta^{18}\text{O}_p$ values

4.2.1 Soils

Tamburini et al. (2012, 2014) have shown that the HCl-P fraction in the upper and most biologically active part of the oldest soil of a chronosequence was formed by secondary minerals precipitated from Pi that were previously cycled by biota. The close-to-equilibrium $\delta^{18}\text{O}_p$ values measured in the agricultural soil samples (samples 10 and 11) can be viewed as evidence that biological cycling had a large impact on the HCl-P fraction (which represents roughly half of the Pt) in these samples. The soil P is supplied by the parent material and fertilizations and is redistributed throughout the soil profile via biological activities. As the parent material is low in Pt, P_{Olsen} and Po, the high concentrations observed in the topsoil sample (see Table 3), as in other cultivated soils, is the long-term result of the accumulation and transformation of P favoring a high P bioavailability, which supports the assumption of an intense biological activity.

The most common fertilizers used in the area have $\delta^{18}\text{O}_p$ values that are higher (16.2-22.5‰) or lower (9.5‰; Guyard 2010) than the values for the studied agricultural soils. Even if we cannot exclude that the observed equilibrium values were only the result of a mixing between the parent material (close to +15.9‰) and a lighter fertilizer input, we assume that the fertilizer has little influence because the fertilizer inputs are almost totally cycled over the long-term (> 1 year) and the amount applied is too small (2-3% of the HCl-P soil stock) over the short-term (one season).

The high and above-equilibrium $\delta^{18}\text{O}_p$ value in the wetland surface soil (sample 9) can also reflect the impact of biota, e.g. plants. The $\delta^{18}\text{O}_p$ value for this sample (18.3‰) cannot be

explained by the mixing of any of the potential input materials present in the area (i.e. bank material or cultivated soil), as +18.3‰ is too far outside the range of the values (11.8 and 15.8‰) recorded by these input materials. The only source with a sufficiently high $\delta^{18}\text{O}_\text{p}$ value would be plant-derived phosphate. Phosphate in plants is enriched in ^{18}O , with the $\delta^{18}\text{O}_\text{p}$ values ranging from 28 to 31‰ (Pfahler et al., 2013; Tamburini et al., 2014). This suggests that the extracted fraction might partially originate from the release of P from plant litter, either directly or via decomposition. A significant fraction of this enriched P pool that is not taken up by microbes (Liang and Blake, 2006; Oberson and Joner, 2005) might have precipitated or been sorbed into the soil mineral pool. Interestingly, this sample had a particularly high Po concentration. Preferential sorption/desorption reactions with soil minerals could also have helped increase the $\delta^{18}\text{O}_\text{p}$ value for this sample. However, previous studies have reported both negative and positive fractionation associated with sorption (Jaisi et al., 2010; Melby et al., 2013b). In any case, this effect is probably too small (<1 ‰) to justify the observed value by itself.

When compared to the equilibrium values calculated using the average river conditions (the values in brackets in Table 3, Fig. 2b), the subsoil/embankment sample and the wetland topsoil are no longer found within the equilibrium range. The situation is different for the agricultural topsoil sample, the actual $\delta^{18}\text{O}_\text{p}$ value of which falls within the equilibrium range, meaning that under these conditions the oxygen isotopic composition alone would not help to distinguish between a sedimentary P- PO_4 pool inherited from this soil and in-stream recycled P- PO_4 .

4.2.2 River sediments upstream and downstream from the wastewater treatment plant discharge

The September sampling showed that the input of P from wastewater was marked by a threefold increase in the SRP concentration in the river water sampled upstream and

downstream from the discharge point (Table 1), accompanied by a drastic change in the P_t , P_i and P_{Olsen} concentrations in the corresponding sediments (Table 3). No changes were detected in the P_o content, meaning that the impact of the point discharge on this pool in the bed sediments was negligible (Dorioz et al., 1998a). Interestingly, despite the small distance (< 100 m) separating the upstream and downstream sites, the $\delta^{18}O_p$ values were also distinct, with a difference of 3‰. Upstream from the WWTP discharge point, the P pools in the sediment are mainly impacted by sources comprising a mixture of riverbanks and agricultural soil erosion (materials similar to samples 12 and 1, 2). The $\delta^{18}O_p$ value (12.2‰) and P speciation for sample 4 were intermediate between those of these source materials.

The two September samples (4 and 5, immediately upstream and downstream from the WWTP) deviate from the isotopic equilibrium values (by -1.5 and +1.4‰, respectively; Fig. 2b), suggesting the existence of conditions that prevent the complete biological turnover of P at both sites (also see Gooddy et al., 2016). Under the assumption that biological equilibration is the dominant process during sewage treatment (Gruau et al., 2005), the $\delta^{18}O_p$ value for dissolved P_i from the WWTP would vary between 13.0 and 14.8‰ using equation 2 (confirmed by the value of 14.0‰ measured by Guyard, 2010), meaning that the higher $\delta^{18}O_p$ value downstream (15.8‰) cannot be interpreted as the result of a mixing between wastewater derived P and P from the upstream sediments. One possible explanation for this could be that part of the P_i injected into the river by the WWTP is rapidly (within a few meters from the injection point) and continuously sorbed onto the sediment particles, as also hypothesized by Dorioz et al. (1998a), and that the heavier isotopologues are preferentially sorbed. However, there is no consensus regarding the associated fractionation (Jaisi et al., 2010; Melby et al., 2013b). An alternative explanation would be that the upstream sample 4 represents a local anomaly, and the signature of the downstream sediments, sample 5, derives from local riverbank erosion. Sample 12 can be considered as representative of riverbank material and its $\delta^{18}O_p$ value is comparable

to that of sample 5. However, this hypothesis is unlikely as it does not explain the drastic change in P speciation observed in the sediments downstream from the WWTP (see Table 3). Only the locality situated immediately downstream from the WWTP discharge point could be sampled in March 2012 (sample 6). The SRP, Pt, Pi and P_{Olsen} concentrations were as high as those found in sample 5, which was collected in September, and much higher than the concentrations measured further upstream, at location 3, once again providing evidence for the strong impact of the WWTP discharge on the river and sediment P pool. The $\delta^{18}\text{Op}$ value for sample 6 was similar to the September sample (sample 5), although it was close to the calculated isotopic equilibrium. The lack of a reference value equivalent to sample 4 in March means that it is not possible to conclude whether the observed value resulted from the complete equilibration of the sedimentary P- PO_4 pool or if it was instead produced by selective sorption processes as we suggest for sample 5. A repeated seasonal sampling of the sediments immediately upstream and downstream from the WWTP could possibly clarify this point.

4.2.3 Head and outlet sediments

The possible mixing of isotopically different sediment source materials coupled with differences in the sediment retention time complicates the interpretation of the $\delta^{18}\text{Op}$ values for the riverbed sediments. Under conditions of low biological activity and a short residence time, the isotopic signature of the sediments is expected to represent primary source signatures. In this context, an isotopic disequilibrium would indicate the existence of factors that limit the in-stream P cycling, intended as uptake by the biota, intracellular equilibration and subsequent release. As judged from the present data, a limiting situation for in-stream P recycling could have been prevalent in the head of the Redon River catchment, where strong out-of-equilibrium $\delta^{18}\text{Op}$ signatures have been recorded (see Table 3 and Fig. 2b). This could be first related to the physical conditions in this part of the catchment - high-energy stream (total renewal of the

sediments with each storm flow) and a short distance from the sources, i.e. riverbanks - which both determined the short residence times. The oligotrophic state of the running water ($\text{SRP} < 10 \mu\text{g P/l}$, $\text{NO}_3 < 0.5 \text{ mg N/l}$) and the low P bioavailability of the riverbed sediments (as indicated by low P_{Olsen} concentrations) could contribute to limit the development of benthic biota and thus P uptake and subsequent release. For the same reasons the contribution of extracellular mineralization of organic P is expected to be negligible. The lack of an isotopic difference and the small differences in the Pt, Pi and Po content between the riverbank sample (12) and the sediment sample (3) collected immediately downstream further confirms the direct filiation between the source and the sediment.

The hypothesis that a rapid transfer does not significantly modify the $\delta^{18}\text{O}_\text{p}$ signature is further supported by the disequilibrium signature recorded in the streamlet samples 1 and 2 and their Pi, Po and P_{Olsen} concentrations, as all of these parameters are intermediate between the values found in the analyzed deep and topsoil samples. This suggests i) a mixed origin for these two sediment samples derived from both superficial soils and ditch banks or field embankments; and ii) that the biological activity was not intense or not long enough to equilibrate the original P oxygen isotope signatures, consistently with the expected very short residence time of the sediments in the streamlets (stream order of 1).

The situation was different for the sediments that were collected closer to the river outlet or in the lake (stream order higher than 3). Close-to-equilibrium values were found in these samples. A corresponding increase in the SRP concentration in running water (a factor of 8 between upstream and the outlet; see Table 1), as the result of the cumulative anthropogenic inputs in the river, was also observed.

The equilibrium values observed in the lowest part of the Redon River suggest the occurrence of processes that are able to efficiently recycle the sedimentary P- PO_4 . This hypothesis would be consistent with the decrease in the $\delta^{18}\text{O}_\text{p}$ values observed during the September campaign

downstream from the WWTP discharge point, i.e. between samples 5 and 8. There was a 3‰ decline in the $\delta^{18}\text{O}_\text{P}$ value between sample 5, which was collected immediately downstream from the WWTP discharge point, and sample 8, which was collected approximately 2 km further downstream. This decrease was accompanied by a strong narrowing of the gap between the measured and calculated equilibrium $\delta^{18}\text{O}_\text{P}$ values. Whereas this gap was +1.4‰ for sample 5, it was reduced to < 0.4‰ for sample 8 (see Table 3). The isotopic equilibration inferred by these data would likely occur in the sediments, given that in this kind of torrential rivers most of the biological activity essentially takes place in the benthic system. Equilibration could be further promoted by the longer residence time of the sediments in the downstream part of the river (Doriot et al., 1989). It may have been also driven by the enhanced bio-availability of P in the lower part of the river, notably downstream from the WWTP, as evidenced by the sharp increase in the P_{Olsen} concentration starting from this point (see Table 3, Figs. 2a and 2b). Most likely, the overall process should be viewed more as the integration of a succession of partial equilibrium occurring over different storage stages under similar equilibrium conditions, as it is unlikely that the bulk of the P-PO_4 could be completely cycled by the benthic biota during a single low-flow period. However, we lack information about the real residence time of the sediments and the P recycling rates in the Redon River. Considering that the measured $\delta^{18}\text{O}_\text{P}$ values for the sediments collected near the river outlet (samples 7 and 8) are identical within error to those measured in the upstream streamlet sediments (samples 1 and 2) and are within the range of the values found in the possible sediment sources (see Fig. 2a), we cannot entirely exclude that the $\delta^{18}\text{O}_\text{P}$ values in these downstream sediments are simply source-inherited isotopic values. More direct evidence for the real residence time of the sediments and the ability of aquatic biota to efficiently recycle P-PO_4 in the downstream part of the river is needed to decide between these two hypotheses.

4.3 Potential and limits of the oxygen isotopes tool

We hypothesize that is possible to trace the potential sources of sediment particulate-P-PO₄ using oxygen isotopes, assuming a limited number of potential sources and under conditions of low biological activity and a short residence time in the river network. In the studied torrential system, it appeared possible to trace these sources in the head of the catchment (only two sources, rapid and high-energy transfer, limiting conditions for the biological activity). It would be also possible at the catchment scale, during winter high-flow periods, when the sediments that originate from banks and agricultural soil erosion are rapidly transferred as suspended matter and without significant interaction with point source discharge and biological activity. In this case, the $\delta^{18}\text{O}_\text{p}$ value would primarily be the result of a two end-members mixing between the cultivated soil (values close to 11-14‰) and moraine material (values close to 15-16‰). However, the isotopic tracing is expected to be less straightforward during the summer and early fall periods, as the storm flows that occur at that time primarily mobilize riverbed sediments. Therefore, it could become more difficult to perform a quantitative assessment of the original particulate-P sources using $\delta^{18}\text{O}_\text{p}$ due to an overprinting of the sediment isotopic signature induced by the high biological activity. During summer low flows, an even stronger overprinting of the original isotopic signature is expected to occur in the downstream part of the river, due to the input of anthropogenic, bioavailable P, a longer sediment residence time and increased biological activity. This being the case, $\delta^{18}\text{O}_\text{p}$ could then instead be used as a tool to assess the in-stream biological turnover of the P.

5. Conclusion

- In this study, we successfully adapted Tamburini et al.'s method (2010) for soil for use with river sediments to provide the first assessment of the variability in $\delta^{18}\text{Op}$ from HCl-extracted P in river sediments along a gradient of increasing anthropogenic impact.
- We observed a contrast between the $\delta^{18}\text{Op}$ signature and the speciation of the sediments collected in the head and at the outlet of the catchment. The sediments that were collected upstream have isotopic signatures that are far from isotopic equilibrium and within the range of the signatures found in the possible sediment sources (i.e. riverbank material), while those collected at the outlet showed $\delta^{18}\text{Op}$ values close to isotopic equilibrium. We related this difference to the contrast between the highly dynamic character (short residence time) of the transfer system in the head of the catchment (preservation of the isotopic signature of the source material in the sediments) and the “quieter” sediment transport system in the lower catchment, which, along with the input of the bioavailable anthropogenic P, likely fostered the recycling of P by benthic biota, leading to an equilibration in the $\delta^{18}\text{O}$ signatures. However, with the available data, we cannot exclude that the signatures at the outlet were the result of the mixing of sediments with higher (upstream/wetland sediments) and lower signatures (agricultural soil erosion).
- Uncertainties remain about the calculation of the equilibrium window. For aquatic environments, the duration of the in-stream equilibration processes should be taken into account, together with the perturbation caused by changes in the hydrological regime. This would require an experimental approach for a proper estimation.
- In terms of the quantitative tracing of potential particulate-P sources, future investigations should focus on winter high flows, when rapid sediment transfer maximally reduces the possibility of biological interactions. On the other hand, the sensitivity of $\delta^{18}\text{Op}$ to biological processes opens up new promising perspectives for studying in-stream processes. In

situ labelling experiments could help develop these studies by providing unique information about the biological uptake, turnover and/or organic P mineralization.

- More insights about the way and extent the biological activity controls P speciation in the soil and sediments could be possibly provided by the development of refined extraction methods that are able to selectively and quantitatively extract other P pools like the one associated with Al and Fe hydroxides, P_{Olsen} and P_o or finer fractions (e.g. $< 50 \mu\text{m}$, $< 2 \mu\text{m}$).

Acknowledgments

This work was funded by the EFPA department of INRA. The authors thank Philippe Quetin, Pascal Perney and Danielle Lacroix (INRA, UMR CARTELE) for their precious help and Jerome Lazzarotto and Liliane Savoye for their advice and fruitful discussions. The authors acknowledge the EDYTEM Laboratory and in particular J.P. Jenny for providing the sediment samples from Lake Geneva.

References

- Al-Charideh, A.R., 1988. Contribution à l'étude hydrologique du bassin du Redon, Haute-Savoie : relation pluie-débit et quelques aspects isotopiques (PhD Thesis). Université Pierre et Marie Curie, Paris.
- Amelung, W., Antar, P., Kleeberg, I., Oelmann, Y., Lücke, A., Alt, F., Lewandowski, H., Pätzold, S., Barej, J.A.M., 2015. The $\delta^{18}\text{O}$ signatures of HCl-extractable soil phosphates: methodological challenges and evidence of the cycling of biological P in arable soil: The $\delta^{18}\text{O}$ signatures of HCl-extractable soil phosphates. *European Journal of Soil Science* 66, 965–972. doi:10.1111/ejss.12288
- Blake, R.E., O'Neil, J.R., Surkov, A.V., 2005. Biogeochemical cycling of phosphorus: insights from oxygen isotope effects of phosphoenzymes. *American Journal of Science* 305, 596–620.
- Burrus, D., Thomas, R.L., Dominik, J., Vernet, J.-P., 1989. Recovery and concentration of suspended solids in the upper rhone river by continuous flow centrifugation. *Hydrol. Process.* 3, 65–74. doi:10.1002/hyp.3360030107
- Chang, S.J., Blake, R.E., 2015. Precise calibration of equilibrium oxygen isotope fractionations between dissolved phosphate and water from 3 to 37°C. *Geochimica et Cosmochimica Acta* 150, 314–329. doi:10.1016/j.gca.2014.10.030

- CIPEL, 1988. Rapport de l'étude des pollutions diffuses dans le bassin lémanique. Commission Internationale pour la Protection des Eaux du Léman, Lausanne.
- Davies, C.L., Surridge, B.W.J., Gooddy, D.C., 2014. Phosphate oxygen isotopes within aquatic ecosystems: Global data synthesis and future research priorities. *Science of The Total Environment* 496, 563–575. doi:10.1016/j.scitotenv.2014.07.057
- Dorioz, J.M., Cassell, E.A., Orand, A., Eisenman, K.G., 1998a. Phosphorus storage, transport and export dynamics in the Foron River watershed. *Hydrol. Process.* 12, 285–309. doi:10.1002/(SICI)1099-1085(199802)12:2<285::AID-HYP577>3.0.CO;2-H
- Dorioz, J.M., Ferhi, A., 1994. Pollution diffuse et gestion du milieu agricole: Transferts compares de phosphore et d'azote dans un petit bassin versant agricole. *Water Research* 28, 395–410. doi:10.1016/0043-1354(94)90277-1
- Dorioz, J.M., Pelletier, J.P., Benoit, P., 1998b. Physico-chemical properties and bioavailability of particulate phosphorus of various origin in a watershed of lake Geneva. *Water Research* 2, 275–286.
- Dorioz, J.M., Pilleboue, E., Ferhi, A., 1989. Dynamique du phosphore dans les bassins versants: Importance des phenomenes de retention dans les sediments. *Water Research* 23, 147–158. doi:10.1016/0043-1354(89)90038-9
- Elsbury, K.E., Paytan, A., Ostrom, N.E., Kendall, C., Young, M.B., McLaughlin, K., Rollog, M.E., Watson, S., 2009. Using Oxygen Isotopes of Phosphate To Trace Phosphorus Sources and Cycling in Lake Erie. *Environmental Science & Technology* 43, 3108–3114. doi:10.1021/es8034126
- Ensign, S.H., McMillan, S.K., Thompson, S.P., Piehler, M.F., 2006. Nitrogen and phosphorus attenuation within the stream network of a coastal, agricultural watershed. *J. Environ. Qual.* 35, 1237–1247. doi:10.2134/jeq2005.0341
- Fares, F., Fardeau, J.C., Jacquin, F., 1974. Quantitative survey of organic phosphorus in different soil types. *Phosphorus in agriculture*.
- Feuillade, M., Dorioz, J.M., 1992. Enzymatic release of phosphate in sediments of various origins. *Water Research* 26, 1195–1201. doi:10.1016/0043-1354(92)90180-C
- Goldhammer, T., Brunner, B., Bernasconi, S.M., Ferdelman, T.G., Zabel, M., 2011. Phosphate oxygen isotopes: Insights into sedimentary phosphorus cycling from the Benguela upwelling system. *Geochimica et Cosmochimica Acta* 75, 3741–3756. doi:10.1016/j.gca.2011.04.006
- Gooddy, D.C., Lapworth, D.J., Bennett, S.A., Heaton, T.H.E., Williams, P.J., Surridge, B.W.J., 2016. A multi-stable isotope framework to understand eutrophication in aquatic ecosystems. *Water Research* 88, 623–633. doi:10.1016/j.watres.2015.10.046
- Gruau, G., Legeas, M., Riou, C., Gallacrier, E., Martineau, F., Hénin, O., 2005. The oxygen isotope composition of dissolved anthropogenic phosphates: a new tool for eutrophication research? *Water Research* 39, 232–238. doi:10.1016/j.watres.2004.08.035
- Guyard, N., 2010. Contribution des méthodes isotopiques à la détermination des origines du phosphore apporté au Lac Léman (Master Thesis). Université de Savoie.
- Hamid, S., Dray, M., Ferhi, A., Dorioz, J.M., Normand, M., Fontes, J.C., 1989. Etude des transferts d'eau à l'intérieur d'une formation morainique dans le bassin du Léman-Transferts d'eau dans la zone non saturée. *Journal of Hydrology* 109, 369–385.
- Hsieh, J.C.C., Chadwick, O.A., Kelly, E.F., Savin, S.M., 1998. Oxygen isotopic composition of soil water: Quantifying evaporation and transpiration. *Geoderma* 82, 269–293. doi:10.1016/S0016-7061(97)00105-5
- Jarvie, H.P., Neal, C., Williams, R.J., Neal, M., Wickham, H.D., Hill, L.K., Wade, A.J., Warwick, A., White, J., 2002. Phosphorus sources, speciation and dynamics in the lowland eutrophic River Kennet, UK. *Science of The Total Environment, Water quality functioning of lowland permeable catchments: inferences from an intensive study of the River Kennet and upper River Thames* 282–283, 175–203. doi:10.1016/S0048-9697(01)00951-2

- Jordan-Meille, L., 1998. Mode de transfert du phosphore d'origine diffuse dans un petit bassin rural lémanique (PhD Thesis). EPFL, Lausanne.
- Kleinman, P.J.A., Sharpley, A.N., McDowell, R.W., Flaten, D.N., Buda, A.R., Tao, L., Bergstrom, L., Zhu, Q., 2011. Managing agricultural phosphorus for water quality protection: principles for progress. *Plant Soil* 349, 169–182. doi:10.1007/s11104-011-0832-9
- Kronvang, B., Vagstad, N., Behrendt, H., Bøgestrand, J., Larsen, S.E., 2007. Phosphorus losses at the catchment scale within Europe: an overview. *Soil Use and Management* 23, 104–116. doi:10.1111/j.1475-2743.2007.00113.x
- Liang, Y., Blake, R.E., 2006. Oxygen isotope signature of Pi regeneration from organic compounds by phosphomonoesterases and photooxidation. *Geochimica et Cosmochimica Acta* 70, 3957–3969. doi:10.1016/j.gca.2006.04.036
- Longinelli, A., Nuti, S., 1973. Revised phosphate-water isotopic temperature scale. *Earth and Planetary Science Letters* 19, 373–376. doi:10.1016/0012-821X(73)90088-5
- Markel, D., Kolodny, Y., Luz, B., Nishri, A., 1994. Phosphorus cycling and phosphorus sources in Lake Kinneret: tracing by oxygen isotopes in phosphate. *Israel Journal of Earth Sciences* 43, 165–178.
- McLaughlin, K., Cade-Menun, B.J., Paytan, A., 2006a. The oxygen isotopic composition of phosphate in Elkhorn Slough, California: A tracer for phosphate sources. *Estuarine, Coastal and Shelf Science* 70, 499–506. doi:10.1016/j.ecss.2006.06.030
- McLaughlin, K., Paytan, A., Kendall, C., Silva, S., 2006b. Oxygen isotopes of phosphatic compounds—Application for marine particulate matter, sediments and soils. *Marine Chemistry* 98, 148–155. doi:10.1016/j.marchem.2005.09.004
- Melby, E.S., Soldat, D.J., Barak, P., 2013. Biological decay of ^{18}O -labeled phosphate in soils. *Soil Biology and Biochemistry* 63, 124–128. doi:10.1016/j.soilbio.2013.03.020
- Mellander, P.-E., Jordan, P., Shore, M., Melland, A.R., Shortle, G., 2015. Flow paths and phosphorus transfer pathways in two agricultural streams with contrasting flow controls. *Hydrol. Process.* 29, 3504–3518. doi:10.1002/hyp.10415
- Moorleghem, C.V., Schutter, N.D., Smolders, E., Merckx, R., 2013. The bioavailability of colloidal and dissolved organic phosphorus to the alga *Pseudokirchneriella subcapitata* in relation to analytical phosphorus measurements. *Hydrobiologia* 709, 41–53. doi:10.1007/s10750-013-1442-8
- Nicoud, G., Coddet, E., Blavoux, B., Dray, M., 1993. Les complexes détritiques de marge glaciaire active du Bas Chablais oriental : implications hydrogéologiques. *Quaternaire* 4, 69–76. doi:10.3406/quate.1993.1994
- Oberson, A., Jöner, E.J., 2005. Microbial turnover of phosphorus in soil, in: *Organic Phosphorus in the Environment*. CABI, Wallingford, pp. 133–164.
- Olsen, S.R., Watanabe, F.S., Cosper, H.R., Larson, W.E., Nelson, L.B., 1954. Residual phosphorus availability in long-term rotations on calcareous soils. *Soil Science* 78, 141–152.
- Pacini, N., Gächter, R., 1999. Speciation of riverine particulate phosphorus during rain events. *Biogeochemistry* 47, 87–109. doi:10.1007/BF00993098
- Pfahler, V., Dürr-Auster, T., Tamburini, F., M. Bernasconi, S., Frossard, E., 2013. ^{18}O enrichment in phosphorus pools extracted from soybean leaves. *New Phytologist* 197, 186–193. doi:10.1111/j.1469-8137.2012.04379.x
- Pistocchi, C., Silvestri, N., Rossetto, R., Sabbatini, T., Guidi, M., Baneschi, I., Bonari, E., Trevisan, D., 2012. A Simple Model to Assess Nitrogen and Phosphorus Contamination in Ungauged Surface Drainage Networks: Application to the Massaciuccoli Lake Catchment, Italy. *Journal of Environment Quality* 41, 544. doi:10.2134/jeq2011.0302
- Pistocchi C., Tamburini F., Savoye L., Sebilo M., Baneschi I., Lacroix D., Perney P., Dorioz J.M., 2014. Développement d'une méthode d'extraction et purification des phosphates à partir

- de matrices sédimentaires pour l'analyse isotopique de leur oxygène. *Le Cahier des Techniques de l'INRA* 81,1.
- Poulenard, J., Dorioz, J.-M., Elsass, F., 2008. Analytical Electron-Microscopy Fractionation of Fine and Colloidal Particulate-Phosphorus in Riverbed and Suspended Sediments. *Aquatic Geochemistry* 14, 193–210. doi:10.1007/s10498-008-9032-5
- Poulenard, J., Perrette, Y., Fanget, B., Quetin, P., Trevisan, D., Dorioz, J.M., 2009. Infrared spectroscopy tracing of sediment sources in a small rural watershed (French Alps). *Science of The Total Environment* 407, 2808–2819. doi:10.1016/j.scitotenv.2008.12.049
- Strahler A.N. 1957. Quantitative Analysis of Watershed Geomorphology. *Transactions American Geophysical Union* 28, 913-920. doi: 10.1029/TR038i006p00913
- Saunders, W.M.H., Williams, E.G., 1955. Observations on the Determination of Total Organic Phosphorus in Soils. *Journal of Soil Science* 6, 254–267. doi:10.1111/j.1365-2389.1955.tb00849.x
- Sharpley, A.N., Gburek, W.J., Folmar, G., Pionke, H.B., 1999. Sources of phosphorus exported from an agricultural watershed in Pennsylvania. *Agricultural Water Management* 41, 77–89. doi:10.1016/S0378-3774(99)00018-9
- Tamburini, F., Bernasconi, S.M., Angert, A., Weiner, T., Frossard, E., 2010. A method for the analysis of the $\delta^{18}\text{O}$ of inorganic phosphate extracted from soils with HCl. *European Journal of Soil Science* 61, 1025–1032. doi:10.1111/j.1365-2389.2010.01290.x
- Tamburini, F., Pfahler, V., Bünemann, E.K., Guelland, K., Bernasconi, S.M., Frossard, E., 2012. Oxygen Isotopes Unravel the Role of Microorganisms in Phosphate Cycling in Soils. *Environmental Science & Technology* 46, 5956–5962. doi:10.1021/es300311h
- Tamburini, F., Pfahler, V., von Sperber, C., Frossard, E., Bernasconi, S.M., 2014. Oxygen Isotopes for Unraveling Phosphorus Transformations in the Soil–Plant System: A Review. *Soil Science Society of America Journal* 78, 38. doi:10.2136/sssaj2013.05.0186dgs
- Turner, B.L., Cade-Menun, B.J., Condon, L.M., Newman, S., 2005. Extraction of soil organic phosphorus. *Talanta, Analysis of Phosphorus in Environmental and Agricultural Samples* 66, 294–306. doi:10.1016/j.talanta.2004.11.012
- Von Sperber, C., Kries, H., Tamburini, F., Bernasconi, S.M., Frossard, E., 2014. The effect of phosphomonoesterases on the oxygen isotope composition of phosphate. *Geochimica et Cosmochimica Acta* 125, 519–527. doi:10.1016/j.gca.2013.10.010
- Von Sperber, C., Tamburini, F., Brunner, B., Bernasconi, S.M., Frossard, E., 2015. The oxygen isotope composition of phosphate released from phytic acid by the activity of wheat and *Aspergillus niger* phytase. *Biogeosciences* 12, 4175–4184. doi:10.5194/bg-12-4175-2015
- Walling, D.E., Collins, A.L., Stroud, R.W., 2008. Tracing suspended sediment and particulate phosphorus sources in catchments. *Journal of Hydrology, Characterization and apportionment of nutrient and sediment sources in catchments* 350, 274–289. doi:10.1016/j.jhydrol.2007.10.047
- Wang, D., Levine, S.N., Meals, D.W., Hoffmann, J.P., Drake, J.C., Cassell, E.A., 1999. Importance of Instream Nutrient Storage to P Export from a Rural, Eutrophic River in Vermont, USA, in: O.nley, T., L.nley, P. (Eds.), *Lake Champlain in Transition: From Research Toward Restoration*. American Geophysical Union, pp. 205–223.
- Weiner, T., Mazeh, S., Tamburini, F., Frossard, E., Bernasconi, S.M., Chiti, T., Angert, A., 2011. A method for analyzing the $\delta^{18}\text{O}$ of resin-extractable soil inorganic phosphate. *Rapid Communications in Mass Spectrometry* 25, 624–628. doi:10.1002/rcm.4899
- Zohar, I., Shaviv, A., Young, M., Kendall, C., Silva, S., Paytan, A., 2010. Phosphorus dynamics in soils irrigated with reclaimed waste water or fresh water — A study using oxygen isotopic composition of phosphate. *Geoderma* 159, 109–121. doi:10.1016/j.geoderma.2010.07.002

Table and Figure captions

Table 1. Characteristics of the sediment sampling sites.

Table 2. Characteristics of the soil sampling sites.

Table 3. P speciation and oxygen isotope composition.

Figure 1. Redon River catchment and sampling sites; the capital letters refer to the sampling locations, the numbers are the sample code.

Figure 2. Oxygen isotope composition ($\delta^{18}\text{O}_p$) and deviation from isotopic equilibrium. The gray circles represent potential source samples, white circles represent riverbed sediment samples, and $\text{Pols} = \text{P}_{\text{Olsen}}/\text{P}_t$ in % (note that the deviation of the potential sources from equilibrium in this image have been calculated according to the average within-river conditions).

Figure 3. Conceptual model describing the evolution of $\delta^{18}\text{O}_p$ for the riverbed sediments in the studied torrential system.

ACCEPTED MANUSCRIPT

Table 1 Characteristics of sediment sampling sites

Site	SO	Sample code	Sampling date	Q	EC	SRP	Altitude	Drainage area	Land use of the drainage basin	Main sources of stream P
				L/s	mS/cm	mg P/L	m a.s.l	ha		
A	1	1	September 3, 2012	<1	543	0.0012	576	4	Cultivated land	NPS (erosion of cultivated soils and tiches)
B	1	2	September 3, 2012	1	557	0.009	577	17	Cultivated land	NPS (erosion of cultivated soils and tiches)
C	2	3	March 26, 2012	35	598	0.008	696	510	Steep forest	NPS (natural: mainly river bank erosion)
D	3	4	September 3, 2012	55	7203	0.027	510	2310	Forest and cultivated land	NPS (natural, agricultural)
E	3	5	March 26, 2012	105	654	0.056	508	2311	Forest and cultivated land	NPS (natural, agricultural) + PS (WWTP)
		6	September 3, 2012	55	753	0.092				
F	4	7	March 26, 2012	340	597	0.062	405	3300	Forest and cultivated land	NPS (natural, agricultural) + PS (WWTP)
		8	September 3, 2012	85	975	0.067				

SO = Strahler stream order (Strahler, 1957)

Q = water discharge

EC = electric conductivity

SRP = soluble reactive phosphorus concentration in water, measured on filtered samples (0.45 μ m) by the molybdate-blue method (Murphy and Riley, 1962)

NPS = Non-point source, PS = Point source

WWTP = waste-water treatment plant

Table 2 Characteristics of soil sampling sites

Sample code	Date of sampling	Location	Soil depth cm	Altitude m a.s.l	Sampling conditions
9	September 3, 2012	Wetland topsoil	-1 to -5	552	Organic soil, saturated
10	September 3, 2012	Cultivated topsoil ^a	-10 to -30	558	Cambisol on moraine deposit, dry
11	September 3, 2012	Cultivated soil (embankment) ^a	-60 to -70	558	Cambisol on moraine deposit, saturated
12	September 3, 2012	Riverbank in a forested upland area	-	980	Würm moraine deposits, saturated
^a samples collected within an agricultural sub-catchment of 33 ha (Figure 1)					

Table 3. P speciation and oxygen isotopes composition

Sample code	Sample type	Pt	P _{Olsen}	Pi	Po	P _{Olsen} /Pt ^a	$\delta^{18}\text{O}_p$	St. Dev.	n	Deviation from equilibrium (‰) ^b
		mgP/kg	mgP/kg	mgP/kg	mgP/kg	%	‰	‰	-	‰
1	River bed sediments ^d	791	14	401	180	2.4	12.5	0.3	3	- 2.4
2		751	20	323	208	3.8	13.3 ^s	0.3	2	- 1.6
3		528	8	411	78	1.7	15.5	0.6	4	+ 1.3
4		609	18	398	202	2.9	12.2 ^s	0.4	2	- 1.5
5		1018	65	586	194	8.4	15.8 ^s	0.7	2	+ 1.4
6		962	67	470	166	10.6	15.2	0.3	3	< 0.4
7		734	26	439	138	4.5	13.8	0.4	2	< 0.4
8		736	21	324	90	5.2	12.8	0.1	3	- 0.6
9	Wetland soil	1163	32	179	383	5.7	18.3 ^s	0.2	2	+ 2.1 (+ 3.6) ^c
10	Agric. topsoil	1256	43	631	450	6.2	14.0	0.4	2	< 0.4 (< 0.4) ^c
11	Embankment/ deep soil	755	5	363	76	1.1	11.8	0.4	2	- 0.7 (- 2.0) ^c
12	Riverbank (moraine)	511	3	305	36	0.7	15.9	0.2	2	+1.9 (+1.2) ^c
13	Lake	459	15	365	121	3.5	12.9	0.3	2	< 0.4
14	sediments	539	17	420	70	3.1	13.9	0.5	2	+ 0.8

^a As a sum of Pi and Po fractions^b See section 2.3.2 and equation 2^c Values not in brackets are calculated according to the temperature and water isotopes conditions at the sampling site, the values in brackets are calculated according to the “within-river” conditions^d Sediment samples are ranked according to the corresponding drained area

Figure 1. Redon River catchment and sampling sites, capital letters refer to sampling locations, numbers to sample code

Figure 2. Oxygen isotopes composition $\delta^{18}\text{Op}$ and deviation from isotopic equilibrium. Grey circles represent potential sources samples, white circles represent riverbed sediments samples, $P_{\text{ols}} = P_{\text{Olsen}}/P_t$ in % (note that deviation from equilibrium of potential sources in this picture have been calculated according to average within-river conditions)

Highlights

- We provided the first assessment of the oxygen isotopic composition of HCl-extracted phosphate in river sediments
- We observe a contrast in the oxygen isotope composition between upstream and downstream sediments
- This contrast can be related to anthropogenic impacts on the river system and to the sediment residence time
- We have shown that in some situations it is possible to trace particulate-P sources of sediments with this isotopic tool