

# Early Paleozoic intracontinental orogeny in the Yunkai domain, South China Block: New insights from field observations, zircon U–Pb geochronological and geochemical investigations

Chaolei Yan, Liangshu Shu, Michel Faure, Yan Chen, Cheng Li

► **To cite this version:**

Chaolei Yan, Liangshu Shu, Michel Faure, Yan Chen, Cheng Li. Early Paleozoic intracontinental orogeny in the Yunkai domain, South China Block: New insights from field observations, zircon U–Pb geochronological and geochemical investigations. *Lithos*, Elsevier, 2017, 268-271, pp.320-333. <10.1016/j.lithos.2016.11.013>. <insu-01427240>

**HAL Id: insu-01427240**

**<https://hal-insu.archives-ouvertes.fr/insu-01427240>**

Submitted on 26 Jun 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



1 **Early Paleozoic intracontinental orogeny in the Yunkai Domain,**  
2 **(South China Block): new insights from field observations, zircon U–**  
3 **Pb geochronological and geochemical investigations**

4 **Chaolei Yan<sup>a</sup>, Liangshu Shu<sup>a\*</sup>, Faure Michel<sup>b</sup>, Yan Chen<sup>b</sup>, Cheng Li<sup>a</sup>**

5 <sup>a</sup>State Key Laboratory for Mineral Deposits Research, School of Earth Sciences and Engineering,  
6 Nanjing University, 164 Xianlin Road, Nanjing 210023, China

7 <sup>b</sup>Institut des Sciences de la Terre d'Orléans, UMR 7327, CNRS, Université d'Orléans, 45071  
8 Orléans, France

9 \* Corresponding author E-mail: [lsshu@nju.edu.cn](mailto:lsshu@nju.edu.cn)

10 **ABSTRACT**

11 Debate on whether the Early Paleozoic tectono–magmatic event in South China is  
12 related to a subduction–collision or an intracontinental orogen has been lasted for  
13 decades within the geoscience community. This study deals with LA–ICP–MS zircon  
14 U–Pb ages, whole–rock chemistry, rare earth elements, trace elements and Hf isotopes  
15 from granitoid samples collected in the Yunkai domain in order to better constrain the  
16 Early Paleozoic tectonic evolution of the South China Block. The weighted mean  
17  $^{206}\text{Pb}/^{238}\text{U}$  ages for eight samples range from 426 Ma to 443 Ma, representing the  
18 crystallization ages of the magma. Fourteen samples were analyzed for geochemistry,  
19 all of which are characterized by a peraluminous signature with A/CNK values greater  
20 than 1.0. The REE geochemistry reveals enrichment in light rare earth element.  
21 LREE/HREE values range from 2.81 to 30.36 and  $(\text{La}/\text{Yb})_{\text{N}}$  vary from 1.23 to 55.14  
22 (mean of 14 analyses is 14.69). All the samples exhibit distinct negative Ba, Sr and Nb  
23 anomalies and enrichment in Rb, Th, U and Pb. Hf isotopic analyses indicate negative  
24  $\epsilon_{\text{Hf}}(t)$  values mainly ranging from -3 to -12, corresponding to two model age  
25 distributing from 1637 Ma to 2208 Ma. The geochemical analyses indicate that the  
26 Silurian granitic magmas in the Yunkai domain were derived from partial melting of  
27 crustal materials with little or no input of mantle source. These new data support the

28 intracontinental subduction model already proposed to account for the Early Paleozoic  
29 tectonic, metamorphic and magmatic event of South China.

30

31 **Keywords:** Early Paleozoic Granitoids; U–Pb geochronology; Geochemistry; Hf  
32 isotope; Yunkai domain; South China

33

## 34 **1. Introduction**

35 The Cathaysia and Yangtze blocks amalgamated in the Neoproterozoic (Fig. 1),  
36 forming the South China Block (SCB), and its subsequent Phanerozoic tectonic  
37 framework is an important constituent of tectonic evolution of Asia (Cawood et al.,  
38 2013; Li et al., 2008a, b, 2009; Rong et al., 2010; Song et al., 2015; Zhang and Zheng,  
39 2013; Zhang et al., 2011; Zhao and Cawood, 2012). Furthermore, as recognized since  
40 1920's (Grabau, 1924), the Middle Devonian terrigenous rocks unconformably  
41 covering Early Paleozoic folded rocks and granitoids, document an Early Paleozoic  
42 tectono–magmatic event widespread in the SE part of the South China Block (Fig. 1).  
43 This pre–Devonian orogeny has been improperly referred to as the "Caledonian  
44 orogeny", however as precise time constraints are now available (Song et al., 2015;  
45 Wang et al., 2007), and because this belt is unrelated to the true Caledonian belt of  
46 Norway and N America, this term should be abandoned. The Early Paleozoic orogeny  
47 of SE China is characterized by: i) the regional absence of Silurian strata, ii) the  
48 unconformity between middle Devonian coarse clastic sequence and Ordovician  
49 marine flysch sequence, iii) a greenschist to amphibolite facies metamorphism coeval  
50 with a ductile deformation, and iv) the occurrence of numerous S–type granitic plutons  
51 (BGMRFJ, 1985; BGMRGX, 1985; BGMRHN, 1988; BGMRJX, 1984; BGMRZJ,  
52 1989; Charvet, 2013; Charvet et al., 2010; Faure et al., 2009; Li et al., 2010; Shu, 2012;  
53 Shu et al., 2008b, 2015; Yao et al., 2013, 2014). The geodynamic significance of this  
54 belt, whether as a collisional orogen or as an intracontinental one, has been debated  
55 since a long time. Guo et al. (1989) argued for the presence of Early Paleozoic  
56 ophiolites in the Yunkai and Wuyi massifs of the SE part of the South China Block.  
57 However, in the past decades, several investigations indicated that the previously

58 proposed oceanic subduction and collision models do not properly account for the Early  
59 Paleozoic lithological and tectonic features of this part of the SCB (Charvet et al., 2010;  
60 Faure et al., 2009; Shu et al., 2014, 2015; Song et al., 2015; Wang et al., 2007, 2011,  
61 2013b; Zhang et al., 2011). The main facts arguing against a collisional model are the  
62 lack of ophiolites, accretionary complexes and magmatic arc. Indeed, ophiolitic  
63 gabbros, once considered as Early Paleozoic, are in fact Neoproterozoic in age, ca. 850–  
64 800 Ma (Li et al., 2005; Shu et al., 2006, 2011). Moreover, most of the "basaltic rocks"  
65 interlayered in the Early Paleozoic strata have been reassessed as meta–greywacke (Shu  
66 et al., 2008a, 2014). Granitoid is an efficient rock to understand the tectonic evolution  
67 of the continental crust (e.g., Pearce et al., 1984; Pitcher, 1983). Some studies have dealt  
68 with the Early Paleozoic granitoids of the South China Block. However, precise  
69 chronological and geochemical data related to their petrogenesis are rare (e.g., Deng et  
70 al., 2012; Li et al., 2010; Liu et al., 2010; Shu et al., 2008a; Wang et al., 2010, 2011;  
71 Xia et al., 2014; Xu et al., 2011; Yang et al., 2010). This study presents zircon U–Pb  
72 geochronology, bulk geochemistry and Hf isotope analysis of granitoids from the  
73 Yunkai domain, providing new insights into the crustal evolution of the SE part of the  
74 SCB in Early Paleozoic.

75

## 76 **2. Geological setting**

### 77 ***2.1 The general framework of SCB***

78 The southern part of the South China Block (Fig. 1) also referred to as “the South  
79 China Fold Belt” is separated by the Neoproterozoic Jiangnan Orogen from the Yangtze  
80 Block (Grabau, 1924; Guo et al., 1989; Li, 1997; Li et al., 2009, 2012; Shu and Charvet,  
81 1996; Shu et al., 2006, 2014; Wang et al., 2007). This NE–SW trending belt is a  
82 complex area that experienced several successive tectonic, metamorphic and magmatic  
83 events, namely: i) a Neoproterozoic (Tonian–Cryogenian, ca. 850 Ma) collision; ii) a  
84 Late Neoproterozoic (Cryogenian–Ediacarian, ca. 820–690 Ma) rifting event marked  
85 by volcanoclastic sedimentation and bimodal magmatism; iii) an Early Paleozoic  
86 orogeny marked by a Devonian angular unconformity, and iv) a Middle Triassic  
87 orogeny marked by a Late Triassic unconformity (BGMRGX, 1985; BGMRJX, 1984;

88 [Faure et al., 2009](#); [Lin et al., 2008](#); [Shu, 2012](#)).

89 The Neoproterozoic Jiangnan orogenic belt in the southeastern margin of the  
90 Yangtze Block is a NW–ward subduction–collision belt developed during the  
91 amalgamation of the Cathaysia and Yangtze blocks ([Shu, 2012](#); [Yao et al., 2014](#)).  
92 Ophiolite, I–type granite, rhyolite, basalt and gabbro, dated at ca. 1000–855 Ma, crop  
93 out along the Shaoxing–Jiangshan–Guilin fault zone ([Li et al., 2009](#); [Shu, 2012](#); [Yao et  
94 al., 2014](#)). The amalgamation resulted in the collision of island arcs with the Yangtze  
95 Block and the closure of a back–arc basin. After the Middle Neoproterozoic orogeny,  
96 from ca. 820 to 690 Ma, the South China block experienced a rifting event coeval with  
97 a bimodal magmatism ([Li et al., 2003, 2005](#); [Shu, 2006](#); [Shu et al., 2011](#); [Wang and Li,  
98 2003](#); [Wang et al., 2006](#)). In Cambrian times, the southern part of the rift was a littoral–  
99 neritic depositional environment whereas during the early–middle Ordovician period,  
100 this area was dominated by a neritic–bathyal setting ([Rong et al., 2010](#); [Shu et al., 2014](#)).  
101 In the late Ordovician, the SE part of the South China Block underwent a  
102 sedimentological change to a littoral silico–clastic environment, along with the  
103 initiation of uplift processes ([BGMRFJ, 1985](#); [BGMRJX, 1984](#); [Shu, 2012](#); [Shu et al.,  
104 2014](#)).

105 During the Silurian, extensive folding, thrusting, metamorphism and anatexis  
106 developed. The emplacement of numerous granitic plutons represents the end of the  
107 orogeny. These features are well recorded in the Wuyi, Jinggang, Nanling and Yunkai  
108 areas ([Figs. 1 and 2](#)). The maximum shortening can reach up to 67% in the Jinggang  
109 and Wuyi belts ([Charvet et al., 2010](#); [Shu, 2012](#); [Shu et al., 2008a, 2015](#)). Fold axes  
110 strike are dominantly E–W. The kinematic analysis in the Jinggang and Wuyi domains  
111 shows that the ductile shearing was directed to the S or SE, however, a northwestward  
112 vergence may develop in the northwestern part of the belt. Concerning the Early  
113 Paleozoic orogeny, two pre–Devonian litho–tectonic units have been identified: 1) A  
114 slate unit, composed of the Sinianto Ordovician marine sandy–muddy rocks, which  
115 experienced a low greenschist facies metamorphism before the intrusion of Silurian  
116 granitoids; 2) A metamorphic unit, comprising Neoproterozoic mica schist, amphibolite,  
117 paragneiss and orthogneiss, and locally Paleoproterozoic amphibolites, gneisses and

118 gneissic granites (Faure et al., 2009; Shu, 2012; Shu et al., 2014; Yu et al., 2009).  
119 However, this orogeny is not well developed in the northwestern part of the South China  
120 Block where the Early Paleozoic strata did not experience any significant  
121 metamorphism, and only underwent slight brittle deformation, together with weak  
122 magmatism. From a geodynamic point of view, the Early Paleozoic orogeny of the  
123 South China Block was interpreted as the consequence of a rift closure from the Late  
124 Ordovician to Early Silurian. Therefore, the orogeny corresponds to an intracontinental  
125 event accommodated by the continental subduction of the southern part of the rift below  
126 its northern one (Faure et al., 2009). It is worth to note that the rift closure area that  
127 corresponds to a crustal scar instead of an ophiolitic suture does not coincide with the  
128 Neoproterozoic suture but is located within the Cathaysia Block. The upper part of the  
129 belt (i.e. the slate unit) is a fold-and-thrust belt limited at its base by a ductile  
130 décollement localized in the Sinian system. The lower part of the belt (i.e. the  
131 metamorphic unit underlying the basal décollement) was characterized by deep burial  
132 giving rise to an amphibolite facies metamorphism (Zhao and Cawood, 1999). During  
133 the exhumation of this lower part, those metamorphic rocks experienced retrogression  
134 and partial melting at ca. 444–420 Ma represented by migmatite and granitoid (Faure  
135 et al., 2009).

136 From the Middle Devonian to Early Carboniferous, quartz sandstone, feldspathic  
137 sandstone, conglomerate, and siltstone intercalated with chert, limestone and bioclastic  
138 limestone were deposited unconformably on the Early Paleozoic sequences (BGMRFJ,  
139 1985; BGMRGX, 1985; BGMRJX, 1984; Shu et al., 2008b, 2015).

140 A Late Triassic regional unconformity implies a Middle Triassic orogeny (e.g.,  
141 Chu et al., 2012; Faure et al., 2009; Lin et al., 2008; Shu, 2012; Wang et al., 2013b).  
142 The Triassic orogens are widespread around the margins and inside the South China  
143 Block. The Early Paleozoic metamorphic rocks were intensely reworked by the Triassic  
144 events.

## 145 ***2.2 Geology of the Yunkai domain***

146 The NE trending Yunkai domain consists of a wide variety of plutonic,  
147 sedimentary and metamorphic rocks (Fig. 1; BGMRGX, 1985; Faure et al., 2016; Lin

148 [et al., 2008](#)). In the study area, located in the NW part of the Yunkai domain, several  
149 stratigraphic units are exposed: 1) a set of low greenschist facies and folded  
150 Neoproterozoic sedimentary rocks, comprising slate, sandy slate and conglomerate; 2)  
151 weakly metamorphosed and folded Early Paleozoic sequences mainly composed of  
152 metasandstone and metamudstone; 3) Post–Ordovician sedimentary rocks ([Fig. 2](#)).

153 In the Yunkai domain, Early Paleozoic biotite–muscovite granite and  
154 monzogranite are well exposed. These plutons are heterogeneously deformed with an  
155 unfoliated core to a gneissic, sometimes mylonitic, margin ([BGMRGX, 1985](#)). In the  
156 section, a progressive reduction in grain size from medium–coarse to fine grained can  
157 be observed from the pluton core to margin. Meanwhile, Early and Middle Triassic  
158 peraluminous plutons are also distributed in this area ([Lin et al., 2008](#); [Wang et al.,](#)  
159 [2007](#)).

160 Our field observations indicate that the bulk architecture results of NE–SW and  
161 E–W trending upright and overturned folds ([Fig. 2](#)) with NW and N vergences,  
162 implying two tectonic deformation stages corresponding to Neoproterozoic and Early  
163 Paleozoic, respectively ([Faure et al., 2016](#); [Lin et al., 2008](#)).

### 164 **3. Sample description**

165 Petrographic features and GPS locations of the analyzed samples are listed in  
166 [Table 1](#). Sample 1571 is a fine to medium grained–massive two–mica granite composed  
167 of ca. 40% K–feldspar, 35% quartz, 10% plagioclase, 7% muscovite, 5% biotite and  
168 other accessories([Fig. 3A and 3F](#)) ([Table 1](#)). Sample 1582 of monzogranite is mainly  
169 composed of 40% quartz, 35% K–feldspar, 15% albite, 5% muscovite and 5% other  
170 accessory minerals ([Table 1](#)). It is located in the boundary of the pluton, displaying a  
171 weak mineral fabric. Sample 1586 is a medium–grained massive K–feldspar muscovite  
172 granite with 45% K–feldspar, 40% quartz, 8% plagioclase, 5% muscovite and other  
173 accessories ([Fig. 3B and 3G](#)) ([Table 1](#)). From the pluton core to margin, a progressive  
174 reduction in grain size from medium–coarse to fine–grained can be observed. Sample  
175 1601 exhibits a distinct granitic texture, with about 35% K–feldspar, 30% quartz, 20%  
176 biotite, 5~8% albite, 5% muscovite and other accessories ([Fig. 3C and 3H](#)) ([Table 1](#)).  
177 Sample 1621 of the muscovite granite is mainly composed of 40% quartz, 38% K–

178 feldspar, 12% plagioclase, 7% muscovite and 3% other accessory minerals (Table 1),  
179 polysynthetic twin and cross hatched twin are well developed in the plagioclase and  
180 microcline, respectively (Fig. 3D). Sample 1628 of the biotite-rich proto-mylonitic  
181 granite with feldspar augen shows about 40% K-feldspar, 30% quartz, 15~18% biotite,  
182 10% plagioclase and other accessories (Table 1), in which the K-feldspar phenocrysts  
183 are commonly deformed (Fig. 3E and 3I). Sample 1630 is a well-foliated gneissic  
184 biotite granite composed of 45% microcline, 25% biotite, 15% quartz, 10% plagioclase  
185 and 5% other accessory minerals (Table 1). Sample 1633 also shows a gneissosity, but  
186 weaker than in Sample 1630. It is mainly composed of 40% microcline, 25% quartz,  
187 20% biotite, 10% plagioclase and 5% other accessories. The dome-shaped granitic  
188 pluton displays obvious zoning with gneissic granite in the middle, and mylonitic  
189 granite around the margin.

190 In this study, five granite samples (1571, 1582, 1586, 1601, 1621), one mylonitic  
191 granite (1628) and two gneissose granites (1630, 1633) were collected for zircon U-Pb  
192 dating and Hf isotope analysis (see locations in Fig. 2). Fourteen rock samples (1571,  
193 1571-1, 1582, 1586, 1586-1, 1601, 1601-1, 1601-2, 1601-3, 1621, 1621-1, 1628,  
194 1630, 1633) were used for whole-rock geochemical study.

195

#### 196 **4. Analytical methods**

197 Whole-rock major element contents were analyzed by ARL-9900 X-ray  
198 fluorescence spectrometer (XRF) at the Testing Center of Shandong Bureau of China  
199 Metallurgical Geology Bureau. The uncertainties reported in this study are 2% for  
200 major elements. Trace elements and rare earth elements (REE) were measured by  
201 Inductively Coupled Plasma- Mass Spectrometry (ICP-MS) (Finnigan Element II) at  
202 the Testing Center of Shandong Bureau of China Metallurgical Geology Bureau and  
203 ALS Chemex (Guangzhou) co., Ltd, respectively. International standards were used to  
204 define the analytical precision and accuracy throughout the analytical processes for  
205 ICP-MS. The uncertainties are 5 percent for trace elements. For the detailed analytical  
206 procedure we refer to those documented in Gao et al. (2003).

207 Zircons were separated from the crushed rocks using heavy liquid and magnetic



208 techniques and then handpicked under a binocular microscope. The zircon grains were  
209 mounted in epoxy resin and polished to expose the approximate center of grains, and  
210 then coated with gold. Cathodoluminescence (CL) images of the zircons were obtained  
211 using a JEOL JXA8230 electron probe microanalyzer at the Testing Center of Shandong  
212 Bureau of China Metallurgical Geology Bureau.

213 The laser ablation (LA)–ICP–MS analysis of zircon U–Pb isotopic compositions  
214 was performed at the State Key Laboratory for Mineral Deposits Research, Nanjing  
215 University, using an Agilent 7500a ICP–MS system attached to New Wave 213 nm  
216 laser ablation system with an in–house sample cell. Samples were analyzed in runs of  
217 ca. 15 analyses including five zircon standards and 10 sample points. The details of the  
218 analytical procedure are documented in [Jackson et al. \(2004\)](#). The U–Pb fractionation  
219 was corrected using zircon standard GEMOC GJ–1 with a  $^{207}\text{Pb}/^{206}\text{Pb}$  age of  $601\pm 12$   
220 Ma and the accuracy was controlled using the zircon standard Mud Tank with an age  
221 of  $735\pm 12$  Ma ([Black and Gulson, 1978](#)). The U–Pb ages were calculated from the raw  
222 signal data using the software Glitter (ver.4.4). Because the  $^{204}\text{Pb}$  could not be measured  
223 owing to low signal and interference from  $^{204}\text{Hg}$  in the gas supply, common lead  
224 correction was carried out using the EXCEL program common Pb correction ([Andersen,](#)  
225 [2002](#)). For zircons older than 1000 Ma, because of large amounts of radiogenic Pb, the  
226  $^{207}\text{Pb}/^{206}\text{Pb}$  age is more reliable than  $^{206}\text{Pb}/^{238}\text{U}$ , whereas for zircons younger than 1000  
227 Ma, as a result of the low content of radiogenic Pb and uncertainty of common Pb  
228 correction, the  $^{206}\text{Pb}/^{238}\text{U}$  age is more reliable ([Griffin et al., 2004](#)).

229 Zircon Hf isotopic composition was analyzed by Neptune MC–ICP–MS, which is  
230 a double focusing multi–collector ICP–MS and has the capability of high mass  
231 resolution measurements in a multiple collector mode. During laser ablation analyses,  
232 the isobaric interference of  $^{176}\text{Lu}$  on  $^{176}\text{Hf}$  is negligible due to the extremely low  
233  $^{176}\text{Lu}/^{177}\text{Hf}$  value in zircon (normally  $<0.002$ ). However, the interference of  $^{176}\text{Yb}$  on  
234  $^{176}\text{Hf}$  must be intensively corrected since the contribution of  $^{176}\text{Yb}$  to  $^{176}\text{Hf}$ . This method  
235 can provide an accurate correction of the  $^{176}\text{Yb}$  interference on  $^{176}\text{Hf}$  ([Kemp et al., 2006](#)).  
236 During analysis, an isotopic ratio of  $^{176}\text{Yb}/^{172}\text{Yb} = 0.5887$  was applied. Standard zircon  
237 91500 was used for the external correction, with a  $^{176}\text{Hf}/^{177}\text{Hf}$  value of  $0.282300 \pm 8$

238 (2 $\sigma$ ). The detailed analytical procedure is similar to the description by Yuan et al. (2008).  
239 Initial  $^{176}\text{Hf}/^{177}\text{Hf}$  values were calculated based on Lu decay constant of  $1.865\text{E}-11$   
240 (Scherer et al., 2001). Model ages were calculated under the assumption that the  
241  $^{176}\text{Lu}/^{177}\text{Hf}$  of average crust is 0.015, and the  $^{176}\text{Hf}/^{177}\text{Hf}$  and  $^{176}\text{Lu}/^{177}\text{Hf}$  ratios of  
242 chondrite and depleted mantle at the present are 0.282772 and 0.0332, 0.28325 and  
243 0.0384, respectively (Blichert-Toft and Albarede, 1997). The model ages (TDM)  
244 provide only a minimum age for the source material of the magma from which the  
245 zircons crystallized.

246

## 247 **5. Analytical results**

### 248 **5.1 Major and trace element compositions**

249 The analytical results of major and trace elements of 14 representative samples are  
250 given in Table 2. Samples 1571, 1571-1, 1586, 1586-1, 1601-2, 1621, 1621-1, 1628,  
251 1630, 1633 exhibit  $\text{SiO}_2$  contents ranging from 69.91 to 77.46 wt%, and plot in the  
252 granite field in the total alkali-silica diagram, displaying  $(\text{Na}_2\text{O}+\text{K}_2\text{O})$  versus  $\text{SiO}_2$   
253 (TAS) (Fig. 4A). However, Samples 1582, 1601, 1601-1 and 1601-3 plot in the field  
254 of quartz diorite, yielding  $\text{SiO}_2$  contents between 65.03 and 69.67 wt%. As shown by a  
255 plot of  $\text{Al}/(\text{Na}+\text{K})$  versus  $\text{Al}/(\text{Ca}+\text{Na}+\text{K})$  (Fig. 4B), all samples fall in the peraluminous  
256 area with high A/CNK ( $>1.1$ ), except Samples 1571, 1571-1 and 1586 which indicate  
257 slightly lower A/CNK (1.08, 1.08 and 1.05, respectively). According to Figure 4C, most  
258 of the samples belong to the high-K calc-alkaline affinity, while only Sample 1582  
259 plots into the calc-alkaline field.

260 The most intuitive character of all samples analyzed for geochemistry is that they  
261 display similar chondrite-normalized steep rare earth (REE) element patterns revealing  
262 obvious enrichment in light rare earth element (LREE) with respect to heavy rare earth  
263 element (HREE) (Fig. 5A). LREE/HREE values range from 2.81 to 30.36, and  
264  $(\text{La}/\text{Yb})_N$  vary from 1.23 to 55.14 (the mean of 14 analyses is 14.69) (Fig. 5A, Table2).  
265 All the samples mark distinct negative Eu anomalies ( $\text{Eu}/\text{Eu}^*$  value of 0.06-0.59, the  
266 mean of 14 analyses is 0.4). On the primitive mantle-normalized spider diagrams (Fig.  
267 5B), all samples exhibit strongly negative Ba, Sr and Nb anomalies and the enrichment

268 of Rb, Th, U and Pb. Nb/Ta values range from 3.76 to 23.22 (average 9.50), consistent  
269 with the geochemical features of crustal derived granite (Corfu et al., 2003; Hoskin and  
270 Schaltegger, 2003; Pearce, 1996; Pearce et al., 1984).

### 271 **5.2 Zircon U–Pb ages**

272 Typical Cathodoluminescence (CL) images of zircons are presented in Figure 7.  
273 U–Pb results are listed in Table 3 and graphically illustrated in Figure 8. Zircons  
274 analyzed in this study range in length from 60 to 160  $\mu\text{m}$ , with length/width ratio ranges  
275 from 2:1 to 4:1. Most of zircon grains display oscillatory zoning (Fig. 7) and high Th/U  
276 values (average 0.80; Table 3), indicating their magmatic origin (Corfu et al., 2003;  
277 Hoskin and Schaltegger, 2003).

278 Two hundred and twenty seven zircon U–Pb ages are obtained, which fall in the  
279 range of 409–1306 Ma. However, most of the data indicate ages between 409 to 460  
280 Ma. These data plotted on concordia diagrams yield eight groups of weighted mean  
281  $^{206}\text{Pb}/^{238}\text{U}$  ages as  $426 \pm 3\text{Ma}$  (Sample 1571, 19 values out of total 25 data);  $439 \pm 3$   
282  $\text{Ma}$  (Sample 1582, 21 values out of total 32 data);  $436 \pm 3\text{Ma}$  (Sample 1586, 18 values  
283 out of total 25 data);  $443 \pm 3\text{Ma}$  (Sample 1601, 18 values out of total 26 data);  $430 \pm$   
284  $3\text{Ma}$  (Sample 1621, 23 values out of total 30 data);  $435 \pm 3\text{Ma}$  (Sample 1628, 23  
285 values out of total 33 data);  $431 \pm 3\text{Ma}$  (Sample 1630, 19 values out of total 29 data)  
286 and  $429 \pm 3\text{Ma}$  (Sample 1633, 18 values out of total 27 data) (Fig. 8). The above results  
287 indicate a Silurian age for these granitoids.

### 288 **5.3 In situ Hf isotopes**

289 More than half of the U–Pb dated zircons were chosen for in–situ Hf isotopic  
290 analysis. The Hf analyses were executed near the fields used for U–Pb dating spots. For  
291 purpose of discussing the Hf isotopic evolution history, the initial  $^{176}\text{Hf}/^{177}\text{Hf}$  values  
292 and  $\epsilon\text{Hf}(t)$  were calculated using the zircon  $^{206}\text{Pb}/^{238}\text{U}$  ages. The results of the Hf  
293 isotopic analyses are presented in Table 4, and the  $\epsilon\text{Hf}(t)$  versus U–Pb age diagram is  
294 illustrated in Figure 9.

295 Lu–Hf isotopic results for the eight granitoid samples are similar. All samples  
296 show negative  $\epsilon\text{Hf}(t)$  values, ranging from -0.80 to -38.96. For the individual plutons,  
297 the ranges are -4.44 to -17.51 (average of -7.60 for Sample 1571), -1.58 to -4.56

298 (average of -3.00 for Sample 1582), -3.00 to -38.96 (average of -15.42 for Sample 1586),  
299 -0.80 to -25.17 (average of -11.06 for Sample 1601), -4.48 to -11.55 (average of -7.25  
300 for Sample 1621), -1.70 to -15.20 (average of -7.92 for Sample 1628), -4.35 to -12.02  
301 (average of -7.67 for Sample 1630), -5.06 to -10.68 (average of -6.99 for Sample 1633),  
302 respectively. Correspondingly, on the  $\epsilon_{\text{Hf}}(t)$  versus U–Pb age plot, the two model ages  
303 ( $T_{\text{DM}2}$ ) mainly concentrate on 1481–2208 Ma, while sparsely distribute in 2208–3843  
304 Ma (Fig. 9). These results indicate that all of the analyzed granitic rocks are derived  
305 from the partial melting of Paleoproterozoic continental basement rocks. 14 groups of  
306 two-model-age data greater than 2.3 Ga may indicate that the magma has been  
307 contaminated by ancient crustal material to a certain extent. In addition, all the isotopic  
308 data are plotted under the CHUR line, suggesting that the involvement of a mantle  
309 component in the granitic magma was negligible.

310

## 311 **6. Discussion**

### 312 *6.1 Crystallization age of the granitoids*

313 The 227 zircons from all eight granitoid samples from the Yunkai domain yield  
314 similar weighted mean U–Pb ages between 426 and 443 Ma (Fig. 8, Table 3), indicative  
315 of a Silurian crystallization age. Many works have been undertaken to investigate the  
316 formation age of the Early Paleozoic granites in the South China Block (Li et al., 2010;  
317 Shu et al., 1999, 2008a, 2008b; Song et al., 2015; Wang et al., 2007, 2013b; Zhang et  
318 al., 1998; Zhao et al., 2013; Zhong et al., 2013). For example, the granitoids for the  
319 Xuefeng, Guidong, Zhuguang, Wugong and Wuyi Mountains display zircon U–Pb ages  
320 of 410–430 Ma, 422–427 Ma, 414–434 Ma, 428–462 Ma, 431–441 Ma, respectively  
321 (Chu et al., 2012; Li, 1990, 1994; Lou et al., 2006; Xu et al., 2011) (Fig. 1). Also,  
322  $^{40}\text{Ar}/^{39}\text{Ar}$  age can be utilized to trace the evolution process of the plutons (Kay et al.,  
323 2011; Makshev et al., 2004; McDougall and Harrison, 1999; Reynolds et al., 1981). Xu  
324 et al. (1992) reported two synkinematic phengites from the mylonites that yield  
325  $^{40}\text{Ar}/^{39}\text{Ar}$  plateau age of 428 Ma from the Dexing–Shexian shear zone in the northern  
326 margin of the Yangtze Block. Wang et al. (2007) showed that the synkinematic phengite  
327 of the mylonitic rocks from the Nanfeng–Yingtian shear zone at the northern margin of

328 the northern Wuyi domain yields  $^{40}\text{Ar}/^{39}\text{Ar}$  plateau at 423–426 Ma. In the Jiuling Mts,  
329 biotite and muscovite from mylonites yield  $^{40}\text{Ar}/^{39}\text{Ar}$  weighted plateau ages ranging  
330 from 379 Ma to 468 Ma (Chu and Lin, 2014). Lastly, Shu et al. (2015) indicated that  
331 the  $^{40}\text{Ar}/^{39}\text{Ar}$  data on newly grown biotites from deformed K–granites and biotite schist  
332 occurring along the Shaoxing–Pingxiang fault zone show pseudoplateau ages of 428–  
333 433 Ma. These  $^{40}\text{Ar}/^{39}\text{Ar}$  ages reflect the regional deformation time that can be  
334 interpreted as the contemporary period of the granitic magmatism. In addition, the  
335 timing of the magmatic activities can be constrained by geological occurrence. To the  
336 east of Hezhou, Cambrian strata are intruded by granite (section A–B in Fig. 2).  
337 Moreover, the Silurian and lower Devonian sequences are absent in this area, along  
338 with the pre–Devonian strata unconformably covered by the middle Devonian sequence.  
339 According to the geological literatures (BGMRGX, 1985; Faure et al., 2009; Shu et al.,  
340 2008a; Wang et al., 2007), it was probably related to a doming induced by thermal event.  
341 Therefore, the field observations support the view that the Early Paleozoic magmatic  
342 event in the Yunkai domain occurred during the Silurian.

343

## 344 ***6.2 Source and petrogenesis of the Early Paleozoic granitoids***

345 A number of studies have addressed the origin and petrogenesis of the Early  
346 Paleozoic granites in the South China block (e.g., Shu et al., 1999, 2015; Wang et al.,  
347 2007, 2010, 2011; Zhang et al., 2011; Zhao et al., 2013). Song et al. (2015) showed that  
348 Early Paleozoic granites collected from the Jiangnan Orogen and the eastern part of the  
349 SCB display  $\epsilon\text{Hf}(t)$  values of -1.44 to -36.84 with two stage model ages of Hf isotope  
350 range from 1.3 Ga to 3.7 Ga, which indicates that the Early Paleozoic granites were  
351 derived from crustal source without or with a small input of mantle materials.  
352 Additionally, the Early Paleozoic granites from Xuefeng, Jinggang and Yunkai domains  
353 show Sr–Nd isotopic compositions with initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios of 0.70924–0.72935,  
354 negative  $\epsilon\text{Nd}(t)$  values of -4.7 to -11.5 and Proterozoic Nd model ages (Wang et al.,  
355 2013b, and references therein). The results of this study indicate that the samples  
356 collected from the Yunkai area also support crustal sources for the genesis of the  
357 granitic magmas. Geochemically, the granite samples are K–feldspar megacryst rich,

358 and display high total alkali and alumina contents, which all plot in the peraluminous  
359 field (Fig. 4B). Geochemically, in the tectonic discrimination diagrams (Fig. 6), most  
360 of the granitoid samples plot in the post–collision field. The geochemical signatures  
361 indicate that the Early Paleozoic granitic magmas represent typical anatectic products  
362 of continental crust. They were generated during the Silurian post–orogenic collapse  
363 event, developed after the intracontinental thickening event. Isotopically, all 125  
364 isotopic data are under the CHUR line (Fig. 9). This feature agrees with the melting of  
365 a continental basement with little or no input of juvenile crust (Song et al., 2015). Two–  
366 stage continental crust model ages of the analyzed zircon grains show a concentration  
367 of 1481–2208 Ma ages (Fig. 9), which reveals that some Paleoproterozoic rocks  
368 probably exist beneath the South China Block. Moreover, 13 groups of  $\epsilon\text{Hf}(t)$  values  
369 less than -15 reveal that Archean material may exist beneath the South China Block or  
370 not far away.

371 Recently, some mafic rocks have been dated at 409–434 Ma. The Taoyuan  
372 hornblende gabbro complex, located to the north of the Wugongshan domain, was  
373 formed by magma mixing and mingling, whose mafic member originated from a  
374 metasomatized lithospheric mantle during the Silurian (Zhong et al., 2013). These  
375 authors interpreted this rock to be generated from the post–orogenic collapse stage of  
376 an intracontinental tectonic regime. Moreover, several gabbroic plutons have been  
377 identified in the Yunkai area (Wang et al., 2013a). It is proposed that the generation of  
378 the source of the mafic magmas was located within Mesoproterozoic to Early  
379 Neoproterozoic asthenospheric mantle, metasomatized by paleosubduction–modified  
380 wedge derived fluids during asthenospheric upwelling in the Silurian, resulting in the  
381 melting of the continental lithospheric mantle. Similar mechanism is well inspired by  
382 de Jong et al. (2015) and references therein. According to the above lines of evidence,  
383 it seems that the granitic magmas in this study area were generated by partial melting  
384 of continental crust without contribution of juvenile crustal components.

385 Generally, the granites can be used to explore petrogenesis and tectonic settings  
386 (Abdallah et al., 2007; Castro, 2014; Foden et al., 2015; John and Blundy, 1993; Shu et  
387 al., 2008a; Song et al., 2015; Vigneresse, 2014; Wang et al., 2007). Field observation

388 shows that the granite structures were gradually changed from the center to the margin  
389 of the plutons. Foliation characterized by the preferred orientation of quartz, feldspar  
390 and mica is common at the pluton margin (Fig. 3). Regional-scale folding during the  
391 Early Paleozoic was widespread together with extensive thrusting and ductile shearing.  
392 Various fold types, such as overturned, chevron, recumbent, and sheath, thrust sheets,  
393 and mylonitic shear zones are widely developed in the pre-Devonian units (Charvet et  
394 al., 2010; Li et al., 2010; Shu et al., 1997; Xu et al., 2011; Yao et al., 2011). The intense  
395 folding and shortening accommodated by intracontinental subduction might raise the  
396 temperature of parent rocks (Shu, 2006; Song et al., 2015; Zhang et al., 2011).  
397 Furthermore, additional heat would be provided by radioactive crustal elements  
398 concentrated during the ductile deformation. Thus the heat budget would trigger partial  
399 melting of the continental crust able to generate the Early Paleozoic granitic magma.  
400 Also, dehydration melting of the hydrous minerals, such as micas, probably plays  
401 important roles in the partial melting process. After its burial, the thickened crust was  
402 quickly eroded and recovered its normal thickness. The metamorphosed continental  
403 rocks were gradually exhumed and decompressed under an extensional tectonic setting,  
404 resulting in the rise of the granitic magma and its emplacement in the upper crust (Faure  
405 et al., 1996).

406

### 407 ***6.3 Tectonic setting of the South China Block in the Early Paleozoic***

408 A large-scale tectono-magmatic event took place in the southeastern South China  
409 Block during the Silurian (Charvet et al., 2010; Faure et al., 2009; Shu et al., 2014;  
410 Wang et al., 2007). But the tectonic setting of the Early Paleozoic magmatism is still  
411 controversial. Guo et al. (1989) proposed that a subduction zone was developed along  
412 the Zhenghe-Dapu section in the Sinian-Late Ordovician period, resulting in the arc-  
413 continent collision and the formation of a large-scale fold belt. According to the  
414 geochemical features of the granites in the Yunkai belt, Peng et al. (2006, 2016a, 2016b)  
415 suggested an ocean-continent subduction-collision and a post-collisional extension-  
416 delamination-underplating model for the South China Block in the Early Paleozoic.  
417 Similarly, a subduction-related mechanism in the northern margin of the Yunkai

418 domain during the Early Paleozoic was assumed by [Qin et al. \(2013\)](#) owing to the arc  
419 magmatic geochemical characters of the Hudong plutonic complex.

420 Nevertheless, the field geological evidences are always the first hand and most  
421 important information concerning the tectonic setting. The Ordovician orogenic event  
422 (ca. 470–460 Ma), referred to as the Grampian Phase in the British Isles or Taconian  
423 Phase in western New England ([McKerrow et al., 2000](#)), is generally accepted as being  
424 due to collision of the rifted Laurentian margin with a continent facing arc. The model  
425 was put forward on the basis of relatively clear field evidence of deformation, ophiolite  
426 obduction and stratigraphy in the orogen ([Dalziel, 1997](#); [Dewey and Shackleton, 1984](#);  
427 [Dewey and Ryan, 1990](#); [Van Staal et al., 1998](#)). Comparing to the classical collisional  
428 belts, however, evidence for oceanic basin, ophiolite suite and subduction complex  
429 associated with an Early Paleozoic subduction is absent in the South China Block.  
430 Contemporaneous arc volcanics and high pressure metamorphic rocks have so far never  
431 been documented in South China. Turbidites with Bouma sequences are also lacking in  
432 spite of widely developed thick flysch sedimentary successions ([BGMRFJ, 1985](#);  
433 [BGMRGX, 1985](#); [BGMRJX, 1984](#); [Wang et al., 2010](#); [Shu, 2012](#); [Shu et al., 2014](#)).  
434 Consequently, it is unreasonable to interpret the Early Paleozoic tectonics of South  
435 China as related to an oceanic subduction and a subsequent collision. On the contrary,  
436 an intracontinental orogeny accommodated by a continental subduction is likely (e.g.,  
437 [Charvet et al., 2010](#); [Chen et al., 2010](#); [Faure et al., 2008](#); [Shu et al., 2008a](#); [Song et al.,](#)  
438 [2015](#)).

439 In the present study, the lithofacies paleogeography of South China provides  
440 constraints on the source direction of thick siliciclastic sediments in the Southeast basin  
441 that were mainly derived from the southeast, such northwestward sedimentary transport  
442 is also supported by several palaeocurrent indicators preserved in the Cambrian-  
443 Ordovician strata, indicating that a paleoplate probably existed to the southeast of SE  
444 SCB ([Ren, 1964](#); [Ren et al., 1990](#); [Rong et al., 2010](#); [Shu, 2012](#); [Shu et al., 2008a](#); [Wu](#)  
445 [et al., 2010](#)). Regional-scale folding during the Early Paleozoic is widespread in the  
446 South China Fold Belt ([BGMRFJ, 1985](#); [BGMRGX, 1985](#); [BGMRJX, 1984](#); [Jahn et](#)  
447 [al., 1990](#); [Shu et al., 1991](#); [Song et al., 2015](#)). Moreover, kinematic studies in the South



448 China Block show a fan-shaped thrust pattern, with top-to-the-northwest in the  
449 northwestern part of SE South China Block and top-to-the-southeast in the  
450 southeastern part of SE South China Block (Faure et al., 2009; Lin et al., 2008, 2011;  
451 Shu et al., 2014). Such double-vergent thrusting may develop in intraplate settings,  
452 associating with syntectonic magmatism, like Petermann and Alice Springs belts in  
453 central Australia (Haines et al., 2001; Raimondo et al., 2009, 2010, 2014; Wade et al.,  
454 2005) or the Middle Triassic Xuefengshan belt in the central part of South China (Chu  
455 et al., 2012).

456 It is widely accepted that the SE SCB has experienced crustal thickening and  
457 anatexis in the Early Paleozoic (Wang et al., 2010; Zeng et al., 2008). Continent-  
458 continent collision may cause thickening of the continental crust over a large wide zone  
459 (1,000 km), with far-field continental shortening, and last for 50 million years. Such  
460 zones are characterized by a wide orogenic plateau, with surrounding and internal  
461 basins bordered by thrust belts (Dewey, 2005).

462 According to the available geological investigations, therefore, we propose that  
463 the Early Paleozoic tectono-magmatic event in SE SCB is probably related to the  
464 continent-continent collision. A Suspected East China Sea Block subducted  
465 northwestward beneath the SE SCB together with the NW SCB subducted  
466 southeastward beneath the SE SCB (Fig. 10D), resulting in the crustal thickening and  
467 intracontinental deformation which may be caused by the stress generated at plate  
468 boundaries and stress transmission with the lithosphere acting as an effective stress  
469 guide (Aitken et al., 2013; Gorczyk and Vogt, 2015; Raimondo et al., 2014).  
470 Subsequently, regional extension occurred in response to crustal thickening at an earlier  
471 stage of the orogeny (Dewey, 1988; Strachan, 1994). The clockwise metamorphic P-T  
472 paths in Chencai, Wuyi and Yunkai regions indicate that the SE SCB experienced  
473 isothermal decompression (Li et al., 2010; Wang et al., 2012; Zhao and Cawood, 2012),  
474 representing the progressive release of stress and rapid crustal denudation after the  
475 crustal thickening and uplift. Coincidentally, the Sr-Nd isotopic compositions of Early  
476 Paleozoic granites from the South China Block are similar to those of Caledonian,  
477 Hercynian tectonic belts and other classical collisional belts (Jahn, 2004; Jahn et al.,

478 2014), which probably represent the similar dynamic mechanism during the  
479 decompression and remelting of the crustal sources.

480

#### 481 *6.4 Geodynamic evolution of the South China Block from the Neoproterozoic to* 482 *Early Paleozoic*

483 A possible spatial and temporal geodynamic evolution model accounting for the  
484 Neoproterozoic to Early Paleozoic geological events is proposed in the following.

485 The Jiangnan orogen is a collisional belt developed during the assembly of the  
486 South China block (Charvet et al., 2010; Li et al., 2002; Shu et al., 2011; Yan et al.,  
487 2015; Yao et al., 2012; Zhao and Cawood, 2012). Previous studies have shown that the  
488 Cathaysia and Yangtze blocks collided at ca. 860 Ma along the Shaoxing–Pingxiang–  
489 Guilin fault zone (Shu et al., 2015; Yao et al., 2014) (Figs. 1 and 10A). During the Late  
490 Neoproterozoic, a regional–scale rifting event affected the South China Block, resulting  
491 in the development of fault–bounded basins. The bimodal volcanic rocks  
492 contemporaneous to the rifting were dated at 820–690 Ma (Wang and Li, 2003) (Fig.  
493 10B). During the Cryogenian–Ediacarian (Sinian) to Early Paleozoic, from 690 to 460  
494 Ma, the southeastern part of the South China Block was in a depositional stage.  
495 Stratigraphic sequences in this period are characterized by graptolite bearing sandy–  
496 muddy rock associations, overlying the rift deposits. Proximal deposits, coarse grain  
497 sandstone, conglomerate and ripple–mark structures suggest a littoral–shelf–slope  
498 depositional environment (Shu, 2012; Shu et al., 2014) (Fig. 10C).

499 During the Late Ordovician and Early Silurian, the tectonic framework of South  
500 China was mainly controlled by a regional compression characterized by km–scale  
501 folds and thrusts, the development of southeast–directed décollement shearing,  
502 amphibolite facies metamorphism. During the late to post–thickening evolution, crustal  
503 melting characterized by migmatites and granitic plutons took place (Charvet et al.,  
504 2010; Faure et al., 2009; Shu, 2006, 2012; Shu et al., 2008a, 2015; Song et al., 2015;  
505 Wang et al., 2013b; Zhang et al., 2011). These structural, metamorphic and magmatic  
506 features were related to the northwest–ward continental subduction of the southeastern  
507 part of the Neoproterozoic rift. This continental area can be considered as a part of the

508 inferred East China Sea Block proposed since a long time (Ren, 1964; Ren et al., 1990)  
509 (Fig. 10D). Subsequently, during the Devonian, the South China Block was in a stable  
510 sedimentary environment.

511

## 512 **7. Conclusions**

513 According to the zircon geochronology, and geochemical study of granitoids in  
514 the NW part of the Yunkai domain, we reach the following conclusions:

515 (1) The analyzed plutons mostly crystallized during 426–443 Ma, suggesting an  
516 Early Paleozoic magmatic event in the Yunkai domain;

517 (2) All granites are peraluminous and represent typical anatectic products of  
518 continental crust, mostly plot in the post–collision field in tectonic  
519 discrimination diagrams. Hf isotopic data indicate that parent magma was  
520 derived from Paleoproterozoic–Mesoproterozoic crustal components with  
521 little or no input of mantle sources;

522 (3) The intracontinental orogeny already proposed to account for the Early  
523 Paleozoic plutonic event in the South China Block is well supported by our  
524 new investigations.

525

## 526 **Acknowledgments**

527 We wish to express our gratitude to the editor Sun-Lin Chung, the reviewers Koen  
528 de Jong and Yuejun Wang and the anonymous reviewer for their very helpful and  
529 constructive suggestions that greatly improved this paper. We also express our  
530 appreciation to Dr. Yao J.L. for the help during whole studying process of this study  
531 and to Wang J.Q. and Song M.J. for their assistance during the work of zircon dating  
532 and Hf isotopic analysis. We acknowledge the financial supports provided by The  
533 Major State Research Development Program of China (Grant No. 2016YFC0600202),  
534 National Natural Science Foundation of China (Nos. 41330208, 41572200) and State  
535 Key Laboratory for Mineral Deposits Research (Nanjing University) (ZZKT–201603).

536

## 537 **References**

538 Abdallah, N., Liégeois, J.P., De Waele, B., Fezaa, N., Ouabadi, A., 2007. The  
539 Temaguessine Fe–cordierite orbicular granite (Central Hoggar, Algeria): U–Pb  
540 SHRIMP age, petrology, origin and geodynamical consequences for the late Pan–  
541 African magmatism of the Tuareg shield. *Journal of African Earth Sciences* 49,  
542 153–178.

543 Aitken, A.R.A., Raimondo, T., Capitanio, F.A., 2013. The intraplate character  
544 of supercontinent tectonics. *Gondwana Research* 24, 807–814.

545 Andersen, T., 2002. Correction of common Pb in U–Pb analyses that do not report <sup>204</sup>Pb.  
546 *Chemical Geology* 192, 59–79.

547 BGMRFJ (Bureau of Geology and Mineral Resources of Fujian Province),  
548 1985. *Regional Geology of Fujian Province*. Geological Publishing House, Beijing,  
549 pp. 7–564 (in Chinese with English abstract).

550 BGMRGX (Bureau of Geology and Mineral Resources of Guangxi Province),  
551 1985. *Regional Geology of Guangxi Autonomous Region*. Geological Publishing  
552 House, Beijing, pp. 1–853 (in Chinese with English Abstract).

553 BGMRHN (Bureau of Geology and Mineral Resources of Hunan Province),  
554 1988. *Regional Geology of Hunan Province*. Geological Publishing House, Beijing,  
555 6–664 (in Chinese with English Abstract).

556 BGMRJX (Bureau of Geology and Mineral Resources Jiangxi Province),  
557 1984. *Regional Geology of Jiangxi Province*. Geology Publishing House, Beijing,  
558 pp. 2–725 (in Chinese with English Abstract).

559 BGMRZJ (Bureau of Geology and Mineral Resources Jiangxi Province),  
560 1989. *Regional Geology of Jiangxi Province*. Geology Publishing House, Beijing,  
561 pp. 5–575 (in Chinese with English Abstract).

562 Black, L.P., Gulson, B.L., 1978. The age of the Mud Tank carbonatite, Strangways  
563 Range, Northern Territory. Bureau of Mineral Resources *Journal of Australian*  
564 *Geology and Geophysics* 3, 227–232.

565 Blichert–Toft, J., Albarede, F., 1997. The Lu–Hf isotope geochemistry of chondrites and  
566 the evolution of the mantle–crust system. *Earth and Planetary Science Letters* 148,  
567 243–258.

568 Castro, A., 2014. The off-crust origin of granite batholiths. *Geoscience Frontiers* 5, 63–  
569 75.

570 Cawood, P.A., Wang, Y.J., Xu, Y.J., Zhao, G.C., 2013. Locating South China in Rodinia  
571 and Gondwana: A fragment of greater India lithosphere? *Geology* 41 (8), 903–906.

572 Charvet, J., 2013. The Neoproterozoic–Early Paleozoic tectonic evolution of the South  
573 China Block: an overview. *Journal of Asian Earth Sciences* 74, 198–209.

574 Charvet, J., Shu, L.S., Faure, M., Choulet, F., Wang, B., Lu, H.F., Breton N.L.,  
575 2010. Structural development of the Lower Paleozoic belt of South China: Genesis  
576 of an intracontinental orogen. *Journal of Asian Earth Sciences* 39, 309–330.

577 Chen, X., Zhang, Y.D., Fan, J.X., Cheng, J.F., Li, Q.J., 2010. Ordovician graptolite–  
578 bearing strata in southern Jiangxi with a special reference to the Kwangsi  
579 Orogeny. *Science China Earth Science* 53, 1602–1610.

580 Chu, Y., Lin, W., Faure, M., Wang, Q.C., Ji, W.B., 2012. Phanerozoic tectonothermal  
581 events of the Xuefengshan Belt, central South China: Implications from U–Pb age  
582 and Lu–Hf determinations of granites. *Lithos* 150, 243–255.

583 Chu, Y., Lin, W., 2014. Phanerozoic polyorogenic deformation in southern Jiuling  
584 Massif, northern South China block: Constraints from structural analysis and  
585 geochronology. *Journal of Asian Earth Sciences* 86, 117–130.

586 Corfu, F., Hanchar, J.M., Hoskin, P.W.O., Kinny, P., 2003. Atlas of zircon textures.  
587 *Reviews in Mineralogy and Geochemistry* 53, 469–500.

588 Cox, K.G., Bell, J.D., Pankhurst, R.J., 1979. *The Interpretation of Igneous*  
589 *Rocks*. George Allen & Unwin, London (450p).

590 Dalziel, I.W.D., 1997. OVERVIEW: Neoproterozoic–Paleozoic geography and  
591 tectonics: Review, hypothesis, environmental speculation. *Geological Society of*  
592 *America Bulletin* 109 (1), 16–42.

593 De Jong, K., Han, S., Ruffet, G., 2015. Fast cooling following a Late Triassic  
594 metamorphic and magmatic pulse: implications for the tectonic evolution of the  
595 Korean collision belt. *Tectonophysics* 662, 271–290.

596 Deng, P., Ren, J.S., Ling, H.F., Shen, W.Z., Sun, L.Q., Zhu, B., Tan, Z.Z., 2012.

597 SHRIMP zircon U–Pb ages and tectonic implications for Indosinian granitoids of  
598 southern Zhuguangshan granitic composite, South China. *Chinese Science*  
599 *Bulletin* 57 (13), 1542–1552.

600 Dewey, J.F., 1988. Extensional collapse of orogens. *Tectonics* 7 (6), 1123–1139.

601 Dewey, J.F., 2005. Orogeny can be very short. *Proceedings of the National*  
602 *Academy of Sciences* 102 (43), 15286–15293.

603 Dewey, J.F., Shackleton, R.M., 1984. A model for the evolution of the Grampian tract  
604 in the early Caledonides and Appalachians. *Nature* 312, 115–121.

605 Dewey, J.F., Ryan, P.D., 1990. The Ordovician evolution of the South Mayo trough,  
606 western Ireland. *Tectonics* 9 (4), 887–901.

607 Faure, M., Sun, Y., Shu, L.S., Charvet, J., 1996. Extensional tectonics within a  
608 subductiontype orogen: the case study of the Wugongshan dome (Jiangxi Province,  
609 Southeastern China). *Tectonophysics* 263, 77–106.

610 Faure, M., Lin, W., Monié, P., Meffre, S., 2008. Palaeozoic collision between the North  
611 and South China blocks. Triassic intracontinental tectonics, and the problem of the  
612 ultrahigh-pressure metamorphism. *Comptes Rendus Geoscience* 340 (2), 139–150.

613 Faure, M., Shu, L.S., Wang, B., Charvet, J., Choulet, F., Monié, P., 2009.  
614 Intracontinental subduction: a possible mechanism for the Early Paleozoic Orogen  
615 of SE China. *Terra Nova* 21, 360–368.

616 Faure, M., Lin, W., Chu, Y., Lepvrier, C., 2016. Triassic tectonics of the southern margin  
617 of the South China Block. *Comptes Rendus Geoscience* 348, 5–14.

618 Foden, J., Sossi, P.A., Wawryk, C.M., 2015. Fe isotopes and the contrasting  
619 petrogenesis of A-, I- and S-type granite. *Lithos* 212–215, 32–44.

620 Gao, J.F., Lu, J.J., Lai, M.Y., Lin, Y.P., Pu, W., 2003. Analysis of trace elements in rock  
621 samples using HR-ICP-MS. *Journal of Nanjing University (Natural Sciences)* 39,  
622 844–850 (in Chinese with English abstract).

623 Gorczyk, W., Vogt, K., 2015. Tectonics and melting in intra-continental settings.  
624 *Gondwana Research* 27, 196–208.

625 Grabau, A.W., 1924. *Stratigraphy of China, Part I, Paleozoic and Older*. Geological  
626 Survey of China, Beijing. 528 pp.

627 Griffin, W.L., Belousova, E.A., Shee, S.R., Pearson, N.J., O'Reilly, S.Y., 2004.  
628        Archaean crustal evolution in the northern Yilgarn Craton: U–Pb and Hf–isotope  
629        evidence from detrital zircons. *Precambrian Research* 131, 231–282.

630 Guo, L.Z., Shi, Y.S., Ma, R.S., Dong, H.G., Yang, S.F., 1989. The pre–Devonian  
631        tectonic patterns and evolution of south China. *Journal of Southeast Asian Earth  
632        Sciences* 3, 87–93.

633 Haines, P.W., Hand, M., Sandiford, M., 2001. Palaeozoic synorogenic sedimentation in  
634        central and northern Australia: A review of distribution and timing with  
635        implications for the evolution of intracontinental orogens. *Australian Journal of  
636        Earth Sciences* 48 (6), 911–928.

637 Hoskin, P.W.O., Schaltegger, U., 2003. The composition of zircon and igneous and  
638        metamorphic petrogenesis. *Reviews in mineralogy and geochemistry* 53 (1), 27–  
639        62.

640 Jahn, B.M., Zhou, X.H., Li, J.L., 1990. Formation and tectonic evolution of  
641        southeastern China and Taiwan: isotopic and geochemical constraints.  
642        *Tectonophysics* 183, 145–160.

643 Jahn, B.M., 2004. The Central Asian Orogenic Belt and growth of the continental crust  
644        in the Phanerozoic. *Geological Society, London, Special Publications* 226 (1), 73–  
645        100.

646 Jahn, B.M., Usuki, M., Usuki, T., Chung, S.L., 2014. Generation of Cenozoic granitoids  
647        in Hokkaido (Japan): Constraints from zircon geochronology, Sr–Nd–Hf isotopic  
648        and geochemical analyses, and implications for crustal growth. *American Journal  
649        of Science* 314 (2), 704–750.

650 Jackson, S.E., Pearson, N.J., Griffin, W.L., Belousova, E.A., 2004. The application of  
651        laser ablation microprobe–inductively coupled plasma–mass spectrometry to in  
652        situ U–Pb zircon geochronology. *Chemical Geology* 211, 47–69.

653 John, B., Blundy, J., 1993. Emplacement–related deformation of granitoid magmas,  
654        southern Adamello Massif, Italy. *Geological Society of America Bulletin* 105,  
655        1517–1541.

656 Kemp, A.I.S., Hawkesworth, C.J., Paterson, B.A., Kinny, P.D., 2006. Episodic growth  
657 of the Gondwana supercontinent from hafnium and oxygen isotopes in zircon.  
658 Nature 439, 580–583.

659 Kay, S.M., Coira, B., Wörner, G., Kay, R.W., Singer, B.S., 2011. Geochemical, isotopic  
660 and single crystal  $^{40}\text{Ar}/^{39}\text{Ar}$  age constraints on the evolution of the Cerro Galán  
661 ignimbrites. Bulletin of Volcanology 73 (10), 1487–1511.

662 Li, S.Z., Santosh, M., Zhao, G.C., Zhang, G.W., Jin, C., 2012. Intracontinental  
663 deformation in a frontier of super–convergence: A perspective on the tectonic  
664 milieu of the South China Block. Journal of Asian Earth Sciences 49, 313–329.

665 Li, W.X., Li, X.H., Li Z.X., 2005. Neoproterozoic bimodal magmatism in the Cathaysia  
666 Block of South China and its tectonic significance. Precambrian Research 136,  
667 51–66.

668 Li, W.X., Li, X.H., Li, Z.X., Lou, F.S., 2008a. Obduction–type granites within the NE  
669 Jiangxi Ophiolite: Implications for the final amalgamation between the Yangtze  
670 and Cathaysia Blocks. Gondwana Research 13, 288–301.

671 Li, X.H., 1990. The magmatic period and crustal activity of Wanyangshan–  
672 Zhuguangshan compound granite batholith. Science China 7, 747–755 (in Chinese).

673 Li, X.H., 1994. A comprehensive U–Pb, Sm–Nd, Rb–Sr and  $^{40}\text{Ar}$ – $^{39}\text{Ar}$   
674 geochronological study on Guidong Granodiorite, southeast China: Records of  
675 multiple tectonothermal events in a single pluton. Chemical Geology 115 (3), 283–  
676 295.

677 Li, X.H., 1997. Timing of the Cathaysia block formation: Constraints from SHRIMP  
678 U–Pb zircon geochronology. Episodes 20, 188–192.

679 Li, X.H., Li, W.X., Li, Z.X., Lo, C.H., Wang, J., Ye, M.F., Yang, Y.H., 2009.  
680 Amalgamation between the Yangtze and Cathaysia blocks in South China:  
681 constraints from SHRIMP U–Pb zircon ages, geochemistry and Nd–Hf isotopes of  
682 the Shuangxiwu volcanic rocks. Precambrian Research 174 (1–2), 117–128.

683 Li, Z.X., Li, X.H., Zhou, H.W., Kinny, P.D., 2002. Grenvillian continental collision in  
684 south China: New SHRIMP U–Pb zircon results and implications for the config–  
685 uration of Rodinia. Geology 30, 163–166.



- 686 Li, Z.X., Li, X.H., Kinny, P.D., Wang, J., Zhang, S., Zhou, H.W., 2003. Geochronology  
687 of Neoproterozoic syn–rift magmatism in the Yangtze Craton, South China and  
688 correlations with other continents: evidence for a mantle superplume that broke up  
689 Rodinia. *Precambrian Research* 122, 85–109.
- 690 Li, Z.X., Bogdanova, S.V., Collins, A.S., Davidson, A., De Waele, B., Ernst, R.E.,  
691 Fitzsimons, I.C.W., Fuck, R.A., Gladkochub, D.P., Jacobs, J., Karlstrom, K.E., Lu,  
692 S., Natapov, L.M., Pease, V., Pisarevsky, S.A., Thrane, K., Vernikovsky, V., 2008b.  
693 Assembly, configuration, and break–up history of Rodinia: A synthesis.  
694 *Precambrian Research* 160, 179–210.
- 695 Li, Z.X., Li, X.H., Wartho, J.A., Clark, C., Li, W.X., Zhang, C.L., Bao, C.M., 2010.  
696 Magmatic and metamorphic events during the Early Paleozoic Wuyi–Yunkai  
697 orogeny, southeastern South China: new age constraints and pressure–temperature  
698 conditions. *Geological Society of America Bulletin* 122 (5/6), 772–793.
- 699 Lin, W., Wang, Q.C., Chen, K., 2008. Phanerozoic tectonics of south China block: New  
700 insights from the polyphase deformation in the Yunkai massif. *Tectonics* 27 (6),  
701 TC6004.
- 702 Lin, W., Faure, M., Lepvrier, C., Chen, Z.C., Chu, Y., Wang, Q.C., Vuong, N.V., Tich,  
703 V.V., 2011. The Early Mesozoic thrust and folds sheet structure along the southern  
704 margin of South China Block and its geodynamic. *ScientiaGeologicaSinica* 46 (1),  
705 134–145.
- 706 Liu, R., Zhou, H.W., Zhang, L., Zhong, Z.Q., Zeng, W., Xiang, H., Jin, S., Lu, X.Q., Li,  
707 C.Z., 2010. Zircon U–Pb ages and Hf isotope compositions of the Mayuan  
708 migmatite complex, NW Fujian Province, Southeast China: constraints on the  
709 timing and nature of a regional tectonothermal event associated with the  
710 Caledonian orogeny. *Lithos* 119, 163–180.
- 711 Lou, F.S., Shen, W.Z., Wang, D.Z., Shu, L.S., Wu, F.J., Zhang, F.R., Yu, J.H., 2006.  
712 Zircon U–Pb Isotopic Chronology of the Wugongshan Dome Compound Granite  
713 in Jiangxi Province. *ActaGeologicaSinica* 79 (5), 636–644 (in Chinese with  
714 English Abstract).
- 715 MaksaeV, V., Munizaga, F., McWilliams, M., Fanning, M., Mathur, R., Ruiz, J.,

716 Zentilli, M., 2004. New Chronology for El Teniente, Chilean Andes, from U–Pb,  
717  $^{40}\text{Ar}/^{39}\text{Ar}$ , Re–Os and Fission–Track Dating: Implications for the evolution of a  
718 Supergiant Porphyry Cu–Mo Deposit. *Andean Metallogeny: New Discoveries,*  
719 *Concepts Update.* Society of Economic Geologists, Special Publication 11, 15–54.

720 Maniar, P.D., Piccoli, P.M., 1989. Tectonic discrimination of granitoids. *Geological*  
721 *Society of America Bulletin* 101 (5), 635–643.

722 McDonough, W.F., Sun, S.S., 1995. Composition of the Earth. *Chemical Geology* 120,  
723 223–253.

724 McDougall, I., Harrison, T.M., 1999. *Geochronology and Thermochronology by the*  
725  *$^{40}\text{Ar}/^{39}\text{Ar}$  Method.* Oxford University Press on Demand.

726 McKerrow, W.S., Niocaill, C.M., Dewey, J.F., 2000. The Caledonian orogeny redefined.  
727 *Journal of the Geological Society* 157 (6), 1149–1154.

728 Pearce, J.A., 1996. Sources and setting of granitic rocks. *Episodes* 19 (4), 120–125.

729 Pearce, J.A., Harris, N.B., Tindle, A.G., 1984. Trace element discrimination diagrams  
730 for the tectonic interpretation of granitic rocks. *Journal of Petrology* 25 (4), 956–  
731 983.

732 Peng, S.B., Jin, Z.M., Liu, Y.H., Fu, J.M., He, L.Q., Cai, M.H., Wang, Y.B., 2006.  
733 Petrochemistry, chronology and tectonic setting of strong peraluminous anatectic  
734 granitoids in Yunkai Orogenic Belt, western Guangdong Province, China. *Journal*  
735 *of China University of Geosciences* 17 (1), 1–12.

736 Peng, S.B., Liu, S.F., Lin, M.S., Wu, C.F., Han, Q.S., 2016a. Early  
737 Paleozoic Subduction in Cathaysia (I): New Evidence from Nuodong Ophiolite.  
738 *Earth Science* 41 (5), 765–778 (in Chinese with English abstract).

739 Peng, S.B., Liu, S.F., Lin, M.S., Wu, C.F., Han, Q.S., 2016b. Early Paleozoic  
740 Subduction in Cathaysia (II): New Evidence from the Dashuang High Magnesian-  
741 Magnesian Andesite. *Earth Science* 41 (6), 931–947 (in Chinese with English  
742 abstract).

743 Pitcher, W.S., 1983. Granite: typology, geological environment and melting  
744 relationships, in M. P. Atherton, and C. D. Gribble, eds. *Migmatites, Melting and*  
745 *Metamorphism.* Nantwich: Shiva, 277–287.

746 Qin, X.F., Wang Z.Q., Hu, G.A., Cao, J., Feng, Z.H., 2013. Geochronology and  
747 geochemistry of Hudong gneissic composite pluton in the junction of Guangdong  
748 and Guangxi provinces: Implications for Early Paleozoic tectono–magmatism  
749 along the northern margin of Yunkai massif. *Acta Petrologica Sinica* 29 (9), 3115–  
750 3130.

751 Raimondo, T., Collins, A.S., Hand, M., Walker–Hallam, A., Smithies, R.H., Evins, P.M.,  
752 Howard, H.M., 2009. Ediacaran intracontinental channel flow. *Geology* 37 (4),  
753 291–294.

754 Raimondo, T., Collins, A.S., Hand, M., Walker–Hallam, A., Smithies, R.H., Evins, P.M.,  
755 Howard, H.M., 2010. The anatomy of a deep intracontinental orogen. *Tectonics*  
756 29 (4), TC4024.

757 Raimondo, T., Hand, M., Collins, W.J., 2014. Compressional intracontinental orogens:  
758 Ancient and modern perspectives. *Earth–Science Reviews* 130, 128–153.

759 Ren, J.S., 1964. A preliminary study on pre–Devonian geotectonic problems of  
760 southeastern China. *Acta Geologica Sinica* 44(4), 418–431.

761 Ren, J.S., Chen, T.Y., Niu, B.G., 1990. The tectonics and mineralization of continental  
762 lithosphere in the East China and adjacent region: Beijing, Science Press, 205 p.  
763 (in Chinese with English abstract).

764 Reynolds, P.H., Zentilli, M., Muecke, G.K., 1981. K–Ar and  $^{40}\text{Ar}/^{39}\text{Ar}$  geochronology  
765 of granitoid rocks from southern Nova Scotia: its bearing on the geological  
766 evolution of the Meguma Zone of the Appalachians. *Canadian Journal of Earth*  
767 *Sciences* 18 (2), 386–394.

768 Rollinson, H., 1993. Using geochemical data: evaluation, presentation, interpretation.  
769 Longman, Singapore, pp. 1–56.

770 Rong, J.Y., Zhan, R.B., Xu, H.G., Huang, B., Yu, G.H., 2010. Expansion of the  
771 Cathaysian Oldland through the Ordovician–Silurian transition: Emerging  
772 evidence and possible dynamics. *Science in China Series D: Earth Sciences* 53 (1),  
773 1–17.

774 Scherer, E., Munker, C., Mezger, K., 2001. Calibration of the Lutetium–Hafnium clock.  
775 *Science* 293, 683–687.

776 Shu, L.S., 2006. Pre-Devonian tectonic evolution of South China: from Cathaysia  
777 Block to Caledonian period folded Orogenic Belt. *Geological Journal of China*  
778 *Universities (Earth Sci.)* 12 (4), 418–431 (in Chinese with English abstract).

779 Shu, L.S., 2012. An analysis of principal features of tectonic evolution in South China  
780 Block. *Geological Bulletin of China* 31(7), 1035–1053 (in Chinese with English  
781 abstract).

782 Shu, L.S., Charvet, J., 1996. Kinematic and geochronology of the Proterozoic  
783 Dongxiang–Shexian ductile shear zone (Jiangnan region), South China.  
784 *Tectonophysics* 267 (1–4), 291–302.

785 Shu, L.S., Charvet, J., Shi, Y.S., Faure, M., Cluzel, D., Guo, L.Z., 1991. Structural  
786 analysis of the Nanchang–Wanzaisinistral ductile shear zone (Jiangnan region,  
787 South China). *Journal of Southeast Asian Earth Sciences* 6 (1), 13–23.

788 Shu, L.S., Lu, H.F., Charvet, J., Faure, M., 1997. Kinematic study of the northern  
789 marginal fault zone of Wuyishan, South China. *Geological Journal of China*  
790 *Universities* 3 (3), 282–292 (in Chinese with English abstract).

791 Shu, L.S., Lu, H.F., Jia, D., Charvet, J., Faure, M., 1999. Study of the  $^{40}\text{Ar}/^{39}\text{Ar}$  Isotopic  
792 Age For the Early Paleozoic Tectonothermal Event In the Wuyishan Region, South  
793 China. *Journal of Nanjing University (Natural Sciences)* 35 (6), 668–674 (in  
794 Chinese with English abstract).

795 Shu, L.S., Faure, M., Jiang, S.Y., Yang, Q., Wang, Y.J., 2006. SHRIMP zircon U–Pb  
796 age, litho and biostratigraphic analyses of the Huaiyu Domain in South China–  
797 Evidence for a Neoproterozoic orogen, not Late Paleozoic–Early Mesozoic  
798 collision. *Episodes* 29(4), 244–252.

799 Shu, L.S., Yu, J.H., Jia, D., Wang, B., Shen, W.Z., Zhang, Y.Q., 2008a. Early Paleozoic  
800 orogenic belt in the eastern segment of South China. *Geological Bulletin of China*  
801 27 (10), 1581–1593 (in Chinese with English abstract).

802 Shu, L.S., Faure, M., Wang, B., Zhou, X.M., Song, B., 2008b. Late Paleozoic–Early  
803 Mesozoic Geological Features of South China: Response to the Indosinian  
804 Collision Event in Southeast Asia. *Comptes Rendus Geosciences* 340 (2–3), 151–  
805 165.

806 Shu, L.S., Faure, M., Yu, J.H., Jahn, B.M., 2011. Geochronological and geochemical  
807 features of the Cathaysia block (South China): New evidence for the  
808 Neoproterozoic breakup of Rodinia: *Precambrian Research* 187 (3–4), 263–276.

809 Shu, L.S., Jahn, B.M., Charvet, J., Santosh, M., Wang, B., Xu, X.S., Jiang, S.Y., 2014.  
810 Early Paleozoic depositional environment and intracontinental orogeny in the  
811 Cathaysia Block (South China): implications from stratigraphic, structural,  
812 geochemical and geochronologic evidence. *American Journal of Science* 314(1),  
813 154–186.

814 Shu, L.S., Wang, B., Cawood P.A., Santosh, M., Xu, Z.Q., 2015. Early Paleozoic and  
815 early Mesozoic intraplate tectonic and magmatic events in the Cathaysia Block,  
816 South China. *Tectonics* 34 (8), 1600–1621.

817 Song, M.J., Shu, L.S., Santosh, M., Li, J.Y., 2015. Late Early Paleozoic and Early  
818 Mesozoic intracontinental orogeny in the South China Craton: Geochronological  
819 and geochemical evidence. *Lithos* 232, 360–374.

820 Strachan, R.A., 1994. Evidence in North-East Greenland for Late Silurian-Early  
821 Devonian regional extension during the Caledonian orogeny. *Geology* 22 (10),  
822 913–916.

823 Sun, S.S., McDonough, W.F., 1989. Chemical and isotopic systematics of oceanic  
824 basalts: implication for mantle composition and processes. In: Saunders, A.D.,  
825 Norry, M.J. (Eds.), *Magmatism in the Ocean Basins* Geological Society 42. Special  
826 Publications, London, pp. 313–345.

827 Van Staal, C.R., Dewey, J.F., Niocaill, C.M., McKerrow, W.S., 1998. The Cambrian-  
828 Silurian tectonic evolution of the northern Appalachians and British Caledonides:  
829 history of a complex, west and southwest Pacific-type segment of Iapetus.  
830 Geological Society, London, Special Publications 143 (1), 197–242.

831 Vignerresse, J.L., 2014. Textures and melt–crystal–gas interactions in granites.  
832 *Geoscience Frontiers* 6 (5), 635–663.

833 Wade, B.P., Hand, M., Barovich, K.M., 2005. Nd isotopic and geochemical constraints  
834 on provenance of sedimentary rocks in the eastern Officer Basin, Australia:  
835 Implications for the duration of the intracratonic Petermann Orogeny. *Journal of*

836 the Geological Society 162, 513–530.

837 Wang, J., Li, Z.X., 2003. History of Neoproterozoic rift basins in South China:  
838 Implications for Rodinia break-up. *Precambrian Research* 122 (1–4), 141–158.

839 Wang, X.L., Zhou, J.C., Qiu, J.S., Zhang, W.L., Liu, X.M., Zhang, G.L., 2006. LA-  
840 ICP-MS U-Pb zircon geochronology of the Neoproterozoic igneous rocks from  
841 Northern Guangxi, South China: implications for petrogenesis and tectonic  
842 evolution. *Precambrian Research* 145(1–2), 111–130.

843 Wang, Y.J., Fan, W.M., Zhao, G.C., Ji, S.C., Peng, T.P., 2007. Zircon U-Pb  
844 geochronology of gneissic rocks in the Yunkai massif and its implications on the  
845 Caledonian event in the South China Block. *Gondwana Research* 12 (4), 404–416.

846 Wang, Y.J., Zhang, F.F., Fan, W.M., Zhang, G.W., Chen, S.Y., Cawood, P.A., Zhang,  
847 A.M., 2010. Tectonic setting of the South China Block in the Early Paleozoic:  
848 Resolving intracontinental and ocean closure models from detrital zircon U-Pb  
849 geochronology. *Tectonics* 29 (6), TC6020, doi:10.1029/2010TC002750.

850 Wang, Y.J., Zhang, A.M., Fan, W.M., Zhao, G.C., Zhang, G.W., Zhang, Y.Z., Zhang,  
851 F.F., Li, S.Z., 2011. Crustal anatexis in an intracontinental regime: Geochemical,  
852 zircon U-Pb geochronological and Hf isotopic fingerprints from the middle  
853 Paleozoic (Kwangian) gneissoid granites in the eastern South China Block.  
854 *Lithos* 127, 239–260.

855 Wang, Y.J., Wu, C.M., Zhang, A.M., Fan, W.M., Zhang, Y.H., Zhang, Y.Z., Peng, T.P.,  
856 Yin, C.Q., 2012. Kwangian and Indosinian reworking of the eastern South China  
857 Block: Constraints on zircon U-Pb geochronology and metamorphism of  
858 amphibolites and granulites. *Lithos* 150, 227–242.

859 Wang, Y.J., Zhang, A.M., Fan, W.M., Zhang, Y.H., Zhang, Y.Z., 2013a. Origin of  
860 paleosubduction-modified mantle for Silurian gabbro in the Cathaysia Block:  
861 geochronological and geochemical evidence. *Lithos* 160, 37–54.

862 Wang, Y.J., Fan, W.M., Zhang, G.W., Zhang, Y.H., 2013b. Phanerozoic tectonics of the  
863 South China Block: Key observations and controversies. *Gondwana Research* 23,  
864 1273–1305.

865 Wu, L., Jia, D., Li, H.B., Deng, F., Li, Y.Q., 2010. Provenance of detrital zircons from

866 the late Neoproterozoic to Ordovician sandstones of South China: implications for  
867 its continental affinity. *Geological Magazine* 147, 974–980.

868 Xia, Y., Xu, X.S., Zou, H.B., Liu, L., 2014. Early Paleozoic crust–mantle interaction  
869 and lithosphere delamination in South China Block: Evidence from geochronology,  
870 geochemistry, and Sr–Nd–Hf isotopes of granites. *Lithos* 184, 416–435.

871 Xu, B., Guo, L.Z., Shi, Y.S., 1992. The Proterozoic Terranes and Multiphase Collision–  
872 Orogens in Anhui–Zhejiang–Jiangxi Area. Geological Publishing House, Beijing,  
873 pp. 1–112 (in Chinese with English abstract).

874 Xu, X.B., Zhang, Y.Q., Shu, L.S., Jia, D., 2011. La–ICP–MS U–Pb and  $^{40}\text{Ar}/^{39}\text{Ar}$   
875 geochronology of the sheared metamorphic rocks in the Wuyishan: Constraints on  
876 the timing of Early Paleozoic and Early Mesozoic tectono–thermal events in SE  
877 China. *Tectonophysics* 501 (1–4), 71–86.

878 Yan, C.L., Shu, L.S., Santosh, M., Yao, J.L., Li, J.Y., Li, C., 2015. The Precambrian  
879 tectonic evolution of the western Jiangnan Orogen and western Cathaysia Block:  
880 Evidence from detrital zircon age spectra and geochemistry of clastic rocks.  
881 *Precambrian Research* 268, 33–60.

882 Yang, D.S., Li, X.H., Li, W.X., Liang, X.Q., Long, W.G., Xiong, X.L., 2010. U–Pb and  
883  $^{40}\text{Ar}$ – $^{39}\text{Ar}$  geochronology of the Baiyunshan gneiss (central Guangdong, south  
884 China): constraints on the timing of Early Paleozoic and Mesozoic tectonothermal  
885 events in the Wuyun (Wuyi–Yunkai) Orogen. *Geological Magazine* 147, 481–496.

886 Yao, J.L., Shu, L.S., Santosh, M., 2011. Detrital zircon U–Pb geochronology, Hf–  
887 isotopes and geochemistry–New clues for the Precambrian crustal evolution of  
888 Cathaysia Block, South China. *Gondwana Research* 20, 553–567.

889 Yao, J.L., Shu, L.S., M. Santosh, Li, J.Y., 2012. Precambrian crustal evolution of the  
890 South China Block and its relation to supercontinent history: Constraints from U–  
891 Pb ages, Lu–Hf isotopes and REE geochemistry of zircons from sandstones and  
892 granodiorite. *Precambrian Research* 208–211, 19–48.

893 Yao, J.L., Shu, L.S., M. Santosh, Xu, Z.Q., 2013. Palaeozoic metamorphism of the  
894 Neoproterozoic basement in NE Cathaysia: Zircon U–Pb ages, Hf isotope and

895 whole rock geochemistry from the Chencai Group. *Journal of the Geological*  
896 *Society, London* 171 (2), 281–297.

897 Yao, J.L., Shu, L.S., M. Santosh, Zhao, G.C., 2014. Neoproterozoic arc-related mafic–  
898 ultramafic rocks and syn-collision granite from the western segment of the  
899 Jiangnan Orogen, South China: Constraints on the Neoproterozoic assembly of the  
900 Yangtze and Cathaysia Blocks. *Precambrian Research* 243, 39–62.

901 Yu, J.H., Wang, L.J., O'Reilly, S.Y., Griffin, W.L., Zhang, M., Li, C.Z., Shu, L.S., 2009.  
902 A Paleoproterozoic orogeny recorded in a long-lived cratonic remnant (Wuyishan  
903 terrane), eastern Cathaysia Block, China. *Precambrian Research* 174, 347–363.

904 Yuan, H.L., Gao, S., Dai, M.N., Zong, C.L., Günther, D., Fontaine, G.H., Liu, X.M.,  
905 DiWu, C.R., 2008. Simultaneous determinations of U–Pb age, Hf isotopes and  
906 trace element compositions of zircon by excimer laser ablation quadrupole and  
907 multiple collector ICP–MS. *Chemical Geology* 247 (1–2), 100–118.

908 Zeng, W, Zhang, L, Zhou, H.W., Zhong Z.Q., Xiang, H., Liu, R., 2008. Caledonian  
909 reworking of Paleoproterozoic basement in the Cathaysia Block: Constraints from  
910 zircon U-Pb dating, Hf isotopes and trace elements. *Chinese Science Bulletin* 53,  
911 895–904.

912 Zhang, S.B., Zheng, Y.F., 2013. Formation and evolution of Precambrian continental  
913 lithosphere in south China. *Gondwana Research* 23, 1241–1260.

914 Zhang, Y., Shu, L.S., Chen, X.Y., 2011. Study of geochemistry, geochronology and  
915 petro-genesis of the Early Paleozoic granitic plutons in the central-southern  
916 Jiangxi Province. *Science China (Earth Sciences)* 54(10), 1492–1510.

917 Zhang, Z.L., Yuan, H.H., Nan, Y., 1998. Whole-grain zircon evaporation for age of  
918 Luoyu Formation, Yunkai Group. *Journal of Mineralogy and Petrology* 18 (1), 85–  
919 90 (in Chinese with English abstract).

920 Zhao, G.C., Cawood P.A., 1999. Tectonothermal evolution of the Mayuan assemblage  
921 in the Cathaysia Block: New evidence for Neoproterozoic collisional-related  
922 assembly of the South China Craton. *American Journal of Science* 299 (4), 309–  
923 339.

924 Zhao, G.C., Cawood, P.A., 2012. Precambrian geology of China. *Precambrian Research*



925 222, 13–54.  
926 Zhao, K.D., Jiang, S.Y., Sun, T., Chen, W.F., Ling, H.F., Chen, P.R., 2013. Zircon U–Pb  
927 dating, trace element and Sr–Nd–Hf isotope geochemistry of Paleozoic granites in  
928 the Miao'ershan–Yuechengling batholith, South China: Implication for  
929 petrogenesis and tectonic–magmatic evolution. *Journal of Asian Earth Sciences*  
930 74, 244–264.  
931 Zhong, Y.F., Ma, C.Q., Zhang, C., Wang, S.M., She, Z.B., Liu, L., Xu, H.J., 2013.  
932 Zircon U–Pb age, Hf isotopic compositions and geochemistry of the Silurian  
933 Fengdingshan I–type granite pluton and Taoyuan mafic–felsic complex at the  
934 southeastern margin of the Yangtze block. *Journal of Asian Earth Sciences* 74, 11–  
935 24.

936  
937

#### 938 **Figure captions**

939 **Figure 1.** Tectonic outline of China and the distribution map of Early Paleozoic granites  
940 in the South China Block (Shu et al., 2015; Song et al., 2015), the numbers 1–5 are  
941 corresponding to the Xuefeng, Guidong, Zhuguang, Wugong and Wuyi, respectively.

942 **Figure 2.** Geological sketch map of the Yunkai domain with cross sections A–B and C–  
943 D (BGMRGX, 1985).

944 **Figure 3.** Representative field photos and photomicrographs. (A) Photomicrograph of  
945 Sample 1571 (two–mica granite) in Yonghe of Lianshan county; (B) Photomicrograph  
946 of Sample 1586 (muscovite granite) in Xindi of Wuzhou city.; (C) Photomicrograph of  
947 Sample 1601 (two–mica granite) in Songguangling of Cenxi city; (D) Photomicrograph  
948 of Sample 1621 (muscovite granite) in Beijie of Xinyi city; (E) Photomicrograph of  
949 Sample 1628 (biotite mylonitic granite) in Lingshan of Rongxian county; (F) Field  
950 photograph of Sample 1571; (G) Field photograph of Sample 1586; (H) Field  
951 photograph of Sample 1601; (I) Field photograph of Sample 1628. Mineral  
952 abbreviations: Kfs, K–feldspar; Mc, microcline; Pl, plagioclase; Ms, muscovite; Bt,  
953 biotite; Qtz, quartz.

954 **Figure 4.** Geochemical features of the granitoids from the Yunkai domain. (A)  
955 (Na<sub>2</sub>O+K<sub>2</sub>O) versus SiO<sub>2</sub> diagram (Cox et al., 1979); (B) Al/(Na+K) versus  
956 Al/(Ca+Na+K) diagram (Maniar and Piccoli, 1989); (C) SiO<sub>2</sub> versus K<sub>2</sub>O diagram  
957 (Rollison, 1993).

958 **Figure 5.** Distribution of (A) rare earth elements and (B) trace elements for the samples  
959 derived from the Yunkai domain. The normalization values for (A) and (B) are cited  
960 from Sun and McDonough (1989) and McDonough and Sun (1995), respectively.

961 **Figure 6.** Tectonic discrimination diagrams for the Early Paleozoic granites from the  
962 Yunkai domain. (A) Rb versus (Y+Nb) diagram (after Pearce, 1996; Pearce et al., 1984);  
963 (B) Rb versus (Yb+Ta) diagram (after Pearce, 1996; Pearce et al., 1984).

964 **Figure 7.** Cathodo-luminescence images of zircons from the granitoids of the Yunkai  
965 domain, attached with analyzed locations and U–Pb ages.

966 **Figure 8.** U–Pb concordia plots for the zircon grains from eight granitoid samples.

967 **Figure 9.** Plot of Epsilon Hf(t) versus U–Pb age of zircons from eight granitoid samples.

968 **Figure 10.** Geodynamic evolution models for the South China Block from  
969 Neoproterozoic to Early Paleozoic. (A) Collision of Yangtze and Cathaysia blocks (Shu,  
970 2012; Yao et al., 2014); (B) Rifting stage of the Yangtze–Cathaysia Block (Shu, 2012;  
971 Yao et al., 2014); (C) Stable depositional stage of the SE South China Block (Ren, 1964;  
972 Shu et al., 2014, 2015); (D) Intraplate Orogeny stage of the SE South China Block (Shu,  
973 2012; Shu et al., 2014, 2015). SCB: South China Block. Depth is not in proportion.

974

## 975 **Tables**

976 **Table 1** GPS locations of the samples and petrological descriptions.

977 **Table 2** Major and trace element data for representative samples from the Yunkai  
978 domain.

979 **Table 3** U–Pb data of the zircons from eight granitoid samples from the Yunkai domain.

980 **Table 4** Hf isotope analyses of the zircons from eight granitoid samples from the Yunkai  
981 domain.