

# Constraints on Titan's atmospheric conductivity and buried ocean depth, disclosed by the Schumann resonance

Christian Béghin, Michel Hamelin, C. Sotin, R.C. Whitten, Jean-Jacques Berthelier, Réjean Grard, F. Simões

► **To cite this version:**

Christian Béghin, Michel Hamelin, C. Sotin, R.C. Whitten, Jean-Jacques Berthelier, et al.. Constraints on Titan's atmospheric conductivity and buried ocean depth, disclosed by the Schumann resonance. EPSC-DPS Joint Meeting 2011, Oct 2011, Nantes, France. <<http://meetings.copernicus.org/epsc-dps2011/home.html>>. <insu-01555780>

**HAL Id: insu-01555780**

**<https://hal-insu.archives-ouvertes.fr/insu-01555780>**

Submitted on 4 Jul 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Constraints on Titan's atmospheric conductivity and buried ocean depth, disclosed by the Schumann resonance

C. Béghin (1), M. Hamelin (2), C. Sotin (3), R.C. Whitten (4), J.J. Berthelier (2), R. Grard (5) and F. Simões (6)

(1) LPC2E-CNRS-Université d'Orléans, 3A, Ave. Recherche Scientifique, 45071 Orléans Cedex 2, France

(2) LATMOS-IPSL, Université Pierre et Marie Curie, 4, place Jussieu, 75252 Paris cedex 05, France

(3) Jet Propulsion Laboratory and California Institute of Technology, Pasadena, CA 91109, USA

(4) SETI Institute, 189 Bernardo Ave., Mountain View, CA 94043, USA

(5) RSSD, ESA-ESTEC, European Space Agency, Keplerlaan 1 2200 AG Noordwijk, The Netherlands

(6) NASA-GSFC, Heliophysics Science Division, Greenbelt, MD 20771, USA

### Abstract

After six years of a thorough data analysis of the data collected by the Permittivity, Wave and Altimetry (PWA/HASI) experiment during the descent of the Huygens Probe through Titan's atmosphere in January 2005, we report the major findings inferred from the measurements of low frequency waves and atmospheric conductivity. The observations display a Schumann resonance trapped within Titan's atmospheric cavity. In this presentation, we describe the characteristics of the observed mode, that allow us to constrain the parameters of the ionospheric cavity and to infer the presence of a conductive water-ammonia ocean buried below the surface, at a likely depth of  $70 \pm 10$  km.

### 1. Introduction

On Earth, the Schumann resonance is triggered by lightning activity and it was anticipated that a similar observation on Titan would reveal the presence of a reflecting inner surface requisite to sustain such a resonance in addition to a sharp ionospheric boundary. However, given the absence of any acknowledged lightning activity on Titan (Fischer and Gurnett, 2011), we interpret the observed signal at 36 Hz, as the second spherical harmonic of the Titan's cavity, triggered and sustained by intense electric currents induced in the ionosphere through the Saturn's magnetospheric plasma flow.

### 2. Model and PWA measurements

After ruling out several possible artefacts (Béghin et al., 2009) the observed signal is presently unambiguously interpreted as the second harmonic of a Longitudinal Section Magnetic (LSM<sub>2</sub>) mode of atypical Schumann resonance in an inhomogeneous and radially layered cavity. Although the triggering

current sources and the ground conductivity are strongly different from conditions encountered for the terrestrial TEM modes, the analytical expression of the modal equation of the eigenmodes can be written in a similar way, as a function of a small number of cavity parameters, such as

$$\omega_l = \frac{c}{a} \left[ l(l+1) \frac{h_1 + z_c / \text{Re}\epsilon_c}{h_1 + z_c + \zeta \ln(1/4k^2\zeta^2)} \right]^{1/2} \quad (1)$$

$$Q = \left[ \frac{2z_c\delta / \text{Re}\epsilon_c + \pi\zeta/2}{h_1 + z_c / \text{Re}\epsilon_c} + \frac{\pi\zeta/2}{h_1 + z_c + \zeta \ln(1/4k^2\zeta^2)} \right]^{-1} \quad (2)$$

where  $\omega_l$  is the eigenfrequency for the harmonic  $l$  (here  $l = 2$ ),  $Q$  is the quality factor,  $c$  and  $k$  are the free-space light velocity and the wave vector, respectively,  $a$  is the Titan's radius (2575 km),  $h_1$  is the altitude of the ionospheric conduction boundary,  $z_c$  and  $\epsilon_c$  are the subsurface crust thickness and permittivity, respectively,  $\delta$  is the loss tangent of the crust material, and  $\zeta$  is the scale height of ionospheric conductivity.

As on Earth, the altitude of the conduction boundary  $h_1$  is nearly the region of maximum signal strength (Fig. 1), i.e. between 90-100 km here, so that we may keep only four unknown parameters ( $z_c$ ,  $\epsilon_c$ ,  $\delta$  and  $\zeta$ ) to be cross-constrained by the measured quantities  $\omega_l$  and  $Q$ . The most likely cross-constraint area for the three parameters of the crust is shown in Fig. 2. The value of the ice thickness determined by the present study strongly suggests that the ice crust lies over an ammonia-rich ocean, which is consistent with either convection or conduction processes. We find indeed that a crust thicker than 80 km would be unstable to convection processes unless the viscosity at the base of the crust would be larger than  $2.10^{16}$  Pa.s, which suggests a temperature less than 220 K. On the other hand, a convective 40-km thick crust cannot be in equilibrium with the internal heating, which suggests a crust growing at the expense of the ocean.




Figure 1. Experimental data. ELF Electric Field and Electron Conductivity were measured with sensors mounted on two deployable booms. A 36 Hz resonance with a level of few  $\text{mVm}^{-1}\text{Hz}^{-1/2}$  and a high Quality Factor ( $Q = 6$ ) was clearly identified throughout the descent. A peak of conductivity, attributed to Galactic Cosmic Rays, was observed at around 60 km.




Figure 2. Cross-constrained shadowed area for the ocean depth versus the crust permittivity ( $2 < \text{Re } \epsilon_c < 3$ ) and the ionospheric scale height of conductivity ( $3.2 < \zeta < 5$ ).

### 3. Summary

Our interpretation of the Titan's Schumann resonance complies with a conduction ionospheric boundary lying at  $\sim 100$  km, and a 60 km thermally conductive crust with an ocean temperature of 200 K (Fig.3) or an 80 km convective crust with a minimum viscosity of  $5 \times 10^{15}$  Pa.s.




Figure 3. Heat power in equilibrium with the crust thickness ( $60 < z_c < 80$  km) for a temperature of the crust-ocean interface  $< 260$  K.

### References

- [1] Béghin, C., et al.: New insights on Titan's plasma-driven Schumann resonance inferred from Huygens and Cassini data, *Planet. Space Sci.*, 57, 1872-1888, 2009.
- [2] Fischer, G. and Gurnett, D.A.: The search for Titan lightning radio emissions, *Geophys. Res. Lett.*, 38, L08206, doi:10.1029/2011GL047316, 2011.