

A review of the Late Jurassic–Early Cretaceous charophytes from the northern Aquitaine Basin in south-west France

Roch-Alexandre Benoit, Didier Néraudeau, Carles Martin-Closas

▶ **To cite this version:**

Roch-Alexandre Benoit, Didier Néraudeau, Carles Martin-Closas. A review of the Late Jurassic–Early Cretaceous charophytes from the northern Aquitaine Basin in south-west France. Cretaceous Research, 2017, 79, pp.199-213. 10.1016/j.cretres.2017.07.009 . insu-01574653

HAL Id: insu-01574653


https://insu.hal.science/insu-01574653

Submitted on 16 Aug 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript



A review of the Late Jurassic–Early Cretaceous charophytes from the northern Aquitaine Basin in south-west France

Roch-Alexandre Benoit, Didier Neraudeau, Carles Martín-Closas

PII: S0195-6671(17)30121-0

DOI: 10.1016/j.cretres.2017.07.009

Reference: YCRES 3658

To appear in: *Cretaceous Research*

Received Date: 13 March 2017

Revised Date: 5 July 2017

Accepted Date: 17 July 2017

Please cite this article as: Benoit, R.-A., Neraudeau, D., Martín-Closas, C., A review of the Late Jurassic–Early Cretaceous charophytes from the northern Aquitaine Basin in south-west France, *Cretaceous Research* (2017), doi: 10.1016/j.cretres.2017.07.009.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

A review of the Late Jurassic-Early Cretaceous charophytes from the northern Aquitaine Basin in south-west France

Roch-Alexandre BENOIT [1,2], Didier NERAUDEAU[1] and Carles MARTÍN-CLOSAS[2*]

[1] Université Rennes 1, Géosciences, CNRS UMR 6118, campus de Beaulieu bât. 15, 263 avenue du Général Leclerc, 35042 Rennes cedex, France

[2] Departament de Dinàmica de la Terra i l'Oceà, Facultat de Geologia, Universitat de Barcelona-UB, Marti i Franquès s/n, 08028 Barcelona, Catalonia, Spain.

* Corresponding author

Abstract

Late Jurassic and Early Cretaceous charophyte assemblages from the northern part of the Aquitaine Basin in south-west France are reviewed here to understand their palaeoecological, palaeobiogeographical and biostratigraphic features. Three sites were studied: the Tithonian-lower Berriasian of Chassiron, and the Berriasian of Cherves-de-Cognac and Angeac-Charente. Abundant porocharaceans, less abundant clavatoraceans and scarce characeans recorded in Cherves-de-Cognac and Angeac-Charente indicate that brackish water environments were substituted by freshwater environments eastwards. The occurrence of *Clavator grovesii* var. *grovesii* and morphotypes intermediate with *C. grovesii* var. *discordis* in the same areas is significant from a biostratigraphic viewpoint, since these species belong to the Maillardii, Incrassatus and Nurrensis European charophyte biozones, representing the Berriasian. This observation refutes a previous dating of the Angeac-Charente site and

highlights the absence of Hauterivian-Barremian records in northern Aquitaine, which is in contrast to the more complete Lower Cretaceous record in southern Aquitaine. These contrasting records could be due to differences in the available sedimentary space produced by the opening of the Bay of Biscay during the Barremian.

Keywords

Charophyta; Freshwater; Tithonian; Berriasian; Charentes basin

Introduction

The Aquitaine Basin in south-west France is among the lesser known European basins in terms of its Lower Cretaceous stratigraphic and palaeontological records. However, its location at the boundary between the Iberian and European plates makes it highly significant in understanding the rift processes that occurred in the Late Jurassic and Early Cretaceous during the opening of the North Atlantic Ocean and the Gulf of Biscay. In this sense, the biostratigraphy of the few Lower Cretaceous outcrops in the northern part of the Aquitaine Basin is fundamental in elucidating the basin's evolution and how it compares with the southern part of the Aquitaine Basin.

Deak and Combaz (1967) were the first to find palynological indicators of the "Wealden", i.e., Lower Cretaceous non-marine strata, in the southern part of the north Aquitaine Basin. These observations were also made in the north-western part by Lanceron (1976), Platel and Moreau (1977) and Platel (1978). Much later, a number of studies devoted to the biostratigraphy of the Cretaceous northern Aquitaine Basin showed the occurrence of the Berriasian based on the presence of ostracodes, charophytes and dinoflagellates (Colin et

al., 2004; El Albani et al., 2004), as well as the occurrence of the Albian and Cenomanian (Néraudeau et al., 2002) and the Turonian (Néraudeau et al., 2016) based on palynological evidence. Moreover, a recent study indicated the possible occurrence of Hauterivian-Barremian deposits in Angeac-Charente (Néraudeau et al., 2012).

The present study is an in-depth review of the charophyte assemblages from three localities, Chassiron in the Oléron Island, Cherves-de-Cognac and Angeac-Charente (Fig. 1), which have been only partly studied in the past. The results significantly advance the biostratigraphic understanding of the Upper Jurassic and Lower Cretaceous of the northern Aquitaine Basin, refuting some of the previous biostratigraphic findings and improving the comparison with southern Aquitaine, the so-called Arzacq-Mauléon Basin.

---------- Please insert Fig. 1 near here ----------

Geological setting and stratigraphy

The Aquitaine Basin was a rift basin during most of the Mesozoic. Its structure was affected by the formation of the oceanic crust between Europe and North America from the latest Callovian to the early Tithonian and the subsequent opening of the Gulf of Biscay during the Barremian, the basin forming along the transform plate boundary at the north of the Iberian plate (Schettino and Scotese, 2002; Schnyder et al., 2012; Tugend et al., 2015).

The northern part of the Aquitaine Basin, sometimes referred to as “Charentes Basin”, is limited by normal faults with a north-west-south-east direction (Platel, 1980). The Upper Jurassic and Lowermost Cretaceous deposits predominantly contain carbonates, with evaporites at its edges (Platel, 1980; Cubaynes et al., 1989) and clayey or conglomeratic bone beds (Néraudeau et al., 2012; Vullo et al., 2012, 2014). The Lower Cretaceous record is

extremely limited in the northern Aquitaine Basin, especially for the Berriasian to Aptian (Fig. 2), while the uppermost Albian deposits are better represented (Néraudeau et al., 2002). A more complete sedimentary record of marine facies is present for the Cenomanian transgression, starting with detrital beds followed by limestone deposits.

In the southern part of the Aquitaine Basin (also called Arzacq-Mauléon Basin), a more complete Lower Cretaceous record provides information about the Berriasian, Hauterivian-Barremian and Aptian, such as the frequent marine deposits to the east and increasing non-marine influences to the west (Peybernès and Combes, 1994). An important transgression occurred during the Barremian that was interrupted by emersion episodes (Martín-Closas and Peybernès, 1987; Combes et al., 1998). Winnock (1973) explained the differences between the northern and southern parts of the Aquitaine Basin to be due to the reactivation of a Triassic fault structure ("flexure celtaquitaine") that separated the two areas and enhanced the subsidence to the south during the Early Cretaceous. Later, chalk sedimentation occurred until the end of the Late Cretaceous, representing the last occurrence of marine sediments.

---------- Please insert Fig. 2 near here ----------

From a palaeogeographical viewpoint, it is assumed that the Aquitaine Basin was a shallow and narrow gulf opening onto the Atlantic domain during the latest Jurassic and earliest Cretaceous (Cubaynes et al., 1989; Hantzpergue and Lafaurie, 1994; Schnyder et al., 2012). During the Kimmeridgian, the Aquitaine Basin was linked to the Paris Basin by the Poitou threshold in its north-eastern part, but this connection ceased before the Tithonian (Thierry, 2000; Schnyder et al., 2012) due to the regressive trend starting in the latest Kimmeridgian (Hantzpergue, 1989). According to Thierry (2000) and Blakey (2011), some

connections with the western Tethys persisted in the Early Cretaceous through the Corbières-Provence platform to the south-east and the Dauphiné Basin to the north-east.

The Jurassic and Cretaceous from northern Aquitaine have been studied as early as the beginning of the 19th century by Fleuriau de Bellevue (1801) and d'Archiac (1837), among others, who concluded that the Lower Cretaceous was not represented in that part of the basin, a hypothesis that had been widely accepted until recently (Platel, 1980; Néraudeau et al., 2012). In the two last decades, a number of studies have provided evidence that the Lower Cretaceous is indeed represented (Schnyder, 2003; Billon-Bruyat, 2003; Colin et al., 2004; Néraudeau et al., 2002, 2012; Vullo et al., 2014). Thus, the present study aimed to perform a detailed review of the Lower Cretaceous records in three different areas of the "Charentes Basin", Chassiron (Oléron Island) in the western part, Cherves-de-Cognac in the central part and Angeac-Charente in the eastern part (Fig. 1).

Chassiron (Oléron Island)

The stratigraphic section studied at Chassiron is composed of four informal members (Fig. 3). The first, up to 24.2 metres (m) thick, is composed of yellowish to grey irregular nodular limestone with thin marl intercalations. These are thought to represent open-marine platform facies (Schnyder et al., 2012). However, there is already evidence of episodic emersion (e.g., mudcracks) 18 m above the base. Furthermore, the presence of charophytes between 20 and 24.2 m from the base suggests increasing non-marine influences (Schnyder et al., 2012). This member is thought to range from the upper Kimmeridgian (Autissiodorensis Zone) to the lower Tithonian (Gigas Zone), according to Hantzpergue et al. (2004).

The second member, 20.5 m thick, begins with a conglomerate composed of a clay matrix and white calcareous clasts (bed 141). It is topped with 0.4-m-thick clay containing abundant conifer wood (Schnyder et al., 2012; Vullo et al., 2014). The upper part of the member is formed of 19 m of blue to black clay and marls, while the bottom part contains an important accumulation of vertebrate remains (Vullo et al., 2014). Charophyte gyrogonites were found in bed 148 (Fig. 3) and documented by Martín-Closas et al. (2008), who attributed them to the Tithonian. A previously unreported charophyte species from the same bed is described herein.

The third member, with a thickness of 23 m, is composed of limestone beds at the base (6 m), followed by 4 m of alternating limestone and clay, with brackish bivalves (e.g., *Mactra*) and diverse marine species (such as corals, brachiopods, echinoids, bivalves, benthic foraminifera, ostracods and *Rhizocorallium* burrows). This member corresponds to sedimentation occurring in a shallow lagoon and other coastal environments, with normal salinity that has been occasionally subjected to storms (Schnyder et al., 2012).

The fourth member of this series, which is 13 to 14 m thick, is composed of finely laminated and marly peloidal limestone, showing dinosaur footprints and numerous mudcrack-bearing surfaces (Moreau et al., submitted). Evaporitic beds (including calcite recrystallised after gypsum), halite pseudomorphs and fenestrae have been observed in some parts. These are the shallowest deposits in this stratigraphic section (Schnyder et al., 2012).

Members 2 to 4 have been assimilated into Purbeckian-type facies, with a strong terrestrial influence (Schnyder et al., 2012; Colombié et al., 2012; Vullo et al., 2014). The dating of the marine beds from this section was performed using calcareous nannofossils and dinoflagellates (Schnyder et al., 2012), ammonites (Hantzpergue et al., 2004), ostracods (Donze, 1960; Malz, 1966) and foraminifera (Bousquet, 1967), which gave a date ranging

149 from the late Kimmeridgian (Autissiodorensis Zone) to late-early Tithonian or early
150 Berriasian.
151
152 Cherves-de-Cognac
153
154 In the Champblanc quarry at Cherves-de-Cognac, two lithological units were
155 identified, each with a thickness of 15 m (Colin et al., 2004). The basal unit (U1) is composed
156 of alternating layers of fibrous gypsum and black clay laminites, sometimes with stromatolitic
157 features (Colin et al., 2004; El Albani et al., 2004), corresponding to an environment of tidal
158 flats in a restricted lagoon under an arid climate (El Albani et al., 2004).
159 The second unit (U2) shows alternating marlstone laminites and gypsum that
160 progressively gives way to fossilerous marly limestones in the upper parts (Colin et al., 2004;
161 El Albani et al., 2004). This unit indicates both marine and freshwater influences in an
162 estuarine context (El Albani et al., 2004). It contains a 1-m-thick bone bed that bears a
163 particularly rich vertebrate fauna including fish, crocodilians and chelonians (Colin et al.,
164 2004). Charophyte specimens were found in beds C32 to C36, C38 and C41, including the
165 bone bed (Fig. 3). Colin et al. (2004) undertook a preliminary study of these charophytes and
166 here, we will review the flora in depth. The Cherves-de-Cognac study area lies directly over
167 Tithonian marine limestone (El Albani et al., 2004).
168
169 Angeac-Charente
170
171 The Angeac-Charente study area is located between the cities of Cognac and
172 Angoulême. It is composed of 7 characteristic beds, An7 to An1 from the base to the top. An7
173 and An6 are composed of red clay and irregularly stratified grey limestone, respectively

(Néraudeau et al., 2012). An5 does not crop out entirely, but is at least 0.7 m thick and contains green clay without macrofossils. Some grey-green sub-angular limestone blocks are observed in various places, suggesting reworking (Néraudeau et al., 2012). An4 is composed of blue-grey lignitic clay, which is generally 0.5 to 1 m thick and rich in plant remains (e.g., wood, cuticles and seeds). This bed yielded most of the large and well-preserved vertebrate remains, as well as woody axes, described in previous studies (Néraudeau, 2011; Néraudeau et al., 2012; Allain et al., 2014). Numerous calcareous clasts occur at the top of An4, and the delimitation with An3 is unclear (Néraudeau et al., 2012). An3 is 0.2 to 0.4 m thick, and composed of conglomeratic calcareous deposits with a blue-grey grainstone matrix and whitish limestone clasts. Diverse vertebrate remains (crocodiles, dinosaurs, turtles and fishes), plant debris (wood and cuticles), charophyte fructifications, ostracods and insect coprolites (termites) have been recorded in this bed (Colin et al., 2011; Néraudeau et al., 2012). An2 contains yellow to grey calcareous deposits with an irregular stratification that is perhaps linked to dinosaur trampling. It has a thickness ranging from 0.2 to 0.4 m, and grades from marlstone to limestone laterally and vertically (Néraudeau et al., 2012). This bed supplied some of the charophyte remains studied herein. Finally, An1 at the top of the section, with a thickness generally ranging from 0.1 to 0.2 m, is composed of white to greyish fine sand that is rich in fish remains such as scales, teeth and bones (Néraudeau et al., 2012).

Charophytes were found in bed An4, in the interbed between An4 and An3 (An3-4) and in An2 (Fig. 3). They were previously studied by Néraudeau et al. (2012) and here we will review the assemblage in depth.

---------- Please insert Fig. 3 near here ----------

Material and methods

199

200 Sample collection

201

202 The charophyte samples from Chassiron (Oléron Island) were collected and lent by D.
203 Gendry (University of Rennes 1, France) and J.-P. Colin. The samples from Chassiron came
204 from level 148 (Cha148-coll. D.G.) of member 2, as described by Schnyder et al. (2012) and
205 Vullo et al. (2014). This also corresponds to the bed 1004-A in which Schnyder (2003)
206 characterised a high concentration of vertebrate remains (Vullo et al., 2014).

207 The samples from Cherves-de-Cognac were collected by J.-P. Colin and lent to C.
208 Martín-Closas for this study. They were collected from the beds C32 to C36, C38 and C41, as
209 described in Rees et al. (2013) and Pouech et al. (2015), corresponding to the "bone bed" of
210 the site (Colin et al., 2004; El Albani et al., 2004).

211 The Angeac-Charente samples were obtained from palaeontological excavations in
212 2010. Samples from beds An1 (An1-SA), An2 (An2-SA) and An4 (An4-SA) were obtained
213 from the first excavation (ANG1-02-2010), while samples from An3-4 (An3-4-SB) were
214 collected from the second excavation (ANG2-02-2010), located about 20 m north of the site
215 of the first excavation (Néraudeau et al., 2012).

216

217 Sample preparation

218

219 Samples were disaggregated in a solution of water and hydrogen peroxide (100
220 volumes), washed and then sieved through 200, 500 and 1000 µm mesh sieves. In some cases,
221 anhydrous sodium carbonate was added to deflocculate the clay. Before sieving, sieves were
222 submerged in a solution of methylene blue to mark the residues of previous samples and then
223 washed. Charophyte remains (thalli, gyrogonites and utricles) were manually picked out under

a Wild M5A stereomicroscope at 40x magnification. When necessary, an ultrasonic cleaner was used. Measurements were performed with the software Motic Images Plus 2.0 ML in a Motic BA310 stereomicroscope. Selected fructifications were photographed with a Quanta 200 scanning electron microscope at the Scientific and Technical Services of the Universitat de Barcelona (UB). The gyrogonite and utricle specimens were stored in the Universities of Rennes (France) and Barcelona (Catalonia, Spain). Specifically all figured specimens, labelled IGR-PAL are stored in the collections of the University of Rennes (France).

Systematic palaeontology

Division Charophyta MIGULA, 1897

Class Charophyceae SMITH, 1938

Order Charales LINDLEY, 1836

Family Porocharaceae GRAMBAST, 1962 emend. SCHUDACK, 1993

Genus *Porochara* (MÄDLER, 1955) emend. SCHUDACK, 1986

Porochara westerbeckensis (MÄDLER, 1952) MÄDLER, 1955

Fig. 4 (A-E).

1952 *Aclistochara westerbeckensis* nov. spec. – Mädler, p. 28-29, pl. B, fig. 20-25.

1955 *Porochara westerbeckensis* nov. comb. – Mädler, p. 271.

Material. About 250 specimens from An3-4-SB and around 120 specimens from Cherves-de-Cognac samples.

Description. Medium-sized gyrogonites, subprolate to prolate in shape. Specimens from Angeac (An3-4-SB) have a size ranging from 400 to 620 µm in height (mean: 505.7 µm) and 285 to 463 µm in width (mean: 385.3 µm). The isopolarity index (ISI) is above 120 for the majority of the gyrogonites (between 112 and 151, with a mean of 131.5). The number of convolutions (NC) is variable, generally 9 (8-10). The apical pore (mean: 48.34 µm) is wider than the basal pore (mean: 32.51 µm) and are very similar in shape, but the apical pore tends to be star shaped or circular and the basal pore pentagonal.

Remarks. Twelve smaller specimens were found in this population (height: 400-471 µm; width: 285-331 µm; ISI: 120-151; and NC: 8-12) that could be accommodated within *Porochara fusca* (MÄDLER, 1952) MÄDLER, 1955. However, Mojon (1989a) pointed out that the polymorphism of the porocharacean populations of the Upper Jurassic and Lower Cretaceous was high, and interpreted this in terms of palaeoecology.

Distribution. *Porochara westerbeckensis* has been found in north-west Germany from the upper Oxfordian to the lower Berriasian (Schudack, 1990, 1993), in south-east France and Switzerland from the Oxfordian/ Kimmeridgian (Mojon, 1989b), in Spain from the Kimmeridgian (Brenner, 1976) to the Barremian (Martín-Closas, 2000), in Portugal from the Kimmeridgian (Grambast-Fessard and Ramalho, 1985) and in Russia from the Kimmeridgian and Tithonian (Shaïkin, 1976).

---------- Please insert Fig. 4 near here ----------

Genus *Latochara* MÄDLER, 1955 emend. FEIST in FEIST and CUBAYNES, 1984

***Latochara latitruncata* (PECK, 1937) MÄDLER, 1955**

Fig. 4 (F-I).

1937 *Aclistochara latitruncata* PECK, n. sp. – PECK, p. 89, pl. 14, figs 1-4

1955 *Latochara latitruncata* (PECK) nov. comb. – MÄDLER, p. 271

1957 *Latochara latitruncata* (PECK) – PECK, p. 32-33, pl. 5, figs. 7, 21-23

1957 *Latochara collina* PECK, n. sp. – PECK, p. 33, pl. 5, figs 1-4.

1957 *Latochara concinna* PECK, n. sp. – PECK, p. 34, pl. 5, figs. 5-6, 8-9

Material. About 200 specimens from the Chassiron samples (Level 148b).

Description. Small- to medium-sized gyrogonites, 326-424 µm high (mean: 388.65 µm) and 254-330 µm wide, with an ISI from 115 to 142 (mean: 128.77). These gyrogonites are subspheroidal to ellipsoidal in shape with an apical neck. The number of convolutions varies from 10 to 13. The spiral cells are flat to concave, with a carinated suture. When approaching the apical part, the spiral cells are wider in the periapical zone and then narrow up to the apical neck. In a few specimens, a periapical depression is present. The apical pore is very small (hardly visible under the stereomicroscope) and the basal plate cannot be observed, but is thought to be multipartite like in the other species of this genus.

Remarks. Peck (1957) noticed some similarities between *L. latitruncata* and *L. concinna*, and suggested also that *L. collina* was a subspecies of *L. latitruncata*. Schudack (1993) and Martín-Closas et al. (2008) regarded the three species as synonymous.

Distribution. This species has been identified in the Morrison Formation (Kimmeridgian to Tithonian in age) in the United States of America (Peck, 1957), from the

lower Tithonian of north-west Germany (Schudack, 1990, 1993) and in the Tithonian from Ukraine (Shaïkin, 1967). Martín-Closas et al. (2008) also described this species from the same area as that studied here.

Family Characeae (RICHARD. ex C.A. AGARDH, 1824) emend. MARTÍN-CLOSAS and SCHUDACK, 1991

Genus *Mesochara* GRAMBAST, 1962

Mesochara* gr. *voluta sensu Martín-Closas (2000)

Mesochara harrisii (MÄDLER, 1952) SHAÏKIN, 1967

Fig. 4 (J-P).

1952 *Tolypella harrisii* nov. spec. – Mädler, p. 31-32, pl. B, fig.31-35

1952 *Tolypella amoena* nov. spec. – Mädler, p. 34-35, pl. B, fig. 43-49

1952 *Tolypella minuta* nov. spec. – Mädler, p. 35-36, pl. B, fig. 50-52

1967 *Mesochara harrisii* SHAÏKIN nov. comb. – Shaïkin, p. 47

Material. 100 gyrogonites from Chassiron (level 148), 25 specimens from An3-4-SB and a few from An2 from Angeac-Charente, and 15 specimens from Cherves-de-Cognac.

Description. The specimens from Chassiron provide the only abundant population of this species in the northern Aquitaine Basin. They present small gyrogonites ranging from 239 to 341 µm in height and 190 to 312 µm in width, with a spheroidal to prolate shape (ISI ranging from 100 to 142). The spiral cells are concave to almost flat and join at the apex without any modification. The base and apex are generally rounded, but are slightly pointed in some cases. The number of convolutions in lateral view ranges between 8 and 11, generally 9, with a mean width of 36.75 µm at the equator. Some of the specimens from Angeac are

slightly larger (353 to 403 µm high and 303 to 361 µm wide) and more spheroidal (ISI, 108–127). A basal plate cannot be observed, but is probably unicellular, based on gyrogonites of the same species from other European areas.

Remarks. Peck (1957), when describing some specimens of "*Mesochara*" *voluta* that are similar to those found in Charentes, did not separate the specimens with a slightly pointed apex and/or base from those with a rounded apex and/or base. He also noticed a strong similarity with another species, *Tolypella minuta* MÄDLER, 1952 from Germany dating to the Kimmeridgian, which now corresponds to *Mesochara harrisii* (MÄDLER, 1952) SHAÏKIN, 1967. Martín-Closas (2000) assigned *M. harrisii* and *M. voluta* to the same morphogroup (*M.* gr. *voluta)* and remarked that the type population of *M. voluta*, included only 3 gyrogonites, which is an insufficient number to properly characterise a fossil charophyte species.

The difference in size between the two morphotypes present in the An3-4 samples could be the result of ecological variations. Vicente et al. (2016) observed that the different genera of Mesozoic characeans, including *Mesochara*, *Tolypella* and *Microchara*, present the smallest morphotypes in shallow, temporary and turbid ponds of floodplains. Recent data indicate that this might be due to the short life cycles adopted in temporary ponds in response to the very warm and exceedingly well illuminated growth conditions (Sanjuan et al., 2017).

It is important to note here that the basal plate is required to distinguish between the gyrogonites of genus *Mesochara* and the genus *Tolypella* (Martín-Closas et al., 2009). The basal plate is formed by three cells (only two are calcified) in *Tolypella* and is unicellular in *Mesochara*. Therefore, the attribution of the specimens to the correct genera was subject to uncertainty in this study.

Distribution. This species has been observed in north-west Germany, from the Upper Oxfordian to the Berriasian-Valanginian? (Schudack, 1990, 1993), in Spain, from the Kimmeridgian (Brenner, 1976; Schudack, 1987), and Berriasian to Barremian (Martín-Closas and Grambast-Fessard, 1986; Schudack, 1989; Martín-Closas, 2000), in Russia from the Kimmeridgian and Tithonian (Shaïkin, 1967, 1976) and in China from the Lower and Upper Cretaceous (Wang, 1965; Hao et al., 1983; Liu and Wu, 1985). In France, the species has been recorded from the Tithonian (Martín-Closas et al., 2008) and the Berriasian in the Aquitaine Basin (Colin et al., 2004; El Albani et al., 2004; Néraudeau et al., 2012), and in the lowermost Aptian of the Northern Subalpine Chains (Martín-Closas et al., 2009). In Japan, *M. harrisii* has been reported in the lower part of the Kitadani Formation that is thought to date back to the Barremian (Kubota, 2005).

Family Clavatoraceae PIA, 1927

Subfamily Clavatoroidae GRAMBAST, 1969 emend. MARTÍN-CLOSAS 1989 ex SCHUDACK, 1993

Genus *Nodosoclavator* MASLOV, 1963 emend. GRAMBAST, 1966

***Nodosoclavator bradleyi* (HARRIS, 1939) comb. nov. GRAMBAST, 1969**

Fig. 5 (A-H).

1939 *Clavator bradleyi* sp. nov. – HARRIS, p. 53-54, pl. XVI, fig. 1, 3-5, 7.

1969 *Nodosoclavator bradleyi* (HARRIS) nov. comb. – GRAMBAST, p. 881, pl. 33, fig. 5.

Material. About 300 specimens in sample An3-4-SB and a few in sample An2 from Angeac-Charente, 1 from Cherves-de-Cognac and 4 specimens from the Chassiron samples.

Description. Medium-sized gyrogonites, often 500 to 650 µm in height and 300 to 415 µm in width, bottle-shaped with a long apical neck and subprolate to prolate (ISI, 124-180). The basal plate has not been observed. The utricle is mostly formed by the nodular layer, with the nodules located on the spiral cell suture. The nodular layer is sometimes partially covered by 5 to 7 individual bract cells, digitated in shape, which are not organised in any particular symmetry and vary in height (from one-third to two-thirds of the gyrogonite).

Remarks. Although the basal plate could not be observed in this species, it has been found to be single celled in several other related species from Clavatoraceae (Clavatoroideae: *Hemiclavator adnatus* (MARTÍN-CLOSAS and GRAMBAST-FESSARD, 1986) SCHUDACK, 1993 and *Clavator harrisii* PECK, 1941; Atopocharoideae: *Atopochara trivolvis* var. *trivolvis* PECK, 1938), as shown by Martín-Closas (1988, 2000). Some of the specimens from Chassiron and Angeac-Charente are well preserved and enabled us to see the features of the outer layer, which is really rare.

Distribution. This species has been identified by Colin et al. (2004) and El Albani et al. (2004) in Cherves-de-Cognac (dating back to the Berriasian) and by Néraudeau et al. (2012) in Angeac-Charente. Outside France, it has been observed in the Lower Saxony basin of Germany from the lower Tithonian (Schudack, 1993) and in the palustrine areas of the Iberian Chain in Spain from the Tithonian to Barremian (Martín-Closas, 2000). In the United States, *N. bradleyi* has been reported to occur in the Cedar Mountain Formation (Utah) from the upper Berriasian and lower Valanginian and more rarely in the Lakota Formation, South Dakota (Martín-Closas et al., 2013), where it had already been identified by Peck (1957).

---------- Please insert Fig. 5 near here ----------

Genus *Clavator* REID and GROVES, 1916 emend. MARTÍN-CLOSAS, 1989 ex SCHUDACK, 1993

Clavator grovesii var. *grovesii* (HARRIS, 1939) nov. comb. MARTIN-CLOSAS, 1996

Fig. 6 (A-G).

1939 *Clavator grovesii* sp. nov. – Harris, p. 46-53, pl. 10 fig. 1-12, pl.11-12, pl. 17, fig. 8-13.

1962 *Flabellochara grovesii* (HARRIS) nov. comb. – Grambast, p. 69.

1993 *Clavator grovesii grovesii* (HARRIS, 1939) n. comb. – Schudack, p. 76-77, fig. 35; pl. 9, figs 8-14.

1996 *Clavator grovesii* var. *grovesii* (HARRIS, 1939) n. comb. – Martín-Closas, p. 278.

Material. Seven specimens from Angeac-Charente (An3-4-SB) and 33 specimens from Cherves-de-Cognac.

Description. Medium-sized utricles, 475 to 650 µm in height and 364 to 533 µm in width, with a bilateral symmetry. They are composed of a structured layer of 3 primary bract cells, one large cell that is opposite the phylloid and two shorter cells in a lateral position and bearing a fan of secondary bract cells. The three primary bract cells are connected to a basal pore. The tips of the lateral bract cells appear as a pore in section. Some utricles (variety 1) show 2 basal cells surrounding the lateral bract cells, while the fan is composed of 5 to 6 flattened cells in the direction of the apex. These cells are fused and ellipsoidal. Other utricles (variety 2) show a small basal cell covering the lateral bract cells, while the fans are similar in structure to the former variety. It is important to note that intermediate forms between these two varieties occur in the same population.

Remarks. The two varieties are part of the *Flabellochara-Clypeator* lineage described by Grambast (1974), later considered by Martín-Closas (1989, 1996,) and Schudack (1993) to represent anagenetic forms of the same evolutionary species *Clavator grovesii*. The first form described above corresponds to *Clavator grovesii* var. *grovesii* (HARRIS, 1939) comb. nov. MARTÍN-CLOSAS, 1996, while the second represents transitional forms between *C. grovesii* var. *grovesii* and *C. grovesii* var. *discordis* (SHAÏKIN, 1976) comb. nov. MARTÍN-CLOSAS, 1996.

It is important to point out the small size of the specimens studied. The utricle size in this species normally ranges from 400 to 850 µm in height and from 350 to 750 µm in width. The small size may be due to ecological variations, as explained previously for *Mesochara harrisii* gyrogonites by Vicente et al. (2016) and for extant *Chara* by Sanjuan et al.(2017).

Distribution. *Clavator grovesii* var. *grovesii* has been observed to occur in England from the Tithonian and Berriasian in the Dorset and Weald Basins (Harris, 1939; Feist et al., 1995), in France (Colin et al., 2004; El Albani et al., 2004; Néraudeau et al., 2012), in the Swiss Jura (Häfeli, 1966; Mojon and Strasser, 1987; Détraz and Mojon, 1989), Spain (Schudack, 1993; Martín-Closas, 2000) and north-west Germany (Schudack, 1993). It has also been found in China from the Berriasian or Valanginian (Wang et al., 1976; and Hao et al., 1983). Martín-Closas et al. (2013) reported on its occurrence in the Berriasian from North America, suggesting that this species was the first clavatoracean to reach a cosmopolitan distribution in a subtropical latitudinal area of the Northern Hemisphere (Martín-Closas, 2015). *C. grovesii* var. *discordis* is also present in Berriasian strata from Spain, often together with *C. grovesii* var. *grovesii* (Schudack, 1993; Martín-Closas, 2000).

---------- Please insert Fig. 6 near here ----------

Discussion

Taphonomy

The charophyte remains from Chassiron generally displayed good preservation. The most significant observation was the occurrence of a number of utricles attached to the same phylloid in *N. bradleyi* (Fig. 5A). In some cases, these specimens still showed parts of the external layer of their utricles. Some calcified parts of clavatoracean axes were also found. These features are indicative of autochthony (El Albani et al., 2004). One part of the charophyte assemblage from Chassiron, especially *M. harrisii* and *L. latitruncata*, is laterally compressed due to diagenetic deformation.

In Cherves-de-Cognac, the clavatoracean utricles (*N. bradleyi* and *C. grovesii*) were well preserved and, like the utricles from Chassiron, some displayed an anatomical connection to the phylloid. Calcified parts of clavatoracean thalli were also found. As above, these features suggest autochthony of the assemblage (El Albani et al., 2004). Characean and porocharacean gyrogonites from Cherves-de-Cognac were also well preserved, but not associated with vegetative remains, indicating that these gyrogonites were gently transported from an adjacent growing area, forming a parautochthonous assemblage, or that the thalli of these species did not calcify (El Albani et al., 2004).

The population from Angeac-Charente (An3-4-SB) presented significant similarities to that from Cherves-de-Cognac. *N. bradleyi* and *C. grovesii* utricles were well preserved and associated with the calcified parts of clavatoracean thalli. This observation is sufficient to justify the autochthony of the assemblage, in contrast to that proposed by Néraudeau et al. (2012). Characean and porocharacean gyrogonites were generally well preserved, but their

thalli were absent. The An2-SA samples contained relatively corroded porocharacean gyrogonites compared to An3-4-SB specimens. This corrosion, in the form of spiral cells with alveolar surfaces, has been attributed to epidiagenetic processes (Martín-Closas and Grambast-Fessard, 1986).

Palaeoecology

During the Late Jurassic and Early Cretaceous, the occurrence of the genus *Latochara* was restricted to a latitudinal area of the boreal realm (Schudack, 1996). Thus, its occurrence in Chassiron represents its southernmost biogeographical distribution of that time (Martín-Closas et al., 2008). Generally, the abundance of porocharaceans in the Upper Jurassic and Cretaceous, particularly that of *Latochara*, indicates brackish water environments such as coastal marshes (Martín-Closas et al., 2008), which is consistent with the data obtained from member 2 of the Chassiron stratigraphic section (Schnyder et al., 2012).

Similarly, assemblages dominated by porocharaceans in the Cretaceous are thought to be indicative of brackish water environments (Martín-Closas and Grambast-Fessard, 1986; Mojon, 1989a, b; El Albani et al., 2004; Villalba-Breva and Martín-Closas, 2013), while clavatoraceans are more abundant and diverse in freshwater facies (Schudack, 1993; El Albani, 2004, Climent et al., 2009). The occurrence of both taxa in Cherves-de-Cognac and Angeac-Charente, with porocharaceans dominating in the former and clavatoraceans in the latter (especially in An3-4 samples), suggests deposition in an increasingly freshwater environment occurring eastwards within the northern Aquitaine Basin. The environment of Angeac-Charente has been postulated to have been a swamp that had occasionally been connected to the sea (Néraudeau et al., 2012); however, the marine influence would have been greater higher up in the stratigraphic section. Villalba-Breva et al. (2012) and Villalba and Martín-Closas (2013) described the occurrence of abundant porocharaceans in the

Maastrichtian of the Vallcebre and Tremp Basins (Catalonia), corresponding to the progradation of a deltaic-estuarine system. The lignite and limestone facies of this unit display some similarities with An3-4 in Angeac-Charente, such as the coal macerals of the vitrinite group, the abundance of conifer remains and the absence of root traces at the base of the coal beds, indicating a parautochthonous accumulation of organic matter rather than a deposit in peat mires or swamps (Villalba-Breva et al. 2012; Villalba-Breva and Martín-Closas, 2013).

Another reason for the occasional marine influence and variation in salinity is the relatively important polymorphism of porocharaceans in the assemblages from Cherves-de-Cognac and Angeac-Charente. Mojon (1989a) highlighted the occurrence of a number of morphotypes within the Berriasian *Porochara* populations found in the Swiss and French Jura mountains. By comparing to the extant species *Lamprothamnium papulosum* (WALLROTH) GROVES, Mojon (1989a) postulated that this variation was linked to seasonal fluctuations of ecological parameters such as salinity and hydrodynamism, with the morphotypes corresponding to different ecophenotypes rather than to different species.

Biostratigraphy

The three study sites enabled us to revisit the biostratigraphic implications of the latest Jurassic and Early Cretaceous charophyte assemblages in the northern part of the Aquitaine Basin (Fig. 7). The assemblages from Chassiron were represented by *L. latitruncata*, *M. harrisii* and rarely *N. bradleyi*. The occurrence of *L. latitruncata* suggests that the assemblages dated from the Kimmeridgian to the late Tithonian, which is consistent with the age obtained from the associated fauna. By contrast, the assemblages from Cherves-de-Cognac and Angeac-Charente consisted of *N. bradleyi* (which was the predominant species in An3-4-SB in Angeac-Charente), *M. harrisii* and *P. westerbeckensis* together with *Clavator grovesii* var. *grovesii* and *C. grovesii* var. *discordis*. The two latter clavatoracean taxa are of

biostratigraphic interest as they form part of the *Flabellochara-Clypeator* anagenetic lineage described by Grambast (1970, 1974). These taxa are typically found in the uppermost Tithonian and Berriasian strata in Europe (Schudack, 1993; Martín-Closas, 2000), within the Maillardi, Incrassatus and Nurrensis charophyte biozones (Riveline et al., 1996). Our findings agree with those of previous studies on samples from Cherves-de-Cognac (Colin et al., 2004; El Albani et al., 2004), but disagree with the date of the Angeac-Charente bone bed proposed by Néraudeau et al. (2012). Using a complex fossil assemblage of terrestrial, freshwater, brackish and marine organisms, Néraudeau et al. (2012) proposed a Hauterivian-Barremian age mainly based on a few specimens of the dinoflagellate *Odontochitina imparilis* (DUXBURY) JAIN and KHOWAJA-ATEEQUZZAMAN and the benthic foraminifer *Trocholina* cf. *odukpaniensis* DESSAUVAGIE. The other fossils were typical of older ages, such as the bryozoans and charophytes, and were considered partly or totally reworked. This dating of the bone bed is refuted here based on the charophyte taphonomic data mentioned above (Fig. 8).

---------- Please insert Fig. 7 near here ----------

Implications for basin evolution

The absence of any Hauterivian-Barremian stratigraphic record in northern Aquitaine compared to the much more complete record of that period in southern Aquitaine, which forms part of the Arzacq-Mauléon Basin, is relevant for analysing the basin's evolution. The creation of sedimentary space in the Iberian and south-eastern French basins has been linked to the "hyperextension" of the continental crust associated with the opening of the Bay of Biscay (Tugend et al., 2015). This process started during the Barremian and lasted until the Early Albian, leading to an increase in the tectonic subsidence of neighbouring basins such as the Maestrat and Cameros Basins in the Iberian Chain, the Arzacq-Mauléon and Organyà

548 Basin in the Pyrenees, the Basque-Cantabrian Range, and the Bay of Biscay-Parentis Basin,
549 including part of Aquitaine (Tugend et al., 2015). However, while this "hyperextension" had a
550 significant effect within the Iberian Plate, reaching up to a thousand kilometres far away from
551 the main rift axis, its influence was comparatively very limited within the European Plate. Our
552 biostratigraphic results confirm that this hyperextension was limited exclusively to the
553 southern part of Aquitaine, its influence disappearing completely around a hundred kilometres
554 north of the main rift zone and thus, failing to reach the Charentes Basin (Fig. 8).
555
556 ---------- Please insert Fig. 8 near here ----------
557
558 **Conclusions**
559
560 The charophyte remains studied in Chassiron, Cherves-de-Cognac and Angeac-
561 Charente provide new data that improve the current knowledge on the Late Jurassic and Early
562 Cretaceous charophyte flora of the northern part of the Aquitaine Basin, yielding further
563 information on the basin's evolution.
564 The Upper Jurassic and Lower Cretaceous northern Aquitaine Basin are uniquely
565 composed of three intervals with freshwater to brackish facies, the Tithonian (in part) from
566 Chassiron, the Berriasian from Cherves-de-Cognac and Angeac-Charente, and the Uppermost
567 Albian from Charente-Maritime. Only the first three areas were studied here. The Tithonian
568 deposits of Chassiron contain a purely brackish water assemblage of charophytes, as reported
569 by Martín-Closas et al. (2008). The assemblage is predominantly composed of *Latochara*
570 *latitruncata*, also containing the rare *Mesochara harrisii* and the newly reported
571 *Nodosoclavator bradleyi*

572 The assemblages of Cherves-de-Cognac and Angeac-Charente are quite similar. The
573 presence of *Porochara westerbeckensis*, *Nodosoclavator bradleyi*, *Clavator grovesii* and
574 *Mesochara harrisii* at different levels, together with sedimentological, taphonomical and
575 palaeontological data, indicate that the environment became increasingly composed of
576 freshwater eastwards during the Berriasian from Cherves-de-Cognac to Angeac-Charente.

577 The association of the clavatoraceans *Clavator grovesii* var. *grovesii* and *Clavator*
578 *grovesii* var. *discordis* in both Cherves-de-Cognac and Angeac-Charente suggests a Berriasian
579 age for the latter area, which disagrees with the Hauterivian-Barremian age proposed by
580 Néraudeau et al. (2012). Indeed, no other Hauterivian-Barremian records have been obtained
581 to date in the northern Aquitaine Basin, which is in contrast to the many found in the southern
582 Aquitaine, i.e., the Arzacq-Mauléon Basin.

583 In terms of basin evolution, the absence of any Hauterivian-Barremian record in
584 northern Aquitaine suggests that the sedimentary space produced as a result of the crustal
585 extension during the opening of the Gulf of Biscay was limited to the southern part of
586 Aquitaine (Arzacq-Mauléon Basin), which was palaeogeographically located close to the
587 main extensional structures. In this sense, northern Aquitaine only represents a minor part of
588 the same basin containing only the Upper Jurassic and Berriasian records linked to the
589 opening of the North Atlantic Ocean.

590

591 Acknowledgements

592 This study was funded by the BIOGEOEVENTS project (CGL2105-69805-P) of the
593 Spanish Ministry of Science (MINECO) and the European fund for Regional Development
594 (EFRD, EU). It was also supported by a grant to RAB in Barcelona. We would like to thank
595 Dr Alejandro Gallardo for his technical help during sample preparation. The English text of
596 the manuscript was revised by Dr Tasneem Ahmed and Dr. Christopher Evans at the

University of Barcelona. The original manuscript was significantly improved by the editor, Eduardo Koutsoukos and the contribution of Ingeborg Soulié-Märsche and Josep Sanjuan during the peer-review process.

References

Agardh, C. A., 1824. Systema Algarum. Lundae Literis Berlingianis, Lund, 312 pp.

Allain, R., Vullo, R., Le Loeuff, J., Tournepiche, J.-F., 2014. European ornithomimosaurs (Dinosauria, Theropoda) : an undetected record. Geologica Acta 12, 2, 127-135.

Archiac, A. d', 1837. Mémoire sur la formation crétacée du Sud-Ouest de la France. Société géologique 2, 157-196.

Billon-Bruyat, J-P., 2003. Les écosystèmes margino-littoraux du Jurassique Terminal et du Crétacé Basal d'Europe Occidentale: biodiversité, biogéochimie et l'événement biotique de la limite Jurassique/Crétacé. [The margino-littoral ecosystems from the Latest Jurassic and Earliest Cretaceous of Western Europe: biodiversity, biogeochemistry and the biotic event of the Jurassic/Cretaceous boundary]. PhD thesis Université de Poitiers, Unpublished.

Blakey, R., 2011. Paleogeographic Map. Colorado Plateau Geosystems, Inc.

Bousquet, P., 1967. Contribution à l'étude stratigraphique et micropaléontologique du Jurassique et du Crétacé de l'île d'Oléron (Charente-Maritime). PhD Thesis Faculté des sciences de l'Université de Paris, France, Unpublished.

Brenner, P., 1976. Ostracoden und Charophyten des spanischen Wealden (Systematik, Oekologie, Stratigraphie, Palaeogeographie). Palaeontographica Abt. A. 152, 113-201.

Climent-Domènech, H., Martín-Closas C., Salas, R., 2009. Charophyte-rich microfacies in the Barremian of the Eastern Iberian Chain (Spain). Facies 55, 387-400.

622 Colin, J.-P., El Albani, A., Fürsich, F. T., Martin-Closas, C., Mazin, J.-M., Billon-Bruyat, J.-
623 P., 2004. Le gisement “Purbeckien” de vertébrés de Cherves-de-Cognac, Charente
624 (SW France): nouvelles données biostratigraphiques. Comptes Rendus Palevol 3, 9-16.
625 Colin, J.P., Néraudeau, D., Nel, A., Perrichot, V., 2011. Termite coprolites (Insecta: Isoptera)
626 from the Cretaceous of western France: a palaeoecological insight. Revue de
627 Micropaléontologie 54, 129-139.
628 Colombié, C., Schnyder, J., Carcel, D., 2012. Shallow-water marl-limestone alternations in
629 the Late Jurassic of Western France: Cycles, storm event deposits or both?
630 Sedimentary Geology 271-272, 28-43.
631 Combes, P.-J., Peybernès, B., Leyreloup, A., F., 1998. Altérites et bauxites, témoins des
632 marges européenne et ibérique des Pyrénées occidentales au Jurassique supérieur-
633 Crétacé inférieur, à l’ouest de la vallé d’Ossau (Pyrénées-Atlantiques, France).
634 Comptes Rendus Académie des Sciences Paris 327, 271-278.
635 Cubaynes, R., Faure, P., Hantzpergue, P., Pélissié, T., Rey, J., 1989. Le Jurassique du
636 Quercy : unités lithostratigraphiques, stratigraphie et organisation séquentielle,
637 évolution sédimentaire. Géologie de la France 3, 33-62.
638 Deak, M.H., Combaz, A., 1967. « Microfossiles organiques » du Wealdien et du Cénomanien
639 dans un sondage de Charente-Maritime. Revue de Micropaléontologie 10, 2, 69-96.
640 Détraz, H., Mojon, P.-O., 1989. Evolution paléogéographique de la marge jurassienne de la
641 Téthys du Tithonique-Portlandien au Valanginien : corrélations biostratigraphique et
642 séquentielle des faciès marins à continentaux. Eclogae geolica Helveticae 82, 1, 37-
643 112.
644 Donze, P., 1960. Les formations du Jurassique terminal de la partie NW de l’île d’Oléron,
645 Charente-Maritime. Travaux du Laboratoire de Géologie de la Faculté des Sciences de
646 Lyon 5, 1-30.

El Albani, A., Fürsich, F. T., Colin, J.-P., Meunier, A., Hochuli, P., Martin-Closas, C., Mazin, J.-M., Billon-Bruyat, J.-P., 2004. Palaeoenvironmental reconstruction of the basal Cretaceous vertebrate bearing in the Northern part of the Aquitaine Basin (SW France): sedimentological and geochemical evidence. Facies 50, 195-215.

Feist, M., Cubaynes, R., 1984. Découverte de Charophytes dans le Lias du Sud de la France. Implications paléoécologiques et phylogénétiques. Comptes Rendus Académie des Sciences Paris 299, 593-596.

Feist, M., Lake, R. D., Wood, C. J., 1995. Charophyte biostratigraphy of the Purbeck and Wealden of southern England. Palaeontology 38, 2, 407-442.

Fleuriau de Bellevue, L. B., 1802. Mémoire sur quelques nouveaux genres de mollusques et de vers lithophages, et sur la faculté qu'ont ces animaux de percer les rochers. Journal de physique, de chimie et d'histoire naturelle 54, 345-355.

Grambast, L., 1962. Classification de l'embranchement des charophytes. Naturalia Monspeliensia, Série botanique 14, 63-86.

Grambast, L., 1966. Remarques sur le genre *Nodosoclavator* MASLOV, emend. (Charophytes). Comptes-rendus Sommaire des Séances de la Société Géologique de France, Séance 20 juin 1966, 7, 269-270.

Grambast, L., 1969. La symétrie de l'utricule chez les Clavatoracées et sa signification phylogénétique. Comptes Rendus Académie des Sciences Paris 269, 878-881.

Grambast, L., 1970. Origine et évolution des *Clypeator* (Charophytes). Comptes Rendus Académie des Sciences Paris 271, 1964-1967.

Grambast, L., 1974. Phylogeny of the Charophyta. Taxon 23, 4, 463-481.

Grambast-Fessard, N., Ramalho, M., 1985. Charophyte du Jurassique supérieur du Portugal. Revue de Micropaléontologie 28, 56-66.

671 Häfeli, C., 1966. Die Jura-Kreide Grenzschichten im Bielerseegebiet (Kt Bern). Eclogae
672 Geologicae Helvetiae 59, 2, 565-596.
673 Hantzpergue, P., 1989. Les ammonites kimméridgiennes du haut-fond d'Europe occidentale :
674 biochronologie, systématique, évolution, paléobiogéographie. Cahiers de
675 paléontologie, 428 pp.
676 Hantzpergue, P., Boullier, A., Colin, J.-P., Abit, D., 2004. Nouveaux repères
677 biostratigraphiques à terebratulidés (Brachiopodes) dans le Tithonien inférieur oust-
678 européen. [New biostratigraphical markers to terebratulids (Brachiopods) from the
679 west-european Lower Tithonian]. Géobios 37, 361-375.
680 Hantzpergue, P., Lafaurie, G., 1994. Les Calcaires lithographiques du Tithonien quercynois:
681 Stratigraphie, paléogéographie et contexte biosédimentaire. [Tithonian lithographic
682 limestones of Quercy area (France): stratigraphy, paleogeography, and biosedimentary
683 context]. Geobios Mémoire Spécial 16, 237-243.
684 Hao, Y., Ruan, P., Song, Q., Yang, G., Cheng, S., Wei, Z., 1983. Middle Jurassic-Tertiary
685 deposits and Ostracoda-Charophyta assemblages of Xining and Minhe basins. Journal
686 Wuhan College Geology 23, 1-210.
687 Harris, T. M., 1939. British Purbeck Charophyta. British Museum oh Natural History,
688 London, 119 pp.
689 Kubota, K., 2005. Charophyte gyrogonites from the Lower Cretaceous Kitadani Formation of
690 the Tetori Group in the Takinamigawa area, Katsuyama City, Fukui Prefecture, central
691 Japan. Paleontological Research 9, 2, 203-213.
692 Lanceron, F., 1976. Contribution à l'étude géologique de la Saintonge dans une optique
693 hydrogéologique. Thèse 3ème cycle Université de Bordeaux I, 184 pp. (unpublished)
694 Lindley, J., 1836. A natural system of Botany. Longman, 2e ed., London, 526 pp.

695 Liu, J., Wu, X., 1985. Charophytes from Tugulu Group of the Junggar Basin. Bulletin
696 Institute Geology Chinese Academy Geological Sciences 11, 139-153. [in Chinese]
697 Mädler, K., 1952. Charophyten aus dem Nordwestdeutschen Kimmeridge. Geologisches
698 Jahrbuch 67, 1-46.
699 Mädler, K., 1955. Zur Taxinomie der tertiären Charophyten. Geologisches Jahrbuch 70, 265-
700 328.
701 Malz, H., 1966. *Rectocythere rugosa*, eine neue Ostracoden-Art aus dem französischen
702 Portlandian. Senckenbergiana Lethaea 47, 405-409.
703 Martín-Closas, C. 1988. Découverte de la plaque basale chez les Clavatoraceae (Charophyta).
704 Implications phylogénétiques. Comptes Rendus de l'Académie des Sciences 306, sér.
705 II, 1131-1136
706 Martín-Closas, C., 1989. Els carofits del Cretaci inferior de les conques perifèriques del bloc
707 de l'Ebre. Tesi doctoral Universitat de Barcelona. 581 pp. (unpublished)
708 Martín-Closas, C., 1996. A phylogenetic system of Clavatoraceae (Charophyta). Review of
709 Palaeobotany and Palynology 94, 259-293.
710 Martín-Closas, C., 2000. Els carofits del Juràssic superior I Cretaci inferior de la Peninsula
711 Ibèrica [Charophytes from the Upper Jurassic and Lower Cretaceous of the Iberian
712 Peninsula-in Catalan with Abridged English Version]. Arxius de les Seccions de
713 Ciències, Institut d'Estudis Catalans 125, 1-304.
714 Martín-Closas, C., 2015. Cosmopolitanism in Northern Hemisphere Cretaceous Charophyta
715 (Clavatoroidae). Palaeogeography, Palaeoclimatology, Palaeoecology 438, 9-23.
716 Martín-Closas, C., Clavel, B., Schroeder, R., Charollais, J., Conrad, M.-A., 2009.
717 Charophytes from Barremian-lower Aptian of the Northern Subalpine Chains and Jura
718 Mountains, France: correlation with associated marine assemblages. Cretaceous
719 Research 30, 49-62.

720 Martín-Closas, C., Colin, J.-P., Billon-Bruyat, J.-P., El Albani, A., Mazin, J.-M., Schnyder, J.,
721 2008. Palaeobiogeography of the genus *Latochara* (fossil Charophyta) in the Upper
722 Jurassic of the Southern Europe. New data from Oléron Island (France). Geologica
723 Croatica 61, 2-3, 177-184.
724 Martín-Closas, C., Grambast-Fessard, N., 1986. Les charophytes du Crétacé Inférieur de la
725 région du Maestrat (Chaine Ibérique-Catalanides, Espagne). Paléobiologie
726 continentale 15, 1-66.
727 Martín-Closas, C., Peybernès, B., 1987. Datation de la transgression éocrétacée dans les
728 pyrénées Basco-Béarnaises à l'aide des charophytes. Geobios 20, 5, 695-700.
729 Martín-Closas, C., Schudack, M. E., 1991. Phylogenetic analysis and systematization of post-
730 paleozoic Charophytes. Bulletin Société botanique de France 138, Actualités Botiques,
731 1, 53-71.
732 Martín-Closas, C., Sames, B., Schudack, M. E., 2013. Charophytes from the Upper Berriasian
733 of the Western Interior Basin of the United States. Cretaceous Research 46, 11-23.
734 Maslov, V. P., 1963. Introduction to the study of fossil charophytes. Akad. Nauk. SSSR
735 Trudy 82, 1-104.
736 Migula, W., 1897. Die Characeen Deutschlands, Österreichs und der Schweiz. In: Rabenhorst,
737 X. (Ed.), Kryptogamic Flora, vol. 5, E. Kummer, Leipzig, 765 pp.
738 Mojon, P.-O., 1989a. Polymorphisme écophénotypique et paléoécologie des porocharacées
739 (charophytes) du Crétacé basal (Berriasien) du Jura Franco-Suisse. Revue
740 Paléobiologie 8, 505-524.
741 Mojon, P.-O., 1989b. Charophytes et ostracodes laguno-lacustres du Jurassique de la
742 Bourgogne (Bathonien) et du Jura Septentrional Franco-Suise (Oxfordien) Remarques
743 sur les discontinuités émersives du Kimméridgien du Jura. Revue de Paléobiologie 3,
744 1-18.

745 Mojon, P.-O., Strasser, A. 1987. Microfaciès, sédimentologie et micropaléontologie du
746 Purbeckien de Bienne (Jura suisse occidental). Eclogae geologica Helveticae 80, 1, 37-
747 58.
748 Moreau, J.D., Néraudeau, D., Vullo. R., Abit, D., Mennecart, B., Schnyder, J.. Late Jurassic
749 dinosaur footprints from Chassiron-La Morelière (Oléron Island, western France).
750 Palaeoecology and Palaeoenvironments (submitted).
751 Néraudeau, D., 2011. Chroniques d'un dinosaure annoncé. La Saintonge Littéraire 97, 28-39.
752 Néraudeau, D., Perrichot, V., Dejax, J., Masure, H., Nel, A., Philippe, M., Moreau, P.,
753 Guillocheau, F., Guyot, T., 2002. Un nouveau gisement à ambre insectifère et à
754 végétaux (Albien terminal probable): Archingeay (Charente-Maritime, France). [A
755 new fossil locality with insects in amber and plants (likely Uppermost Albian):
756 Archingeay (Charente-Maritime, France)]. Geobios 35, 233-240.
757 Néraudeau, D., Allain, R., Ballèvre, M., Batten, D.J., Buffetaut, E., Colin, J.-P., Dabard, M.-
758 P., Daviero-Gomez, V., El Albani, A., Gomez, B., Grosheny, D., Le Loeuff, J.,
759 Leprince, A., Martín-Closas, C., Masure, E., Mazin, J.-M., Philippe, M., Pouech, J.,
760 Tong, H., Tournepiche, J.-F., Vullo, R., 2012. The Hauterivian-Barremian lignitic
761 bone bed of Angeac (Charente, south-west France): Stratigraphical, palaeobiological
762 and palaeogeographical implications. Cretaceous Research 37, 1-14.
763 Néraudeau, D., P, V., Batten, D., Boura, A., Girard, V., Jeanneau, L., B., Nohra, Y., Polette,
764 F., Saint Martin, S., Saint Martin, J.P., Thomas, R., 2016. Upper Cretaceous amber
765 from Vendée, north-western France: age dating and geological, chemical and
766 palaeontological characteristics. Cretaceous Research, in press.
767 Peck, R. E., 1937. Morrison Charophyta from Wyoming. Journal of Paleontolgy 11, 2, 83-90.
768 Peck, R. E., 1938. A new family of Charophyta from the Lower Cretaceous of Texas. Journal
769 of Paleontology 12, 2, 173-176.

770 Peck, R. E., 1941. Lower Cretaceous Rocky Mountain nonmarine microfossils. Journal of
771 Paleontology 15, 285-304.
772 Peck, R. E., 1957. North American Mesozoic Charophyta. Geological Survey Professional
773 Paper 294A, 1-44.
774 Peybernès, B. Combes, P.-J., 1994. Stratigraphie séquentielle du Crétacé basal (intervalle
775 Berriasien-Hauterivien) des Pyrénées centrales et orientales franco-espagnoles.
776 Cretaceous Research 15, 535-546.
777 Pia, J., 1927. Charophyta. In: Hirmer, M. (Ed.]), Handbuch der Paläobotanik, 1, Munich-
778 Berlin, pp. 88-93.
779 Platel, J.-P., 1978. Données récentes sur la connaissance géologique de la région du sud des
780 Charentes. Norois 100, 600-608.
781 Platel, J.-P., 1980. Le Bassin Gypsifère des Charentes. Synthèse des données sur le Purbecko-
782 Wealdien. Proposition d'une méthode de prospection par approche géophysique.
783 Préfecture de la région Poitou-Charentes, commission « argiles et terres réfractaires ».
784 Rapport du Bureau de Recherches Géologiques et Minières 80 SGN 313 PROC,
785 Poitou-Charentes.
786 Platel, J.-P., Moreau, P., 1977. Feuille de Saint-Agnant. Carte géologique de la France à
787 1/50000 682, Bureau de Recherches Géologiques et Minières.
788 Poeuch, J., Mazin, J.-M., Cavin, L., Poyato-Ariza, F. J., 2015. A Berriasian actinopterygian
789 fauna from Cherves-de-Cognac, France: Biodiversity and palaeoenvironmental
790 implications. Cretaceous Research 55, 32-43.
791 Rees, J., Cuny, G., Poeuch, J., Mazin, J.-M., 2013. Non-marine selachians from the basal
792 Cretaceous of Charente, SW France. Cretaceous Research 40, 122-131.
793 Reid, C., Groves, J., 1916. Preliminary report on the Purbeck Characeae. Proceedings of the
794 Royal Society B89, 252-256.

Riveline, J., Berger, J.-P., Feist, M., Martín-Closas, C., Schudack, M., Soulié-Märsche, I., 1996. European Mesozoic-Cenozoic charophyte biozonation. Bulletin Société Géologique France 167, 3, 453-468.

Sanjuan, J., Vicente, A., Flor, N., Cambra, J. Martín-Closas, C., 2017. Effects of light and temperature on *Chara vulgaris* gyrogonite productivity and polymorphism. Palaeoenvironmental implications. Phycologia. 56, 204-212.

Schettino, A., Scotese, C., 2002. Global kinematic constraints to the tectonic history of the Mediterranean region and surrounding areas during the Jurassic and Cretaceous. Reconstruction of the evolution of the Alpian-Himalayan Orogen. Journal of the virtual Explorer 7, 147-166.

Schnyder, J., 2003. Le passage Jurassique/Crétacé: événements instantanés, variations climatiques enregistrées dans les faciès purbeckiens français (Boulonnais, Charentes) et anglais (Dorset). PhD thesis University of Lille 1, 389 p. (Unpublished)

Schnyder, J., Deconinck, J.-F., Baudin, F., Colombié, C., Jan du Chêne, R., Gardin, S., Galbrun, B., De Rafélis, M., 2012. Purbeck beds (Late Jurassic) in the Phare de Chassiron section (Ile d'Oléron, NW Aquitaine Basin, France): Refined age-assignment and long-term depositional sequences. Geobios 45, 485-499.

Schudack, M. E., 1986. Zur Nomenklatur der Gattungen *Porochara* MÄDLER 1955 (syn. *Musacchiella* FEIST and GRAMBAST-FESSARD 1984) und *Feistiella* n. gen. (Charophyta). Paläontogische Zeitung 60, 1-2, 21-27.

Schudack, M. E., 1987. Charophytenflora und fazielle Entwicklung der Grenzschichten mariner Jura/Wealden in den Nordwestlichen Iberischen Ketten (mit Vergleichen zu Asturien und Kantabrien). Paleontographica 204, B, 1-180.

818 Schudack, M. E., 1989. Charophytenfloren aus den unterkretazischen Vertebrate-
819 Fundschichten bei Galve und Una (Ostspanien). Berliner Geowissenschaftliche
820 Abhandlungen, Reihe A 106, 409-443.
821 Schudack, M. E., 1990. Bestandsaufnahme und Localzonierung der Charophyten aus Oberjura
822 und Unterkreide des Nordwestdeutschen Beckens. Berliner Geowissenschaftliche
823 Abhandlungen, Reihe A, 124, 209-245.
824 Schudack, M. E., 1993. Die Charophyten im Oberjura und Unterkreide Westeuropas: Mit
825 einer phylogenetischer Analyse der Gesamtgruppe. Geowissenschaftliche
826 Abhandlungen, Reihe E, 8, 1-209.
827 Schudack, M. E., 1996. Ostracode and Charophyte biogeography in the continental Upper
828 Jurassic of Europe and North America influenced by Plate Tectonics and paleoclimate.
829 In: MORALES, M. [ed.]. The continental Jurassic. Museum of Northern Arizona
830 Bulletin 60, 333-341.
831 Shaïkin, I. M., 1967. Fossil Charophyta of Upper Jurassic deposits of Dniepr-Donets
832 depression. In: Fossil Algae of the USSR. IZDAT NAUK SSSR, pp. 43-47. [In
833 Russian].
834 Shaïkin, I. M., 1976. New data on biostratigraphy of the Jurassic and Cretaceous of the Fore-
835 Dobrogean Trough. Geological Zhurnal 36, 2, 77-86. [In Russian]
836 Smith, G. M., 1938. Botany. In: Algae and Fungi. Charophyceae, 1, Mcgraw Hill, New-York,
837 127 pp.
838 Thierry, J., 2000. Early Kimmeridgian. In: Dercourt, M., et al. (Ed), Atlas Peri-Tethys:
839 Palaeogeographical Maps, pp. 85-97.
840 Tugend, J., Manatschal, G., Kusznir, N.J., 2015. Spatial and temporal evolution of
841 hyperextended rift systems: Implication for the nature, kinematics, and timing of the
842 Iberian-European plate boundary. Geology 43, 15-18.

Vicente, A., Exposito, M., Sanjuan, J., Martín-Closas, C., 2016. Smalll sized charophyte gyrogonites in the Maastrichtian of Coll de Nargo, Eastern Pyrenees: An adaptation to temporary floodplain ponds. Cretaceous Research 57, 443-456.

Villalba-Breva, S., Martín-Closas, C., 2013. Upper Cretaceous paleogeography of the Central Southern Pyrenean Basins (Catalonia, Spain) from microfacies analysis and charophyte biostratigraphy. Facies 59, 319-345.

Villalba-Breva S, Martín-Closas C, Marmi J, Gomez B, Fernández-Marrón MT. 2012. Peat-forming plants in the Maastrichtian coals of the Eastern Pyrenees. Geologica Acta. 10, 189-207.

Vullo, R., Abit, D., Ballèvre, M., Billon-Bruyat, J.-P., Bourgeais, R., Buffetaut, E., Daviero-Gomez, V., Garcia, G., Gomez, B., Mazin, J.-M., Morel, S., Néraudeau, D., Pouech, J., Rage, J.-C., Schnyder, J., Tong, H., 2014. Palaeontology of the Purbeck-type (Tithonian, Late Jurassic) bonebeds of Chassiron (Oléron Island, western France). Comptes Rendus Palevol 13, 421-441.

Vullo, R., Buffetaut, E., Néraudeau, D., Le Loeuff, J., Heil, J.F., Dunand, M., 2012. Le "Mégalosaure" (Dinosauria, Sauropoda) de Saint-Agnant (Charente-Maritime, France): description et origine stratigraphique. Annales de Paléontologie 98, 115-129.

Wang, S., 1965. Mesozoic and Tertiary Charophyta from Jinquan basin of Kansu province. Acta Paleontologica Sinica 13, 485-499.

Wang, Z., Huang, R., Wang, S., 1976. Mesozoic and Cenozoic charophytes from Yunnan. In: Mesozoic fossils from Yunnan, 1, 65-93.

Winnock, E., 1973. Exposé succinct de l'évolution paléogéologique de l'Aquitaine. Bulletin Société Géologique France 1, 7, 15, 5-12.

867 **Figure legends**

868 **Figure 1.** Geological map of the Charentes Basin and locations of the studied sites (modified
869 from Géoportail.fr).

870 **Figure 2.** Stratigraphic framework of the Charentes Basin.

871 **Figure 3.** Stratigraphic columns of Chassiron, Cherves-de-Cognac and Angeac-Charente
872 (modified from Schnyder et al. (2012), Rees et al. (2013) and Néraudeau et al. (2012),
873 respectively).

874 **Figure 4.** Charophyte fructifications from the Upper Jurassic and Lower Cretaceous of the
875 northern part of the Aquitaine Basin. A-E, *Porochara westerbeckensis* from the Berriasian of
876 Angeac-Charente. (A) Lateral view of a gyrogonite from sample An2-SA, IGR-PAL-2779.12;
877 (B) lateral view of a gyrogonite (sample An3-4-SB), IGR-PAL-2779.13; (C) a multipartite
878 basal plate (sample An3-4-SB), IGR-PAL-2779.32; (D) basal view of a gyrogonite (sample
879 An3-4-SB), IGR-PAL-2779.15; and (E) apical view of a gyrogonite (sample An3-4-SB),
880 IGR-PAL-2779.16. F-I, *Latochara latitruncata* from the lower Tithonian of Chassiron (level
881 148). (F,G) Lateral views, IGR-PAL-2779.3 and IGR-PAL-2779.5; (H) basal view, IGR-
882 PAL-2779.6, (I) apical view, IGR-PAL-2779.4. J-P, *Mesochara harrisii* from level 148 of the
883 lower Tithonian of Chassiron (J, M, O) and from level An3-4-SB of the Berriasian of Angeac-
884 Charente (H-L,N,P) . (J) Lateral view of a gyrogonite, IGR-PAL-2779.7; (K) lateral view,
885 IGR-PAL-2779.17; (L) lateral view, IGR-PAL-2779.18; (M) basal view, IGR-PAL-2779.10;
886 (N) basal view, IGR-PAL-2779.20; (O) apical view, IGR-PAL-
887 2779.9; and (P) apical view, IGR-PAL-2779.19.

888 **Figure 5.** Charophyte fructifications from the Upper Jurassic and Lower Cretaceous of the
889 northern part of the Aquitaine Basin. A-H, *Nodosoclavator bradleyi* from level 148 of the
890 lower Tithonian Chassiron (A) and from the Berriasian of Angeac, samples An4 (B) and An3-
891 4-SB (C-F, H) and An2-SA (G). (A) Lateral view of two utricles still attached to the phylloid,

with small bract cells of the structured layer visible, IGR-PAL-2779.1; (B) lateral view of a utricle, with the nodular layer visible, IGR-PAL-2780.5; (C) lateral view of a utricle with the structured layer visible, IGR-PAL-2780.13; (D) apical view of a utricle, IGR-PAL-2779.26; (E) basal view of a utricle, IGR-PAL-2779.21; (F) lateral view of a utricle with the structured layer visible, IGR-PAL-2779.25; (G) lateral view of a gyrogonite with remains of the nodular layer attached to it, IGR-PAL-2779.11; and (H) lateral view of a utricle, IGR-PAL-2779.22.

Figure 6. Charophyte fructifications from the Upper Jurassic and Lower Cretaceous of the northern part of the Aquitaine Basin. A-G, *Clavator grovesii* from the Berriasian of Angeac-Charente (sample An3-4-SB). (A) Lateral view of a utricle, IGR-PAL-2780.6, (B) lateral view of a utricle, IGR-PAL-2779.27, (C) lateral view of a poorly calcified gyrogonite, IGR-PAL-2780.7, (D) lateral view of a utricle, IGR-PAL-2779.28, (E), lateral view of a utricle, IGR-PAL-2780.11, (F) lateral view of a utricle, IGR-PAL-2780.12, and (G) lateral view of a utricle, IGR-PAL-2779.30.

Figure 7. Stratigraphic distribution of species in Chassiron, Cherves-de-Cognac and Angeac-Charente.

Figure 8. Palaeogeographic and structural map of the different rift systems between the European and the Iberian plates during the Late Jurassic-Early Cretaceous, showing the position of the Charentes basin (modified from Tugend et al., 2015). Study area in the Charentes marked with a rectangle.


Table 1. Distribution of charophyte species and their abundance in the samples studied.


ACCEPTED MANUSCRIPT


Lithostratigraphy / Locality / Samples / Species	Chassiron	Cherves-de-Cognac							Angeac-Charente				
	L148	C32	C33	C34	C35	C36	C38	C41	An1	An2	An3	An3-4	An4
Porochara westerbeckensis		●	●		⬤	●			•			⬤	
Latochara latitruncata	⬤												
Mesochara harrisii	●	•			•	•				•		●	
Nodosoclavator bradleyi	•	•								•		⬤	
Clavator grovesii		●		•	•	•						•	

• 0-10 ● 11-50 ⬤ 51-100 ⬤ >100

ACCEPTED MANUSCRIPT



A
B
C
D
E
F
G
H
: 500 µm


A
B
C
D
E
F
G
: 500 µm




ACCEPTED MANUSCRIPT


ACCEPTED MANUSCRIPT




ACCEPTED

ACCEPTED MANUSCRIPT





A
B
C
D
E
F
G
H
I
J
K
L
M
N
O
P
: 500 µm



ACCEPTED MANUSCRIPT

ACCEPTED MANUSCRIPT



ACCEPTED MANUSCRIPT

Highlights

Charophytes from Charente are exclusively Tithonian-Berriasian in age

Former claims of Hauterivian-Barremian records are not confirmed in the Charente basin

The opening of the Biscay basin left no sedimentary record in the Charente basin.

Clavatoraceans dominated Berriasian assemblages over characeans and porocharaceans

Charophyte assemblages show increasing freshwater influence eastwards in the basin

ACCEPTED MANUSCRIPT