

Explorer of Enceladus and Titan (E2 T): Investigating ocean worlds' evolution and habitability in the solar system

Giuseppe Mitri, Frank Postberg, Jason M. Soderblom, Peter Wurz, Paolo Tortora, Bernd Abel, Jason W. Barnes, Marco Berga, Nathalie Carrasco, Athena Coustenis, et al.

▶ **To cite this version:**

Giuseppe Mitri, Frank Postberg, Jason M. Soderblom, Peter Wurz, Paolo Tortora, et al.. Explorer of Enceladus and Titan (E2 T): Investigating ocean worlds' evolution and habitability in the solar system. Planetary and Space Science, 2018, 155, pp.73-90. 10.1016/j.pss.2017.11.001 . insu-01636074


HAL Id: insu-01636074
https://insu.hal.science/insu-01636074

Submitted on 27 Feb 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript



Explorer of Enceladus and Titan (E^2T): Investigating ocean worlds' evolution and habitability in the solar system

Giuseppe Mitri, Frank Postberg, Jason M. Soderblom, Peter Wurz, Paolo Tortora, Bernd Abel, Jason W. Barnes, Marco Berga, Nathalie Carrasco, Athena Coustenis, Jean Pierre Paul de Vera, Andrea D'Ottavio, Francesca Ferri, Alexander G. Hayes, Paul O. Hayne, Jon K. Hillier, Sascha Kempf, Jean-Pierre Lebreton, Ralph D. Lorenz, Andrea Martelli, Roberto Orosei, Anastassios E. Petropoulos, Kim Reh, Juergen Schmidt, Christophe Sotin, Ralf Srama, Gabriel Tobie, Audrey Vorburger, Véronique Vuitton, Andre Wong, Marco Zannoni

PII: S0032-0633(17)30144-7

DOI: 10.1016/j.pss.2017.11.001

Reference: PSS 4418

To appear in: *Planetary and Space Science*

Received Date: 29 April 2017

Revised Date: 11 September 2017

Accepted Date: 1 November 2017

Please cite this article as: Mitri, G., Postberg, F., Soderblom, J.M., Wurz, P., Tortora, P., Abel, B., Barnes, J.W., Berga, M., Carrasco, N., Coustenis, A., Paul de Vera, J.P., D'Ottavio, A., Ferri, F., Hayes, A.G., Hayne, P.O., Hillier, J.K., Kempf, S., Lebreton, J.-P., Lorenz, R.D., Martelli, A., Orosei, R., Petropoulos, A.E., Reh, K., Schmidt, J., Sotin, C., Srama, R., Tobie, G., Vorburger, A., Vuitton, Vé., Wong, A., Zannoni, M., Explorer of Enceladus and Titan (E^2T): Investigating ocean worlds' evolution and habitability in the solar system, *Planetary and Space Science* (2017), doi: 10.1016/j.pss.2017.11.001.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Explorer of Enceladus and Titan (E^2T): Investigating Ocean Worlds' Evolution and Habitability in the Solar System

Giuseppe Mitri [1], Frank Postberg [2], Jason M. Soderblom [3], Peter Wurz [4], Paolo Tortora [5], Bernd Abel [6], Jason W. Barnes [7], Marco Berga [8], Nathalie Carrasco [9], Athena Coustenis [10], Jean Pierre Paul de Vera [11], Andrea D'Ottavio [8], Francesca Ferri [12], Alexander G. Hayes [13], Paul O. Hayne [14], Jon K. Hillier [15], Sascha Kempf [16], Jean-Pierre Lebreton [17], Ralph D. Lorenz [18], Andrea Martelli [8], Roberto Orosei [19], Anastassios E. Petropoulos [14], Kim Reh [14], Juergen Schmidt [20], Christophe Sotin [14], Ralf Srama [21], Gabriel Tobie [1], Audrey Vorburger [4], Véronique Vuitton [22], Andre Wong [14], Marco Zannoni [5]

[1] LPG, Université de Nantes, France
[2] Klaus-Tschira-Laboratory for Cosmochemistry, University of Heidelberg, Germany
[3] Massachusetts Institute of Technology, USA
[4] University of Bern, Switzerland
[5] University of Bologna, Italy
[6] University of Leipzig Germany
[7] University of Idaho, USA
[8] Thales Alenia Space, Italy
[9] LATMOS, France
[10] LESIA, Observ. Paris-Meudon, CNRS, Univ. P. et M. Curie, Univ. Paris-Diderot, France
[11] DLR, Germany
[12] University of Padova -CISAS, Italy
[13] Cornell University, USA
[14] Jet Propulsion Laboratory, California Institute of Technology, USA
[15] University of Kent, UK
[16] University of Colorado, USA
[17] LPC2E, France
[18] JHU Applied Physics Laboratory, USA
[19] INAF, Italy
[20] University of Oulu, Finland
[21] University of Stuttgart, Germany
[22] Univ. Grenoble Alpes, CNRS, IPAG, France

11 September 2017

Manuscript submitted to Planetary Space Science

ACCEPTED MANUSCRIPT

Corresponding Author:

Giuseppe Mitri

Laboratoire de Planetologie et de Geodynamique

Université de Nantes

2 rue de la Houssinière

44322 Nantes, France

Giuseppe.Mitri@univ-nantes.fr

ACCEPTED MANUSCRIPT

ACCEPTED MANUSCRIPT

Abstract

Titan, with its organically rich and dynamic atmosphere and geology, and Enceladus, with its active plume, both harbouring global subsurface oceans, are prime environments in which to investigate the habitability of ocean worlds and the conditions for the emergence of life. We present a space mission concept, the Explorer of Enceladus and Titan (E^2T), which is dedicated to investigating the evolution and habitability of these Saturnian satellites. E^2T is proposed as a medium-class mission led by ESA in collaboration with NASA in response to ESA's M5 Cosmic Vision Call. E^2T proposes a focused payload that would provide in-situ composition investigations and high-resolution imaging during multiple flybys of Enceladus and Titan using a solar-electric powered spacecraft in orbit around Saturn. The E^2T mission would provide high-resolution mass spectrometry of the plume currently emanating from Enceladus' south polar terrain and of Titan's changing upper atmosphere. In addition, high-resolution infrared (IR) imaging would detail Titan's geomorphology at 50–100 m resolution and the temperature of the fractures on Enceladus' south polar terrain at meter resolution. These combined measurements of both Titan and Enceladus would enable the E^2T mission scenario to achieve two major scientific goals: 1) Study the origin and evolution of volatile-rich ocean worlds; and 2) Explore the habitability and potential for life in ocean worlds. E^2T's two high-resolution time-of-flight mass spectrometers would enable resolution of the ambiguities in chemical analysis left by the NASA/ESA/ASI Cassini-Huygens mission regarding the identification of low-mass organic species, detect high-mass organic species for the first time, further constrain trace species such as the noble gases, and clarify the evolution of solid and volatile species. The high-resolution IR camera would reveal the geology of Titan's surface and the energy dissipated by Enceladus' fractured south polar terrain and plume in detail unattainable by the Cassini mission.

Keywords: *Enceladus; Titan; Origin of volatiles; Habitability*

1. Introduction

The NASA/ESA/ASI Cassini-Huygens mission has revealed Titan and Enceladus to be two unique worlds in the Solar System during its thirteen years of observations in the Saturnian system (July 2004 - September 2017). Titan, with its organically rich and dynamic atmosphere and geology, and Enceladus, with its active plume system composed of multiple jets (Waite et al., 2006; Spahn et al., 2006; Porco et al., 2006), both harbouring global subsurface oceans (Iess et al., 2010, 2012, 2014; see also discussion in Sotin et al., 2010), are ideal environments in which to investigate the conditions for the emergence of life and the habitability of ocean worlds as well as the origin and evolution of complex planetary systems. The prime criteria of habitability include energy sources, liquid water habitats, nutrients and a liquid transport cycle to move nutrients and waste (McKay et al., 2008, 2016; Lammer et al., 2009). The best-known candidates in the Solar System for habitability at present meeting these criteria are the ocean worlds in the outer Solar System, which include: Enceladus, Titan, Europa, and Ganymede (Lunine, 2016; Nimmo and Pappalardo, 2016). While the Jovian moons will be thoroughly investigated by the ESA Jupiter Icy moon Explorer (JUICE), Enceladus and Titan, which provide environments that can be easily sampled from orbit in a single mission, are currently not targeted by any future exploration. The joint exploration of these two fascinating objects will allow us to better understand the origin of their organic-rich environments and will give access to planetary processes that have long been thought unique to the Earth.

Titan is an intriguing world that is similar to the Earth in many ways, with its dense nitrogen-methane atmosphere and familiar geological features, including dunes, mountains, seas, lakes and rivers (e.g., Stofan et al., 2007; Lorenz et al., 2006, 2009; Lopes et al., 2007a; Mitri et al., 2010). Titan undergoes seasonal changes similar to Earth, driven by its orbital inclination of 27° and Saturn's approximately 30 year orbit. Exploring Titan then offers the

possibility to study physical processes analogous to those shaping the Earth's landscape, where methane takes on water's role of erosion and formation of a distinct geomorphological surface structure.

Enceladus is an enigma; it is a tiny moon (252 km radius) that harbours a subsurface liquid-water ocean (Iess et al., 2014; McKinnon et al., 2015; Thomas et al., 2016; Čadek et al., 2016), which jets material into space. The eruption activity of Enceladus offers a unique possibility to sample fresh material ejected from subsurface liquid water and understand how exchanges with the interior controls surface activity, as well as to constrain the geochemistry and astrobiological potential of internal oceans on ocean worlds (e.g., Porco et al., 2006). Since the 1997 launch of the Cassini-Huygens mission, there has been great technological advancement in instrumentation that would enable answering key questions that still remain about the Saturnian ocean worlds.

The scientific appeal of Titan and Enceladus has stimulated many previous mission studies (e.g. see reviews by Lorenz, 2000; 2009), which have articulated detailed scientific objectives for post-Cassini scientific exploration (e.g. Mitri et al., 2014a; Tobie et al., 2014). At Titan, in particular, the diversity of scientific disciplines (Dougherty et al., 2009) has prompted the study of a variety of observing platforms from orbiters ("Titan Explorer", Leary et al., 2007; Mitri et al., 2014a), landers for the seas ("Titan-Saturn System Mission – TSSM", Strange et al., 2009; "Titan Mare Explorer - TiME", Stofan et al., 2013, Mitri et al., 2014a), landers for land (Titan Explorer), fixed-wing aircraft ("AVIATR", Barnes et al., 2012), to balloons (Titan Explorer, TSSM and others). Additionally, Enceladus' plume has attracted designs of spacecraft to sample it: "Titan and Enceladus Mission TANDEM" (Coustenis et al., 2009), "Journey to Enceladus and Titan – JET" and "Enceladus Life Finder – ELF" (Reh et al., 2016).

120 We present a space mission concept, the Explorer of Enceladus and Titan (E^2T), which is
121 dedicated to investigating the evolution and habitability of these Saturnian satellites and is
122 proposed as a medium-class mission led by ESA in collaboration with NASA in response to
123 ESA's M5 Cosmic Vision Call. In Section 2 we present the science case for the future
124 exploration of Enceladus and Titan as proposed by the E^2T mission, and Section 3 the science
125 goals for the E^2T mission. In Sections 4 and 5 we discuss the proposed payload and mission
126 and spacecraft configuration necessary to achieve E^2T mission goals.
127

128 **2. Science Case for the Exploration of Enceladus and Titan**

129 Titan, Saturn's largest satellite, is unique in the Solar System with its dense, extensive
130 atmosphere composed primarily of nitrogen (97%) and methane (1.4%) (e.g., Bézard, 2014),
131 and a long list of organic compounds resulting from multifaceted photochemistry that occurs
132 in the upper atmosphere down to the surface (e.g., Israël et al., 2005; Waite et al., 2007;
133 Gudipati et al., 2013; Bézard, 2014). As methane is close to its triple point on Titan, it gives
134 rise to a methane cycle analogous to the terrestrial hydrological cycle, characterized by cloud
135 activity, precipitation, river networks, lakes and seas covering a large fraction of the northern
136 terrain (Figure 1) (e.g., Tomasko et al., 2005; Stofan et al., 2007; Mitri et al., 2007; Lopes et
137 al., 2007a; Hayes et al., 2008).

138 FIGURE 1

139 With an environment that changes on a 29.5 year cycle, it is crucial to study Titan during
140 an entire orbital period. Cassini has investigated Titan over only two seasons: from Northern
141 winter solstice to summer solstice. While ground-based observations, have observed Titan in
142 other seasons, these data are not sufficient to address many of the outstanding questions.
143 Current measurements with Cassini/CIRS show that the chemical content of Titan's
144 atmosphere has significant seasonal and latitudinal variability (Coustenis et al., 2013; 2016);

145 future extended exploration of Titan is necessary to get a full picture of the variations within
146 this complex environment.

147 Titan is the only known planetary body, other than the Earth, with long-standing liquid on
148 its surface, albeit hydrocarbons instead of water; these lakes and seas are likely fed by a
149 combination of precipitation, surface runoff and subsurface alkanofers (hydrocarbon
150 equivalent of aquifers) in the icy crust (Hayes et al., 2008). The presence of radiogenic noble
151 gases in the atmosphere indicates some communication between the surface and the
152 subsurface and is suggestive of water-rock interactions and methane outgassing processes
153 (Tobie et al., 2012), possibly associated with cryovolcanic activity or other exchange
154 processes (Lopes et al., 2007b, 2016; Solomonidou et al., 2014, 2016). The detection of a
155 salty ocean at an estimated 50-80 km depth (Iess et al., 2012; Beghin et al., 2012; Mitri et al.,
156 2014b) and the possible communication between this ocean and the organic-rich surface hint
157 at exciting astrobiological possibilities. While Cassini has provided tantalizing views of the
158 surface with its lakes and seas, dunes, equatorial mountains, impact craters and possible
159 cryovolcanos, the low spatial, spectral, and mass resolution of the Cassini scientific
160 instruments make it difficult to identify morphological features, to quantify geological
161 processes and relationships between different geological units and to monitor changes due to
162 geologic or atmospheric activity. Constraining the level of geological activity on Titan is
163 crucial to understanding its evolution and determining if this ocean world could support
164 abiotic/prebiotic activity.

165 Both Titan and Enceladus possess several, if not all, of the key ingredients necessary for
166 life: an energy source, liquid habitats, nutrients (organic compounds) and a liquid transport
167 cycle to move nutrients and waste (McKay et al., 2008, 2016). While sunlight is a minimal
168 source of energy for solid bodies in the outer Solar System, interior heat sources derived from
169 a rocky core or tidal forces produced by neighbouring satellites and planet can be quite

significant. Most recently, the Cassini INMS has identified molecular hydrogen at the level of 0.4–1.4% in Enceladus' plume (Waite et al., 2017) providing further evidence of water-rock interactions. This suggests that methane formation from CO_2 in Enceladus' subsurface ocean could occur in a similar fashion as it occurs on Earth, where extremophile microbes in hydrothermal sea vents produce methane as a metabolic by-product (McKay et al., 2008). Another compelling discovery is the complex large nitrogen-bearing organic molecules in Titan's upper atmosphere by Cassini (Waite et al., 2007; Coates et al., 2007). The low resolution of the in-situ mass spectrometers on Cassini, however, precludes the determination of the chemical composition of this complex organic matter. In situ exploration with more advanced instruments is required to investigate the prebiotic potential of Titan.

The discovery in 2005 of a plume emanating from multiple jets in Enceladus' south polar terrain is one of the major highlights of the Cassini–Huygens mission (Figure 2) (Dougherty et al., 2006; Porco et al., 2006; Spahn et al., 2006). Despite its small size (10 times smaller than Titan), Enceladus is the most geologically active moon of the Saturnian system. Although geyser-like plumes have been observed on Triton (Soderblom et al., 1990) and more recently transient water vapour activity around Europa has been reported (Roth et al., 2014; Sparks et al., 2016, 2017), Enceladus is the only satellite for which this activity is known to be endogenic in nature. The jets, of which approximately one hundred have been identified (Porco et al., 2014) form a huge plume of vapour and ice grains above Enceladus' south polar terrain and are associated with elevated heat flow along tectonic ridges, called 'tiger stripes'. Enceladus' endogenic activity and gravity measurements indicate that it is a differentiated body, providing clues to its formation and evolution (Iess et al., 2014). Sampling of the plume by Cassini's instruments revealed the presence of water vapour, ice grains rich in sodium and potassium salts (Postberg et al., 2011). Organic materials were observed, both in the gas (Waite et al., 2009) and in the ice grains (Postberg et al., 2008, 2015), and molecular

hydrogen (Waite et al., 2017). The jet sources are connected to a subsurface salt-water reservoir that is likely alkaline in nature and undergoing hydrothermal water-rock interactions (Porco et al., 2006, 2014; Waite et al., 2006, 2009, 2017; Postberg et al., 2009, 2011; Hsu et al., 2011, 2014; Glein et al., 2015). The putative exothermic water-rock interactions on Enceladus could be further constrained by quantifying He and constraining the amount of H_2 in the plume. Gravity, topography and libration measurements demonstrate the presence of a global subsurface ocean (Iess et al., 2014; McKinnon et al., 2015; Thomas et al., 2016; Čadek et al., 2016; Beuthe et al., 2016). The co-existence of organic compounds, salts, liquid water and energy sources on this small moon appear to provide all of the necessary ingredients for the emergence of life by chemoautotrophic pathways (McKay et al., 2008) – a generally held model for the origin of life on Earth in deep sea vents.

FIGURE 2

Titan and Enceladus offer an opportunity to study analogous prebiotic processes that may have led to the emergence of life on Earth. Retracing the processes that allowed the emergence of life on Earth around 4 Ga is a difficult challenge since most traces of the environmental conditions at that time have been erased. It is, therefore, crucial for astrobiologists to find extraterrestrial planetary bodies with similarities to our planet, providing a way to study some of the processes that occurred on the primitive Earth, when prebiotic chemistry was active. The eruption activity of Enceladus offers a possibility to sample fresh material emerging from subsurface liquid water and to understand how exchange processes with the interior control surface activity. It provides us with an opportunity to study phenomena in-situ that has been important in the past on Earth and throughout the outer Solar System.

ACCEPTED MANUSCRIPT

3. Scientific Objectives and Investigations

The proposed E^2T mission has two major goals and several science objectives that would be pursued through Enceladus and Titan investigations. The first scientific goal of the E^2T mission on Enceladus would focus on the origin and evolution of volatile compounds in the plume vapour and icy grains. On Titan the first scientific goal includes two objectives; the first would focus on determining the history and extent of volatile exchange on Titan and the second objective would aim to understand how Titan's surface processes evolved. The second scientific goal on Enceladus would examine the nature of hydrothermal activity and search for evidence of abiotic/prebiotic processes. On Titan, the second scientific goal would aim to discern to what level of complexity abiotic/prebiotic chemistry has evolved. The scientific objectives and investigations of E^2T are discussed in detail in the Sections 3.1 and 3.2; Table 1 details the scientific objectives, the scientific questions, and the measurements requirements (payload, instrument parameters) to address the scientific goals of the E^2T mission.

Enceladus and Titan' investigations would be conducted using the E^2T mission model payload, which consists of three instruments: two time-of-flight mass spectrometers, the Ion and Neutral Gas Mass Spectrometer (INMS) and the Enceladus Icy Jet Analyzer (ENIJA) dust instrument; and a high-resolution infrared (IR) camera, the Titan Imaging and Geology, Enceladus Reconnaissance (TIGER). The scientific payload will be described in Section 4.

TABLE 1

3.1 Origin and Evolution of Volatile-Rich Ocean Worlds in the Saturn System

3.1.1 Chemical Constraints on the Origin and Evolution of Titan and Enceladus

The origin and evolution of Titan's methane still needs to be constrained. It is a key open question whether Titan's methane is primordial likely due to water-rock interactions in Titan's interior during its accretionary phase (Atreya et al., 2006) or else is delivered to Titan during its formation processes (Mousis et al., 2009) or by cometary impacts (Zahnle et al., 1992; Griffith and Zahnle, 1995). On Titan, the Huygens probe detected a small argon abundance (^{36}Ar) and a tentative amount of neon (^{22}Ne) in its atmosphere (Niemann et al., 2005, 2010), but was unable to detect the corresponding abundance of xenon and krypton. The presence of ^{22}Ne (^{36}Ar /^{22}Ne~1) was unexpected, as neon is not expected to be present in any significant amounts in protosolar ices (Niemann et al., 2005, 2010), and may indicate water-rock interactions and outgassing processes (Tobie et al., 2012). The non-detection of xenon and krypton supports the idea that Titan's methane was generated by serpentinization of primordial carbon monoxide and carbon dioxide delivered by volatile depleted planetesimals originating from within Saturn's subnebula (e.g., Atreya et al., 2006). To support a primordial methane source, xenon and krypton both would have to be sequestered from the atmosphere. While xenon is soluble in liquid hydrocarbon (solubility of 10^{-3} at 95 K) and could potentially be sequestered into liquid reservoirs, argon and krypton cannot be sequestered as soluble materials (Cordier et al., 2010). Therefore, the absence of measureable atmospheric krypton requires either sequestration into non-liquid surface deposits, such as clathrates (Mousis et al., 2011), or depletion in the noble gas concentration of the planetesimals (Owen and Niemann, 2009). Unlike Cassini INMS, the E^{2}T INMS has the mass range and the sensitivity to accurately measure xenon. E^{2}T would measure the abundance of noble gases in the upper atmosphere of Titan to discriminate between crustal carbon sequestration and carbon delivery via depleted planetesimals.

The longevity of methane in Titan's atmosphere is still a mystery. The value of ^{12}C/^{13}C in Titan's atmosphere has been used to conclude that methane outgassed ~10^7 years ago (Yung

et al., 1984), and is being lost via photolysis and atmospheric escape (Yelle et al., 2008). It is an open question whether the current methane rich atmosphere is a unique event, in a steady state where methane destruction and replenishment are in balance (Jennings et al., 2009), or is a transient event and is in a non-steady state where methane is being actively depleted or replenished. Indeed, the possibility that Titan did not always possess a methane rich atmosphere seems to be supported by the fact that the amount of ethane on Titan's surface should be larger than the present inventory; though Wilson and Atreya (2009) contend that missing surface deposits may simply be reburied into Titan's crust and Mousis and Schmitt (2008) have shown that it is possible for liquid ethane to react with a water-ice and methane-clathrate crust to create ethane clathrates and release methane. Nixon et al. (2012), however, favours a model in which methane is not being replenished and suggest atmospheric methane duration is likely between 300 and 600 Ma given that Hörst et al. (2008) demonstrated that 300 Ma is necessary to create Titan's current CO inventory and recent surface age estimates based on cratering (Neish and Lorenz, 2012). Mandt et al. (2012) suggests that methane's presence in the atmosphere, assumed here to be due to outgassing, has an upper limit of 470 Ma or else up to 940 Ma if the presumed methane outgassing rate was large enough to overcome $^{12}C/^{13}C$ isotope fractionation resulting from photochemistry and escape. Both the results of Mandt et al. (2012) and Nixon et al. (2012) fall into the timeline suggested by interior models (Tobie et al., 2006) which suggests that the methane atmosphere is the result of an outgassing episode that occurred between 350 and 1350 Ma. On Titan, both simple (methane, ethane and propane) and complex hydrocarbons precipitate out of the atmosphere onto the surface. Measuring the isotopic ratios ($^{14}N/^{15}N$; $^{12}C/^{13}C$; D/H; $^{16}O/^{18}O$; $^{17}O/^{16}O$) and abundances of the simple alkanes (e.g., methane, ethane and propane) would constrain the formation and evolution of the methane cycle on Titan. Further measurements of radiogenic noble gases such as ^{40}Ar and ^{22}Ne, which are typically markers of volatile elements from

Titan's interior can constrain outgassing episodes. Detection of ^{40}Ar and tentatively ^{22}Ne in the atmosphere has provided circumstantial evidence of water-rock interactions and methane outgassing from the interior (Niemann et al., 2010; Tobie et al., 2012). Recent measurements by ground-based observatories including measurements of CO and its carbon and oxygen isotopologues accompanied by the first detection of ^{17}O in Titan and indeed in the outer Solar System by Atacama Large Millimeter/submillimeter Array (ALMA) can be followed up on in more detail by in-situ spectroscopic measurements (Serigano et al., 2016). E^2T would measure the composition and isotopic ratios of Titan's upper atmosphere to determine the age of methane in the atmosphere and characterize outgassing history.

On Enceladus, Cassini measurements by INMS (Waite et al., 2006, 2009, 2017) and UVIS (Hansen et al., 2006, 2008) showed that plume gas consists primarily of water vapour with a few percent other volatiles. In addition to H_2O, as the dominant species, INMS was able to identify CO_2 (0.3-0.8%), CH_4 (0.1-0.3%), NH_3 (0.4-1.3%) and H_2 (0.4-1.4%), in the vapour plume as well as an unidentified species with a mass-to-charge (m/z) ratio of 28, which is thought to be either N_2 or C_2H_4, a combination of these compounds, or CO. The low mass resolution of Cassini INMS is insufficient to separate these species, and the UVIS measurements can only provide upper limits on N_2 and CO abundance. Determining the abundance ratio between these different species is, however, essential to constrain the origin of volatiles on Enceladus and to assess if they were internally reprocessed. A high CO/N_2 ratio, for instance, would suggest a cometary-like source with only a moderate modification of the volatile inventory, whereas a low CO/N_2 ratio would indicate significant internal reprocessing.

In addition to these main volatile species, the possible presence of trace quantities of C_2H_2, C_3H_8, methanol, formaldehyde and hydrogen sulphide has been detected within the INMS data recorded during some Cassini flybys (Waite et al., 2009). Organic species above the

INMS mass range of 99 u are also present but could not be further constrained (Waite et al., 2009). The identification and the quantification of the abundances of these trace species remains very uncertain due to the limitations of the mass spectrometer onboard Cassini.

Except for the measurement of D/H in H_2O on Enceladus (which has large uncertainty, Waite et al., 2009), no information is yet available for the isotopic ratios in Enceladus' plume gas. The E^2T mission would determine the isotopic ratios (D/H, $^{12}C/^{13}C$, $^{16}O/^{18}O$, $^{14}N/^{15}N$) in major gases compounds of Enceladus' plume, as well as $^{12}C/^{13}C$ in organics contained in icy grains. Comparison of gas isotopic ratios (e.g., D/H in H_2O and CH_4, $^{12}C/^{13}C$ in CH_4, CO_2, and CO; $^{16}O/^{18}O$ in H_2O, CO_2, CO; $^{14}N/^{15}N$ in NH_3 and N_2) and with Solar System standards would provide essential constraints on the origin of volatiles and how they may have been internally reprocessed. Simultaneous precise determination of isotopic ratios in N, H, C and O- bearing species in Enceladus' plume and Titan's atmosphere would permit a better determination of the initial reference values and a quantification of the fractionation due to internal and atmospheric processes on both moons.

Noble gases also provide essential information on how volatiles were delivered to Enceladus and if significant exchanges between the rock phase and water-ice phase occurred during Enceladus' evolution. The E^2T mission would be able to determine the abundance of ^{40}Ar, expected to be the most abundant isotope, as well as the primordial (non-radiogenic) argon isotopes, ^{36}Ar and ^{38}Ar. The detection and quantification of ^{36}Ar and ^{38}Ar would place fundamental constraints on the volatile delivery in the Saturn system. A low $^{36}Ar/N_2$ ratio, for instance, would indicate that N_2 on Enceladus is not primordial, like on Titan (Niemann et al., 2010), and that the fraction of argon brought by cometary materials on Enceladus is rather low. In addition to argon, if Ne, Kr, and Xe are present in detectable amounts, E^2T would be able to test whether primordial noble gases on Enceladus were primarily brought by a chondritic phase or cometary ice phase, which has implications for all the other primordial

volatiles. The $^{40}Ar/^{38}Ar/^{36}Ar$ as well as $^{20}N/^{21}Ne/^{22}Ne$ ratios would also allow for testing of how noble gases were extracted from the rocky core. Abundance ratios between Ar/Kr and Ar/Xe, if Kr and Xe are above detection limits, would offer an opportunity to test the influence of clathrate storage and decomposition in volatile exchanges through Enceladus's ice shell.

The origin of methane detected in Enceladus' plume is still uncertain. Methane, ubiquitous in the interstellar medium, was most likely embedded in the protosolar nebula gas. The inflow of protosolar nebular gas into the Saturn subnebula may have trapped methane in clathrates that were embedded in the planetesimals of Enceladus during their formation. Alternatively, methane may have been produced via hydrothermal reactions in Enceladus' interior; a possibility made more evident by the recent discovery of molecular hydrogen in Enceladus' plume (Waite et al., 2017). Mousis et al. (2009) suggests that if the methane of Enceladus originates from the solar nebula, then Xe/H_2O and Kr/H_2O ratios are predicted to be equal to $\sim 7\times10^{-7}$ and 7×10^{-6} in the satellite's interior, respectively. On the other hand, if the methane of Enceladus results from hydrothermal reactions, then Kr/H_2O should not exceed $\sim 10^{-10}$ and Xe/H_2O should range between $\sim 1\times10^{-7}$ and 7×10^{-7} in the satellite's interior. The E^2T mission by performing in situ analysis with high-resolution mass spectrometry of both the vapour and solid phases would quantify the abundance ratios between the different volatile species present in the plume of Enceladus, the isotopic ratios in major species, and the noble gas abundance.

3.1.2 Sources and Compositional Variability of Enceladus' Plume

The detection of salty ice grains (Postberg et al., 2009, 2011), the high solid/vapour ratio (Porco et al., 2006; Ingersoll and Ewald, 2011), and the observations of large particles in the

lower part of the plume (Hedman et al., 2009) all indicate that the plume of Enceladus originates from a liquid source likely from the subsurface ocean rather than from active melting within the outer ice shell (Figure 3). However, the abundance of the major gas species observed by Cassini suggests some contribution from the surrounding cold icy crust should also be considered. If plume gases exclusively originate from a liquid water reservoir, low-solubility species would be more abundant than high-solubility compounds, which is not apparent in the INMS data. The saltiness of the ice grains and the recent detections of nanometer sized silica dust in E-ring particles, which are believed to come from Enceladus (Hsu et al., 2011, 2015), and molecular hydrogen in Enceladus' plume (Waite et al., 2017) all indicate their origin is a location where alkaline high temperature hydrothermal reactions and likely water-rock interactions are occurring.

FIGURE 3

Although the Cassini (Cosmic Dust Analyzer) CDA has constrained knowledge of plume compositional stratigraphy, measurements of the absolute abundance and composition of organics, silicates and salts are poorly constrained given the low spatial resolution (10 km), low mass resolution and limited mass range of the CDA. The E^2T ENIJA is capable of providing a spatial accuracy of better than 100 m, allowing for a precise determination of compositional profiles along the spacecraft trajectory (Srama et al., 2015). The Cassini INMS provided only plume integrated spectra and is not able to separate gas species with the same nominal mass. E^2T INMS' high mass resolution, which is 50 times larger than that of Cassini INMS, would allow for separation of isobaric interferences, for example separating ^{13}C and ^{12}CH and CO and N_2. Determining high-resolution spatial variations in composition is crucial to establish whether the jets are fed by a common liquid reservoir or if jet sources are disconnected, and if the local liquid sources interact with a heterogeneous region in the icy crust. Variations in composition between the solid and gas phases as a function of distance

391 from jet sources can also provide information about how the less volatile species condense on
392 the grains, thus constraining the eruption mechanisms. The E^2T mission would allow for the
393 determination of the compositional distribution between both solid and vapour phases of
394 Enceladus' plume, thus providing crucial constraints on the nature and composition of the jet
395 sources, and on the relative contributions of subsurface liquid reservoirs and the surrounding
396 cold icy crust. Spatial variations in composition within the plume and possible correlations
397 with the jet sources would permit for testing if the volatile compounds originate from a
398 common reservoir and how the less volatile compounds are integrated in the solid particles
399 during the eruption processes.

400

401 *3.1.3 Geological Constraints on Titan's Methane Cycle and Surface Evolution*

402 There is an open question whether Titan's methane-rich atmosphere is being actively
403 replenished, or if methane is being lost and Titan's methane may eventually be depleted
404 (Yung et al., 1984). Cryovolcanism has been suggested as a mechanism by which methane
405 and argon can be transported from Titan's interior to its surface (e.g., Lopes et al., 2013).
406 Cryovolcanic activity may also promote methane outgassing (Tobie et al., 2006); while
407 methane clathrates are stable in Titan's ice shell in the absence of destabilizing thermal
408 perturbations and/or pressure variation, variations in the thermal structure of Titan's outer ice
409 shell during its evolution could have produced thermal destabilization of methane clathrates
410 generating outgassing events from the interior to the atmosphere (Tobie et al., 2006; see also
411 Davies et al., 2016). A number of candidate cryovolcanic features have been identified in
412 Cassini observations (Lopes et al., 2013). E^2T TIGER high-resolution colour images would
413 provide the data needed to determine the genesis of these features. Stratigraphic relationships

and crater counting would provide a means by which the relative ages of these features may be constrained.

A related question to the age of Titan's atmosphere is if Titan's climate is changing. At present, most of the observed liquid methane is located in the north polar region (Aharonson et al., 2009). There have been suggestions, however, that organic seas may have existed in Titan's tropics (Moore and Howard, 2010; MacKenzie et al., 2014), and/or in broad depressions in the south (Aharonson et al., 2009; Hayes et al., 2011). Observations and models suggest Titan's methane distribution varies on seasonal timescales (e.g., Waite et al., 2007; Hayes et al., 2010; Turtle et al., 2011; Coustenis et al., 2013, 2016) or Milankovitch timescales (Aharonson et al., 2009). Alternative models suggest that methane is being depleted and Titan's atmosphere is drying out (Moore and Howard, 2010). High-resolution images of the margins and interiors of these basins would allow us to determine if they once held seas. Identification of impact features or aeolian processes within these basins would help to constrain the timing of their desiccation.

In addition to their inherent scientific interest, Titan's dunes serve as sensitive indicators of climatic evolution (Lorenz et al., 2006; Radebaugh et al., 2008). Larger dune forms take longer to form than smaller dune forms. In Earth's Namib desert, these differing timescales result in large, longitudinal dunes that adhere to the overall wind conditions from the Pleistocene 20,000 years ago, while smaller superposing dunes (sometimes called rake dunes, or flanking dunes) have responded to the winds during our current interglacial and orient ages accordingly (Lancaster et al., 1989). On Titan, E^2T TIGER's superior spatial resolution would resolve these potential smaller dunes on top of the known longitudinal dunes, and would therefore reveal if Titan's recent climate has been stable or if it has changed over the past few Ma. The E^2T mission would provide high-resolution colour imaging of Titan that can be used

438 to characterize candidate cryovolcanic features that could be replenishing Titan's atmosphere
439 and paleo-seas, or dune patterns that evidence changes in Titan's climate.

440 Titan's geology is unique in that liquid and solid organics likely play key roles in many of
441 the observed processes. These processes, in turn, play an important role in modifying these
442 organics, both physically and chemically. Understanding these modification processes is
443 crucial to investigating the complex chemistry occurring on this moon. Furthermore, study of
444 Titan's geology allows us to investigate processes that are common on Earth, but in
445 drastically different environmental conditions, providing a unique way to gain insight into the
446 processes that shaped the Earth and pre-Noachian Mars.

447 Observations of Titan suggest the landscape is significantly modified by liquid organics
448 (e.g., Tomasko et al., 2005; Soderblom et al., 2007; Burr et al., 2013). Fluvial erosion is
449 observed at all latitudes, with a variety of morphologies suggesting a range of controls and
450 fluvial processes (Burr et al., 2013). High-resolution color imaging would provide insight into
451 the nature of this erosion: whether it is predominantly pluvial or sapping in nature and
452 whether it is dominated by mechanical erosion or dissolution. Dissolution processes are also
453 suspected to control the landscape of Titan's labyrinth terrains (Cornet et al., 2015) and may
454 be responsible for the formation of the polar sharp edged depressions (Hayes et al., 2008):
455 E^2T imaging would allow direct testing of these hypotheses.

456 Both fluvial and aeolian processes likely produce and transport sediments on Titan. Dunes
457 are observed across Titan's equator (e.g., Radebaugh et al., 2008; Malaska et al., 2016) while
458 a variety of fluvial sediment deposits have been identified across Titan (e.g., Burr et al., 2013;
459 Birch et al., 2016). Detailed images of the margins of the dune fields would allow us to
460 determine the source and fate of sands on Titan. E^2T images would also help determine
461 whether the observed fluvial features are river valleys or channels (cf. Burr et al., 2013)

462 providing key information in obtaining accurate discharge estimates needed to model
463 sediment transport (Burr et al., 2006). E^2T observation would provide insight into the primary
464 erosion processes acting on crater rims, which likely comprise a mixture of organics and
465 water ice (cf. Neish et al., 2015, 2016). Finally, E^2T images may provide insight into the
466 nature of erosion that exists in Titan's mid-latitudes, a region that shows little variability in
467 Cassini observations.

468 Of great interest in understanding the evolution of Titan's surface is determining the nature
469 of the observed geologic units, including their mechanical and chemical properties. Fluvial
470 processes, the degree to which mechanical versus dissolution dominates and the existence of
471 sapping, reflect the material properties of the surface and therefore can be used as a powerful
472 tool to investigate the properties of the surface. E^2T imaging would also allow us to
473 investigate the strength of the surface materials by constraining the maximum slopes
474 supported by different geologic units. Detailed color and stereo imaging of the boundaries of
475 units would also allow investigation of the morphology, topography, and spectral relationship
476 across unit boundaries. E^2T would take high-resolution color images of Titan that would
477 elucidate the nature of the geological evolution of Titan's organic-rich surface.

478

479 **3.2 Habitability and Potential for Life in Ocean Worlds, Enceladus and Titan**

480 **3.2.1 Evidence for Prebiotic and Biotic Chemical Processes on Titan and Enceladus**

481 Unlike the other ocean worlds in the Solar System, Titan has a substantial atmosphere,
482 consisting of approximately 95% nitrogen and 5% methane with trace quantities of
483 hydrocarbons such as ethane, acetylene, and diacetylene, and nitriles, including hydrogen
484 cyanide (HCN) and cyanogen (C_2N_2). Somewhat more complex molecules such as
485 cyanoacetylene, vinyl and ethylcyanide follow from these simpler units. In Titan's upper

atmosphere, Cassini has detected large organic molecules with high molecular masses over 100 u. In-situ measurements by the Cassini Plasma Spectrometer (CAPS) detected heavy positive ions (cations) up to 400 u (Crary et al., 2009) and heavy negative ions (anions) with masses up to 10,000 u (Coates et al., 2007) in Titan's ionosphere. Whereas Cassini INMS only had the ability to detect cations, E^2T INMS can detect both cations and anions and can do so with much better mass resolution than Cassini-INMS (and a fortiori than Cassini-CAPS). It is thought that these heavy negative ions, along with other heavy molecules found in the upper atmosphere, are likely the precursors of aerosols that make up Titan's signature orange haze, possibly even precipitating to the surface. While the compositions of these molecules are still unknown, their presence suggest a complex atmosphere that could hold the precursors for biological molecules such as those found on Earth. The ability to detect prebiotic molecules in Titan's atmosphere is currently limited by the mass range of the Cassini INMS to the two smallest biological amino acids, glycine (75 u) and alanine (89 u), and the limited mass resolution precludes any firm identification. While Cassini INMS has not detected 75 or 89 u molecules, it has detected positive ions at masses of 76 u and 90 u, which are consistent with protonated glycine and alanine, respectively (Vuitton et al., 2007; Hörst et al., 2012). Experimental results from a Titan atmosphere simulation experiment found 18 molecules that could correspond to amino acids and nucleotide bases (Hörst et al., 2012). The E^2T mission would use high-resolution mass spectrometry to measure heavy neutral and ionic constituents up to 1000 u, and the elemental chemistry of low-mass organic macromolecules and aerosols in Titan's upper atmosphere, and would monitor neutral-ionic chemical coupling processes.

The plume emanating from Enceladus' south pole probably contains the most accessible samples from an extra-terrestrial liquid water environment in the Solar System. The plume is mainly composed of water vapour and trace amounts of other gases (Waite et al., 2017). In

addition, higher molecular weight compounds with masses exceeding 100 u, were detected in the plume emissions (Waite et al., 2009; Postberg et al., 2015). The presence of CO_2, CH_4 and H_2 can constrain the oxidation state of Enceladus' hydrothermal system during its evolution. The minor gas constituents in the plume are indicative of high-temperature oxidation-reduction (redox) reactions in Enceladus' interior possibly a result of decay of short-lived radionucleides (Schubert et al., 2007). In addition, H_2 production and escape may be a result of redox reactions indicative of possible methanogenesis similar to the process occurring in terrestrial submarine hydrothermal vents (McKay et al., 2008; Waite et al., 2017). Further, the high temperatures and H_2 escape may have led to the oxidation of NH_3 to N_2 (Glein et al., 2008). Detection and inventory of reduced and oxidized species in the plume material (e.g., NH_3/N_2 ratio, H_2 abundance, reduced versus oxidized organic species) can constrain the redox state and evolution of Enceladus' hydrothermal system.

Cassini CDA measurements identified three types of grains in the plume and Saturn's E-ring. Type I and Type II grains are both salt-poor (Figure 4). Type I ice grains are nearly pure-water ice while Type II grains also possess silicates and organic compounds and Type III grains are salt-rich (0.5–2.0% by mass) (Postberg et al., 2009, 2011). The salinity of these particles, the high solid/vapor ratio (Porco et al., 2006, Ingersoll and Ewald, 2011) of the plume and observations of large particles in the lower part of the plume (Hedman et al., 2009) indicate that the plume originates from a subsurface liquid source. The Cassini CDA later detected nanometer sized silica dust particles in E-ring stream particles (Hsu et al., 2011, 2015) which indicate that water-rock interactions are likely taking place within this liquid reservoir. Gravity and libration data demonstrated that this liquid reservoir was a subsurface global ocean (Iess et al., 2014; Thomas et al., 2015)

FIGURE 4

Hsu et al. (2015) suggest that the ocean should be convective in order to have silica nanoparticles transported from hydrothermal sites at the rocky core up to the surface of the ocean where they can be incorporated into icy plume grains (Hsu et al., 2015). To confirm this hypothesis of current hydrothermal activity on Enceladus, a direct detection of silica and other minerals within ejected ice grains is required. SiO_2 nano-particles detected in Saturn's E-ring could be much better investigated and quantified by E^2T ENIJA given its high dynamic range (10^6–10^8). By performing high-resolution mass spectrometry of ice grains in Enceladus' plume, the E^2T mission would characterize the composition and abundance of organics, salts and other minerals embedded in ice grains, as messengers of rock/water interactions. It would also search for signatures of on-going hydrothermal activities from possible detection of native He and further constrain recent measurements of native H_2 found in Enceladus' plume (Waite et al., 2017).

3.2.2 Physical Dynamics in Enceladus' Plume and Links to Subsurface Reservoirs

The total heat emission at Enceladus' tiger stripes is at least 5 GW - possibly up to 15 GW, (Howett et al., 2011), and in some of the hot spots where jets emanate, the surface temperatures are as high as 200 K (Goguen et al., 2013). Cassini observations show that the plume is made up of approximately 100 discrete collimated jets as well as a diffuse distributed component (Hansen et al., 2008, 2011; Postberg et al., 2011; Porco et al., 2014). The majority of plume material can be found in the distributed diffuse portion of the plume, which likely originates from elongated fissures along Enceladus' tiger stripes while only a small portion of gas and grains are emitted from the jets (Hansen et al., 2011; Postberg et al., 2011). CDA measurements demonstrate that the majority of salt-poor grains tend to be ejected through the jets and at faster speeds while larger salt-rich grains tend to be ejected more

559 slowly through the distributed portion of the plume (Postberg et al., 2011). The ice to vapour
560 ratio can constrain how Enceladus' plume material is formed and transported to the surface.
561 For example, ice/vapour ratios > 0.1–0.2 would exclude plume generation mechanisms that
562 require a large amount of ice grains to be condensed from vapour (Porco et al., 2006;
563 Ingersoll and Pankine, 2011). However, this ratio is poorly constrained with estimates ranging
564 from 0.05 (Schmidt et al., 2008) to 0.4 (Porco et al., 2006) to 0.35–0.7 (Ingersoll and Ewald,
565 2011). E^2T high-resolution IR images and ENIJA can help constrain this important ratio.
566 Cassini ISS images used to track plume brightness variation, which is proportional to the
567 amount of grains in the plume, with the orbital position of Enceladus found more ice grains
568 are emitted when Enceladus is near its farthest point from Saturn (apocenter). It is not
569 understood if the plume vapour has such a variation. This temporal variation of the plume
570 indicates that it is tidally driven but could also be due to possible physical libration (Hurford
571 et al., 2009; Kite et al., 2016). Most recently, Kite et al. (2016) has suggested that the tiger
572 stripe fissures are interspersed with vertical pipe-like tubes with wide spacing that extend
573 from the surface to the subsurface water. This mechanism allows tidal forces to turn water
574 motion into heat, generating enough power to produce eruptions in a sustained manner.
575 TIGER can provide high spatial resolution thermal emissions maps to constrain the amount of
576 energy dissipated between the tiger stripes. The E^2T mission would use high resolution IR
577 imaging of the south polar terrain and mass spectra of the grains to provide new details of its
578 surface and constrain the links between plume activity, subsurface reservoirs and deep
579 hydrothermal processes.

580

581

582

3.2.3 Geological Evidence for Interior-Surface Communication on Titan

Geological processes such as tectonism and cryovolcanism indicate communication between the surface and subsurface. While Titan's surface offers a wealth of geological processes, the Cassini data lack the resolution needed in which to constrain the detailed nature of these processes, and thus to understand the extent that Titan's surface may be chemically interacting with its water-rich interior. Also of great importance to habitability are the transient H_2O melt sheets and flows (e.g., Soderblom et al., 2010) associated with impacts. On Titan, several features with volcanic landforms, lengthy flows, tall mountains, large caldera-like depressions, have been identified as possible cryovolcanic sites but could also possibly be due to other endogenic processes (Lopes et al., 2016; Solomonidou et al., 2016). At present, the Hotei Regio flows and the Sotra Patera region, which includes Sotra Patera, an elliptical deep depression on Titan, Mohini Fluctus, a lengthy flow feature, and Doom and Erebor Montes, two volcanic edifices, are considered to host the strongest candidates for cyrovolcanism on Titan (Lopes et al., 2013; Solomonidou et al., 2014, 2016).

A variety of mountainous topography has been observed on Titan (Radebaugh et al., 2007; Cook-Hallett et al., 2015). The observed morphologies of many of Titan's mountains suggest contractional tectonism (Mitri et al., 2010; Liu et al., 2016). This is somewhat surprising, however, since tectonic landforms observed on other ocean worlds and icy satellites in the outer solar system appear to be extensional in nature. Understanding Titan's tectonic regime would, thus not only provide insight into the transport of material between surface and the interior, but also into the evolution of the other ocean worlds. We would test the hypothesis that Titan's mountains are formed by contraction by mapping the faults driving mountain formation in topographic context. The shape of the fault outcrop draped against topography would allow us to measure the dips of faults, which would be ~30 degrees to the horizontal for compressive mountains and ~60 degrees for extensional mountains.

In addition to cryovolcanism and tectonism, which may transport water to Titan's surface, impact craters likely have created transient liquid-water environments on Titan's surface. Because of Titan's dense atmosphere, models suggest that melt sheets and flows associated with impact craters may remain liquid for 10^4–10^6 years (Thompson and Sagan, 1992; Artemieva and Lunine, 2005), though the stability of such melts is questioned (Senft and Stewart, 2011; Zahnle et al., 2014) and detailed imaging of the floors of young craters is needed to constrain these models. Titan offers numerous pathways for interaction between its organic-rich surface and liquid water. E^2T would provide high-resolution mapping (30 m/pixel with DTM vertical resolution of 10 m) that would offer the ability to distinguish cryovolcanic features and to investigate the morphology of Titan's mountains and impact craters.

4. Scientific Payload

The Explorer of Enceladus and Titan (E^2T) has a focused payload that would provide in-situ mass spectrometry and high-resolution imaging of Enceladus' south polar terrain and plume, and Titan's upper atmosphere and surface, from a solar-electric powered spacecraft in orbit around Saturn. The in-situ measurements of Titan's upper atmosphere would be acquired during 17 flybys with an altitude as low as 900 km. At Enceladus, in-situ measurements would be conducted during 6 flythroughs of the plume and flybys of the south polar terrain at altitudes between 50 and 150 km. At the closest approach the velocity of the S/C with respect to Enceladus surface is ~5 km/s and with respect to Titan surface is ~7 km/s. Imaging data will be collected during inbound and outbound segments of each flyby.

The E^2T mission model payload consists of three science instruments: two time-of-flight mass spectrometers, the Ion and Neutral gas Mass Spectrometer (INMS) and the Enceladus

Icy Jet Analyzer (ENIJA); and a high-resolution infrared camera, Titan Imaging and Geology, Enceladus Reconnaissance (TIGER). Two instruments in the E^2T payload were proposed to be provided by ESA member states. INMS would be provided by the Swiss Space Office (SSO) and ENIJA would be provided by the German Aerospace Center (DLR). NASA proposed to provide the third instrument, TIGER. The characteristics of the science payload are shown in Table 2. In addition, a Radio Science Experiment (RSE), not necessarily requiring specific hardware on-board the spacecraft – thus regarded as an "experiment of opportunity", was considered for further study. The RSE would improve the current determination of the gravity fields of Enceladus and Titan, by using the radio links between the E^2T spacecraft and Earth to better constrain their internal structure. The proposed main funding agency of the RSE is the Italian Space Agency (ASI).

TABLE 2

4.1 Ion and Neutral Mass Spectrometer (INMS)

The Ion and Neutral Mass Spectrometer (INMS) is a reflectron time-of-flight mass spectrometer that would record mass spectra of neutral and ionised gases during flybys and fly-throughs of Enceladus' plume and Titan's upper atmosphere. During flybys and flythroughs of Saturn's E-ring, Enceladus' plume and of Titan's upper atmosphere, INMS would record mass spectra of neutral, positive and negative ionized gases within the mass range 1 – 1000 u/e with a mass resolution at 5000 m/Δm (50%) and a sensitivity (detection threshold) of 1 cm^{-3} (~10^{-16} mbar) in a 5 s measurement cadence (Abplanalp et al., 2010; Wurz et al., 2012). The Cassini INMS (Ion and Neutral Mass Spectrometer) is limited to the detection of low mass species of 1–99 u and is therefore not able to measure the high-mass molecular species abundant in Enceladus' plume and in Titan's upper atmosphere. Further,

656 E^2T INMS' high mass resolution ($m/\Delta m = 5000$), 50 times larger than Cassini INMS, allows
657 for the separation of isobaric interference such as occur at masses 16, 20, 28, and 40 u/e
658 which is possible based on heritage of RTOF/ROSINA from the Rosetta mission (Scherer et
659 al., 2006; Wurz et al., 2015) as shown in Figure 5.

660 FIGURE 5

661 Cassini INMS only had the ability to measure positively charged ions. In-situ
662 measurements by the Cassini Plasma Spectrometer (CAPS) detected heavy negative ions in
663 Titan's ionosphere and negative water group ions in Enceladus' plume (Coates et al., 2010).
664 E^2T INMS would record 10,000 mass spectra per second and accumulate these for a pre-set
665 period, allowing for a time resolution of measurements in the range from 0.1 to 300 s. The
666 maximum INMS time resolution at Enceladus is 0.1 s for recording a mass spectrum
667 (corresponding to a spatial resolution of ~0.5 km for a S/C velocity of 5 km/s) to resolve
668 small-scale structure in the plume; the maximum used INMS resolution at Titan is 5 s
669 (corresponding to a spatial resolution of ~35 km for a S/C velocity of 7 km/s).

670 The E^2T INMS has heritage based on P-BACE instrument (Abplanalp et al., 2009.), RTOF
671 instrument of ROSINA (Scherer et al., 2006) on Rosetta and on the gas-chromatograph-
672 neutral gas mass spectrometer prototype which will be used in upcoming lunar exploration
673 conducted by the Russian Space Agency in 2019 (Hofer et al., 2015; Wurz et al., 2015). Most
674 recently, a similar instrument, the Neutral gas and Ion Mass spectrometer (NIM) instrument
675 of the Particle Environment Package (PEP) consortium, is developed for the JUICE mission
676 of ESA.

677

678

679 **4.2 Enceladus Icy Jet Analyzer (ENIJA)**

680 Enceladus Icy Jet Analyzer (ENIJA) is optimized to search for prebiotic molecules and
681 biogenic compounds in objects with high dust fluxes and number densities such as occur in
682 Enceladus' plume or cometary comae. ENIJA consists of two time-of-flight mass
683 spectrometer subsystems and a software-enabled high flux detector (HFD) that runs in parallel
684 to the spectrometers. The high flux detector measurement mode will map the dynamical
685 profile (number density and size distribution) of Enceladus' ice jets. Compared to the Cassini
686 CDA (Cosmic Dust Analyzer), ENIJA has a 40 times better mass resolution, a 100 times
687 better maximum flux and sensitivity, and a spatial resolution that is 50 times better (Srama et
688 al., 2004, 2015; Postberg et al., 2009). Moreover, the twin-detector instrument will acquire
689 mass spectra of both cations and anions created upon ice particle impact, simultaneously,
690 whereas CDA only measured cations. During flythroughs of Enceladus plume and in the E
691 ring, ENIJA provides time-of-flight mass spectra for ice particles with a mass range of 1–
692 2000 u with a mass resolution (m/Δm) of 970 between 23-2000 u. ENIJA acquires time-of-
693 flight mass spectra of individual ice grains by impact ionization and analyses the generated
694 anions and cations with two separate time of flight spectrometer sub-units. In this way ENIJA
695 is able to quantify organic compounds at <10 ppm; additionally some polar organic species,
696 such as most amino acids, can be quantified well below 1 ppm (Figure 6). With a dynamic
697 range up to 10^8, inorganic trace components with sub-ppm concentrations from Enceladus'
698 ocean, can now be investigated simultaneously with the more abundant mineral species.
699 ENIJA is not a newly developed instrument; rather it is an optimization and miniaturization of
700 flight-proven hardware. ENIJA has heritage based on Giotto-Particle Impact Analyzer (PIA),
701 Stardust Cometary Interstellar Dust Analyzer (CIDA), Cassini's' Cosmic Dust Analyzer
702 (CDA), Rosetta COmetary Secondary Ion Mass Analyzer (COSIMA) and Europa-SUrface
703 Dust mass Analyzer (SUDA).

ACCEPTED MANUSCRIPT

704 FIGURE 6

705

706 **4.3 Titan Imaging and Geology, Enceladus Reconnaissance (TIGER)**

707 Titan Imaging and Geology, Enceladus Reconnaissance (TIGER) is a near infrared (NIR)
708 camera designed to acquire high resolution images of Titan and Enceladus. TIGER would
709 observe Titan at 30–100 m/pixel in three wavelengths, 1.3, 2, and 5 µm and Enceladus
710 emissions at 1 m/pixel at two wavelengths, 5 and 5.3 µm. Images acquired by TIGER would
711 enable investigation of Titan's geology, hydrology, and compositional variability and study of
712 the composition and kinematics of Enceladus' jets and plumes. The TIGER band passes are
713 selected to match with Titan's atmospheric transmission windows (Lemmon et al., 1993;
714 Sotin et al., 2005) to enable direct ground observations using reflected sunlight and to
715 measure thermal emission from Enceladus. The 5 µm images are subject to virtually no
716 scattering from Titan's atmospheric aerosols, allowing diffraction limited images achieving
717 spatial resolutions an order of magnitude better than Cassini observations (Clark et al., 2010;
718 Soderblom et al., 2012; Barnes et al., 2014) and would be highly sensitive to organic
719 composition (Clark et al., 2010; Barnes et al., 2014). TIGER has the capability to image Titan
720 at Huygens DISR resolution (Figure 7). At Enceladus, the 5 and 5.3 µm observations would
721 measure thermal emission of surfaces as cold as 130 K and would provide temperature maps
722 of the surface at Cassini-ISS image scales. A fine steering mirror (FSM) would be employed
723 to select and track regions of interest during the flyby and compensate for spacecraft jitter
724 allowing for longer exposures and better signal-to-noise ratio (SNR). Digital time-delay
725 integration (TDI) would also be employed, as needed, during closest approach when the
726 ground speed is highest.

727 FIGURE 7

728 While the TIGER instrument is a new design, it utilizes high-heritage subsystems with a
729 Technology Readiness Level (TRL) larger of 6. The TIGER Focal Plane Array (FPA) is from
730 Teledyne Imaging Sensors and utilizes the same HgCdTe detector and H2RG qualified for
731 James Webb Space Telescope (JWST) Near Infrared Camera (NIRCam) (Rieke et al., 2005),
732 Near Infrared Spectrometer (NIRSpec) (Rauscher et al., 2004), and Fine Guidance Sensor
733 (FGS) (Doyon et al., 2012) instruments. TIGER also implements FPA readout electronics that
734 have been qualified extensively for the JWST instruments (Loose et al., 2006) and are used by
735 the Euclid Near Infrared Spectrometer and Photometer (NISP) (Maciaszek et al., 2014).

736

737 **4.4 Radio Science Experiment**

738 Gravity field measurements are powerful tools to constrain the interior structure of the planets
739 and satellites and to assess mass anomalies, providing information on the internal dynamics
740 and evolution. The observable quantities used by gravity science experiments are obtained by
741 means of spacecraft tracking at microwave frequencies from a terrestrial ground antenna. The
742 eight gravity flybys of Titan conducted by the Cassini mission (not including the T122 flyby,
743 recently completed in August 2016) yielded sufficient information to obtain a robust
744 estimation of the degree-3 static gravity field, plus the fluid Love number k_2 (Iess et al., 2010,
745 2012); however, this was not the case for Enceladus (Rappaport et al., 2008; Iess et al., 2014),
746 where only the degree-2 static gravity field and J_3 were observed, and the tidal response to
747 Saturn's gravity was not detected.

748 In Enceladus' south polar terrain a larger time-variation of the gravity field with respect to
749 the global solution of the time variation of the gravity field is expected because the ice shell
750 thickness is anticipated to be locally thin in that area. A gravity science experiment, based on
751 radio tracking and precise spacecraft multiarc orbit determination (see e.g. Tortora et al.,

752 2016), would determine the local solution of the gravity field of Enceladus at the south polar
753 terrain thus allowing the determination of the thickness variation at the south polar regions
754 and constraining the mechanical properties (viscosity) of the ice overlying the outer ice shell.
755 The expected tidal deformation is characterized by a pattern more complex than the standard
756 degree-two pattern, with a strong amplification of the tidal fluctuation in the south polar
757 terrain (Behounkova et al., 2017).

758 For Titan gravity science, a desirable objective would be the improve of Cassini's results
759 in terms of reduction of the uncertainty in the fluid Love number k_2. In particular, the
760 geophysics objectives relative to the subsurface liquid ocean extent, shell thickness and shell
761 viscosity would be improved if the formal uncertainty of k_2 could be reduced down to values
762 in the range 0.05-0.01. The feasibility of such result strongly depends on the actual geometry
763 of Titan's flyby (in particular: altitude at pericenter and mean anomaly of Titan around Saturn
764 at the time of closest approach with the E^2T spacecraft), in addition to the stability of the radio
765 link (preferably at Ka-band, to reduce the detrimental effect of dispersive media).

766 The availability of a two-way (uplink and downlink) radio system would also allow
767 expanding the long series of Titan's occultations carried out by Cassini's radio science
768 subsystem to probe Titan's neutral and ionized atmosphere (Coustenis et al., 2016; Kliore et
769 al., 2008, 2011; Schinder et al., 2012, 2015). The main objective would be the
770 characterization of Titan's atmospheric structure (pressure, density, and total electron content
771 profiles, versus altitude) at different latitude/longitudes.

772

773 **5. Proposed Mission and Spacecraft Configuration**

774 The baseline scenario for the E^2T mission is a solar electric powered spacecraft (S/C), in orbit
775 around Saturn, performing multiple flybys of Titan and Enceladus. The baseline includes a

shared launch on the Ariane 6.4 with a co-manifest of estimated launch mass of ~2636 kg to geosynchronous transfer orbit (GTO), with a forecasted shared launch opportunity in 2030. The Ariane 6.4 with four solid rocket boosters is scheduled to debut in 2020–2021. The upper stage reignition capability of the Ariane 6.4 will enable a GTO/escape dual launch. The mass estimate for the co-manifest places it in the lower end of the Intermediate Class (2,500-4,200 kg) of commercial satellites (FAA, 2015). After a transfer from GTO to a hyperbolic escape trajectory, E^2T S/C would pursue a gravity assist flyby of the Earth to help propel itself to the Saturn system. The cruise phase from Earth to Saturn would be 6 years long. The E^2T tour in the Saturn system would be 3.5 years long. The E^2T tour consists of 6 flybys of Enceladus above the south polar terrain with a flight altitude range between 50 to 150 km, and 17 flybys of Titan at a reference flight altitude ranging from 1500 down to 900 km.

5.1 Transfer Orbit to Saturn

The nominal transfer to Saturn uses a $1:1^+$ resonant Earth flyby trajectory, as shown in Table 3 and Figure 8. Electric propulsion is used to strongly increase the hyperbolic flyby speed at the Earth, and to provide a subsequent energy boost to reduce the total flight time to 6 years to Saturn orbit insertion (SOI). After separation from the co-manifested satellite and the SYstème de Lancement Double Ariane (SYLDA) carrier, the upper stage will be re-ignited near perigee to impart approximately 1.2 km/s ΔV to the E^2T S/C, putting it on a hyperbolic escape trajectory with a characteristic energy (C_3) of 9.6 km^2/s^2. Thanks to the use of electric propulsion and an Earth gravity assist, the declination of the asymptote can vary between ±5 degrees without significant propellant mass penalty. Similarly, as shown in Table 4, there is negligible performance variation over a 21-day launch period. The data in Table 3 is based on a proposed launch from January-May 2030 during which there is little variation in flight time or required propellant mass. Lunar flybys on the escape trajectory were not considered, but

could be used to effectively lower the required escape ΔV, which in turn would allow for a heavier co-manifested satellite.

FIGURE 8

TABLE 3

TABLE 4

5.2 Saturn Tour

The duration of the tour from SOI through to the end of the 17-flyby Titan phase is about 3.5 years. A sample tour is shown in Figure 9. Like Cassini, end-of-mission spacecraft disposal is Saturn impact, by means of three to four additional Titan flybys.

The tour is divided in a first Enceladus science phase and in a second Titan science phase. The S/C should perform at least 6 flybys of Enceladus above the south polar terrain and at least 17 flybys of Titan. To prevent contamination from Titan's organics for Enceladus science, E^2T S/C will perform close flybys of Enceladus at the beginning of the tour (Enceladus science phase); distant flybys of Titan would be performed during the initial tour phase. After the main Enceladus phase, close flybys of Titan with atmospheric sampling would be performed (Titan science phase).

FIGURE 9

During Enceladus science phase, E^2T would provide in-situ sampling of the plume at a minimum altitude from Enceladus' surface ranging between 50 and 150 km using INMS and ENIJA. At the closest approach, the velocity of the S/C with respect to Enceladus' surface will be approximately 5 km/s. E^2T would provide high-resolution imaging of Enceladus

surface with the TIGER camera. During Enceladus flybys the observations of INMS, ENIJA and TIGER would be performed simultaneously. During this phase observations of Titan's surface using TIGER are scheduled during distant flybys.

During the Titan science phase, E^2T would provide in-situ sampling of the upper atmosphere at a minimum altitude from Titan surface as low as 900 km using INMS. At the closest approach the velocity of the S/C with respect to Titan's surface will be approximately 7 km/s. E^2T will provide high-resolution imaging of Titan's surface with TIGER camera. During Titan flybys the observations of INMS and TIGER can be performed simultaneously.

5.3 Spacecraft Design and Structure

The proposed E^2T spacecraft (S/C) design is the result of the harmonisation of several high-TRL key space technologies already developed on behalf of several ESA space exploration missions, such as: ExoMars 2016, JUICE, BepiColombo and Rosetta. The proposed E^2T spacecraft architecture would be derived from the ESA ExoMars Trace Gas Orbiter (TGO) developed for the 2016 mission. The proposed E^2T architecture is in compliance with the Ariane 6.4 launcher size constraints. Table 5 summarizes its main technical characteristics. The E^2T spacecraft baseline configuration is depicted in Figure 10.

TABLE 5

FIGURE 10

The Solar Arrays (SA) are the main S/C electrical power source, whereas the S/C main battery provides the emergency backup one. The E^2T four panel solar arrays are based on space-qualified Thales Alenia Space SolarBus W51 technologies and GaAs/Ge Low Intensity Low Temperature (LILT) solar cells technology used in the ESA JUICE mission. The Solar

Electric Propulsion System (SEPS) is based on four (3+1 for redundancy) QinetiQ's T6 gridded ion thruster engines, whereas two chemical propulsion systems have been also implemented for S/C manoeuvres and Attitude and Orbital Control System (AOCS) purposes. The propulsion systems' designs have been directly derived from different mission heritages, such as the two ESA missions GOCE and BepiColombo, and the NASA/ESA/ASI Cassini mission. The heritage of systems and subsystems from previous ESA missions requiring limited technological development enables a relatively low cost for the industrial development. The E^2T S/C design was performed by Thales Alenia Space (TAS-I, Torino, Italy)

6. Summary

The previous decades have seen a revolution in our understanding of ocean worlds thanks in part to the Cassini-Huygens mission. Recent space missions to the outer Solar System have shown that liquid water once thought to be confined to Earth is quite common in the Solar System. The finding of subsurface oceans on Titan and Enceladus increases the possibility of extra-terrestrial habitability since life as we know it requires water, energy and nutrients (McKay et al., 2008, 2016). The geological activity represented by the eruptions forming Enceladus' plume at its south polar terrain and the organic-rich atmosphere and geological landscape, including lakes and seas, on Titan provide an ideal laboratory to investigate processes that may have been operational in Earth's early history.

Scientific investigations at both Titan and Enceladus as well as terrestrial experiments replicating conditions found on the icy moons have hinted at the real possibility of prebiotic activity on these icy ocean worlds. The ability to detect prebiotic molecules in Titan's atmosphere is currently limited by the mass range of the Cassini INMS to the two smallest

biological amino acids, glycine (75 u) and alanine (89 u), and the limited mass resolution precludes any firm identification. However, the recent detection of prebiotic chemicals (glycine and phosphorus) in comet Churyumov/Gerasimenko (Altwegg et al., 2016) makes it likely that prebiotic molecules could be also present in Enceladus' plumes. Experimental results from a Titan atmosphere simulation experiment found 18 molecules that could correspond to amino acids and nucleotide bases (Hörst et al., 2012). The E^2T mission would use high-resolution mass spectrometry to measure heavy neutral and ionic constituents up to 1000 u, and the basic chemistry of low-mass organic macromolecules and aerosols in Titan's upper atmosphere, and would monitor neutral-ionic chemical coupling processes.

Investigations of Enceladus' plume emissions has yielded tantalizing information which are suggestive of prebiotic/biotic activity, including the detection of higher molecular weight compounds with masses exceeding 100 u, (Waite et al., 2009; Postberg et al., 2015) and minor gas constituents in the plume that are indicative of high-temperature oxidation-reduction (redox) reactions in Enceladus' interior. Recent detection of molecular hydrogen in Enceladus' plume by Cassini could indicate processes that are similar in nature to processes that occur in terrestrial submarine hydrothermal vents which emit large quantities of molecular hydrogen due to water-rock interactions. In these hydrothermal vents, molecular hydrogen serves as a fuel that supports both abiotic and biotic production of organic molecules that have been detected in Enceladus' plume (Waite et al., 2015, 2017). Detection and inventory of reduced and oxidized species in the plume material (e.g., NH_3/N_2 ratio, H_2 abundance, reduced versus oxidized organic species) such as proposed by the E^2T mission could constrain the redox state and evolution of Enceladus' hydrothermal system.

Another key question still to be resolved is to understand how the plume is generated. E^2T's imaging system could provide high spatial resolution thermal emissions maps to constrain the amount of energy dissipated between the tiger stripes. The E^2T mission would

use high resolution infrared imaging of the south polar terrain and mass spectra of the grains to provide new data of its surface to constrain the links between plume activity, subsurface reservoirs and deep hydrothermal processes.

While Titan's surface offers a wealth of geological processes, the Cassini data lack the resolution necessary to constrain the detailed nature of these processes. E^2T would provide high-resolution mapping (30 m/pixel with DTM vertical resolution of 10 m) that would offer the ability to distinguish geological features and morphology. Further, the high-resolution imaging of Titan's surface could improve understanding of the extent that Titan's organic-rich surface may be chemically interacting with its water-rich interior. By combining in-situ chemical analysis of Titan's atmosphere and Enceladus' plume with observations of Enceladus' plume dynamics and Titan's surface geology, E^2T will be able to address fundamental science questions that would relate to content, origin and evolution of ocean worlds. The science that a mission such as E^2T can provide can also answer questions about Earth's early history as well as help to predict the physical and chemical properties of exoplanets beyond the Solar System.

Acknowledgments

We thank two anonymous reviewers for their constructive remarks and suggestions.

ACCEPTED MANUSCRIPT

REFERENCES

Abplanalp et al. 2010. An optimised compact electron impact ion storage source for a time-of-flight mass spectrometer International Journal of Mass Spectrometry 294, 33-39.

Abplanalp, D., Wurz, P., Huber, L., Leya, I., Kopp, E., Rohner, U. Wieser, M., Kalla, L., Barabash, S. 2009. A neutral gas mass spectrometer to measure the chemical composition of the stratosphere Advances in Space Research, 44, pp. 870-878

Aharonson, O., Hayes, A.G., Lunine, J.I., Lorenz, R.D., Allison, M.D., Elachi, C. 2009. An asymmetric distribution of lakes on Titan as a possible consequence of orbital forcing. Nature Geoscience 2, 851-854.

Altwegg, K., H. Balsiger, A. Bar-Nun, J.-J. Berthelier, A. Bieler, P. Bochsler, C. Briois, U. Calmonte, M. Combi, H. Cottin, J. De Keyser, B. Fiethe, S.A. Fuselier, S. Gasc, T.I. Gombosi, K.C. Hansen, M. Hässig, A. Jäckel, E. Kopp, A. Korth, L. Le Roy, U. Mall, B. Marty, O. Mousis, T. Owen, H. Rème, M. Rubin, T. Sémon, C.-Y. Tzou, J.H. Waite, and P. Wurz, Prebiotic chemicals - amino acid and phosphorus - in the coma of comet 67P/Churyumov-Gerasimenko, Science Advances 2:e1600285 (2016) 5pp.

Artemieva, N., Lunine, J.I. 2005. Impact cratering on Titan II. Global melt, escaping ejecta, and aqueous alteration of surface organics. Icarus 175, 522-533.

Atreya, S.K., Adams, E.Y., Niemann, H.B., Demick-Montelara, J.E., Owen, T.C., Fulchignoni, M., Ferri, F., Wilson, E.H. 2006. Titan's methane cycle. Planetary and Space Science 54, 1177-1187.

Barnes, J.W., Sotin, C., Soderblom, J.M., Brown, R.H., Hayes, A.G., Donelan, M., Rodriguez, S., Le Mouélic, S., Baines, K.H. and McCord, T.B., 2014. Cassini/VIMS observes rough surfaces on Titan's Punga Mare in specular reflection. Planetary Science, 3(1), p.1.

Barnes, J., et al. 2012. AVIATR—Aerial Vehicle for In-Situ and Airborne Titan Reconnaissance, Experimental Astronomy, Vol. 33, No. 1, pp. 55–127. doi:10.1007/s10686-011-9275-9

Beghin, C., Randriamboarison, O., Hamelin, M., Karkoschka, E., Sotin, C., Whitten, R.C., Berthelier, J.J., Grard, R., Simoes, F. 2012. Analytic theory of Titan's Schumann resonance: Constraints on ionospheric conductivity and buried water ocean. Icarus 218, 1028-1042.

Behounkova, M., Soucek, O., Hron, J. and Cadek O. 2017. Plume activity and tidal deformation on Enceladus influenced by faults and variable ice shell thickness, Astrobiology, 17, doi:10.1089/ast.2016.1629.

Beuthe, M., Rivoldini, A., Trinh, A. 2016. Enceladus's and Dione's floating ice shells supported by minimum stress isostasy. Geophysical Research Letters 43, 10.

Bèzard, B. 2014. The methane mole fraction in Titan’s stratosphere from DISR measurements during the Huygens probe’s descent. Icarus 242, 64-73.

Birch, S.P.D., and 20 colleagues 2017. Geomorphologic mapping of titan's polar terrains: Constraining surface processes and landscape evolution. Icarus 282, 214-236.

Burr, D.M., Emery, J.P., Lorenz, R.D., Collins, G.C., Carling, P.A. 2006. Sediment transport by liquid surficial flow: Application to Titan. Icarus 181, 235-242.

Burr, D.M., Drummond, S.A., Cartwright, R., Black, B.A., Perron, J.T. 2013. Morphology of fluvial networks on Titan: Evidence for structural control. Icarus 226, 742-759.

Čadek, O., and 10 colleagues 2016. Enceladus's internal ocean and ice shell constrained from Cassini gravity, shape, and libration data. Geophysical Research Letters 43, 5653-5660.

Clark, R.N., Curchin, J.M., Barnes, J.W., Jaumann, R., Soderblom, L., Cruikshank, D.P., Brown, R.H., Rodriguez, S., Lunine, J., Stephan, K. and Hoefen, T.M., 2010. Detection

961 and mapping of hydrocarbon deposits on Titan. Journal of Geophysical Research: Planets,
962 115(E10).
963 Coates, A.J., Crary, F.J., Lewis, G.R., Young, D.T., Waite, J.H., Sittler, E.C. 2007. Discovery
964 of heavy negative ions in Titan's ionosphere. Geophysical Research Letters 34, L22103.
965 Coates, A.J., Jones, G.H., Lewis, G.R., Wellbrock, A., Young, D.T., Crary, F.J., Johnson,
966 R.E., Cassidy, T.A., Hill, T.W. 2010. Negative ions in the Enceladus plume. Icarus 206,
967 618-622.
968 Cook-Hallett, C., Barnes, J.W., Kattenhorn, S.A., Hurford, T., Radebaugh, J., Stiles, B.,
969 Beuthe, M. 2015. Global contraction/expansion and polar lithospheric thinning on Titan
970 from patterns of tectonism. Journal of Geophysical Research (Planets) 120, 1220-1236.
971 Cordier, D., Mousis, O., Lunine, J.I., Lebonnois, S., Lavvas, P., Lobo, L.Q., Ferreira, A.G.M.
972 2010. About the Possible Role of Hydrocarbon Lakes in the Origin of Titan's Noble Gas
973 Atmospheric Depletion. The Astrophysical Journal 721, L117-L120.
974 Cornet, T., Cordier, D., Bahers, T.L., Bourgeois, O., Fleurant, C., Mouelic, S.L., Altobelli, N.
975 2015. Dissolution on Titan and on Earth: Toward the age of Titan's karstic landscapes.
976 Journal of Geophysical Research (Planets) 120, 1044-1074.
977 Coustenis, A. and 114 co-authors, TanDEM: Titan and Enceladus Mission, Experimental
978 Astronomy, 23, 893-946, 2009
979 Coustenis, A., Lunine, J., Lebreton, J.-P., Matson, D., Erd, Ch., Reh, K., Beauchamp, P.,
980 Lorenz, R., Waite, H., Sotin, Ch., Gurvits, L., Hirtizg, M., 2009. Earth-based support for
981 the Titan Saturn Mission. Earth Moon and Planets 105, 135-142, DOI: 10.1007/s11038-
982 009-9308-9.
983 Coustenis, A., Bampasidis, G., Achterbergh, R. K., Lavvas, P., Nixon, C. A., Jennings, D. E.,
984 Teanby, N. A., Vinatier, S., Flasar, F. M., Carlson, R. C., Orton, G., Romani, P. N.,

985 Guardique, E. A., 2013. Evolution of the stratospheric temperature and chemical
986 composition over one Titanian year. Astrophys. J. 799, 177, 9p.
987 Coustenis, A., Jennings, D. E., Achterbergh, R. K., Bampasidis, G., Lavvas, P., Nixon, C. A.,
988 Teanby, N. A., Anderson, C. M., Flasar, F. M., 2016. Titan's temporal evolution in
989 stratospheric trace gases near the poles. Icarus 270, 409-420.
990 Crary, F.J., Magee, B.A., Mandt, K., Waite, J.H., Westlake, J., Young, D.T. 2009. Heavy
991 ions, temperatures and winds in Titan's ionosphere: Combined Cassini CAPS and INMS
992 observations. Planetary and Space Science 57, 1847-1856.
993 Davies, A.G., Sotin, C., Choukroun, M., Matson, D.L., Johnson, T.V. 2016. Cryolava flow
994 destabilization of crustal methane clathrate hydrate on Titan. Icarus 274, 23-32.
995 Dougherty, M.K., Khurana, K.K., Neubauer, F.M., Russell, C.T., Saur, J., Leisner, J.S.,
996 Burton, M.E. 2006. Identification of a Dynamic Atmosphere at Enceladus with the Cassini
997 Magnetometer. Science 311, 1406–1409.
998 Dougherty, M.K., Esposito, L.W., Krimigis, S.M. 2009. Saturn from Cassini-Huygens. Saturn
999 from Cassini-Huygens .
1000 Doyon, R., Hutchings, J., Beaulieu, M., Albert, L., Lafrenière, D., Willott, C., Touahri, D.,
1001 Rowlands, N., Maszkiewicz, M., Fullerton, A.W. and Volk, K. 2012. The JWST fine
1002 guidance sensor (FGS) and near-infrared imager and slitless spectrograph (NIRISS). In
1003 Proc. SPIE (Vol. 8442, p. 84422R).
1004 Federal Aviation Administration (FAA) Commercial Space Transportation (AST) and the
1005 Commercial Space Transportation Advisory Committee (COMSTAC) 2015 Commercial
1006 Space Transportation Forecast April 2015
1007 Glein, C.R., Zolotov, M.Y., Shock, E.L. 2008. The oxidation state of hydrothermal systems
1008 on early Enceladus. Icarus 197, 157-163.

ACCEPTED MANUSCRIPT

1009 Glein, C.R., Baross, J.A., Waite, J.H. 2015. The pH of Enceladus' ocean. Geochimica et
1010 Cosmochimica Acta 162, 202-219.
1011 Goguen, J.D., and 12 colleagues 2013. The temperature and width of an active fissure on
1012 Enceladus measured with Cassini VIMS during the 14 April 2012 South Pole flyover.
1013 Icarus 226, 1128-1137.
1014 Griffith, C.A., Zahnle, K. 1995. Influx of cometary volatiles to planetary moons: The
1015 atmospheres of 1000 possible Titans. Journal of Geophysical Research 100, 16907-16922.
1016 Gudipati, M. S., Jacovi, R., Couturier-Tamburelli, I., Lignell, A., Allen, M. 2013.
1017 Photochemical activity of Titan’s low-altitude condensed haze. Nature Communications.
1018 4:1648, DOI: 10.1038/ncomms2649.
1019 Hansen, C.J., Esposito, L., Stewart, A.I.F., Colwell, J., Hendrix, A., Pryor, W., Shemansky,
1020 D., West, R. 2006. Enceladus' Water Vapor Plume. Science 311, 1422-1425.
1021 Hansen, C.J., Esposito, L.W., Stewart, A.I.F., Meinke, B., Wallis, B., Colwell, J.E., Hendrix,
1022 A.R., Larsen, K., Pryor, W., Tian, F. 2008. Water vapour jets inside the plume of gas
1023 leaving Enceladus. Nature 456, 477-479.
1024 Hansen, C.J., and 10 colleagues 2011. The composition and structure of the Enceladus plume.
1025 Geophysical Research Letters 38, L11202.
1026 Hayes, A., and 13 colleagues 2008. Hydrocarbon lakes on Titan: Distribution and interaction
1027 with a porous regolith. Geophysical Research Letters 35, L09204.
1028 Hayes, A.G., and 11 colleagues 2010. Bathymetry and absorptivity of Titan's Ontario Lacus.
1029 Journal of Geophysical Research (Planets) 115, E09009.
1030 Hayes, A.G., and 14 colleagues 2011. Transient surface liquid in Titan's polar regions from
1031 Cassini. Icarus 211, 655-671.

1032 Hedman, M.M., Nicholson, P.D., Showalter, M.R., Brown, R.H., Buratti, B.J., Clark, R.N.
1033 2009. Spectral Observations of the Enceladus Plume with Cassini-Vims. The
1034 Astrophysical Journal 693, 1749-1762.
1035 Hofer, L., and 10 colleagues 2015. Prototype of the gas chromatograph-mass spectrometer to
1036 investigate volatile species in the lunar soil for the Luna-Resurs mission. Planetary and
1037 Space Science 111, 126-133.
1038 Hörst, S.M., Vuitton, V., Yelle, R.V. 2008. Origin of oxygen species in Titan's atmosphere.
1039 Journal of Geophysical Research (Planets) 113, E10006.
1040 Hörst, S.M., and 12 colleagues 2012. Formation of Amino Acids and Nucleotide Bases in a
1041 Titan Atmosphere Simulation Experiment. Astrobiology 12, 809-817.
1042 Howett, C.J.A., Spencer, J.R., Pearl, J., Segura, M. 2011. High heat flow from Enceladus'
1043 south polar region measured using 10-600 cm-1 Cassini/CIRS data. Journal of Geophysical
1044 Research (Planets) 116, E03003.
1045 Hsu, H.-W., Postberg, F., Kempf, S., Trieloff, M., Burton, M., Roy, M., Moragas-
1046 Klostermeyer, G., Srama, R. 2011. Stream particles as the probe of the dust-plasma-
1047 magnetosphere interaction at Saturn. Journal of Geophysical Research (Space Physics)
1048 116, A09215.
1049 Hsu, H.-W., and 14 colleagues 2015. Ongoing hydrothermal activities within Enceladus.
1050 Nature 519, 207-210.
1051 Hurford, T.A., Bills, B.G., Helfenstein, P., Greenberg, R., Hoppa, G.V., Hamilton, D.P. 2009.
1052 Geological implications of a physical libration on Enceladus. Icarus 203, 541-552.
1053 Iess, L., Rappaport, N.J., Jacobson, R.A., Racioppa, P., Stevenson, D.J., Tortora, P.,
1054 Armstrong, J.W., Asmar, S.W. 2010. Gravity Field, Shape, and Moment of Inertia of
1055 Titan. Science 327, 1367.

1056 Iess, L., Jacobson, R.A., Ducci, M., Stevenson, D.J., Lunine, J.I., Armstrong, J.W., Asmar,
1057 S.W., Racioppa, P., Rappaport, N.J., Tortora, P. 2012. The Tides of Titan. Science 337,
1058 457.
1059 Iess, L., and 10 colleagues 2014. The Gravity Field and Interior Structure of Enceladus.
1060 Science 344, 78-80.
1061 Ingersoll, A.P., Pankine, A.A. 2010. Subsurface heat transfer on Enceladus: Conditions under
1062 which melting occurs. Icarus 206, 594-607.
1063 Ingersoll, A.P., Ewald, S.P. 2011. Total particulate mass in Enceladus plumes and mass of
1064 Saturn's E ring inferred from Cassini ISS images. Icarus 216, 492-506.
1065 Israël, G., and 21 colleagues 2005. Complex organic matter in Titan's atmospheric aerosols
1066 from in situ pyrolysis and analysis. Nature 438, 796-799.
1067 Jennings, D.E., Romani, P.N., Bjoraker, G.L., Sada, P.V., Nixon, C.A., Lunsford, A.W.,
1068 Boyle, R.J., Hesman, B.E., McCabe, G.H. 2009. 12C/13C Ratio in Ethane on Titan and
1069 Implications for Methane's Replenishment. Journal of Physical Chemistry A 113, 11101-
1070 11106.
1071 Kite, E.S., Rubin, A.M. 2016. Sustained eruptions on Enceladus explained by turbulent
1072 dissipation in tiger stripes. Proceedings of the National Academy of Science 113, 3972-
1073 3975.
1074 Kliore, A.J., Nagy, A.F., Cravens, T.E., Richard, M.S., Rymer, A.M. 2011. Unusual electron
1075 density profiles observed by Cassini radio occultations in Titan's ionosphere: Effects of
1076 enhanced magnetospheric electron precipitation? Journal of Geophysical Research: Space
1077 Physics, 116, A11318.
1078 Kliore, A.J. et al. 2008. First results from the Cassini radio occultations of the Titan
1079 ionosphere. Journal of Geophysical Research: Space Physics 113 (9), A09317

1080 Lammer, H., and 16 colleagues 2009. What makes a planet habitable?. Astronomy and
1081 Astrophysics Review 17, 181-249.
1082 Lancaster, N. 1989. The Namib Sand Sea: Dune Forms, Processes, and Sediments. A.A.
1083 Balkema, Rotterdam, 200 pp.
1084 Leary, J., Jones, C., Lorenz, R., Strain, R. D., and Waite, J. H., Titan Explorer Flagship
1085 Mission Study, Johns Hopkins Univ., Applied Physics Lab., Laurel, MD, Aug. 2007; also
1086 http://www.lpi.usra.edu/opag/Titan_Explorer_Public_Report.pdf, Jan. 2008 [retrieved
1087 2015].
1088 Lemmon, M.T., Karkoschka, E. and Tomasko, M. 1993, Titan's Rotaion: Surface Features
1089 Observed. Icarus, 103, pp. 329-332.
1090 Liu, Z.Y.C., Radebaugh, J., Harris, R.A., Christiansen, E.H., Neish, C.D., Kirk, R.L., Lorenz,
1091 R.D. 2016. The tectonics of Titan: Global structural mapping from Cassini RADAR. Icarus
1092 270, 14-29.
1093 Loose, M., Beletic, J., Garnett, J. and Muradian, N., 2006, June. Space qualification and
1094 performance results of the SIDECAR ASIC. In SPIE Astronomical Telescopes+
1095 Instrumentation (pp. 62652J-62652J). International Society for Optics and Photonics.
1096 Lopes, R.M.C., and 15 colleagues 2007a. The Lakes and Seas of Titan. EOS Transactions 88,
1097 569-570.
1098 Lopes, R.M.C., and 43 colleagues 2007b. Cryovolcanic features on Titan's surface as revealed
1099 by the Cassini Titan Radar Mapper. Icarus 186, 395-412.
1100 Lopes, R.M.C., and 15 colleagues 2013. Cryovolcanism on Titan: New results from Cassini
1101 RADAR and VIMS. Journal of Geophysical Research (Planets) 118, 416-435.
1102 Lopes, R.M.C., Malaska, M.J., Solomonidou, A., LeGall, A., Janssen, M.A., Neish, C.,
1103 Turtle, E.P., Birch, S.P.D., Hayes, A.G., Radebaugh, J., Coustenis, A., Schoenfeld, A.,
1104 Stiles, B.W., Kirk, R.L., Mitchell, K.L., Stofan, E.R., Lawrence, K.J., and the Cassini

RADAR Team 2016. Nature, Distribution, and Origin of Titan's Undifferentiated Plains ("Blandlands"). Icarus, 270, 162-182.

Lorenz, R.D., and 39 colleagues 2006. The Sand Seas of Titan: Cassini RADAR Observations of Longitudinal Dunes. Science 312, 724-727.

Lorenz, R. D. 2000. Post-Cassini Exploration of Titan: Science Rationale and Mission Concepts, Journal of the British Interplanetary Society, Vol. 53, pp. 218–234.

Lorenz, R. D. 2009. A Review of Titan Mission Studies, Journal of the British Interplanetary Society, 62, 162-174.

Lunine, J.I. 2017. Ocean worlds exploration. *Acta Astronautica* 131, 123-130.

Maciaszek, T., Ealet, A., Jahnke, K., Prieto, E., Barbier, R., Mellier, Y., Costille, A., Ducret, F., Fabron, C., Gimenez, J.L. and Grange, R., 2014, August. Euclid near infrared spectrophotometer instrument concept and first test results at the end of phase B. In SPIE Astronomical Telescopes+ Instrumentation (pp. 91430K-91430K). International Society for Optics and Photonics.

MacKenzie, S.M., and 10 colleagues 2014. Evidence of Titan's climate history from evaporite distribution. Icarus 243, 191-207.

Malaska, M.J., Lopes, R.M., Hayes, A.G., Radebaugh, J., Lorenz, R.D., Turtle, E.P. 2016. Material transport map of Titan: The fate of dunes. Icarus 270, 183-196.

Mandt, K.E., and 18 colleagues 2012. Ion densities and composition of Titan's upper atmosphere derived from the Cassini Ion Neutral Mass Spectrometer: Analysis methods and comparison of measured ion densities to photochemical model simulations. Journal of Geophysical Research (Planets) 117, E10006.

McKay, C.P., Porco, C., Altheide, T., Davis, W.L., Kral, T.A. 2008. The Possible Origin and Persistence of Life on Enceladus and Detection of Biomarkers in the Plume. Astrobiology 8, 909-919.

McKay, C.P. 2016. Titan as the Abode of Life. Life, 6, 8.

McKinnon, W.B. 2015. Effect of Enceladus's rapid synchronous spin on interpretation of Cassini gravity. Geophysical Research Letters 42, 2137-2143.

Mitri, G., Showman, A.P., Lunine, J.I., Lorenz, R.D. 2007. Hydrocarbon lakes on Titan. Icarus 186, 385-394.

Mitri, G., Bland, M.T., Showman, A.P., Radebaugh, J., Stiles, B., Lopes, R.M.C., Lunine, J.I., Pappalardo, R.T. 2010. Mountains on Titan: Modeling and observations. Journal of Geophysical Research (Planets) 115, E10002.

Mitri, G., and 16 colleagues 2014a. The exploration of Titan with an orbiter and a lake probe. Planetary and Space Science 104, 78-92.

Mitri, G., Meriggiola, R., Hayes, A., Lefevre, A., Tobie, G., Genova, A., Lunine, J.I., Zebker, H. 2014b. Shape, topography, gravity anomalies and tidal deformation of Titan. Icarus 236, 169-177.

Moore, J.M., Howard, A.D. 2010. Are the basins of Titan's Hotei Regio and Tui Regio sites of former low latitude seas?. Geophysical Research Letters 37, L22205.

Mousis, O., Schmitt, B. 2008. Sequestration of Ethane in the Cryovolcanic Subsurface of Titan. The Astrophysical Journal 677, L67.

Mousis, O., and 10 colleagues 2009. Clathration of Volatiles in the Solar Nebula and Implications for the Origin of Titan's Atmosphere. The Astrophysical Journal 691, 1780-1786.

Mousis, O., Lunine, J.I., Picaud, S., Cordier, D., Waite, J.H., Jr., Mandt, K.E. 2011. Removal of Titan's Atmospheric Noble Gases by Their Sequestration in Surface Clathrates. The Astrophysical Journal 740, L9.

Neish, C.D., Lorenz, R.D. 2012. Titan’s global crater population: A new assessment. Planetary and Space Science 60, 26-33.

1155 Neish, C.D., and 14 colleagues 2015. Spectral properties of Titan's impact craters imply
1156 chemical weathering of its surface. Geophysical Research Letters 42, 3746-3754.
1157 Neish, C.D., Molaro, J.L., Lora, J.M., Howard, A.D., Kirk, R.L., Schenk, P., Bray, V.J.,
1158 Lorenz, R.D. 2016. Fluvial erosion as a mechanism for crater modification on Titan. Icarus
1159 270, 114-129.
1160 Niemann, H.B., and 17 colleagues 2005. The abundances of constituents of Titan's
1161 atmosphere from the GCMS instrument on the Huygens probe. Nature 438, 779-784.
1162 Niemann, H.B., Atreya, S.K., Demick, J.E., Gautier, D., Haberman, J.A., Harpold, D.N.,
1163 Kasprzak, W.T., Lunine, J.I., Owen, T.C., Raulin, F. 2010. Composition of Titan's lower
1164 atmosphere and simple surface volatiles as measured by the Cassini-Huygens probe gas
1165 chromatograph mass spectrometer experiment. Journal of Geophysical Research (Planets)
1166 115, E12006.
1167 Nimmo, F., Pappalardo, R.T. 2016. Ocean worlds in the outer solar system, J. Geophys. Res.,
1168 121,1378-1399.
1169 Nixon, C.A., and 12 colleagues 2012. Isotopic Ratios in Titan's Methane: Measurements and
1170 Modeling. The Astrophysical Journal 749, 159.
1171 Owen, T., Niemann, H.B. 2009. The origin of Titan's atmosphere: some recent advances.
1172 Philosophical Transactions of the Royal Society of London Series A 367, 607-615.
1173 Porco, C.C., and 24 colleagues 2006. Cassini Observes the Active South Pole of Enceladus.
1174 Science 311, 1393-1401.
1175 Porco, C., DiNino, D., Nimmo, F. 2014. How the Geysers, Tidal Stresses, and Thermal
1176 Emission across the South Polar Terrain of Enceladus are Related. The Astronomical
1177 Journal 148, 45.

1178 Postberg, F., Kempf, S., Hillier, J.K., Srama, R., Green, S.F., McBride, N., Grun, E. 2008.
1179 The E-ring in the vicinity of Enceladus. II. Probing the moon's interior. The composition of
1180 E-ring particles. Icarus 193, 438-454.
1181 Postberg, F., Kempf, S., Rost, D., Stephan, T., Srama, R., Trieloff, M., Mocker, A., Goerlich,
1182 M. 2009. Discriminating contamination from particle components in spectra of Cassini's
1183 dust detector CDA. Planetary and Space Science 57, 1359-1374.
1184 Postberg, F., Kempf, S., Schmidt, J., Brilliantov, N., Beinsen, A., Abel, B., Buck, U., Srama,
1185 R. 2009. Sodium salts in E-ring ice grains from an ocean below the surface of Enceladus.
1186 Nature 459, 1098-1101.
1187 Postberg, F., Schmidt, J., Hillier, J., Kempf, S., Srama, R. 2011. A salt-water reservoir as the
1188 source of a compositionally stratified plume on Enceladus. Nature 474, 620-622.
1189 Postberg, F., Khawaja, N., Hsu, H.W., Sekine, Y., Shibuya, T. 2015. Refractory Organic
1190 Compounds in Enceladus' Ice Grains and Hydrothermal Activity. AGU Fall Meeting
1191 Abstracts.
1192 Radebaugh, J., Lorenz, R.D., Kirk, R.L., Lunine, J.I., Stofan, E.R., Lopes, R.M.C., Wall,
1193 S.D., the Cassini Radar Team 2007. Mountains on Titan observed by Cassini Radar. Icarus
1194 192, 77-91.
1195 Radebaugh, J., and 15 colleagues 2008. Dunes on Titan observed by Cassini Radar. Icarus
1196 194, 690-703.
1197 Rappaport, N.J., Iess, L., Wahr, J., Lunine, J.I., Armstrong, J.W., Asmar, S.W., Tortora, P., Di
1198 Benedetto, M., Racioppa, P., 2008. Can Cassini detect a subsurface ocean in Titan from
1199 gravity measurements? Icarus 194, 711–720.
1200 Rauscher, B.J., Figer, D.F., Regan, M.W., Boeker, T., Garnett, J., Hill, R.J., Bagnasco, G.,
1201 Balleza, J., Barney, R., Bergeron, L.E. and Brambora, C., 2004, October. Detectors for the
1202 James Webb Space Telescope near-infrared spectrograph. In SPIE Astronomical

Telescopes+ Instrumentation (pp. 710-726). International Society for Optics and Photonics.

Reh, K., Spilker, L., Lunine, J.I., Waite, J.H., Cable, M.L., Postberg, F. and Clark, K., 2016, March. Enceladus Life Finder: The search for life in a habitable Moon. In Aerospace Conference, 2016 IEEE (pp. 1-8)

Rieke, M.J., Kelly, D. and Horner, S., 2005, August. Overview of James Webb Space Telescope and NIRCam's Role. In Optics & Photonics 2005 (pp. 590401-590401). International Society for Optics and Photonics.

Roth, L., Saur, J., Retherford, K.D., Strobel, D.F., Feldman, P.D., McGrath, M.A., Nimmo, F. 2014. Transient Water Vapor at Europa's South Pole. Science 343, 171-174.

Scherer, S., K. Altwegg, H. Balsiger, J. Fischer, A. Jäckel, A. Korth, M. Mildner, D. Piazza, H. Rème, and P. Wurz, A novel principle for an ion mirror design in time-of-flight mass spectrometry, Int. Jou. Mass Spectr. 251 (2006) 73–81.

Schinder, P.J. et al. 2015. A numerical technique for two-way radio occultations by oblate axisymmetric atmospheres with zonal winds. Radio Science 50 (7), pp. 712-727.

Schinder, P.J. et al. 2012. The structure of Titan's atmosphere from Cassini radio occultations: Occultations from the Prime and Equinox missions. Icarus, 221, 1020-1031.

Schmidt, J., Brilliantov, N., Spahn, F., Kempf, S. 2008. Slow dust in Enceladus' plume from condensation and wall collisions in tiger stripe fractures. Nature 451, 685-688.

Schubert, G., Anderson, J.D., Travis, B.J., Palguta, J. 2007. Enceladus: Present internal structure and differentiation by early and long-term radiogenic heating. Icarus 188, 345-355.

Senft, L.E., Stewart, S.T. 2011. Modeling the morphological diversity of impact craters on icy satellites. Icarus 214, 67-81.

1227 Serigano, J., Nixon, C.A., Cordiner, M.A., Irwin, P.G.J., Teanby, N.A., Charnley, S.B.,
1228 Lindberg, J.E. 2016. Isotopic Ratios of Carbon and Oxygen in Titan's CO using ALMA.
1229 The Astrophysical Journal 821, L8.
1230 Soderblom, J.M., Brown, R.H., Soderblom, L.A., Barnes, J.W., Jaumann, R., Le Mouélic, S.,
1231 Sotin, C., Stephan, K., Baines, K.H. and Buratti, B.J., Clark, R.N., Nicholson, P.D., 2010.
1232 Geology of the Selk crater region on Titan from Cassini VIMS observations. Icarus,
1233 208(2), pp.905-912.
1234 Soderblom, J.M., Barnes, J.W., Soderblom, L.A., Brown, R.H., Griffith, C.A., Nicholson,
1235 P.D., Stephan, K., Jaumann, R., Sotin, C., Baines, K.H. and Buratti, B.J., 2012. Modeling
1236 specular reflections from hydrocarbon lakes on Titan. Icarus, 220(2), pp.744-751.
1237 Soderblom, L.A., Becker, T.L., Kieffer, S.W., Brown, R.H., Hansen, C.J., Johnson, T.V.,
1238 Kirk, R.L., Shoemaker, E.M., Cook, A.F. 1990. Triton's geyser-like plumes - Discovery
1239 and basic characterization. Science 250, 410-415.
1240 Soderblom, L.A., and 26 colleagues 2007. Correlations between Cassini VIMS spectra and
1241 RADAR SAR images: Implications for Titan's surface composition and the character of
1242 the Huygens Probe Landing Site. Planetary and Space Science 55, 2025-2036.
1243 Solomonidou, A., Hirtzig, M., Coustenis, A., Bratsolis, E., Le Mouélic, S., Rodriguez, S.,
1244 Stephan, K., Drossart, P., Sotin, C., Jaumann, R., Brown, R.H., Kyriakopoulos, K., Lopes,
1245 R.M.C., Bampasidis, G., Stamatelopoulou-Seymour, K., Moussas, X. (2014). Surface
1246 albedo spectral properties of geologically interesting areas on Titan. J. of Geophys. Res.,
1247 119, 1729-1747.
1248 Solomonidou, A., A. Coustenis, M. Hirtzig, S. Rodriguez, K. Stephan, R.M.C. Lopes, P.
1249 Drossart, C. Sotin, S. Le Mouélic, K. Lawrence, E. Bratsolis, R. Jaumann and R.H. Brown,
1250 (2016). Temporal variations of Titan's surface with Cassini/VIMS, Icarus 270, 85-99.

1251 Sotin, C. et al., 2005. Release of volatiles from a possible cryovolcano from near-infrared
1252 imaging of Titan. Nature, 435, 786-798.
1253 Sotin, C., Mitri, G., Rappaport, N., Schubert, G., Stevenson, D. 2010. Titan's Interior
1254 Structure. Titan from Cassini-Huygens 61.
1255 Spahn, F., and 15 colleagues 2006. Cassini Dust Measurements at Enceladus and Implications
1256 for the Origin of the E Ring. Science 311, 1416-1418.
1257 Sparks, W. B., Hand, K. P. , McGrath, M. A., Bergeron, E., Cracraft, M., and Deustua, S. E.,.
1258 2016. Probing for Evidence of Plumes on Europa with HST/STIS. Ap. J. 829:121.
1259 Sparks, W.B., Schmidt, McGrath, M. A., Hand, K. P., Spencer, J. R., Cracraft, M., and
1260 Deustua, S. E, 2017. Active Cryovolcanism on Europa?, Ap. J. Lett., 839:L18.
1261 Srama, R., and 43 colleagues 2004. The Cassini Cosmic Dust Analyzer. Space Science
1262 Reviews 114, 465–518
1263 Srama, R., and 17 colleagues 2015. Enceladus Icy Jet Analyzer (ENIJA): Search for life with
1264 a high resolution TOF-MS for in situ characterization of high dust density regions.
1265 European Planetary Science Congress 2015, EPSC2015-769.
1266 Stofan, E.R., and 37 colleagues 2007. The lakes of Titan. Nature 445, 61-64.
1267 Stofan, E., Lorenz, R., Lunine, J., Bierhaus, E., Clark, B., Mahaffy, P., and Ravine, M.,
1268 TiME—The Titan Mare Explorer, IEEE Aerospace Conference, IEEE Publ., Piscataway,
1269 NJ, March 2013, pp. 1–10.
1270 Strange, N., Spilker, T., Landau, D., Lam, T., Lyons, D. and Guzman, J., 2009. Mission
1271 Design for the Titan Saturn System Mission Concept. Advances in the Astronautical
1272 Sciences, 135(2), pp.919-934.
1273 Thomas, P.C., Tajeddine, R., Tiscareno, M.S., Burns, J.A., Joseph, J., Loredo, T.J.,
1274 Helfenstein, P., Porco, C. 2016. Enceladus's measured physical libration requires a global
1275 subsurface ocean. Icarus 264, 37-47.

1276 Thompson, W.R., Sagan, C. 1992. Organic chemistry on Titan: Surface interactions.
1277 Symposium on Titan 338.
1278 Tobie, G., Teanby, N.A., Coustenis, A., Jaumann, R., Raulin, F., Schmidt, J., Carrasco, N.,
1279 Coates, A.J., Cordier, D., De Kok, R. and Geppert, W.D., 2014. Science goals and mission
1280 concept for the future exploration of Titan and Enceladus. Planetary and Space Science,
1281 104, pp.59-77.
1282 Tobie, G., Lunine, J.I., Sotin, C. 2006. Episodic outgassing as the origin of atmospheric
1283 methane on Titan. Nature 440, 61-64.
1284 Tobie, G., Gautier, D., Hersant, F. 2012. Titan's Bulk Composition Constrained by Cassini-
1285 Huygens: Implication for Internal Outgassing. The Astrophysical Journal 752, 125.
1286 Tomasko, M.G., and 39 colleagues 2005. Rain, winds and haze during the Huygens probe's
1287 descent to Titan's surface. Nature 438, 765-778.
1288 Tortora, P., Zannoni, M., Hemingway, D., Nimmo, F., Jacobson, R.A., Iess, L., Parisi, M.
1289 2016. Rhea gravity field and interior modeling from Cassini data analysis. Icarus 264, 264-
1290 273.
1291 Turtle, E.P., and 13 colleagues 2011. Rapid and Extensive Surface Changes Near Titan’s
1292 Equator: Evidence of April Showers. Science 331, 1414.
1293 Vuitton, V., Yelle, R.V., McEwan, M.J. 2007. Ion chemistry and N-containing molecules in
1294 Titan's upper atmosphere. Icarus 191, 722-742.
1295 Yelle, R.V., Cui, J., Müller-Wodarg, I.C.F. 2008. Methane escape from Titan's atmosphere.
1296 Journal of Geophysical Research (Planets) 113, E10003.
1297 Yung, Y.L., Allen, M., Pinto, J.P. 1984. Photochemistry of the atmosphere of Titan -
1298 Comparison between model and observations. The Astrophysical Journal Supplement
1299 Series 55, 465-506.

1300 Waite, J.H., and 13 colleagues 2006. Cassini Ion and Neutral Mass Spectrometer: Enceladus
1301 Plume Composition and Structure. Science 311, 1419-1422.
1302 Waite, J.H., Young, D.T., Cravens, T.E., Coates, A.J., Crary, F.J., Magee, B., Westlake, J.
1303 2007. The Process of Tholin Formation in Titan's Upper Atmosphere. Science 316, 870.
1304 Waite, J.H., Jr., and 15 colleagues 2009. Liquid water on Enceladus from observations of
1305 ammonia and ^{40}Ar in the plume. Nature 460, 487-490.
1306 Waite, J. H. et al. 2006. Cassini ion and neutral mass spectrometer: Enceladus plume
1307 composition and structure. Science, 311(5766), 1419-1422.
1308 Waite, J.H., and 12 colleagues 2017. Cassini finds molecular hydrogen in the Enceladus
1309 plume: Evidence for hydrothermal processes. Science, 356 (6334), 155-159.
1310 Wilson, E.H., Atreya, S.K. 2009. Titan's Carbon Budget and the Case of the Missing Ethane.
1311 Journal of Physical Chemistry A 113, 11221-11226.
1312 Wurz, P., Abplanalp, D., Tulej, M., Lammer, H. 2012. A neutral gas mass spectrometer for
1313 the investigation of lunar volatiles. Planetary and Space Science 74, 264-269.
1314 Wurz, P., and 18 colleagues 2015. Solar wind sputtering of dust on the surface of 67P
1315 Churyumov-Gerasimenko. Astronomy and Astrophysics 583, A22.
1316 Zahnle, K., Pollack, J.B., Grinspoon, D., Dones, L. 1992. Impact-generated atmospheres over
1317 Titan, Ganymede, and Callisto. Icarus 95, 1-23.
1318 Zahnle, K.J., Korycansky, D.G., Nixon, C.A. 2014. Transient climate effects of large impacts
1319 on Titan. Icarus 229, 378-391.

ACCEPTED MANUSCRIPT

ACCEPTED MANUSCRIPT

1320 TABLES AND FIGURES

ACCEPTED MANUSCRIPT

ACCEPTED MANUSCRIPT

Table 1. Scientific objectives and questions, and measurement requirements (payload, instrument parameters required and performance) of E^2T.

GOAL A: ORIGIN AND EVOLUTION OF VOLATILE-RICH OCEAN WORLDS, ENCELADUS AND TITAN						
Science Objectives	**Scientific Questions**	**Measurement Description**	**Measurement Requirements**			
			Payload	**Parameter**	**Requirement**	**Performance**
A.1: Are Enceladus' volatile compounds primordial or have they been re-processed and if so, to what extent?	Are major volatiles in the plume material primordial (as opposed to geochemical or biological) and, if so, how were they delivered?	Quantify the abundance of radiogenic and non-radiogenic noble gases (40/38/36 Ar, 4/3He, 20/22Ne, 128-136Xe) from mass spectra of plume vapour phases	INMS	Dynamic range Resolution Sensitivity	6/8 decades 5000 m/Δm 10^1 #/cm^3	6/10 decades 5000 m/Δm 10^0 #/cm^3
		Determine the extent of molecular complexity in the plume up to 1000 amu	INMS	Dynamic range Resolution Sensitivity	6/8 decades >300 m/Δm 10^1 #/cm^3	6/10 decades 5000 m/Δm 10^0 #/cm^3
		Measure isotopic ratios (^{12}C/^{13}C, D/H, ^{14}N/^{15}N, ^{16}O/^{18}O) in major species from mass spectra of plume vapour, including allowing the separation of species with isobaric interferences for example, ^{12}CH from ^{13}C	INMS	Dynamic range Resolution Sensitivity	6/8 decades >2500-3000 m/Δm 10^3 #/cm^3	6/8 decades 5000 m/Δm 10^0 #/cm^3
		Measure isotopic ratio ^{12}C/^{13}C from mass spectra of condensed organics in icy grains	ENIJA	Dynamic range Mass range Resolution Sensitivity	3 decades 12–100m/z 100 m/Δm 1 ppm	6/8 decades 1–2000m/z 1000m/Δm 0.1 ppm
		Determine the abundance ratio between major C-, N-, O-, S-bearing compounds from mass spectra of plume vapour and solid phases, including allowing the separation of vapour-phase species with isobaric interferences for example, CO from N_2	INMS	Dynamic range Resolution Sensitivity	6/8 decades 3000 m/Δm 10^3 #/cm^3	6/10 decades 5000 m/Δm 10^0 #/cm^3
			ENIJA	Dynamic range Mass range Resolution Sensitivity	5 decades 12–200 m/z 200 m/Δm 1 ppm	6/8 decades 1–2000m/z 1000m/Δm 0.1 ppm

ACCEPTED MANUSCRIPT

		Do Enceladus' volatile compounds originate from a unique reservoir and, if so, how is their distribution between solid and vapour phases affected by eruption processes?	Determine the compositional profile of major species through the plume (both volatile and non-volatile) from mass spectra of plume vapour and condensed phases in icy grains	INMS	Dynamic range Resolution Sensitivity	6/8 decades 300 m/Δm 10^3 #/cm^3	6/10 decades 5000 m/Δm 10^0 #/cm^3
				ENIJA	Dynamic range Mass range Resolution Sensitivity Spatial resolution	5 decades 7–500 m/z 500 m/Δm 1 ppm 1000 m	6/8 decades 1–2000m/z 1000m/Δm 0.1 ppm 100 m
	A.2: What is the history and extent of volatile exchange on Titan?	How were Titan's volatiles delivered, and were they internally processed?	Quantify the abundance of noble gases in the atmosphere	INMS	Dynamic range Resolution Sensitivity	6/8 decades 3000–5000 m/Δm 10^3 #/cm^3	6/10 decades 5000 m/Δm 10^0 #/cm^3
			Measure and compare isotopic ratios: $^{12}C/^{13}C$ and D/H in the atmospheric main constituents	INMS	Dynamic range Resolution Sensitivity	6 decades 3000 m/Δm 10^5 #/cm^3	6 decades 5000 m/Δm 10^0 #/cm^3
		What is the age of Titan's methane cycle, and is there evidence for renewal from internal outgassing?	Measure and compare isotopic ratios: ($^{14}N/^{15}N$; $^{12}C/^{13}C$; D/H; $^{16}O/^{18}O$) in the upper atmosphere	INMS	Dynamic range Resolution Sensitivity	6/8 decades 3000 m/Δm 10^3 #/cm^3	6/10 decades 5000 m/Δm 10^0 #/cm^3
			Constrain the genesis of candidate cryovolcanoes and cryovolcanic flows	TIGER	Coverage Scale Wavelength DTM vertical resol.	Sotra, Tui, & Hotei 50 m/pixel 3 colours 20 m	Sotra, Tui, & Hotei 30–50 m/pixel 3 colours 15–20 m
		How does Titan's surface reflect climate change?	Determine whether seas have existed in the tropics and/or south polar region	TIGER	Coverage Scale Wavelength DTM vertical resol.	Regions within Hotei, Tui, Xanadu & the putative south-polar basins 100m/pix 3 colour N/A	Regions within Hotei, Tui, Xanadu & the putative south-polar basins 100m/pix 3 colour N/A
			Investigate fine-scale dune patterns and how they reorient or breakup	TIGER	Coverage Scale Wavelength DTM vertical resol.	4 dune regions 50m/pix 3 colour N/A	4 dune regions 30–50m/pix 3 colour N/A

	A.3: How has Titan's organic-rich surface evolved?	How have liquid organics modified Titan's surface?	Investigate the morphology of river networks and channels	TIGER	Coverage Scale Wavelength DTM vertical resol.	3 drainages 30m/pix 3 colour 15 m	3 drainages 25–30m/pix 3 colour 10–15 m
			Determine whether the labyrinth terrains and sharp edged depressions are formed by dissolution	TIGER	Coverage Scale Wavelength DTM vertical resol.	2 regions 100m/pix 3 colour 40 m	2 regions 50–100m/pix 3 colour 20–40 m
		To what degree are sediments produced and transported by fluvial and eolian processes?	Investigate the margins of the dune fields to identify sources and sinks of sand	TIGER	Coverage Scale Wavelength DTM vertical resol.	4 dune regions 50m/pix 3 colour N/A	4 dune regions 30–50m/pix 3 colour N/A
			Determine the primary erosion processes that degrade impact crater rims	TIGER	Coverage Scale Wavelength DTM vertical resol.	4 impacts 100m/pix 3 colour N/A	4 impacts 50–100m/pix 3 colour N/A
			Investigate the fine-scale geologic processes that modify the plains	TIGER	Coverage Scale Wavelength DTM vertical resol.	3 regions 50m/pix 3 colour N/A	3 regions 30–50m/pix 3 colour N/A
		How do the mechanical and chemical properties (i.e., composition) vary between the observed geologic units on Titan?	Determine how fluvial erosion varies among geologic units	TIGER	Coverage Scale Wavelength DTM vertical resol.	5 regions 50m/pix 3 colour N/A	5 regions 30–50m/pix 3 colour N/A
			Constrain the maximum slopes and topography supported by different geologic units	TIGER	Coverage Scale Wavelength DTM vertical resol.	5 regions 100m/pix 3 colour 40m	5 regions 50–100m/pix 3 colour 20–40m
			Investigate the morphology, topography, and spectral relationships across unit boundaries	TIGER	Coverage Scale Wavelength DTM vertical resol.	5 boundaries 100m/pix 3 colour 40m	5 boundaries 50–100m/pix 3 colour 20–40m

ACCEPTED MANUSCRIPT

GOAL B: HABITABILITY AND POTENTIAL FOR LIFE IN OCEAN WORLDS, ENCELADUS AND TITAN						
Science Objectives	**Scientific Questions**	**Measurement Description**	**Instrument**	**Measurement Requirements**		
				Parameter	**Requirement**	**Performance**
B.1: Is Enceladus' aqueous interior an environment favourable to the emergence of life?	What is the nature of hydrothermal activity on Enceladus and how does it connect with the plume activity?	Detect and inventory reduced and oxidized species in the plume material (e.g., NH_3/N_2 ratio, H_2 abundance, reduced versus oxidized organic species)	INMS	Dynamic range Resolution Sensitivity	6/8 decades 3000m/Δm 10^3 #/cm^3	6/10 decades 5000m/Δm 10^0 #/cm^3
			ENIJA	Dynamic range Mass range Resolution Sensitivity	4 decades 7–200 m/z 200 m/Δm 1 ppm	6/8 decades 1–2000m/z 1000m/Δm 0.1 ppm
		Characterize composition and abundance of salts and other minerals as messengers of rock/water interaction from mass spectra of icy grains	ENIJA	Dynamic range Mass range Resolution Sensitivity	6 decades 7–200 m/z 200 m/Δm 1 ppm	6/8 decades 1–2000m/z 1000m/Δm 0.1 ppm
		Determine spatial variations in plume composition (vapour and icy grains) and correlate with source activity	ENIJA	Dynamic range Mass resolution Resolution Sensitivity Spatial Resolution	4 decades 7–500 m/z 500 m/Δm 1 ppm 1000 m	6/8 decades 1–2000m/z 1000m/Δm 0.1 ppm 100 m
			TIGER	Coverage Scale Wavelength DTM vertical res.	2 cross-stripe tracks 5m/pix 5 & 5.3 µm N/A	2 cross-stripe tracks 2.5–5m/pix 5 & 5.3 µm N/A
		Determine how much energy the jets dissipate	TIGER	Coverage Scale Wavelength DTM vertical res.	2 night-time cross-stripe tracks 5m/pix 5 & 5.3 µm N/A	2 cross-stripe tracks 2.5–5m/pix 5 & 5.3 µm N/A

ACCEPTED MANUSCRIPT

	Is there evidence that biotic/prebiotic processes are operating in Enceladus' interior?	Compare mass distribution of organic molecules to that expected from abiotic synthesis (e.g., Poisson versus non-Poisson) in the mass distribution of organic molecules	INMS	Dynamic range Resolution Sensitivity	6/8 decades 3000m/Δm 10^3 #/cm^3	6/10 decades 5000m/Δm 10^0 #/cm^3
			ENIJA	Dynamic range Mass range Resolution Sensitivity	6 decades 26-1000 m/z 500 m/Δm 1 ppm	6/8 decades 1-2000m/z 1000m/Δm 0.1 ppm
		Search for abnormal isotopic ratio in organics in plume vapour, including allowing the separation of species with isobaric interferences for example, ^{12}CH from ^{13}C	INMS	Dynamic range Resolution Sensitivity	6/8 decades 2500-3000 m/Δm 10^3 #/cm^3	6/10 decades 5000 m/Δm 10^0 #/cm^3
		Search for amino acids and abnormal ^{12}C/^{13}C ratio in organics that may have been captured in the ice matrix of solid plume material	ENIJA	Dynamic range Mass range Resolution Sensitivity	5 decades 12–500m/z 500 m/Δm 1 ppm	6/8 decades 1–2000m/z 1000m/Δm 0.1 ppm
B.2: To what level of complexity has prebiotic chemistry evolved in the Titan system?	What chemical processes control the formation and evolution of complex organics in Titan's atmosphere?	Measure the elemental chemistry of low mass organic molecules in Titan's atmosphere	INMS	Dynamic range Resolution Sensitivity	6/8 decades 3000m/Δm 10^1 #/cm^3	6/10 decades 5000m/Δm 10^0 #/cm^3
		Measure organic macromolecules and ioni species (cations and anions) in order to monitor chemical coupling processes between ions and neutral species	INMS	Dynamic range Resolution Sensitivity	6/8 decades 3000m/Δm 10^1 #/cm^3	6/10 decades 5000m/Δm 10^0 #/cm^3
		Characterize the structure of haze layers, along with their spatial and temporal variations	TIGER	Coverage Scale	10 temporally spaced full-disk images 1km/pix	10 temporally spaced full-disk images 500m–1km/pix

				Wavelength DTM vertical resol.	3 colours N/A	3 colours N/A
		Determine the cloud distribution and morphology, including estimates of their top altitudes and vertical extent	TIGER	Coverage Scale Wavelength DTM vertical resol.	10 temporally spaced full-disk images 1km/pix 3 colours N/A	10 temporally spaced full-disk images 500m–1km/pix 3 colours N/A
		Investigate the morphology and compositional variability of candidate cryovolcanoes	TIGER	Coverage Scale Wavelength DTM vertical resol.	Sotra, Tui & Hotei 50m/pix 3 colours 20m	Sotra, Tui & Hotei 30–50m/pix 3 colours 15–20m
	To what extent does Titan's organic-rich surface chemically communicate with its water-rich interior?	Investigate the morphology of Titan's mountains to look for evidence of compressional vs extensional tectonics	TIGER	Coverage Scale Wavelength DTM vertical resol.	3 mountains 100 m/pix 3 colours 40 m	3 mountains 50–100 m/pix 3 colours 20–40 m
		Investigate melt sheets and flows associated with impact craters	TIGER	Coverage Scale Wavelength DTM vertical resol.	2 larger fresh craters 50 m/pix 3 colours N/A	2 larger fresh craters 30–50 m/pix 3 colours N/A
		Measure the abundance of radiogenically derived noble gases (e.g., ^{40}Ar, ^{21}Ne) and helium	INMS	Dynamic range Resolution Sensitivity	6/8 decades 3000m/Δm 10^1 #/cm^3	6/10 decades 5000m/Δm 10^0 #/cm^3

76 **Table 2.** Summary of Instrument Characteristics.

Instrument/ Experiment (Proposed Agency)	Mass (kg)	Peak Power (W)	TRL	Science Contribution
INMS (SSO)	6.2	34	6	1) Analysis of elemental, molecular and isotopic composition of neutral and ionic gas phase constituents in a mass-to-charge range of 1–1000 u in Saturn's E-ring, Enceladus' plume and Titan's upper atmosphere 2) Search for spatial variations in composition and correlate with jet sources.
ENIJA (DLR)	6.5	19.2	5–6	1) Analysis of elemental, molecular and isotopic composition of solid phase constituents in a mass-to-charge range of 1–2000 u of Enceladus' plume / E-ring 2) Measure fluxes at high impact rates up to 108 $s^{-1}m^{-2}$ to map the dynamical profile (number density, ejection speeds and size distribution) of Enceladus' ice jets
TIGER (NASA)	45	30	5–6	1) Detailed analysis of the geology of Titan's surface at 30–100 m/pixel and of Enceladus' plume sources at 1 m/pixel 2) Measure of the thermal emission from Enceladus's south polar terrain, at 1 m/pixel
Total	57.7	83.2		

77

78

79

ACCEPTED MANUSCRIPT

80 **Table 3.** Nominal interplanetary trajectory

Departure	
Launch date to GTO	April 2030
Hyperbolic Injection date	27 April 2030
Injection ΔV from upper stage (km/s)	1.20
Injected Mass (kg)	4056
C_3 (km^2/s^2)	9.58
Declination of escape asymptote (deg)	5
Cruise	
Earth flyby date	20 August 2031
Earth flyby altitude (km)	500
Earth flyby V-infinity (km/s)	9.28
S/C mass at Earth flyby	3637
Arrival	
Saturn arrival date	26 April 2036
Flight time (years)	6.0
Arrival V-infinity (km/s)	5.72
Arrival declination (deg)	19.0
SOI ΔV (m/s)	569
Range at SOI (R_S)	1.05
Orbital period (day)	150
Apoapsis range (R_S)	180
S/C mass after SOI (kg)	2753

81

82

ACCEPTED MANUSCRIPT

83 **Table 4.** Launch period performance, fixed arrival date.

Launch Date	Nominal –10 days	Nominal (27Apr2030)	Nominal+10days
C_3 (km^2/s^2)	9.58	9.58	9.58
Xe propellant (kg)	750	747	752
SOI ΔV (m/s)	563	569	571

84

85

ACCEPTED MANUSCRIPT


86 **Table 5.** E^2T spacecraft technical characteristics.

E^2T Spacecraft main technical characteristics		**Heritage**
Spacecraft Dimensions	• Main S/C body: 2974mm × 2860mm × 2589mm with 4 Solar Array wings of an overall area of 160 m^2. • SEPS cylindrical section: 600mm radius × 970mm height with 3+1 QinetiQ T6-Ion Thruster Engine of 7.5 kW each one.	ExoMars BepiColombo
Overall Launch mass	• 4056 kg (including 58 kg of science payload), dry mass 1876 kg	
Propulsion Architecture	The E^2T S/C propulsion architecture is based on the integration of three different propulsion systems: • Solar Electrical Propulsion System (SEPS) based on 3+1 QinetiQ T6 gridded Ion-Thruster engines (1 for redundancy) • Main Bi-propellant propulsion system for S/C manoeuvres purposes • Hydrazine Mono-propellant propulsion system of 16 RCS Thruster (8+8 in hot redundancy) for Attitude and Orbital Control System (AOCS) purposes.	ExoMars BepiColombo
Power Architecture	The E^2T Electrical Propulsion System (EPS) architecture proposed is based on: • 4 Solar Array Wings able to sustain an End of Life (EOL) power demand of 620 W (including margins) with a reference solar constant of 15 W/m^2 at 9.14 AU, with a Sun-Aspect-Angle of 0°. • One 31 kg Li-Ion battery of 3691 Wh total capacity and able to sustain a peak power load of demand of 700 W during the forecast eclipses. • One secondary battery installed on SEPS module for 100 V High-Voltage Sun-regulated power bus stability purposes.	JUICE Rosetta BepiColombo
Baseline payload	• Ion and Neutral Mass Spectrometer (INMS) • Enceladus Icy Jet Analyser (ENIJA) • Titan Imaging and Geology, Enceladus Reconnaissance (TIGER) high-resolution infrared camera	

87

88

ACCEPTED MANUSCRIPT



89

90 **Figure 1.** Cassini SAR mosaic images of the north polar region showing Kraken, Ligeia and Punga

91 Maria (from Mitri et al., 2014a).

92

ACCEPTED MANUSCRIPT



Figure 2. Plume emanating from multiple jets in Enceladus' south polar terrain (NASA/JPL/Space Science Institute).

ACCEPTED MANUSCRIPT





Figure 3. Enceladus' internal structure inferred from gravity, topography and libration measurement provided by Cassini mission. A global subsurface ocean is present under the outer ice shell. The ice shell is believed to be a few kilometers thin at the south polar region where the center of the geological activity is with the formation of the plume formed by multi-jets (NASA/JPL-Caltech).


ACCEPTED MANUSCRIPT



124 **Figure 4.** Composition of salt-poor (Type I and II) and salt-rich (Type III) particles in
125 Saturn's E-ring and Enceladus' plume as measured by Cassini CDA instrument (Postberg et
126 al., 2009).

127

ACCEPTED MANUSCRIPT



140 **Figure 5.** Separation of isobaric interference such as occur at 16, 20, 28, and 40 u/e for the
141 indicated species is possible based on heritage of RTOF/ROSINA from the Rosetta mission
142 (Scherer et al., 2006; Wurz et al., 2015).

143

ACCEPTED MANUSCRIPT




Figure 6. Laser dispersion mass spectrum of aspartic acid (12 ppm), glutamic acid (12 ppm), and arginine (8 ppm) dissolved in a salt-water matrix simulating Enceladus' ocean composition. The complex amino acids are detectable in comparable quantities to glycine (15 ppm, not shown), even though the spectrum has a mass resolution 3 times less than ENIJA's. S/N ratio of this laser dispersion spectrum is comparable to a much lower analysed concentration in ENIJA spectra (≤1 ppm). Most un-annotated mass lines are due to salt-water cluster ions. By co-adding multiple ice grain spectra the S/N ratio can be further improved leading to an ENIJA detection limit of 10–100 ppb for most amino acids in ice grains formed from Enceladus' ocean water.

ACCEPTED MANUSCRIPT




Figure 7. The surface of Titan imaged by DISR camera during the Huygens probe descend. TIGER has the capability to image Titan at Huygens DISR resolution (ESA/NASA/JPL/University of Arizona).

ACCEPTED MANUSCRIPT





Figure 8. Interplanetary transfer to Saturn. Red arrows indicate electric propulsion thrust.

ACCEPTED MANUSCRIPT



191

192 **Figure 9.** Sample tour with two period- and inclination-management Titan flybys followed
193 by a science phase with 6 Enceladus flybys and 17 Titan flybys (Inertial representation).

194

ACCEPTED MANUSCRIPT




Figure 10. E^2T baseline proposed configuration of the spacecraft. Left panel shows an enlarged view of the S/C and right panel shows a close-up view of the spacecraft.

ACCEPTED MANUSCRIPT

GOAL A: ORIGIN AND EVOLUTION OF VOLATILE-RICH OCEAN WORLDS, ENCELADUS AND TITAN						
Science Objectives	**Scientific Questions**	**Measurement Description**	**Measurement Requirements**			
			Payload	**Parameter**	**Requirement**	**Performance**
A.1: Are Enceladus' volatile compounds primordial or have they been re-processed and if so, to what extent?	Are major volatiles in the plume material primordial (as opposed to geochemical or biological) and, if so, how were they delivered?	Quantify the abundance of radiogenic and non-radiogenic noble gases (40/38/36 Ar, 4/3He, 20/22Ne, 128-136Xe) from mass spectra of plume vapour phases	INMS	Dynamic range Resolution Sensitivity	6/8 decades 5000 m/Δm 10^1 #/cm^3	6/10 decades 5000 m/Δm 10^0 #/cm^3
		Determine the extent of molecular complexity in the plume up to 1000 amu	INMS	Dynamic range Resolution Sensitivity	6/8 decades >300 m/Δm 10^1 #/cm^3	6/10 decades 5000 m/Δm 10^0 #/cm^3
		Measure isotopic ratios (^{12}C/^{13}C, D/H, ^{14}N/^{15}N, ^{16}O/^{18}O) in major species from mass spectra of plume vapour, including allowing the separation of species with isobaric interferences for example, ^{12}CH from ^{13}C	INMS	Dynamic range Resolution Sensitivity	6/8 decades >2500-3000 m/Δm 10^3 #/cm^3	6/8 decades 5000 m/Δm 10^0 #/cm^3
		Measure isotopic ratio ^{12}C/^{13}C from mass spectra of condensed organics in icy grains	ENIJA	Dynamic range Mass range Resolution Sensitivity	3 decades 12–100m/z 100 m/Δm 1 ppm	6/8 decades 1–2000m/z 1000m/Δm 0.1 ppm
		Determine the abundance ratio between major C-, N-, O-, S-bearing compounds from mass spectra of plume vapour and solid phases, including allowing the separation of vapour-phase species with isobaric interferences for example, CO from N_2	INMS	Dynamic range Resolution Sensitivity	6/8 decades 3000 m/Δm 10^3 #/cm^3	6/10 decades 5000 m/Δm 10^0 #/cm^3
			ENIJA	Dynamic range Mass range Resolution Sensitivity	5 decades 12–200 m/z 200 m/Δm 1 ppm	6/8 decades 1–2000m/z 1000m/Δm 0.1 ppm

		Do Enceladus' volatile compounds originate from a unique reservoir and, if so, how is their distribution between solid and vapour phases affected by eruption processes?	Determine the compositional profile of major species through the plume (both volatile and non-volatile) from mass spectra of plume vapour and condensed phases in icy grains	INMS	Dynamic range Resolution Sensitivity	6/8 decades 300 m/Δm 10^3 #/cm^3	6/10 decades 5000 m/Δm 10^0 #/cm^3
				ENIJA	Dynamic range Mass range Resolution Sensitivity Spatial resolution	5 decades 7–500 m/z 500 m/Δm 1 ppm 1000 m	6/8 decades 1–2000m/z 1000m/Δm 0.1 ppm 100 m
	A.2: What is the history and extent of volatile exchange on Titan?	How were Titan's volatiles delivered, and were they internally processed?	Quantify the abundance of noble gases in the atmosphere	INMS	Dynamic range Resolution Sensitivity	6/8 decades 3000–5000 m/Δm 10^3 #/cm^3	6/10 decades 5000 m/Δm 10^0 #/cm^3
			Measure and compare isotopic ratios: $^{12}C/^{13}C$ and D/H in the atmospheric main constituents	INMS	Dynamic range Resolution Sensitivity	6 decades 3000 m/Δm 10^5 #/cm^3	6 decades 5000 m/Δm 10^0 #/cm^3
		What is the age of Titan's methane cycle, and is there evidence for renewal from internal outgassing?	Measure and compare isotopic ratios: ($^{14}N/^{15}N$; $^{12}C/^{13}C$; D/H; $^{16}O/^{18}O$) in the upper atmosphere	INMS	Dynamic range Resolution Sensitivity	6/8 decades 3000 m/Δm 10^3 #/cm^3	6/10 decades 5000 m/Δm 10^0 #/cm^3
			Constrain the genesis of candidate cryovolcanoes and cryovolcanic flows	TIGER	Coverage Scale Wavelength DTM vertical resol.	Sotra, Tui, & Hotei 50 m/pixel 3 colours 20 m	Sotra, Tui, & Hotei 30–50 m/pixel 3 colours 15–20 m
		How does Titan's surface reflect climate change?	Determine whether seas have existed in the tropics and/or south polar region	TIGER	Coverage Scale Wavelength DTM vertical resol.	Regions within Hotei, Tui, Xanadu & the putative south-polar basins 100m/pix 3 colour N/A	Regions within Hotei, Tui, Xanadu & the putative south-polar basins 100m/pix 3 colour N/A
			Investigate fine-scale dune patterns and how they reorient or breakup	TIGER	Coverage Scale Wavelength DTM vertical resol.	4 dune regions 50m/pix 3 colour N/A	4 dune regions 30–50m/pix 3 colour N/A

	A.3: How has Titan's organic-rich surface evolved?	How have liquid organics modified Titan's surface?	Investigate the morphology of river networks and channels	TIGER	Coverage Scale Wavelength DTM vertical resol.	3 drainages 30m/pix 3 colour 15 m	3 drainages 25–30m/pix 3 colour 10–15 m
			Determine whether the labyrinth terrains and sharp edged depressions are formed by dissolution	TIGER	Coverage Scale Wavelength DTM vertical resol.	2 regions 100m/pix 3 colour 40 m	2 regions 50–100m/pix 3 colour 20–40 m
		To what degree are sediments produced and transported by fluvial and eolian processes?	Investigate the margins of the dune fields to identify sources and sinks of sand	TIGER	Coverage Scale Wavelength DTM vertical resol.	4 dune regions 50m/pix 3 colour N/A	4 dune regions 30–50m/pix 3 colour N/A
			Determine the primary erosion processes that degrade impact crater rims	TIGER	Coverage Scale Wavelength DTM vertical resol.	4 impacts 100m/pix 3 colour N/A	4 impacts 50–100m/pix 3 colour N/A
			Investigate the fine-scale geologic processes that modify the plains	TIGER	Coverage Scale Wavelength DTM vertical resol.	3 regions 50m/pix 3 colour N/A	3 regions 30–50m/pix 3 colour N/A
		How do the mechanical and chemical properties (i.e., composition) vary between the observed geologic units on Titan?	Determine how fluvial erosion varies among geologic units	TIGER	Coverage Scale Wavelength DTM vertical resol.	5 regions 50m/pix 3 colour N/A	5 regions 30–50m/pix 3 colour N/A
			Constrain the maximum slopes and topography supported by different geologic units	TIGER	Coverage Scale Wavelength DTM vertical resol.	5 regions 100m/pix 3 colour 40m	5 regions 50–100m/pix 3 colour 20–40m
			Investigate the morphology, topography, and spectral relationships across unit boundaries	TIGER	Coverage Scale Wavelength DTM vertical resol.	5 boundaries 100m/pix 3 colour 40m	5 boundaries 50–100m/pix 3 colour 20–40m

GOAL B: HABITABILITY AND POTENTIAL FOR LIFE IN OCEAN WORLDS, ENCELADUS AND TITAN						
Science Objectives	**Scientific Questions**	**Measurement Description**	**Instrument**	**Measurement Requirements**		
				Parameter	**Requirement**	**Performance**
B.1: Is Enceladus' aqueous interior an environment favourable to the emergence of life?	What is the nature of hydrothermal activity on Enceladus and how does it connect with the plume activity?	Detect and inventory reduced and oxidized species in the plume material (e.g., NH_3/N_2 ratio, H_2 abundance, reduced versus oxidized organic species)	INMS	Dynamic range Resolution Sensitivity	6/8 decades 3000m/Δm 10^3 #/cm^3	6/10 decades 5000m/Δm 10^0 #/cm^3
			ENIJA	Dynamic range Mass range Resolution Sensitivity	4 decades 7–200 m/z 200 m/Δm 1 ppm	6/8 decades 1–2000m/z 1000m/Δm 0.1 ppm
		Characterize composition and abundance of salts and other minerals as messengers of rock/water interaction from mass spectra of icy grains	ENIJA	Dynamic range Mass range Resolution Sensitivity	6 decades 7–200 m/z 200 m/Δm 1 ppm	6/8 decades 1–2000m/z 1000m/Δm 0.1 ppm
		Determine spatial variations in plume composition (vapour and icy grains) and correlate with source activity	ENIJA	Dynamic range Mass resolution Resolution Sensitivity Spatial Resolution	4 decades 7–500 m/z 500 m/Δm 1 ppm 1000 m	6/8 decades 1–2000m/z 1000m/Δm 0.1 ppm 100 m
			TIGER	Coverage Scale Wavelength DTM vertical res.	2 cross-stripe tracks 5m/pix 5 & 5.3 µm N/A	2 cross-stripe tracks 2.5–5m/pix 5 & 5.3 µm N/A
		Determine how much energy the jets dissipate	TIGER	Coverage Scale Wavelength DTM vertical res.	2 night-time cross-stripe tracks 5m/pix 5 & 5.3 µm N/A	2 cross-stripe tracks 2.5–5m/pix 5 & 5.3 µm N/A

		Is there evidence that biotic/prebiotic processes are operating in Enceladus' interior?	Compare mass distribution of organic molecules to that expected from abiotic synthesis (e.g., Poisson versus non-Poisson) in the mass distribution of organic molecules	INMS	Dynamic range Resolution Sensitivity	6/8 decades 3000m/Δm 10^3 #/cm^3	6/10 decades 5000m/Δm 10^0 #/cm^3
				ENIJA	Dynamic range Mass range Resolution Sensitivity	6 decades 26-1000 m/z 500 m/Δm 1 ppm	6/8 decades 1-2000m/z 1000m/Δm 0.1 ppm
			Search for abnormal isotopic ratio in organics in plume vapour, including allowing the separation of species with isobaric interferences for example, ^{12}CH from ^{13}C	INMS	Dynamic range Resolution Sensitivity	6/8 decades 2500-3000 m/Δm 10^3 #/cm^3	6/10 decades 5000 m/Δm 10^0 #/cm^3
			Search for amino acids and abnormal ^{12}C/^{13}C ratio in organics that may have been captured in the ice matrix of solid plume material	ENIJA	Dynamic range Mass range Resolution Sensitivity	5 decades 12–500m/z 500 m/Δm 1 ppm	6/8 decades 1–2000m/z 1000m/Δm 0.1 ppm
	B.2: To what level of complexity has prebiotic chemistry evolved in the Titan system?	What chemical processes control the formation and evolution of complex organics in Titan's atmosphere?	Measure the elemental chemistry of low mass organic molecules in Titan's atmosphere	INMS	Dynamic range Resolution Sensitivity	6/8 decades 3000m/Δm 10^1 #/cm^3	6/10 decades 5000m/Δm 10^0 #/cm^3
			Measure organic macromolecules and ioni species (cations and anions) in order to monitor chemical coupling processes between ions and neutral species	INMS	Dynamic range Resolution Sensitivity	6/8 decades 3000m/Δm 10^1 #/cm^3	6/10 decades 5000m/Δm 10^0 #/cm^3
			Characterize the structure of haze layers, along with their spatial and temporal variations	TIGER	Coverage Scale	10 temporally spaced full-disk images 1km/pix	10 temporally spaced full-disk images 500m–1km/pix

ACCEPTED MANUSCRIPT

					Wavelength DTM vertical resol.	3 colours N/A	3 colours N/A
			Determine the cloud distribution and morphology, including estimates of their top altitudes and vertical extent	TIGER	Coverage Scale Wavelength DTM vertical resol.	10 temporally spaced full-disk images 1km/pix 3 colours N/A	10 temporally spaced full-disk images 500m–1km/pix 3 colours N/A
		To what extent does Titan's organic-rich surface chemically communicate with its water-rich interior?	Investigate the morphology and compositional variability of candidate cryovolcanoes	TIGER	Coverage Scale Wavelength DTM vertical resol.	Sotra, Tui & Hotei 50m/pix 3 colours 20m	Sotra, Tui & Hotei 30–50m/pix 3 colours 15–20m
			Investigate the morphology of Titan's mountains to look for evidence of compressional vs extensional tectonics	TIGER	Coverage Scale Wavelength DTM vertical resol.	3 mountains 100 m/pix 3 colours 40 m	3 mountains 50–100 m/pix 3 colours 20–40 m
			Investigate melt sheets and flows associated with impact craters	TIGER	Coverage Scale Wavelength DTM vertical resol.	2 larger fresh craters 50 m/pix 3 colours N/A	2 larger fresh craters 30–50 m/pix 3 colours N/A
			Measure the abundance of radiogenically derived noble gases (e.g., ^{40}Ar, ^{21}Ne) and helium	INMS	Dynamic range Resolution Sensitivity	6/8 decades 3000m/Δm 10^{1} #/cm^{3}	6/10 decades 5000m/Δm 10^{0} #/cm^{3}

ACCEPTED MANUSCRIPT

Highlights

- We present a space mission concept, the Explorer of Enceladus and Titan (E^2T), which is dedicated to investigating the evolution and habitability of these Saturnian satellites.

- E^2T is proposed as a medium-class mission led by ESA in collaboration with NASA in response to ESA's M5 Cosmic Vision Call.

- E^2T proposes a focused payload that would provide in-situ composition investigations and high-resolution imaging during multiple flybys of Enceladus and Titan using a solar-electric powered spacecraft in orbit around Saturn.

ACCEPTED MANUSCRIPT