UNIVERSITY OF LEEDS

This is a repository copy of *Asian monsoons and aridification response to Paleogene sea retreat and Neogene westerly shielding indicated by seasonality in Paratethys oysters*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/136029/>

Version: Accepted Version

Article:

Bougeois, L, Dupont-Nivet, G, de Rafélis, M et al. (6 more authors) (2018) Asian monsoons and aridification response to Paleogene sea retreat and Neogene westerly shielding indicated by seasonality in Paratethys oysters. *Earth and Planetary Science Letters*, 485. pp. 99-110. ISSN 0012-821X

<https://doi.org/10.1016/j.epsl.2017.12.036>

(c) 2017, Elsevier B.V. This manuscript version is made available under the CC BY-NC-ND 4.0 license <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Asian monsoons and aridification response to Paleogene sea retreat and westerly shielding indicated by strong seasonality in Paratethys oysters

Laurie Bougeois^{a,*}, Guillaume Dupont-Nivet^{b,c,d}, Marc de Rafélis^e, Julia C. Tindall^f, Jean-Noël Proust^b, Gert-Jan Reichart^g, Lennart J. de Nooijer^g, Zhaojie Guo^d, Cholponbelk Ormukov^h

^a*Ministère de l'Éducation nationale, Académie de Montpellier, France*

^b*Géosciences Rennes, UMR-CNRS 6118, Université de Rennes 1, Rennes, France*

^c*Department of Earth and Environmental Sciences, Potsdam University, Germany*

^d*Key Laboratory of Orogenic Belts and Crustal Evolution, Ministry of Education, Beijing, China*

^e*GET, Observatoire Midi Pyrénées, CNRS, IRD, Université de Toulouse, Toulouse, France*

^f*School of Earth and Environment, University of Leeds, Leeds LS2 9JT, UK*

^g*Department of Marine Geology, Royal Netherlands Institute for Sea Research, Texel, The Netherlands*

^h*Central Asian Institute for Applied Geosciences (CAIAG), Bishkek, T. Frunze rd.73/2, 720027, Bishkek, Kyrgyzstan*

Abstract

Asian climate patterns, characterized by highly seasonal monsoons and continentality, are thought to originate in the Eocene epoch (56 to 34 million years ago - Ma) in response to global climate, Tibetan Plateau uplift and the disappearance of the giant Proto-Paratethys sea formerly extending over Eurasia. The influence of this sea on Asian climate has hitherto not been constrained by proxy records despite being recognized as a major driver by climate models. We report here strongly seasonal records preserved in annual lamina of Eocene oysters from the Proto-Paratethys with sedimentological and numerical data showing that monsoons were not dampened by the sea and that aridification was modulated by westerly moisture sourced from the sea. Hot and arid summers despite the presence of the sea suggest a strong anticyclonic zone at Central Asian latitudes and an orographic effect from the emerging Tibetan Plateau. Westerly moisture precipitating during cold and wetter winters appear to have decreased in two steps. First in response to the late Eocene (34-37 Ma) sea retreat; second by the orogeny of the Tian Shan and Pamir ranges shielding the westerlies after 25 Ma. Paleogene sea retreat and Neogene westerly shielding thus provide two successive mechanisms forcing coeval Asian desertification and biotic crises.

Keywords: Eocene monsoon, aridification, Paratethys sea, Central Asia, seasonality, bivalves

*Corresponding author at:

Email address: laurie.bougeois@gmail.com (Laurie Bougeois)

1. Introduction

Asian climate, governing the livelihood of billions of people, is characterized by high seasonality. In southeastern Asia seasons are expressed by high summer rainfall compared to winter in response to the archetypal monsoonal circulation. In contrast, central continental Asia records extreme seasonal temperature and minimal precipitation (Araguás-Araguás et al., 1998). The origin and past evolution of these seasonal patterns remain poorly constrained although they are key to deciphering their forcing mechanisms and to validate climate model predictions. To understand Asian climate, it is crucial to assess and quantify its past seasonality, in particular during the Eocene epoch (56 to 34 Myr ago), when high atmospheric pCO₂ levels kept global climate in a greenhouse state and the India-Asia collision shaped the paleogeographic features that would define Asian climate patterns (Zachos et al., 2001; Pagani et al., 2005).

During this epoch, the India-Asia collision resulted in the growth of the Tibetan Plateau and Himalayas. These orogenies are traditionally held responsible for the establishment of monsoons and desertification (Guo et al., 2002; France-Lanord et al., 1993) by creating orographic barriers as well as increasing atmospheric circulation through enhanced heat transfer from the plateau surface to the atmosphere (Molnar et al., 2010; Boos and Kuang, 2010; Wu et al., 2012; Liu et al., 2012). These environmental changes during Cenozoic have been called upon to explain major biotic events (Favre et al., 2015) such as the Mongolian Remodelling (Kraatz and Geisler, 2010; Fortelius et al., 2014) and the emergence of grassland and C4 plants (Edwards et al., 2010). Another competing forcing mechanism during this epoch is the strong global climate cooling from a greenhouse to an icehouse state. Models suggest that strong monsoonal circulation was maintained by warmer Eocene conditions (Licht et al., 2014) but subsequent global cooling led to Asian aridification and decreasing monsoonal intensity mostly due to diminished moisture transport (Dupont-Nivet et al., 2007; Licht et al., 2014). We focus here on the influence of the large epicontinental Proto-Paratethys sea that formerly covered Eurasia (Bosboom et al., 2014c). It is also recognized as a major driver by modelling studies suggesting that its presence would dampen the seasonal thermal contrast between the continent and surrounding oceans negating the possibility of intense Eocene monsoons (Ramstein et al., 1997; Zhang et al., 2007). Also the sea potentially provided an important moisture source transported by the westerlies into Asia (Dupont-Nivet et al., 2007; Zhang et al., 2012). The sea fluctuations and retreat may therefore have modulated Asian environments leading to desertification and increasing monsoonal circulation (Ramstein et al., 1997).

In principle, climate proxies of seasonality contrasts from the sedimentary records should enable

to disentangle the respective contributions of forcing mechanisms suggested by climate models. If the sea indeed dampened the ocean-continent thermal contrast, a temperate climate with low seasonality would be expected. In addition, seasonality can be used to discriminate between monsoonal vs. westerly moisture sources because they have opposite precipitation patterns: monsoons are dominated by summer precipitation while westerlies are characterised by winter precipitation. Existing records of aeolian loess-like deposits (Licht et al., 2014), pedogenic stable isotope (Caves et al., 2015) and fossil pollen studies (Dupont-Nivet et al., 2008) point to monsoonal circulations with Asian interior aridity despite the Proto-Paratethys sea presence. However, no quantitative records of Eocene seasonality from this area have been hitherto produced due to the paucity of appropriate records and reliable methods to extract them.

To constrain seasonality, Bougeois et al. (2014, 2016) developed a geochemical high resolution multi-proxy approach (oxygen stable isotopes $\delta^{18}\text{O}$ - and Mg/Ca elemental ratios) on pilot samples from fossil oyster shells of the Proto-Paratethys itself. Here we apply this approach to numerous specimens and complement this geochemical data with sedimentological paleoenvironmental analyses and a coupled atmosphere-ocean general circulation model -GCM- constraining independently oceanic and atmospheric temperatures, precipitation patterns and isotopic composition in the Proto-Paratethys region. Furthermore, the isotopic signatures of Paleogene to Neogene pedogenic carbonates are compared to the marine data to identify the evolution of moisture sources during sea retreat, regional uplift and global cooling.

2. Geological setting

This study focuses on the well-dated Proto-Paratethys sea sediments exposed today in the Tarim and Alai Valley basins over the foothills of the Central Asian (Pamir and Tian Shan) ranges (Bosboom et al., 2014c,a,b) (Fig. 1) that uplifted since the early Miocene (ca. 25 Ma Sobel et al. (2006); Zheng et al. (2015); Blayney et al. (2016)). In the early Eocene (ca. 55 Ma) the sea reached its maximum expansion, from the Tarim basin to the Mediterranean and linked Arctic and paleo-Indian oceans (Dercourt et al., 1993). After this, the sea retreated westward. It was barely reaching the Tarim basin towards the Late Eocene (ca. 37 Ma) and had shrunk to the Caspian sea's present position by the 34 Ma Eocene-Oligocene transition (Bosboom et al., 2014b). The retreat therefore likely results from tectonic deformations in response to the early Eocene India-Asia collision onset (Molnar et al., 2010). However, the eustatic drop associated with global late Eocene cooling that lead to the Antarctic ice sheet at the Eocene-Oligocene transition, probably also contributed to the sea retreat (Bosboom et al., 2014b).

3. Material and methods

3.1. Climatic model simulations

The model used to investigate the Eocene climate is the Hadley Centre General Circulation Model (HadCM3 [Gordon et al., 2000](#)) with isotope tracers incorporated ([Tindall et al., 2009](#)). The model resolution is $3.75^\circ \times 2.5^\circ$, with 19 vertical levels in the atmosphere and 20 levels in the ocean. The simulations used are as described by [Tindall et al. \(2010\)](#) and are based on Early Eocene boundary conditions (see supplementary material for details). However, we note that model boundary conditions for this time are subject to considerable uncertainty. Briefly, CO_2 was set to 1680 ppmv ($6\times$ pre-industrial levels) and was intended to represent the combined radiative forcing from all greenhouse gases (since CH_4 and N_2O were kept as pre-industrial). The land-sea mask and orography is described in [Tindall et al. \(2010\)](#) and was produced using similar methods to [Markwick and Valdes \(2004\)](#). Of particular relevance for our study is that Tibetan plateau is set to a maximum height of ~ 1500 m, the topography of the Tian Shan and Pamir are absent and Central Asia is covered by the Proto-Paratethys with a depth of under 100 m, and water exchange between the Proto-Paratethys and the Indian ocean is possible through a gateway wider than 15 latitudinal degrees. Exchange with the Arctic is more difficult as the Turgai Strait is only one gridbox wide and 80 m deep. As a result there can be no baroclinic flow and limited barotropic flow between the Proto-Paratethys and the Arctic. Globally the Eocene simulation was 14°C warmer and 20% more precipitation than a corresponding pre-industrial simulation (see supplementary material for more information). Here we focus on gridboxes 37.5°N , 71.25°E and 40.0°N 63.25°E for ocean, and 40°N , 75°E for atmosphere, which are the closest area from field study since the position of the sites experienced no statistically significant latitudinal tectonic motion since the time of deposition ([Bougeois et al., 2014](#)).

3.2. Geochemical data for sclerochronology

To quantify seasonal variations of temperature and salinity of the Proto-Paratethys seawater, we applied geochemical incremental analyses on fossil oyster's ligamental area following the multi-proxy methodology developed by [Bougeois et al. \(2014, 2016\)](#).

Oyster sampling was performed with particular attention to: (1) good preservation of a ligamental area large enough for a high resolution infra-annual record through numerous years, (2) ensuring that specimens fossilized in living position in fully marine environments. We focus here to the well-dated Middle Eocene (Lutetian) species ([Bosboom et al., 2014a,b,c](#)) *Ostrea (Turkostrea) strictiplicata* and *Sokolowia*

91 *buhssii*, which lived in subtidal environment (Fig. 2e,f) as attested by sedimentological analyses (see also
92 supplementary material), and that have been shown to provide reliable Mg/Ca results (Bougeois et al.,
93 2016). The oyster shells were sectioned perpendicular to their maximal growth axis and well polished be-
94 fore geochemical analyses. Cross sections of oyster shells reveal large numbers of distinct light and dark
95 growth bands, especially well-expressed in the ligamental area resulting from the typical incremental
96 growth of yearly dark-light couplets (Bougeois et al., 2014). Cathodoluminescence microscopy revealed
97 the annual banding, attesting no diagenesis effect on the calcitic shells.

98 Mg/Ca analyses were performed with Laser Ablation-Inductively Coupled Plasma Mass Spectrometer
99 (LA-ICP MS) at the Department of Earth Sciences in Utrecht University following two parallel transects
100 perpendicularly to the growth direction. Upon checking the consistency of the two parallel transects,
101 their results were averaged such that a single datapoint was obtained for each incremental position, then
102 a moving average on 21 points is calculated to overcome ICP-MS noise (see Bougeois et al., 2014, 2016,
103 for more details).

104 Microsample powders were drilled following growth layers every 100 to 120 μm using a Merchantek
105 MicroMill then analysed for stable isotopes composition using a KIEL-III device coupled online to a
106 Finnigan MAT-253 mass spectrometer at the Department of Earth Sciences in Utrecht University (KY01,
107 AT04) and using a KIEL-IV device coupled to an Isoprime DI-IRMS at the Department of Earth Sciences
108 in Pierre et Marie Curie University (AL02, MS05, AT20, AT19). Internal and international (NBS 19)
109 standards were used for reproducibility. For both mass spectrometers, long-term analytical precision was
110 better than 0.08‰ for $\delta^{18}\text{O}$.

111 From all the specimens shown in Bougeois et al. (2016), we finally selected six specimens where
112 reliable Mg/Ca and $\delta^{18}\text{O}_\text{c}$ were available. We show here only the part of the shells where both proxy
113 were performed (all data sets are provided in supplementary material).

114 3.3. Geochemical data for carbonates sediments

115 Carbonates sediments (bioclastic grainstone to wackstone for marine sediments and carbonaceous pe-
116 dogenic horizons for continental sediments) were sampled at Mine and Aertashi sections in Tarim Basin
117 (China) for stable isotopes analyses (horizon level indicated in supplementary material). Continental
118 sediments selected were carefully chosen from the finest granulometry (mudstones to siltstones) with
119 carbonaceous matrix unaltered and devoid of secondary vein of calcite. To avoid effects of diagenesis, in
120 laboratory we sampled the fresh core of samples.

121 After milling the sediments, we analysed the carbonate fraction using mass spectrometer SIRA 9 at

122 University Pierre and Marie Curie (Paris 6). Internal (white marble Marceau) and international (NBS 19)
123 standards were used for reproducibility. Long-term analytical precision was better than 0.05‰ for $\delta^{13}\text{C}$
124 and 0.1‰ for $\delta^{18}\text{O}$.

125 **4. Eocene Central Asian seasonality and monsoons**

126 *4.1. Paleogene sedimentological facies analyses.*

127 Paleogene sedimentological facies analyses have been performed throughout the Aertashi and Mine
128 sections (Fig. 1) displaying alternation of marine and continental deposits recording several sea incur-
129 sions and subsequent retreats (Fig. 3, 4 and detailed facies associations in supplementary material).
130 Sedimentary facies and fossil assemblages of marine sediments (Fig. 2c-d) are characteristic of shallow
131 marine environments between upper offshore to coastal plain, and typical of warm, carbonate-rich neritic
132 ramps. Tidal flat environments indicate a calm and shallow epicontinental sea prone to record paleo-
133 climate fluctuations. Continental deposits are indicative of flood plains, playa and sabhka environments
134 (Fig 2a-b). Alternations of sandy fluvial channel-fills and flood plain red silty clays with nodular gypsum
135 and desiccation cracks attest for strong seasonal contrasts testifying for successions of floods events and
136 dessication periods typical of semi-arid climates as also indicated by existing marine microfossils and
137 pollen data (Sun and Wang, 2005; Bosboom et al., 2014a,c). A modern analogue is provided by the
138 Persian Gulf, which is directly connected to the Indian Ocean but protected in a wide gulf and subject to
139 semi-arid conditions leading to playa and sabkhas hypersaline deposits (James and Dalrymple, 2010). To
140 quantify the seasonality we explore model and proxy data in the following

141 *4.2. Numerical simulations of Eocene climate.*

142 Eocene climate (Fig. 5b) over the Proto-Paratethys sea region provided by the HadCM3 General
143 Circulation Model shows annual average sea surface temperatures (SST) of 23°C, with average seasonal
144 cycle between 16°C and 34°C. Modelled $\delta^{18}\text{O}$ of sea water surface ($\delta^{18}\text{O}_{\text{sw}}$) values average 0.44‰
145 (SMOW) and, because ocean gridboxes are large and well mixed, show negligible seasonal variability.
146 However, a higher variability is expected in the environments studied here because modelled air temper-
147 atures fluctuate largely between 7°C and 49°C (average of 26°C) and coastal areas are prone to seasonal
148 water balance variations (Goodwin et al., 2001). Modelled precipitation in our study area are strongly
149 seasonal peaking at 38 mm in January and reaching a 0.5 mm minimum in July. This pattern is consistent
150 with previous Eocene model simulations (Zhang et al., 2012) and similar to modern conditions west of
151 the Central Asian ranges (Bukhara site, Fig. 5) subjected to winter westerly precipitation. Stable isotope

composition of precipitation ($\delta^{18}\text{O}_p$) is stable around -6‰ except for a significant increase during the warmest month with $\delta^{18}\text{O}_p = -1.7\text{‰}$ in July. The model winter values are very far from isotopic composition of modern precipitation East and South of the Central Asian ranges (Araguás-Araguás et al. (1998), Hotan site, Fig. 5c). The general contrast between Eocene model and modern precipitation and isotopic seasonality suggests conditions have changed drastically since Eocene times in this area.

4.3. Eocene seasonality revealed by oyster shell geochemistry.

Along the ligamental areas of the shells, the growth bands show Mg/Ca and $\delta^{18}\text{O}$ periodic fluctuations, which are synchronized and anti-correlated (high values of Mg/Ca corresponding to low values of $\delta^{18}\text{O}$ and inversely). These variations with clear banding attest for a well-recorded seasonal pattern and no diagenetic alteration, as also supported by cathodoluminescence analyses (Fig. 6). The primary character of trace element and stable isotope values is attested by measurement reproducibility in these species, which display homogeneous values both within the same sedimentary horizon and in the different sections of Central Asia. According to Bougeois et al. (2014, 2016), we infer quantitatively temperature changes from the chemical oyster shell composition using the relationships calibrated in the modern oyster *Crassostrea gigas* (Mouchi et al., 2013):

$$T(^{\circ}\text{C}) = 3.77 \times \text{Mg/Ca}(\text{mmol/mol}) + 1.88 \quad (1)$$

The SST reconstructed from Mg/Ca are in excellent agreement with temperatures derived from the modelled HadCM3 (Fig. 5a,b). The seasonal temperatures amplitudes based on Mg/Ca ($\Delta\text{SST}=19^{\circ}\text{C}$) is slightly higher to the SST seasonality predicted from HadCM3 ($\Delta\text{SST}=18^{\circ}\text{C}$) and the annual averages ($27\pm 2^{\circ}\text{C}$ according Mg/Ca vs $23\pm 6^{\circ}\text{C}$ according model) are comparable (details in supplementary material). Especially, Mg/Ca-based temperatures lay in-between the SSTs and the surface air temperatures predicted by HadCM3. These amplitudes between air and sea temperature are typically expected in these coastal environments (Goodwin et al., 2001) which further supports the validity of temperatures estimated with Mg/Ca.

We infer further paleoclimate information from the fossil $\delta^{18}\text{O}_c$ using the well established relationship for bivalves linking temperature T ($^{\circ}\text{C}$), $\delta^{18}\text{O}_c$ (‰ VPDB) and $\delta^{18}\text{O}_{\text{sw}}$ (‰ VSMOW) (Anderson and Arthur, 1983):

$$T = 16 - 4.14 \times (\delta^{18}\text{O}_c - \delta^{18}\text{O}_{\text{sw}}) + 0.13 \times (\delta^{18}\text{O}_c - \delta^{18}\text{O}_{\text{sw}})^2 \quad (2)$$

On first approximation a stable $\delta^{18}\text{O}_{\text{sw}}$ of 0.44‰ derived from the modelled HadCM3 (Fig. 5b) is used. The obtained average temperatures ($28\pm 2^{\circ}\text{C}$) are comparable to annual averages of modelled ($23\pm 6^{\circ}\text{C}$)

180 and Mg/Ca ($27 \pm 2^\circ\text{C}$) SST. However, temperature amplitudes reconstructed from $\delta^{18}\text{O}_c$ are considerably
181 lower ($\Delta\text{SST}=9^\circ\text{C}$) than amplitudes of modelled and Mg/Ca SST. Given that $\delta^{18}\text{O}_c$ depends on tempera-
182 ture and $\delta^{18}\text{O}_{\text{sw}}$, this discrepancy is most likely due to stronger seasonal variability in $\delta^{18}\text{O}_{\text{sw}}$ compared
183 to stable values in open water predicted by HadCM3.

184 These seasonal fluctuations in $\delta^{18}\text{O}_{\text{sw}}$ can be directly derived from equation (2) using the shell $\delta^{18}\text{O}_c$
185 and temperatures (T) deduced from Mg/Ca:

$$\delta^{18}\text{O}_{\text{sw}} \simeq \frac{T - 16}{4.14} + \delta^{18}\text{O}_c \quad (3)$$

186 Seawater oxygen isotope compositions thus obtained indicate high seasonal fluctuations with highest
187 $\delta^{18}\text{O}_{\text{sw}}$ during summer months and $\sim 3\text{‰}$ lower values during winter (Fig. 6). These fluctuations are in
188 full agreement with our sedimentological interpretations of oysters living environments at the epiconti-
189 nental sea margin where runoff, precipitation and evaporation have a strong effect on $\delta^{18}\text{O}_{\text{sw}}$. This $\delta^{18}\text{O}_{\text{sw}}$
190 seasonality is also consistent with typical coastal environment influenced by dry summer conditions re-
191 sulting in a negative water balance (increasing $\delta^{18}\text{O}_{\text{sw}}$ and salinity) contrasted with positive water balance
192 during cooler and wetter winter months (decreasing $\delta^{18}\text{O}_{\text{sw}}$ and salinity) (Goodwin et al., 2001, 2010).

193 4.4. Interpretation of seasonality.

194 Our results show that the Eocene Central Asian summer climate was hotter than today and already
195 arid despite the Proto-Paratethys sea presence (Fig. 5b). According to the model, Eocene seasonal air
196 temperature amplitudes ($\Delta T \sim 42^\circ\text{C}$) were higher than today ($\Delta T \sim 32^\circ\text{C}$), and aridity was more sustained in
197 summer with very low precipitation. As a moisture source, the Proto-Paratethys appears to have had little
198 impact on local climate during summer. Most importantly, the high reconstructed summer temperatures
199 imply that the shallow sea did not thermally buffer the Asian interior and delay the onset of monsoonal
200 circulation, as suggested by previous models (Ramstein et al., 1997; Zhang et al., 2007). This may be
201 attributed to overall warmer Eocene global climate imposing a stronger anticyclonic Hadley high pressure
202 cell descending at these latitudes (25 to 45°N) over Central Asia (Zhang et al., 2012). It is also consistent
203 with recent studies showing that high atmospheric $p\text{CO}_2$ levels had more impact on circulation than local
204 paleogeography (Lunt et al., 2016). In addition, the emerging Proto-Tibetan plateau during this period
205 (Molnar et al., 2010), even at low altitude, may have contributed to a stronger Foehn effect during summer
206 months bringing warm and dry air into Central Asia (Fig. 7a). Our results are thus supported by recent
207 model and proxy data suggesting modern-like Asian monsoonal circulation already established as early
208 as Eocene times (Sun and Wang, 2005; Huber and Goldner, 2012; Licht et al., 2014; Caves et al., 2015).

209 In contrast, the observed summer aridity precludes previously proposed pre-Neogene low pressures and
210 humid conditions north of the Tibetan Plateau, as this region would have been sufficiently shielded from
211 Asian monsoon rains at this time and high pressures cell hence fixed to its north (Allen and Armstrong,
212 2012).

213 Furthermore, our results imply enhanced winter over summer Eocene precipitation, which is sup-
214 ported by climate model simulations suggesting a dominant westerly winter moisture source (Tindall
215 et al., 2010; Zhang et al., 2012). Eocene winter air temperature was significantly warmer than today and
216 the source of moisture unshielded by Central Asian ranges. The relatively high $\delta^{18}\text{O}_p$ and precipitation
217 during Eocene winters can thus be interpreted as resulting from winter westerlies bringing moist air from
218 the neighbouring Proto-Paratethys and adjoining seas (Fig. 7b). Reconstructed Eocene seasonality is ac-
219 tually comparable to modern conditions on the Central Asian ranges' western flank exposed to westerlies
220 with enhanced winter precipitation (up to 85 mm/month - Fig. 5c, Bukhara site) (Araguás-Araguás et al.,
221 1998). This sharply contrasts with modern climate patterns on the other side of the Central Asian ranges
222 (Fig. 5c, Kashgar site). There, climate is hyper arid with maximum seasonal rainfall reaching only
223 12 mm/month in late spring and summer. The minimal moisture is typically recycled locally through
224 groundwater evaporation or plant cover transpiration (Araguás-Araguás et al., 1998) resulting in strong
225 seasonal variability off precipitation stable isotope (Fig. 5c).

226 Compared to Eocene, these regions are more arid today with a reversed summer/winter precipitation
227 seasonality pattern. To understand the potential driving factors of these changes from Eocene to the
228 modern climate patterns we investigate below the Eocene to Pliocene moisture evolution.

229 5. Eocene to Pliocene moisture evolution

230 To track the moisture composition through the Cenozoic, we analysed bulk carbon ($\delta^{13}\text{C}$) and oxygen
231 isotopic compositions of Paleogene carbonates of Aertashi and Mine sections (Fig. 3 and 4), which are
232 prone to reflect the isotopic composition of water in which they precipitated. These analyses include
233 marine and pedogenic carbonates (Tab. ?? and ??) and are complemented by the Neogene data provided
234 by Kent-Corson et al. (2009).

235 Stable isotopes in continental vs. marine deposits are fundamentally different. In marine systems,
236 there is a substantial influence of ice volume, temperature and especially in coastal area, a component
237 of runoff on $\delta^{18}\text{O}$. In terrestrial systems, $\delta^{18}\text{O}$ is primarily controlled by the ratio of precipitation to
238 evapotranspiration (Winnick et al., 2014). Similarly, $\delta^{13}\text{C}$ should have fundamentally different values

239 in both marine and terrestrial systems, reflecting different sources of the carbon. The data are therefore
240 interpreted separately.

241 The $\delta^{18}\text{O}$ from bulk marine limestones show a slightly decreasing trend from ca. -2 to -8‰ through-
242 out the Late Paleocene to the Late Eocene. In contrast, terrestrial $\delta^{18}\text{O}$ strongly decrease from ca. -7 to
243 -14‰ from the Eocene to the Miocene (Fig. 8).

244 $\delta^{13}\text{C}$ decreases from 6 to -5 ‰ in Eocene marine limestones. Then $\delta^{13}\text{C}$ increases to 3 ‰ from Late
245 Eocene to Miocene continental pedogenic carbonates (details in supplementary material).

246 The Late Paleocene to Late Eocene decrease in marine $\delta^{13}\text{C}$ is consistent with an increase in runoff
247 and a decrease in fully marine contributions due to the sea retreat. The Eocene to Neogene increase
248 in terrestrial carbonates $\delta^{13}\text{C}$ may be partly related to the coeval mudstones to conglomerates lithologic
249 changes (Kent-Corson et al., 2009). However, the rise is also consistent with regional aridification (Bos-
250 boom et al., 2014b; Sun and Wang, 2005; Quan et al., 2012) suggesting alternatively that it results from
251 a combination of water scarcity increasing the $\delta^{13}\text{C}$ of plant matter (Suits et al., 2005; Diefendorf et al.,
252 2010; Kohn, 2010) and a decrease in plant productivity (Caves et al., 2016) that would reduce the quantity
253 of soil respired CO_2 .

254 The $\delta^{18}\text{O}$ decrease in marine limestones since ca. 55 Ma likely reflects the retreating sea with a shift
255 to more coastal environments increasingly affected by precipitation and runoff (Bosboom et al., 2014a).
256 At the transition from marine to continental deposits associated with the sea retreat out of the Tarim Basin
257 (ca. 37 Ma), overlapping marine limestones to continental carbonates $\delta^{18}\text{O}$ values suggest a gradual tran-
258 sition with continental precipitation being evaporated from the nearby sea. After the marine-continental
259 transition and up to the Pliocene, the $\delta^{18}\text{O}$ decrease must be interpreted in terms of precipitation. These
260 are most likely governed by westerly moisture sources given the predominant winter precipitation indi-
261 cated by the seasonality data above. This corroborates the recent compilation of pedogenic and lacustrine
262 carbonate $\delta^{18}\text{O}$ data across Central Asia also indicating that the westerlies were the dominant source of
263 moisture and therefore must have controlled aridification (Caves et al., 2015). Of the many factors that
264 may have influenced the precipitation $\delta^{18}\text{O}$ decrease, the distance from the source and an orographic
265 rain-shadow effect of the Central Asian ranges probably dominated compared to relatively small ex-
266 pected $\delta^{18}\text{O}$ decrease due to altitude and temperature changes of the site (Araguás-Araguás et al., 1998;
267 Botsyun et al., 2016). Indeed this time interval corresponds to further westward sea retreat (Bosboom
268 et al., 2014c,b) and regional uplift shielding the Tarim Basin from the westerlies (Sobel et al., 2006;
269 Zheng et al., 2015; Blayney et al., 2016; Caves et al., 2016, 2017).

270 Finally, because the sea had already retreated back to the present Caspian Sea location after the
271 Eocene-Oligocene transition (Bosboom et al., 2014c), most of the subsequent isotopic change must be
272 attributed to orographic effects related to the Tian Shan and Pamir uplifts (ca. 25-15 Ma Sobel et al.,
273 2006; Zheng et al., 2015; Blayney et al., 2016). In addition, decreasing $\delta^{18}\text{O}$ may results from a greater
274 contribution of high-elevation precipitation of the local water (Macaulay et al., 2016).

275 6. Conclusions

276 Our results reveal clear and cyclic geochemistry alternations in fossil oyster shells indicating an ex-
277 ceptional preservation suitable for climate proxy reconstruction. These records, in excellent agreement
278 with sedimentology and numerical simulations, enable to constitute the first robust quantitative estimate
279 of seasonality for this area with the following implications.

280 Despite the presence of the Eocene Proto-Paratethys sea, the Asian interior climate was semi-arid and
281 strongly seasonal receiving dominantly winter moisture from the westerlies. Highly seasonal temperature
282 contrasts indicate that the shallow sea did not have a strong dampening effect that may imply monsoonal
283 circulation. This contrasts with previous modelling studies (Ramstein et al., 1997) but confirms recent
284 regional evidence for strong Eocene monsoons (Licht et al., 2014).

285 The sea, however, provided moisture to Central Asia through westerlies during Eocene winters. Our
286 results, thus suggest a two step aridification. The first one related to the Eocene to Oligocene west-
287 ward Proto-Paratethys sea retreat and affecting central to eastern Central Asia (Bosboom et al., 2014a).
288 The subsequent aridification associated with the early Miocene uplift of Pamir and Tian Shan affecting
289 the regions east of these ranges shielding the westerlies and leading to enhanced aridification, recycled
290 precipitation patterns and desertification of Taklamakan, Qaidam and Gobi regions. These two events
291 are consistent with the documented paleo-wind patterns (Licht et al., 2016), and provide respectively a
292 driving mechanisms for the generation of (1) Eocene aeolian loess-like deposits (Licht et al., 2014) in
293 response to the sea retreat, and (2) Mio-Pliocene Loess (Nie et al., 2015) in response to Central Asian
294 ranges orogenies. The past diminution of westerly rather than monsoonal moisture was thus more likely
295 the governing factor of the aridification held responsible for major biotic crisis documented in this area
296 (Kraatz and Geisler, 2010; Fortelius et al., 2014; Edwards et al., 2010).

297 7. Acknowledgements

298 This project was funded by the Netherlands Organization for Scientific Research, the A. v. Humboldt
299 foundation, the Marie Curie Career Integration Grant FP7 CIG grant 294282 'HIRESDAT', the French
300 Ministry of Higher Education and Research, Horizon 2020 ERC grant 649081 'MAGIC', ANR DSP-
301 Tibet and the CaiYuanpei programme of the French ministry of foreign affairs. We thank R. Bosboom, G.
302 Heilbronn, B. Paulet and Y. WEI for their contribution in the field. We thank G. Ramstein for constructive
303 criticisms and suggestions that greatly improved this manuscript. We are grateful to H. de Waard and A.
304 Van Dijk (LA-ICP-MS and stable isotopes analyses in Utrecht University) and to N. Labourdette and C.
305 Pierre (stable isotopes analyses and Micromill access in Pierre et Marie Curie University). We thank J.
306 Caves for constructive reviews and suggestions.

307 References

- 308 Allen, M., Armstrong, H., 2012. Reconciling the Intertropical Convergence Zone, Himalayan/Tibetan
309 tectonics, and the onset of the Asian monsoon system. *Journal of Asian Earth Sciences* 44, 36–47.
- 310 Anderson, T., Arthur, M., 1983. Stable isotopes of oxygen and carbon and their application to sedimen-
311 tologic and paleoenvironmental problems. *Stable isotopes in sedimentary geology* 10, 1–151.
- 312 Araguás-Araguás, L., Froehlich, K., Rozanski, K., 1998. Stable isotope composition of precipitation over
313 southeast Asia. *Journal of Geophysical Research: Atmospheres* (1984–2012) 103, 28721–28742.
- 314 Baker, C., Eischeid, J., Karl, T., Diaz, H., 1995. The quality control of long-term climatological data
315 using objective data analysis, in: *Preprints of AMS ninth conference on applied climatology*, Dallas,
316 TX, pp. 15–20.
- 317 Blayney, T., Najman, Y., Dupont-Nivet, G., Carter, A., Miller, I., Garzanti, E., Sobel, E.R., Rittner, M.,
318 Ando, S., Guo, Z., Vezzoli, G., 2016. Indentation of the Pamirs with respect to the northern margin of
319 Tibet: constraints from the Tarim Basin sedimentary record. *Tectonics* , 2345–2369.
- 320 Boos, W., Kuang, Z., 2010. Dominant control of the South Asian monsoon by orographic insulation
321 versus plateau heating. *Nature* 463, 218–222.
- 322 Bosboom, R., Dupont-Nivet, G., Grothe, A., Brinkhuis, H., Villa, G., Mandic, O., Stoica, M., Huang,
323 W., Yang, W., Guo, Z., Krijgsman, W., 2014a. Linking Tarim Basin sea retreat (west China) and Asian
324 aridification in the late Eocene. *Basin Research* .

325 Bosboom, R., Dupont-Nivet, G., Grothe, A., Brinkhuis, H., Villa, G., Mandic, O., Stoica, M., Kouwen-
 326 hoven, T., Huang, W., Yang, W., et al., 2014b. Timing, cause and impact of the late Eocene stepwise
 327 sea retreat from the Tarim Basin (west China). *Palaeogeography, Palaeoclimatology, Palaeoecology* .
 328 Bosboom, R., Dupont-Nivet, G., Mandic, O., Proust, J., Ormukov, C., 2014c. Late Eocene paleogeog-
 329 raphy of the Proto-Paratethys Sea in Central Asia (NW China, S Kyrgyzstan and SW Tajikistan).
 330 Geological Society of London Special Publications .
 331 Botsyun, S., Sepulchre, P., Risi, C., Donnadieu, Y., 2016. Impacts of Tibetan Plateau uplift on atmo-
 332 spheric dynamics and associated precipitation $\delta^{18}\text{O}$. *Climate of the Past* 12, 1401–1420.
 333 Bougeois, L., De Rafélis, M., Reichart, G.J., de Nooijer, L.J., Dupont-Nivet, G., 2016. Mg/Ca in fossil
 334 oyster shells as palaeotemperature proxy, an example from the Palaeogene of Central Asia. *Palaeo-*
 335 *geography, Palaeoclimatology, Palaeoecology* 441, 611–626.
 336 Bougeois, L., de Rafélis, M., Reichart, G.J., de Nooijer, L.J., Nicollin, F., Dupont-Nivet, G., 2014. A
 337 high resolution study of trace elements and stable isotopes in oyster shells to estimate Central Asian
 338 Middle Eocene seasonality. *Chemical Geology* 363, 200–212.
 339 Caves, J.K., Bayshashov, B.U., Zhamangara, A., Ritch, A.J., Ibarra, D.E., Sjostrom, D.J., Mix, H.T.,
 340 Winnick, M.J., Chamberlain, C.P., 2017. Late Miocene Uplift of the Tian Shan and Altai and Reorga-
 341 nization of Central Asia Climate. *GSA Today* 27.
 342 Caves, J.K., Moragne, D.Y., Ibarra, D.E., Bayshashov, B.U., Gao, Y., Jones, M.M., Zhamangara, A.,
 343 Arzhannikova, A.V., Arzhannikov, S.G., Chamberlain, C.P., 2016. The Neogene de-greening of Cen-
 344 tral Asia. *Geology* 44, 887–890.
 345 Caves, J.K., Winnick, M.J., Graham, S.A., Sjostrom, D.J., Mulch, A., Chamberlain, C.P., 2015. Role
 346 of the westerlies in Central Asia climate over the Cenozoic. *Earth and Planetary Science Letters* 428,
 347 33–43.
 348 Dercourt, J., Ricou, L.E., Vrielynck, B., 1993. *Atlas Tethys, Paleoenvironmental Maps: Explanatory*
 349 *Notes*. Gauthier-Villars.
 350 Diefendorf, A.F., Mueller, K.E., Wing, S.L., Koch, P.L., Freeman, K.H., 2010. Global patterns in leaf
 351 ^{13}C discrimination and implications for studies of past and future climate. *Proceedings of the National*
 352 *Academy of Sciences* 107, 5738–5743.

353 Dupont-Nivet, G., Hoorn, C., Konert, M., 2008. Tibetan uplift prior to the Eocene-Oligocene climate
354 transition: Evidence from pollen analysis of the Xining Basin. *Geology* 36, 987.

355 Dupont-Nivet, G., Krijgsman, W., Langereis, C., Abels, H., Dai, S., Fang, X., 2007. Tibetan plateau
356 aridification linked to global cooling at the Eocene–Oligocene transition. *Nature* 101, 45.

357 Edwards, E.J., Osborne, C.P., Strömberg, C.A., Smith, S.A., Consortium, C.G., et al., 2010. The origins
358 of C4 grasslands: integrating evolutionary and ecosystem science. *science* 328, 587–591.

359 Favre, A., Päckert, M., Pauls, S.U., Jähnig, S.C., Uhl, D., Michalak, I., Muellner-Riehl, A.N., 2015. The
360 role of the uplift of the Qinghai-Tibetan Plateau for the evolution of Tibetan biotas. *Biological Reviews*
361 90, 236–253.

362 Fortelius, M., Eronen, J.T., Kaya, F., Tang, H., Raia, P., Puolamäki, K., 2014. Evolution of Neogene
363 mammals in Eurasia: environmental forcing and biotic interactions. *Annual Review of Earth and*
364 *Planetary Sciences* 42, 579–604.

365 France-Lanord, C., Derry, L., Michard, A., 1993. Evolution of the Himalaya since Miocene time: isotopic
366 and sedimentological evidence from the Bengal Fan. *Geological Society, London, Special Publications*
367 74, 603–621.

368 Goodwin, D.H., Cohen, A.N., Roopnarine, P.D., 2010. Forensics on the half shell: a sclerochronological
369 investigation of a modern biological invasion in San Francisco Bay, United States. *Palaios* 25, 742–
370 753.

371 Goodwin, D.H., Flessa, K.W., Schöne, B.R., Dettman, D.L., 2001. Cross-calibration of daily growth
372 increments, stable isotope variation, and temperature in the Gulf of California bivalve mollusk *Chione*
373 *cortezi*: implications for paleoenvironmental analysis. *Palaios* 16, 387–398.

374 Gordon, C., Cooper, C., Senior, C.A., Banks, H., Gregory, J.M., Johns, T.C., Mitchell, J.F., Wood, R.A.,
375 2000. The simulation of SST, sea ice extents and ocean heat transports in a version of the Hadley
376 Centre coupled model without flux adjustments. *Climate Dynamics* 16, 147–168.

377 Guo, Z., Ruddiman, W.F., Hao, Q., Wu, H., Qiao, Y., Zhu, R.X., Peng, S., Wei, J., Yuan, B., Liu, T.,
378 2002. Onset of Asian desertification by 22 Myr ago inferred from loess deposits in China. *Nature* 416,
379 159–163.

380 Huber, M., Goldner, A., 2012. Eocene monsoons. *Journal of Asian Earth Sciences* 44, 3–23.

381 James, N.P., Dalrymple, R.W., 2010. *Facies models*. Geolog. Assoc. of Canada.

382 Kent-Corson, M.L., Ritts, B.D., Zhuang, G., Bovet, P.M., Graham, S.A., Page Chamberlain, C., 2009.
383 Stable isotopic constraints on the tectonic, topographic, and climatic evolution of the northern margin
384 of the Tibetan Plateau. *Earth and Planetary Science Letters* 282, 158–166.

385 Kohn, M.J., 2010. Carbon isotope compositions of terrestrial C3 plants as indicators of (paleo) ecology
386 and (paleo) climate. *Proceedings of the National Academy of Sciences* 107, 19691–19695.

387 Kraatz, B.P., Geisler, J.H., 2010. Eocene–Oligocene transition in Central Asia and its effects on mam-
388 malian evolution. *Geology* 38, 111–114.

389 Licht, A., Dupont-Nivet, G., Pullen, A., Kapp, P., Abels, H., Lai, Z., Guo, Z., Abell, J., Giesler, D.,
390 2016. Resilience of the Asian atmospheric circulation shown by Paleogene dust provenance. *Nature*
391 *Communications* 7.

392 Licht, A., Van Cappelle, M., Abels, H., Ladant, J.B., Trabucho-Alexandre, J., France-Lanord, C., Don-
393 nadiou, Y., Vandenberghe, J., Rigaudier, T., Lécuyer, C., et al., 2014. Asian monsoons in a late Eocene
394 greenhouse world. *Nature* 513, 501–506.

395 Liu, Y., Wu, G., Hong, J., Dong, B., Duan, A., Bao, Q., Zhou, L., 2012. Revisiting Asian monsoon
396 formation and change associated with Tibetan Plateau forcing: II. Change. *Climate dynamics* 39,
397 1183–1195.

398 Lunt, D.J., Farnsworth, A., Loftson, C., Foster, G.L., Markwick, P., O’Brien, C.L., Pancost, R.D., Robin-
399 son, S.A., Wrobel, N., 2016. Palaeogeographic controls on climate and proxy interpretation. *Climate*
400 *of the Past* 12, 1181–1198.

401 Macaulay, E.A., Sobel, E.R., Mikolaichuk, A., Wack, M., Gilder, S.A., Mulch, A., Fortuna, A.B., Hynek,
402 S., Apayarov, F., 2016. The sedimentary record of the Issyk Kul basin, Kyrgyzstan: climatic and
403 tectonic inferences. *Basin Research* 28, 57–80.

404 Markwick, P.J., Valdes, P.J., 2004. Palaeo-digital elevation models for use as boundary conditions in
405 coupled ocean–atmosphere GCM experiments: a Maastrichtian (late Cretaceous) example. *Palaeo-*
406 *geography, Palaeoclimatology, Palaeoecology* 213, 37–63.

407 Molnar, P., Boos, W., Battisti, D., 2010. Orographic controls on climate and paleoclimate of Asia:
 408 thermal and mechanical roles for the Tibetan Plateau. *Annual Review of Earth and Planetary Sciences*
 409 38, 77–102.

410 Mouchi, V., de Rafélis, M., Lartaud, F., Fialin, M., Verrecchia, E., 2013. Chemical labelling of oyster
 411 shells used for time-calibrated high-resolution Mg/Ca ratios: A tool for estimation of past seasonal
 412 temperature variations. *Palaeogeography, Palaeoclimatology, Palaeoecology* 373, 66–74.

413 Nie, J., Stevens, T., Rittner, M., Stockli, D., Garzanti, E., Limonta, M., Bird, A., Andò, S., Vermeesch,
 414 P., Saylor, J., et al., 2015. Loess plateau storage of northeastern Tibetan plateau-derived Yellow River
 415 sediment. *Nature communications* 6.

416 Pagani, M., Zachos, J.C., Freeman, K.H., Tipple, B., Bohaty, S., 2005. Marked decline in atmospheric
 417 carbon dioxide concentrations during the Paleogene. *Science* 309, 600–603.

418 Quan, C., Liu, Y.S.C., Utescher, T., 2012. Eocene monsoon prevalence over china: a paleobotanical
 419 perspective. *Palaeogeography, palaeoclimatology, palaeoecology* 365, 302–311.

420 Ramstein, G., Fluteau, F., Besse, J., Joussaume, S., 1997. Effect of orogeny, plate motion and land-sea
 421 distribution on Eurasian climate change over the past 30 million years. *Nature* 386, 788–795.

422 Sobel, E.R., Chen, J., Heermance, R.V., 2006. Late Oligocene–Early Miocene initiation of shortening in
 423 the Southwestern Chinese Tian Shan: implications for Neogene shortening rate variations. *Earth and*
 424 *Planetary Science Letters* 247, 70–81.

425 Suits, N.S., Denning, A.S., Berry, J., Still, C., Kaduk, J., Miller, J., Baker, I.T., 2005. Simulation of
 426 carbon isotope discrimination of the terrestrial biosphere. *Global Biogeochemical Cycles* 19.

427 Sun, X., Wang, P., 2005. How old is the Asian monsoon system? *Palaeobotanical records from China.*
 428 *Palaeogeography, Palaeoclimatology, Palaeoecology* 222, 181–222.

429 Tindall, J., Flecker, R., Valdes, P., Schmidt, D., Markwick, P., Harris, J., 2010. Modelling the oxygen
 430 isotope distribution of ancient seawater using a coupled ocean-atmosphere GCM: implications for
 431 reconstructing early Eocene climate. *Earth and Planetary Science Letters* 292, 265–273.

432 Tindall, J., Valdes, P., Sime, L.C., 2009. Stable water isotopes in HadCM3: Isotopic signature of El Niño–
 433 Southern Oscillation and the tropical amount effect. *Journal of Geophysical Research: Atmospheres*
 434 (1984–2012) 114.

- 435 Winnick, M.J., Chamberlain, C.P., Caves, J.K., Welker, J.M., 2014. Quantifying the isotopic ‘continental
436 effect’. *Earth and Planetary Science Letters* 406, 123–133.
- 437 Wu, G., Liu, Y., Dong, B., Liang, X., Duan, A., Bao, Q., Yu, J., 2012. Revisiting Asian monsoon
438 formation and change associated with Tibetan Plateau forcing: I. Formation. *Climate dynamics* 39,
439 1169–1181.
- 440 Zachos, J., Pagani, M., Sloan, L., Thomas, E., Billups, K., 2001. Trends, rhythms, and aberrations in
441 global climate 65 Ma to present. *Science* 292, 686.
- 442 Zhang, Z., Flatøy, F., Wang, H., Bethke, I., Bentsen, M., Guo, Z., 2012. Early Eocene Asian climate
443 dominated by desert and steppe with limited monsoons. *Journal of Asian Earth Sciences* 44, 24–35.
- 444 Zhang, Z., Wang, H., Guo, Z., Jiang, D., 2007. What triggers the transition of palaeoenvironmental
445 patterns in China, the Tibetan Plateau uplift or the Paratethys Sea retreat? *Palaeogeography, Palaeo-*
446 *climatology, Palaeoecology* 245, 317–331.
- 447 Zheng, H., Wei, X., Tada, R., Clift, P.D., Wang, B., Jourdan, F., Wang, P., He, M., 2015. Late Oligocene–
448 Early Miocene birth of the Taklimakan Desert. *Proceedings of the National Academy of Sciences* 112,
449 7662–7667.

Figure 1: Localisation of study area. Modern topographic map of the study area showing the main tectonic features and the localisation of the sampled sedimentary sections (AL: Alai Valley, 39.6°N, 72.4°E; MS: Mine, 39°51'N, 74°32'E; AT: Aertashi, 37°58'N, 76°33'E; KY: Keyliand, 37°27'N, 77°86'E). The position of the sites experienced no statistically significant latitudinal tectonic motion since the time of deposition (Bougeois et al., 2014).

Figure 2: Paleogene depositional environments. Nodular (a.) and massive (b.) gypsum deposits indicate playa, salinas and sabkha environments typical from semi-arid climate (alternation of floods events and dessication periods). Tidal flat with neap-spring tide alternations (c.) and littoral barrier with rippled bioclastic grainstone (d.) indicate a low energy carbonate ramp environment. Fossil oysters lived in subtidal zone where *O. (T.) strictiplicata* (e.) built bioherms (patch reefs) and *S. buhsii* (f.) stood isolated in blue marls with bryozoa, serpulids, echinoids (see frame), foraminiferas and fishes attesting for a more quiet open marine environment.

Figure 3: Sedimentary log of Aertashi section (facies associations are described in supplementary material). Alternation of marine and continental deposits and record several sea incursions and subsequent retreats. Associated carbonate geochemistry analyses ($\delta^{18}\text{O}$, $\delta^{13}\text{C}$, % in CaCO_3) are reported throughout the section.

Figure 4: Sedimentary log of Mine section (facies associations are described in supplementary material). Associated carbonate geochemistry analyses are reported throughout the section.

Figure 5: Seasonal data. Comparison between (a) Eocene proxy data (see details in Fig. 6) and (b) monthly data provided by climatic simulations. (c) Average of monthly modern air temperature and precipitation in Kashgar (China) from 1951 to 1993 (Baker et al., 1995) and in Bukhara (Uzbekistan) from 1982 to 2012 (www.climate-data.org) ; modern oxygen stable isotopic composition of precipitation in Hotan (China) (Araguás-Araguás et al., 1998).

Figure 6: Polished cross sections of *O. (T.) strictiplicata* (3 top samples) and *S. buhsii* (3 bottom samples) revealing annual growth bands (arrows indicate growth direction). Cathodoluminescence microscopy analyses (CL) show no diagenetic alteration. Grey bands correspond to dark bands in the shells which are built during the coldest months of the year. Black lines indicate main paths drilled for $\delta^{18}\text{O}$ analyses. Orange lines indicate transects followed by laser for Mg/Ca analyses. Mg/Ca is given in mmol/mol, $\delta^{18}\text{O}_c$ in ‰ VPDB, $\delta^{18}\text{O}_{sw}$ in ‰ SMOW, temperatures in ° Celsius. Distance is measured from the first drilled micro-sample for $\delta^{18}\text{O}$. Temperatures are estimated from elemental composition (Mouchi et al., 2013) and Apparent temperatures from $\delta^{18}\text{O}_c$ (Anderson and Arthur, 1983), with a constant $\delta^{18}\text{O}_{sw} = 0.44\text{‰}$. See main text for the calculation of $\delta^{18}\text{O}_{sw}$ from Mg-deduced temperature and $\delta^{18}\text{O}_c$.

Figure 7: Eocene paleogeographic maps of Asia showing the interpreted general summer (a) and winter (b) wind patterns 40 Ma ago, according to this study and previous numerical simulations (Huber and Goldner, 2012; Zhang et al., 2012; Licht et al., 2014). The descending branch of the Hadley cell is responsible for a broad and large band of semi-arid to arid climate (Zhang et al., 2012), which is intensified during summer months in Central Asia due to the Foehn effect induced by the emerging Tibetan plateau. In winter, aridity is less strong due to precipitation associated with the westerly winds over the Proto-Paratethys sea that were not yet impeded by the Pamir and Tian Shan.

Figure 8: Eocene to Pliocene stable isotopic composition ($\delta^{13}\text{C}$ and $\delta^{18}\text{O}$) for Tarim Basin carbonaceous sediment from this study (Aertashi and Mine sections, see supplementary material) and Kent-Corson et al. (2009) data (Aertashi section). Crosses show averages and standard deviation for each time interval. Gray arrow shows the trend through geological times.

Table 1: Carbon ($\delta^{13}\text{C}$ in ‰ VPDB) and oxygen ($\delta^{18}\text{O}$ in ‰ VPDB) isotopic composition from marine and pedogenic carbonates in Aertashi section.

Table 2: Carbon ($\delta^{13}\text{C}$ in ‰ VPDB) and oxygen ($\delta^{18}\text{O}$ in ‰ VPDB) isotopic composition from marine and pedogenic carbonates in Mine section.