

Synchronization of the astronomical time scales in the Early Toarcian: A link between anoxia, carbon-cycle perturbation, mass extinction and volcanism

Fatima-Zahra Ait-Itto, Mathieu Martinez, Gregory Price, Abdellah Ait Addi

► To cite this version:

Fatima-Zahra Ait-Itto, Mathieu Martinez, Gregory Price, Abdellah Ait Addi. Synchronization of the astronomical time scales in the Early Toarcian: A link between anoxia, carbon-cycle perturbation, mass extinction and volcanism. Earth and Planetary Science Letters, 2018, 493, pp.1-11. 10.1016/j.epsl.2018.04.007. insu-01777678

HAL Id: insu-01777678 https://insu.hal.science/insu-01777678

Submitted on 18 Jul2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	Synchronization of the astronomical time scales in the Early Toarcian: a link between
2	anoxia, carbon-cycle perturbation, mass extinction and volcanism
3	
4	Fatima-Zahra Ait-Itto ^{a,*} , Mathieu Martinez ^{b,c} , Gregory D. Price ^d , Abdellah Ait Addi ^a
5	
6	^a Cadi Ayyad University, Faculty of Sciences and Techniques, Geosciences and
7	Environmental research's Laboratory, P.O. Box 549, Marrakech, Morocco
8	^b MARUM: Center for Marine Environmental Sciences, Universität Bremen, Leobenerstr.,
9	28359 Bremen, Germany.
10	^c Univ Rennes, CNRS, Géosciences Rennes - UMR 6118, 35000 Rennes, France
11	^d School of Geography, Earth and Environmental Sciences Plymouth University, Plymouth,
12	Devon PL4 8AA, UK
13	

14 Abstract

The Late Pliensbachian-Early Toarcian is a pivotal time in the Mesozoic era, marked by 15 pronounced carbon-isotope excursions, biotic crises and major climatic and oceanographic 16 17 changes. Here we present new high-resolution carbon-isotope and magnetic-susceptibility measurements from an expanded hemipelagic Late Pliensbachian-Early Toarcian section from 18 the Middle Atlas Basin (Morocco). Our new astronomical calibration allows the construction 19 20 of an orbital time scale based on the 100-kyr eccentricity cycle. The Early Toarcian Polymorphum Zone contains 10 to 10.5 repetitions of the 100-kyr eccentricity both in the 21 carbon-isotope and the magnetic-susceptibility data, leading to an average duration of 1.00 \pm 22 0.08 myr. We also show that the Late Pliensbachian-Early Toarcian global carbon-cycle 23 perturbation has an average duration of 0.24 ± 0.02 myr. These durations are comparable to 24 previous astrochronological time scales provided for this time interval in the most complete 25

sections of the Tethyan area, and longer than what has been provided in condensed sections. 26 27 Anchoring this framework on published radiometric ages and astrochronological time scales, we estimate that the carbon-cycle perturbation of the Late Pliensbachian-Early Toarcian 28 29 corresponds with the early phase of the Karoo and Chonke Aike large igneous provinces. Likewise, our new age constraints confirm that the Toarcian oceanic anoxic event is 30 synchronous to the main phase of the Ferrar volcanic activity. Thus, these successive and short 31 32 phases of the volcanic activity may have been at the origin of the successive phases of the mass extinctions observed in marine biotas in the Pliensbachian and Toarcian times. 33

34 Keywords:

35 Astrochronology; Toarcian; carbon cycle; Middle Atlas; oceanic anoxic events

36

37 **1. Introduction**

The Early Toarcian (184.15 – 174.1 Ma) was a time of global warming events (Dera et al., 2011), mass extinctions (Little and Benton, 1995), and pronounced negative carbon-isotope excursions recorded in marine carbonates and organic matter, brachiopods, biomarkers and fossil wood (Hesselbo et al., 2007; Suan et al., 2008a; Ait-Itto et al., 2017). These events coincide with carbonate production demises (Bassoullet and Baudin, 1994; Wilmsen and Neuweiler, 2008) and widespread oceanic anoxia (Jenkyns, 1988; Hesselbo et al., 2007).

Although large amounts of data have been produced, the timing and rhythms of the environmental perturbations are still debated, leading to controversies on the mechanisms at the onset of the climatic changes in the Early Toarcian (e.g. Kemp et al., 2005; Suan et al., 2008b). For instance, Martinez et al (2017) assess the duration of Pliensbachian-Toarcian event (P-To event) at 0.18 to 0.27 myr, while in the Peniche section (Lusitanian Basin, Portugal) the duration of this event is evaluated as 0.05 myr (Suan et al., 2008b). Similarly, the estimates of duration of the Polymorphum Zone vary from 90 kyr to 1.15 myr (Matiolli and Pittet, 2004; Suan et al.,

2008b, Boulila et al., 2014; Huang and Hesselbo, 2014; Ruebsam et al., 2014; Martinez et al., 51 52 2017). The main reason of these differences is the occurrence of four discontinuity events identified by correlations of δ^{13} C and sedimentological features throughout the Tethyan 53 margins, which are due to changes in the sea level observed in the earliest Toarcian (Pittet et 54 al., 2014). The section studied here in Issouka, Morocco, is expanded compared to the 55 56 sedimentary series from the northern Tethyan margin. It provides the opportunity to fill the gap 57 in the time scale of the Toarcian Stage and to establish a detailed chronology of the succession of the environmental disturbances occurring in the Early Toarcian. 58

59

60 2. Geological setting

The geological history of Morocco was influenced by two important events, starting in the early 61 Mesozoic with the opening of the north Atlantic and western Tethys and the collision of Africa 62 and Europe during the middle Cenozoic (Michard, 1976). These events led to the formation of 63 64 fault-bounded basins, which are made up of several smaller depocenters, separated by synsedimentary highs (Studer and Du Dresnay, 1980). The Middle Atlas is one of these small 65 fault-bounded basins (Fig. 1). It constitutes a part of a Meso-Cenozoic intracontinental chain, 66 67 namely the Moroccan Atlas (Michard, 1976). The Middle Atlas of Morocco is structurally dominated by four NE-SW trending anticlines and is mainly constituted of Lower and Middle 68 Jurassic formations (Du Dresnay, 1971). The basin is bounded by the Saïs Plain and the front 69 of the Rifian Nappes in the North, by the Guercif Basin in the northeast, by the Moulouya Plain 70 to the southeast, and by the Hercynian Central Massif in the West. 71

The Middle Atlas Basin is deep in the center and shallows towards the northern and southern basin margins (Du Dresnay, 1971). The study area during the Early Toarcian was located at a palaeolatitude of ~20°N (Bassoulet et al., 1993) (Fig. 1A). The sedimentary evolution and palaeogeographic differentiation is controlled by tectonic activity, combined with

the rate of sedimentation and global eustatic variations (Wilmsen and Neuweiler, 2008). The 76 rapid transition from shallow marine carbonates to hemipelagic marls has been taken to reflect 77 a major deepening phase across the entire Middle and High Atlas area (Wilmsen and Neuweiler, 78 2008). The Pliensbachian-Toarcian transition coincides with a dislocation of the Lower 79 Jurassic carbonate platform (Lachkar et al., 2009; Ait Addi and Chafiki, 2013) with Toarcian 80 deposits dominated by marls lying upon Upper Pliensbachian shallow marine limestones and 81 calcareous marls. This drowning episode is linked to the eustatic sea-level rise of the Early 82 Toarcian described in Europe and Africa (e.g. Hallam, 1997). 83

The biostratigraphy in the Middle Atlas has been established with ammonites (El 84 85 Hammichi et al., 2008) with further biostratigraphic data provided by benthic foraminifera (Bejjaji et al., 2010). The ammonite zonation in the Issouka section is based on the 86 Mediterranean zonation. Notably, the ammonites *Emaciaticeras emaciatum* of the Emaciatum 87 Zone, Dactylioceras polymorphum of the Toarcian Polymorphum Zone and Hildaites in the 88 Semicelatum Zone have been identified (e.g. El Hammichi et al., 2008). Furthermore, the 89 occurrence of the benthic foraminifera Lenticulina sublaevis in the Issouka section is correlated 90 by Bejjaji et al. (2010) to the Emaciatum ammonite zone of the Pliensbachian, whilst 91 Lenticulina bochardi and Lenticulina toarcense are correlated with the Toarcian Polymorphum 92 93 Zone and Lenticulina obonensis with the Serpentinus Zone.

The Issouka section is situated near the village of Issouka, ~25 km southwest of Immouzer Marmoucha, in the Middle Atlas (N33° 26' 55.56"; W4° 20' 33.83"; Fig. 1). The section begins with centimeter thick limestone–marl alternations (Fig. 2). The limestone beds contain a rich ammonite fauna, with also belemnites, echinoids and brachiopods. Foraminifera suggested a Late Pliensbachian age (Bejjaji et al., 2010; Fig. 2A). The early Toarcian succession starts with green marls and marl–limestone alternations, rich in foraminifera, belemnites, echinoids and gastropods. The Toarcian deposits are generally hemipelagic and correspond to
deep marine environments (Bejjaji et al., 2010; Fig. 2B).

102

103 **3. Material and methods**

A total of 430 bulk-rock samples were collected with an even sample distance of 10 cm in the Issouka section. The samples were recovered from up to 15 cm below the surface, to minimize the effects of surface weathering. The sampled interval encompasses the end of the Pliensbachian Stage to the lowermost part of the Levisoni ammonite Zone. Such a range allows the Polymorphum Zone to be entirely sampled. The bulk-rock samples were then powdered using a metal ring grinder and analyzed for stables isotopes, magnetic susceptibility and calcium carbonate contents.

111

112 **3.1.** Carbon isotopes

113 A total of 430 samples were analyzed for stable isotopes at the University of Plymouth. Using 114 200 to 300 micrograms of carbonate, stable isotope data were generated on a VG Optima mass 115 spectrometer with a Gilson autosampler. Isotope ratios were calibrated using NBS19 standards 116 and are given in δ notation relative to the Vienna Pee Dee Belemnite (VPDB). Reproducibility 117 was generally better than 0.1 ‰ for samples and standard material.

118

119 **3.2. Magnetic susceptibility**

120 The magnetic susceptibility (MS) of the 430 powdered samples was measured with a 121 Kappabridge KLY-3. The samples are placed in small plastic cubes of 10 cm³ and then 122 introduced inside the instrument using the "pick-up" unit. We measured the cube empty and 123 then with samples for blank correction. The blank-corrected MS values are normalized to their 124 mass and volume. The resulting values are reported as mass-specific MS (m³.kg⁻¹).

126 **3.3. Calcium carbonate content**

The samples were analyzed for their calcium carbonate content using a Bernard calcimeter at
the Geosciences and Environment Laboratory, Cadi Ayyad University. The values are given
with a precision between 1 and 5% (Lamas et al., 2005).

130

131 **3.4. Spectral analysis**

Prior to spectral analysis, the long-term trend of both series was measured and subtracted from 132 the series to ensure stationarity. The detrend procedures removed the high powers in the 133 134 frequencies close to 0, while not creating new spectral peaks in the low frequencies (see Vaughan et al., 2015). Best-fit linear trend and LOWESS smoothing curves (Cleveland et al., 135 1979) were applied. The exact coefficients depend on each of the series and are detailed in the 136 results. Following the removal of this long-term trend, the multi-taper spectral analyses, using 137 three 2π prolate tapers (2π -MTM) with robust red-noise modelling (Mann and Lees, 1996) 138 modified in Meyers (2014) were applied to evaluate lithological cycles as a possible record of 139 astronomically forced sedimentation following the LOWSPEC approach (Meyers, 2012). The 140 141 significance levels were then calculated assuming a chi-square distribution of the spectral 142 background. We used the time-frequency weighted fast Fourier transforms (T-F WFFT; Martinez et al., 2015) to characterize the evolution of the periods through the sedimentary 143 series. Taner low-pass and band-pass filters (Taner, 2003) were then applied to isolate each 144 145 signal interpreted as an orbital forcing and calculate the duration of the Polymorphum Zone by cycle counting. Orbital tuning procedures are then used to anchor the sedimentary cycles to 146 their corresponding orbital periods, allowing depth-time conversions to be done. Durations are 147 then estimated by calculating the difference of relative ages. Error margins of each of the time 148

intervals include the difference between the average and the calculated durations and theuncertainty in the orbital frequencies.

151

152 **4. Results**

153 **4.1. Carbonate carbon isotopes**

The carbon-isotope profile of the Issouka section shows relatively high values in the 154 Pliensbachian part of the section, ranging from +1.7‰ to +2.7‰ (Fig. 3A). From 5 m to 5.6 m, 155 the δ^{13} C decreases from 2.5% to -0.6%. This abrupt decrease of 3% in the δ^{13} C observed at 156 157 the base of the Toarcian is related to the Pliensbachian-Toarcian event (P-To event; Suan et al., 2008a; Ait-Itto et al., 2017). The δ^{13} C values then remain low from 5 m to 23 m. In this interval, 158 the values show gentle fluctuations from 5 to 17 m. At 17 m, the δ^{13} C abruptly increase by 1‰ 159 before they gently decrease until 23 m to -2‰. The values then progressively increase from 23 160 m to 30 m, changing from -2‰ to 1‰. Above 30 m to the top of the section, the values slightly 161 162 increase from 1 to 2‰.

163 **4.2. Magnetic susceptibility**

The magnetic susceptibility curve (Fig. 3E) firstly shows from the base of the series to level 5.1 m low values, ranging from 9.65×10^{-9} to 3.15×10^{-8} m³.kg⁻¹. This interval corresponds to the limestone-dominated part of the formation of Pliensbachian age. From 5.2 m to the top of the series, MS values increase with an average value 1.04×10^{-7} m³.kg⁻¹. The interval from 27 m to 35 m shows lower values of the MS in a more carbonated interval. Calcium carbonate content and MS values display a strong inverse correlation (r = -0.91, Fig. 4), indicating that the lithology mainly controls the MS variations.

171 **4.3.** Calcium carbonate content

The carbonate content series (Fig. 3D) firstly show high values (>80% on average) from the
base of the series to 5.1 m, in the interval of the limestone beds of the Pliensbachian. From 5.2

m to 8.4 m, the values rapidly decrease from 91% to 8.6%. From 5.2 m to 18 m, the CaCO₃ values remain low with average values around 22%. The transition between the first to the second interval is more progressive than observed in the δ^{13} C series (Fig. 3A). From 18 m to 42.9 m CaCO₃ series show high-amplitude cycles and high-frequency fluctuations from 60% and 20% linked to the limestone-marl alternations (Fig. 3D).

179 **4.4. Spectral analyses**

180 **4.4.1. Carbon-isotope signal**

The δ^{13} C shows a rapid decrease at 5.2 m, and a rapid increase at 16.6 m (Fig. 3A). The series 181 182 was stationarized using a best-fit linear trend from 0 to 5.1 m, a LOWESS regression with a coefficient of 0.6 from 5.2 to 16.5 m and a LOWESS regression with a coefficient of 0.3 from 183 16.6 to 42.9 m. The LOWESS coefficients have been chosen in such a way that they 184 approximatively cover the same length of the section (i.e. ~7 m on average), which represents 185 ca. 20% of the Toarcian part. This choice decreased the high powers at frequencies near 0, 186 187 while not creating spurious spectral peaks in the low frequencies (Vaughan et al., 2015). The three subseries were then standardized (average = 0, standard deviation = 1) and stacked. Two 188 points at 11.7 m and 12 m showing outstandingly low values when detrending may bias the 189 190 spectrum in the high frequencies and have been removed from the series before detrending. The evolutive spectrum of the δ^{13} C series shows a high-power spectral band in the low frequencies 191 evolving having a period of 4.1 to 4.2 m from the base of the series to level 24 m (Fig. 3C). The 192 period of this band then decreases to 3 m, from level 24 m to level 31 m, and stabilizes at 3 m 193 from level 31 m to the top of the series. These changes in the period of the ~4-m peak likely 194 195 reflects variations in the sedimentation rate. These distort the sedimentary record of the orbital cycles by decreasing the *p*-value of the low frequency and flattening the spectrum at high 196 197 frequencies (Weedon, 2003; Kemp, 2016; Martinez et al., 2016). We thus decided to analyze the frequency content of two sub-intervals of the series in which variations in the sedimentation 198

rate are limited. The range of each of the sub-intervals is given in Fig. 3C. We compared the frequencies of the carbon isotopes in each interval with the orbital frequencies calculated for the Toarcian (Waltham, 2015), using the average spectral misfit (ASM; Meyers and Sageman, 2007). The ASM method compares the sedimentary frequencies converted into time to the orbital frequencies and provides a quantitative assessment on the most likely sedimentation rate in each sub-interval (see supplementary materials).

Spectra of Interval 1, from levels 0 to 31 m, show significant periods above the 99%
confidence level (CL) at 4.2 m (frequency: 0.2404 cycles.m⁻¹), and above the 95% CL at 1.2 m
(frequency: 0.8013 cycles.m⁻¹), 0.83 m (frequency: 1.202 cycles.m⁻¹), and 0.31 m (Fig. 5A, C).
Spectra of Interval 2, from levels 24 m to the top of the series show significant peaks above the
95% CL at 3.2 m (frequency: 0.3158 cycles.m⁻¹), 0.63 m (1.579 cycles.m⁻¹) and 0.33 m (Fig. 5B, D).

211 As noted above, various astrochronological frameworks estimate the duration of the Polymorphum Zone from 0.09 myr to 1.15 myr (Mattioli and Pittet, 2004; Suan et al., 2008b; 212 213 Huang and Hesselbo, 2014; Boulila et al., 2014; Martinez et al., 2017; Boulila and Hinnov, 2017), implying a sedimentation rate varying from 400 m/myr to 31 m/myr. When converting 214 m-cycles to time-cycles, the peaks at ~0.3 m have periods below the Milankovitch band 215 216 whatever the sedimentation rate considered, so were excluded from the ASM analysis. Testing ASM over 500 sedimentation rates ranging from 0.4 cm/kyr to 40 cm/kyr leads to a most likely 217 sedimentation rate of 3.9 cm/kyr (39 m/myr) in Interval 1, and to a most likely sedimentation 218 rate of 3.2 cm/kyr (32 m/myr) in Interval 2 (Fig. 5E, F). The H0-significance level at the most 219 220 likely sedimentation rate is below the critical significance level below which we can consider without ambiguity these spectra reflect an orbital forcing. In Interval 1, peaks at 4.2 m, 1.2 m 221 and 0.83 m are respectively due to the 100-kyr eccentricity, the obliquity and the precession 222 cycles. Notice than the peak of 1.2 m leads to an obliquity period which is shorter than expected 223

in the theory (here 29.4 kyr vs. 37.5 kyr in Waltham, 2015). In Interval 2, peaks at 3.2 m and
0.63 m are respectively due to the 100-kyr eccentricity and the precession cycles. In addition,
the peak at 0.33 m has a mean period of 10 kyr, which corresponds to the semi-precession, the
first harmonic of the precession (Berger et al., 2006).

The evolutive spectral analysis indicates the 100-kyr cycle is the most continuous throughout the studied series (Fig. 3C). We filtered the 100-kyr eccentricity recorded in the δ^{13} C series and used it as target cycle for calibrating the series in time. We filtered the 100-kyr cycle using a Taner low-pass filter with a frequency cut of 0.3953 cycles.m⁻¹ and a roll-off rate of 10³⁶. The output filter indicates that the Polymorphum Zone contains 10 repetitions of the 100-kyr eccentricity cycle (Fig. 3B).

234

235 **4.4.2. Magnetic susceptibility**

The MS series is marked by a rapid increase of values from 5.0 m to 6.8 m. The trend of the 236 237 series is then gentler. The series was stationarized using a best-fit linear regression from 0 to 5.2 m and a LOWESS smoothing with a coefficient of 0.2 in the remainder of the series, which 238 represents 7.5 m of the section. The two subseries were then standardized (average = 0, standard 239 240 deviation = 1) and stacked. The evolutive spectrum of the MS series shows a band in the low frequencies evolving from 3.1 m to 3.7 m in the lower part of the series, from the base of the 241 series to level 20 m (Fig. 3G). Then, this period decreases to 2.9 m from level 26 m to the top 242 of the series. 2π -MTM spectra have been generated from Interval 1 (from the base of the series 243 to level 26 m), and from Interval 2 (from 21 m to the top of the series) (Fig. 6). Spectra of 244 245 Interval 1 show spectral peaks over the 99% CL at 3.3 m and 0.38 m, and peaks over the 95% CL at 0.81 m and 0.23 m (Fig. 6A, C). Spectra of Interval 2 show peaks over the 99% CL at 2.9 246 m, 2.0 m, 0.27 m and 0.22 m, and a peak exceeding the 95% CL at 1.0 m (Fig. 6B, D). 247

Assuming from ASM applied on the δ^{13} C spectra that Interval 1 has a mean sedimentation rate of 39 m/myr, the peaks of 3.3 m, 0.81 m and 0.38 m are attributed to the 100-kyr eccentricity, the precession and the semi-precession, respectively. Assuming from ASM that Interval 2 has a mean sedimentation rate of 32 m/myr, the peak of 2.9 m, 1.0 m and 0.27 m are respectively attributed to the 100-kyr eccentricity, the obliquity and the semiprecession (Fig. 6).

In Interval 1, the period of the 100-kyr cycle is lower in the MS series than in the δ^{13} C. On the evolutive spectral analyses, this is particularly obvious in the first 10 m of the series, when the lithology evolves from the limestones of the Pliensbachian to the marls of the Toarcian (Fig. 3). The 100-kyr eccentricity recorded in the MS series was isolated using a Taner lowpass filter with a frequency cut of 0.4070 cycles.m⁻¹ and a roll-off rate of 10³⁶ (Fig. 3). The output filter indicates that the Polymorphum Zone clearly contains ~10.5 repetitions of the 100kyr eccentricity cycle compared to the 10 repetitions observed in the δ^{13} C (Fig. 3).

261

262 **5. Discussion**

5.1. Astrochronology of the the Polymorphum Zone

The δ^{13} C and the MS signals both display 12 complete short eccentricity cycles along the 264 studied section. In the astronomical models computed for the last 20 myr (Laskar et al., 2004), 265 the average duration of 12 consecutive short eccentricity cycles is 1149 ± 63 kyr (2σ), so that 266 the average duration of a short eccentricity cycle 95.8 \pm 5.2 kyr (2 σ). Assuming a constant 267 sedimentation rate between two successive anchor points of the 100-kyr cycle obtained from 268 the filters of the δ^{13} C and the MS signals, the durations calculated of the Polymorphum Zone 269 range from 0.97 myr with the δ^{13} C signal, to 1.02 myr with the MS signal. In the spectra of the 270 untuned series, the period of the 100-kyr cycle is longer in the δ^{13} C than in the MS (Figs. 3, 5, 271 6). The difference is particularly noticeable in the first 10 m (Fig. 3). Below level 5 m, the filter 272

of 100-kyr cycle in the δ^{13} C is inversely correlated to the filter of the 100-kyr in the MS signal. 273 After 5 m, i.e. after the start of the P-To event, the two filters are in phase. The limestone beds 274 275 in the Pliensbachian of the atlasic basins originate from exports from the neritic environments, in which the δ^{13} C was higher than in pelagic environments (Wilmsen and Neuweiler, 2008). In 276 the Pliensbachian, the δ^{13} C is higher in limestone beds. The P-To Event corresponds to a demise 277 in the neritic carbonate production. Fluctuations in the δ^{13} C thus correspond to fluctuations in 278 the pelagic production, which is usually higher during the deposit of marlier intervals, when 279 280 detrital and nutrient increased the primary productivity (Mutterlose and Ruffell, 1999; Mattioli and Pittet, 2004). The demise of the carbonate platforms thus changed the phasing between the 281 MS and the δ^{13} C, which is a source of uncertainty in the calculation of the durations. Another 282 change of phasing occurs at 28 m, where the δ^{13} C and the CaCO₃ content increase. As the 283 phasing between the orbital configuration and the proxies is still unclear, we choose to retain 284 285 the average duration of 1.00 ± 0.08 myr for the Polymorphum Zone (the uncertainty includes the age model and the duration of a short eccentricity period, see supplementary materials). 286

The spectrum of the calibrated δ^{13} C shows a high power at 96 kyr (expected when 287 calibrated to the short eccentricity), 32.4 kyr (main obliquity), 24.1 and 19.7 kyr (precession), 288 and 13.2 to 11.0 kyr (half-precession) (Fig. 7A). In addition, a peak exceeding the 95% 289 confidence level appears at 400 kyr, which corresponds to the long-eccentricity cycle. The 290 291 spectrum of the calibrated MS shows a high power at 95.7 kyr (expected), 65.5 kyr, 31.9 kyr 292 (main obliquity), from 24.4 to 19.1 kyr (precession), and from 12.1 to 10.6 kyr (half precession) (Fig. 7B). In the δ^{13} C and MS signals, the precession cycle has slightly longer periods than 293 294 expected in the theory (Waltham, 2015). This may be the consequence of a precession cycle missing per short eccentricity. In the astronomical solutions, the amplitude of the precession 295 band is modulated by the 405-kyr and the 100-kyr eccentricity cycles, while the amplitude of 296 the 100-kyr cycle is modulated by the 400-kyr cycle (Laskar et al., 2004). The Taner band-pass 297

filter of the precession cycle in the δ^{13} C series was performed with frequency cuts of 3.583 x 298 10⁻² cycles/kyr and 5.417 x 10⁻² cycles/kyr, and a roll-off rate of 10³⁶ (Fig. 8B, E). We calculated 299 the amplitude modulation (AM) of the precession cycles using a Hilbert transform. The 300 spectrum of the AM of the precession filter shows cycles at 340 kyr (near the 405-kyr 301 eccentricity), and at 123 and 82.8 kyr (near the periods of 124 and 95 kyr of the short 302 eccentricity) (Fig. 8F). We filtered the 100-kyr band of the AM of the precession filter using a 303 Taner band-pass filter with frequency cuts of 6.250×10^{-3} cycles/kyr and 1.558×10^{-2} cycles/kyr 304 and a roll-off rate of 10^{36} (Fig. 8F). The AM of this filter shows a cycle at 340 kyr (Fig. 8G). 305 306 Similarly, the spectrum of the AM calculated on the direct filter of the 100-kyr band shows a cycle at 436 kyr (Fig. 8H). Thus, the AM of the precession and the 100-kyr eccentricity show 307 similar patterns as expected in the astronomical models. The filters of the two bands shows 11 308 cycles of short eccentricity within the Polymorphum Zone (Fig. 8C), while the various filters 309 of the 405-kyr cycles in the AM of the precession and the 100-kyr cycles show from 3 to 4 310 311 repetitions of the 405-kyr cycle (Fig. 8A, C, D). The difference between the direct filter and the filter of the AM is explained by the residence time of carbon in the ocean, which creates a delay 312 in the response of the δ^{13} C to the orbital forcing (Laurin et al., 2017). In this context of 313 hemipelagic sediments, with the limestone beds originating from exports from neritic 314 315 environments, the difference may also be due to the change of the carbonate source.

The duration of the Polymorphum Zone from this study is longer than the duration of ~0.8 myr assessed in the Peniche section (Suan et al., 2008b; Huang and Hesselbo, 2014) and the Foum Tillicht section (Martinez et al., 2017). The correlation of these sections led to a duration of a composite Polymorphum Zone of 0.9-1.0 myr, which agrees with the duration calculated here. In addition, the duration calculated here is in close agreement with the duration estimated from Central Italy, in which 11 bundles of marl-limestone alternations have been related to the 100-kyr eccentricity cycle (Mattioli and Pittet, 2004).

The series of condensation affecting the Pliensbachian-Toarcian transition strongly 323 324 affects the duration assessments in the Northern Tethyan margin. For instance, in the Paris Basin, the duration of the Tenuicostatum Zone (equivalent to Polymorphum) varies from 0.09 325 myr to 0.5 myr (Boulila et al., 2014; Ruebsam et al., 2014). Correlations of δ^{13} C curves indicate 326 that these sections are affected by a series of condensation events, leading to an underestimation 327 328 of the duration of the Tenuicostatum Zone (Pittet et al., 2014). Recently, Boulila and Hinnov (2017) suggested that the 100-kyr eccentricity cycles identified in Suan et al. (2008b) and 329 Huang and Hesselbo (2014) are the record of the obliquity cycle. However, this would imply a 330 hiatus of ~800 kyr in the middle of the Tenuicostatum Zone, which is not indicated in the $\delta^{13}C$ 331 data (Pittet et al., 2014). In addition, detailed observations of the sedimentological pattern of 332 Peniche indicate than these cycles correspond to bundles of 4-6 marl-limestone alternations, 333 which is related to the hierarchy in between the short eccentricity and precession (Suan et al., 334 335 2008b). We thus retain here for the Peniche section the astrochronological models from Suan et al. (2008b) and Huang and Hesselbo (2014). 336

337

338 **5.2 Duration of the P-To Event**

The duration of the P-To event varies from ~50 kyr to 0.27 myr from various astrochronological 339 340 models (Suan et al., 2008b; Huang and Hesselbo, 2014; Martinez et al., 2017). Uncertainties are due to the condensation event often observed at the Pliensbachian-Toarcian boundary (Pittet 341 342 et al., 2014) and to the difficulty in defining the upper boundary of the P-To event (Martinez et al., 2017). The P-To event indeed starts with an abrupt decrease of 2 to 3‰ the δ^{13} C and change 343 in lithology from carbonated to detrital sediment (Suan et al., 2008a; Bodin et al., 2010). Then, 344 the δ^{13} C values gently increase, making the definition of an upper limit of the P-To event 345 problematic. Martinez et al. (2017) defined two options to define the upper boundary of the P-346 To event: (i) the end of the decrease of the δ^{13} C and CaCO₃ values; (ii) the local increase of the 347

 δ^{13} C values in the lower part of the Polymorphum Zone, corresponding to a maximum of the 348 filter of the 405-kyr band in the δ^{13} C. In the Issouka section, the filter of the 405-kyr band was 349 calculated on the calibrated series (high frequency cut: 0.005833 cycles/kyr; roll-off rate: 10^{36}) 350 and converted into depths shows that this maximum is at 13.1 m (Fig. 9). We suggest using this 351 352 marker as the end of the P-To event, as it can be observed both in the Central High Atlas and the Middle Atlas. From the δ^{13} C and the MS age models, the duration of the P-To event as 353 defined here varies from 0.23 to 0.25 myr, with an average duration of 0.24 ± 0.02 myr (this 354 uncertainty takes into account the uncertainty of the age model and the uncertainty of the period 355 356 of the eccentricity, see supplementary materials). These durations are in agreement with Option 1 from the Foum Tillicht section, which vary from 0.25 to 0.29 myr, with an average duration 357 of 0.27 myr (Martinez et al., 2017). This duration of the P-To event appears stable in Moroccan 358 359 sections, ranging from 0.23 to 0.29 myr.

360

5.3. A synthetic astrochronology for the Tethyan area in the Early Toarcian

The Polymorphum Zone at Issouka shows 10 to 10.5 repetitions of the 100-kyr eccentricity 362 363 cycles. In comparison, the sedimentary records of Foum Tillicht and Peniche show a total of 364 eight 100-kyr eccentricity cycles for the Polymorphum Zone. Lithological patterns in the Umbria-Marche Basin tend to indicate that the Polymorphum Zone contains ten to eleven 365 repetitions of 100-kyr cycle (Mattioli and Pittet, 2004). Four discontinuities identified in the 366 367 Pliensbachian-Toarcian transition have affected the completeness of the sedimentary record in the Foum Tillicht and the Peniche sections. In particular, at Peniche, the Pliensbachian-Toarcian 368 transition appears condensed, while in Foum Tillicht the top of Polymorphum is missing 369 (Martinez et al., 2017). In Issouka, Foum Tillicht and Peniche, δ^{13} C has been measured at a 370 high resolution and astrochronologies have been produced, making it possible to correlate the 371 sections at a Milankovitch scale. 372

Based on the astrochronological frameworks produced in this study and in Huang and 373 374 Hesselbo (2014) and Martinez et al. (2017), we can correlate these three sections at the eccentricity scale (Fig. 9). Radiometric ages have been anchored on stratigraphic ages at the 375 376 Triassic-Jurassic boundary, in the Hettangian and the Sinemurian (Ruhl et al., 2016). A U-Pb age has been proposed for the uppermost part of the Tenuicostatum Zone at 183.22 ± 0.26 Ma 377 in Southern Peru (Sell et al., 2014). The lack of intercalibration with astrochronology makes it 378 379 difficult to anchor it on our time scale. Conversely, the Triassic-Jurassic boundary have been bracketed at 201.36 ± 0.17 Ma with U-Pb ages from the Pucara Basin (Peru) and the New York 380 Canyon (Nevada, USA), recalibrated applying version 3.0 of EARTHTIME tracer (Wotzlaw et 381 382 al., 2014). The duration of the interval from the base of the Jurassic to the base of the Toarcian have been assessed at 17.21 myr (Ikeda and Tada, 2014), which matches with the time scale 383 suggested in Ruhl et al. (2016) with an uncertainty of \pm one 405-kyr cycle. It positions the base 384 385 of the Toarcian at 184.15 ± 0.58 Ma. The uncertainty retained here corresponds to the sum of uncertainties of astrochronology and radiochronology. 386

387 Other ages are as follows (see also supplementary materials for details of the 388 calculations):

- The average duration of the P-To event is 0.24 ± 0.02 myr, positioning the top of the P-To
event at 183.91 ± 0.60 myr,

- The average duration of the Tenuicostatum/Polymorphum Zone is 1.00 ± 0.08 myr, positioning the top of Tenuicostatum/Polymorphum at 183.15 ± 0.66 Ma. Notice that this age fits with the age of Sell et al. (2014) of 183.22 ± 0.26 Ma for the uppermost part of the Tenuicostatum Zone,

- The base of the T-OAE occurs 0.03 myr after the end of the Polymorphum Zone (Huang and
Hesselbo, 2014; Fig. 9), i.e. 183.12 ± 0.66 Ma,

- The T-OAE encompasses 11 short-eccentricity cycles (Suan et al., 2008b; Huang and Hesselbo, 2014; Thibault et al., 2018), corresponding to a duration of 1.05 ± 0.06 myr (Fig. 9).
The top of the T-OAE is thus dated here at 182.07 ± 0.72 Ma,

400

401 **5.4. Origins of P-To and T-OAE disturbances**

The large negative CIEs observed in the Early Toarcian, the P-To event and the T-OAE, imply 402 403 the injection of large amount of light carbon in the oceanic system (Suan et al., 2015). This has been interpreted as dissociation of methane hydrate, maybe orbitally paced (Kemp et al., 2005), 404 or production of thermogenic methane due to volcanic intrusion during the Karoo-Ferrar 405 406 activity (Svensen et al., 2007). The long duration of the T-OAE (Suan et al., 2008b; Huang and 407 Hesselbo, 2014) implies that the amount of methane needed to maintain the negative excursion 408 for such a long time would exceed the volume of gas hydrate reservoir (Suan et al., 2008b). In 409 addition, the interval of the T-OAE display a positive excursion in Hg/TOC, associated with a volcanic activity (Percival et al., 2015). CA-ID-TIMS U-Pb ages indicate that the Ferrar 410 411 province activity occurred from 182.779 \pm 0.033 Ma and lasted 349 \pm 49 kyr (Burgess et al., 2015). These ages fall in the range of the ages assessed here for the T-OAE (183.12 \pm 0.66 Ma 412 to 182.07 ± 0.72 Ma; Fig. 9). All these observations tend to favour the crucial role of the Karoo-413 414 Ferrar activity on the inception of the T-OAE. Hg/TOC data from Yorkshire and the Peniche sections show a positive excursion within the P-To event, which tend to indicate the 415 concomitance between the P-To event and volcanic activity. The P-To event started 184.15 \pm 416 417 0.58 Ma and ended at 183.91 ± 0.60 Ma (Fig. 9), so that the P-To event appears significantly 418 older than the activity of the Ferrar province. Dating of the early phase of the Karoo province tends to indicate that the Karoo province started earlier than the Ferrar (Burgess et al., 2015; 419 Moulin et al., 2017). Notably the oldest ⁴⁰Ar/³⁹Ar ages of the Karoo activity have been found 420 in the southern part of the Barkly East Formation (South Africa), within the Omega and the 421

Moshesh's Ford units (Moulin et al., 2017 and references therein). Other ages older than 183 422 423 Ma have been reported in the Northern part of the Chonke Aike LIP (South America) in the Rio Negro Province and in the eastern part of the Chubut Province (Féraud et al., 1999). The 424 location and the thicknesses of the formations in which these older ⁴⁰Ar/³⁹Ar and and U/Pb ages 425 have been provided are, to our knowledge, much more limited than that of the main phase of 426 the Karoo-Ferrar activity (~183 Ma). While this main phase is likely to be responsible for the 427 428 inception of the T-OAE, limited volcanic activities in the early phase of the Karoo and Chonke Aike LIP may have affected the environmental and ecological perturbations recorded at the 429 Pliensbachian-Toarcian boundary. 430

431

432 **6.** Conclusions

The carbon isotope and MS variations in the Late Pliensbachian-Early Toarcian, performed in 433 434 the expanded hemipelagic in Issouka section from the Middle Atlas Basin, Morocco, reveal superposed frequencies consistent with Milankovitch forcing (eccentricity, obliquity, and 435 precession). The Polymorphum Zone contains 10 to 10.5 repetitions of the 100-kyr eccentricity 436 cycle, so that its duration is assessed here at 1.00 ± 0.08 myr. The duration of the P-To event is 437 in addition assessed at 0.24 ± 0.02 myr. The recognition of the 100-kyr and the 405-kyr 438 439 eccentricity enabled us to establish a stratigraphic framework between the Northern and the Southern Tethyan margins. Anchoring this framework on published radiometric ages and 440 astrochronological time scales, we could estimate the age of the base of the Toarcian as 184.15 441 \pm 0.58 Ma, the top of the P-To event as 183.91 \pm 0.60 Ma, the base of the Toarcian OAE as 442 183.12 \pm 0.66 Ma, and the top of the Toarcian OAE as 182.07 \pm 0.72 Ma. The age of the 443 Toarcian OAE fits with the ages published for the Ferrar volcanic activity. Conversely, the age 444 of the P-To event seems to correspond with early phase of the Karoo and Chonke Aike activity. 445 Thus, the successive and short phases of the volcanic activity of the Chonke Aike, Karoo and 446

447 Ferrar provinces may have been the smoking gun of the successive phases of the mass448 extinction observed in marine biotas in the Pliensbachian and Toarcian times.

449

450 Acknowledgements

This paper benefitted considerably from 3 constructive reviews. We would like to thank also the IAS Postgraduate Grant Scheme for the financial support that Fatima-Zahra Ait-Itto received in 2016 for making the isotopic analyses at Plymouth University. Mathieu Martinez was supported by ERC Consolidator Grant Earthsequencing.

455

456 **References**

- 457 Ait Addi, A., Chafiki, D., 2013. Sedimentary evolution and palaeogeography of mid Jurassic
 458 deposits of the Central High Atlas, Morocco. J. Afr. Earth Sci. 84, 54–69.
 459 https://doi.org/10.1016/j.jafrearsci.2013.04.002.
- Ait-Itto, F.-Z., Price, G.D., Ait Addi, A., Chafiki, D., Mannani, I., 2017. Bulk-carbonate and
 belemnite carbon-isotope records across the Pliensbachian-Toarcian boundary on the
 northern margin of Gondwana (Issouka, Middle Atlas, Morocco). Palaeogeogr.
 Palaeoclim. Palaeoecol. 466, 128–136. https://doi.org/10.1016/j.palaeo.2016.11.014.
- Bassoullet, J.P., Baudin, F., 1994. Le Toarcien inférieur : une période de crise dans les bassins
 et sur les plate-formes carbonatées de l'Europe du Nord-Ouest et de la Téthys. Geobios.
 17, 645–654. https://doi.org/10.1016/S0016-6995(94)80227-0.
- Bassoulet, J., Elmi, S., Poisson, F., Cecca, F., Belion, Y., Guiraud, R., Baudin, F., 1993. Mid
 Toarcian. In: Dercourt, J., Ricou, L.E., Vrielynck, B. (Eds.), Atlas Tethys,
 Paleoenvironmental Maps: Becip–Franlab, Rueil–Malmaison, France, 63–80.
- Bejjaji, Z., Chakiri, S., Reolid, M., Boutakiout, M., 2010. Foraminiferal biostratigraphy of the
 Toarcian deposits (Lower Jurassic) from the Middle Atlas (Morocco). Comparison with

472 western Tethyan areas. J. Afr. Earth Sci. 57, 154–162.
473 https://doi.org/10.1016/j.jafrearsci.2009.08.002.

- Berger, A., Loutre, M.F., Melice, J.L., 2006. Equatorial insolation: from precession harmonics
 to eccentricity frequencies. Clim. Past 2, 131–136. https://doi.org/10.5194/cp-2-1312006.
- Bodin, S., Mattioli, E., Fröhlich, S., Marshall, J.D., Boutib, L., Lahsini, S., Redfern, J., 2010.
 Toarcian carbon isotope shifts and nutrient changes from the Northern margin of
 Gondwana (High Atlas, Morocco, Jurassic): Palaeoenvironmental implications.
 Palaeogeogr. Palaeocl. 297, 377–390. https://doi.org/10.1016/j.palaeo.2010.08.018.
- Boulila, S., Galbrun, B., Huret, E., Hinnov, L.A., Rouget, I., Gardin, S., Bartolini, A., 2014.
 Astronomical calibration of the Toarcian Stage: Implications for sequence stratigraphy
 and duration of the early Toarcian OAE. Earth and Planet. Sci. Lett. 386, 98–111.
 https://doi.org/10.1016/j.epsl.2013.10.047.
- Boulila, S., Hinnov L.A., 2017. A review of tempo and scale of the early Jurassic Toarcian
 OAE: implications for carbon cycle and sea level variations. Newsl. Stratigr. In press.
 https://doi.org/10.1127/nos/2017/0374.
- Burgess, S.D., Bowring, S.A., Fleming, T.H., Elliot, D.H., 2015. High-precision geochronology
 links the Ferrar large igneous province with early-Jurassic ocean anoxia and biotic crisis.
- 490 Earth Planet. Sci. Lett. 415, 90–99. <u>https://doi.org/10.1016/j.epsl.2015.01.037</u>.
- 491 Cleveland, W.S., 1979. Robust Locally Weighted Regression and Smoothing Scatterplots.
 492 Journal of the American Statistical Association 74, 829-836.
- 493 Dera, G., Brigaud, B., Monna, F., Laffont R., Pucéat, E., Deconinck, J.-F., Pellenard, P.,
 494 Joachimski, M.M., Durlet, C., 2011. Climatic ups and downs in a disturbed Jurassic
 495 world. Geology 39, 215–218. http://dx.doi.org/10.1130/G31579.1.

- Du Dresnay, R., 1971. Extension et développement des phénomènes récifaux Jurassique dans
 le domaine Atlasique Marocain, particulièrement au Lias moyen. Bull. Soc. Geol. Fr.
 13, 46–56. http://dx.doi.org/10.2113/gssgfbull.S7-XIII.1-2.46.
- El Hammichi, F., Benshili, K., Elmi, S., 2008. Les faunes d'Ammonites du Toarcien-Aalénien
 du Moyen Atlas sud-occidental (Maroc). Rev. Paléobiol. 27, 429–447.
- Féraud, G., Alric, V., Fornari, M., Bertrand, H., Haller, M., 1999. 40Ar/39Ar dating of the 501 Jurassic volcanic province of Patagonia: migrating magmatism related to Gondwana 502 subduction. Earth Planet. Sci. 83-96. break-up and Lett. 172. 503 https://doi.org/10.1016/S0012-821X(99)00190-9. 504
- Hallam, A., 1997. Estimates of the amount and rate of the of sea level change across the
 Rhaetian–Hettangian and Pliensbachian–Toarcian boundary (latest Triassic to Early
 Jurassic). J. Geol. Soc. Lond. 154, 773–779.
 http://dx.doi.org/10.1144/gsjgs.154.5.0773.
- Hesselbo, S.P., Jenkyns, H.C., Duarte, L.V., Oliveira, L.C.V., 2007. Carbon–isotope record of
 the Early Jurassic (Toarcian) Oceanic Anoxic Event from fossil wood and marine
 carbonate (Lusitanian Basin, Portugal). Earth Planet. Sci. Lett. 253, 455–470.
 https://doi.org/10.1016/j.epsl.2006.11.009.
- Huang, C., Hesselbo, S.P., 2014. Pacing of the Toarcian Oceanic Anoxic Event (Early Jurassic)
 from astronomical correlation of marine sections. Gondwana Res. 25, 1348–1356.
 https://doi.org/10.1016/j.gr.2013.06.023.
- 516 Ikeda, M., Tada, R., 2014. A 70 million year astronomical time scale for the deep-sea bedded
 517 chert sequence (Inuyama, Japan): Implications for Triassic–Jurassic geochronology.
 518 Earth Planet. Sci. Lett. 399, 30–43. https://doi.org/10.1016/j.epsl.2014.04.031.
- Jenkyns, H.C., 1988. The Early Toarcian anoxic event: stratigraphic, sedimentary and
 geochemical evidence. Am. J. Sci. 288, 101–151.

- Kemp, D.B., Coe, A.L., Cohen, A.S., Schwark, L., 2005. Astronomical pacing of methane
 release in the Early Jurassic period. Nature 437, 396–399.
 http://dx.doi.org/10.1038/nature04037.
- Lachkar, N., Guiraud, M., El Harfi, A., Dommergues, J. L., Dera, G., Durlet, C., 2009. Early
 Jurassic normal faulting in a carbonate extensional basin: characterization of
 tectonically driven platform drowning (High Atlas rift, Morocco). J. Geol. Soc. Lond.
 166, 413–430. http://dx.doi.org/10.1144/0016-76492008-084.
- Lamas, F., Irigaray, C., Oteo, C., Chacón, J., 2005. Selection of the most appropriate method
 to determine the carbonate content for engineering purposes with particular regard to
- 530 marls. Eng. Geol. 81, 32–41. <u>https://doi.org/10.1016/j.enggeo.2005.07.005</u>.
- Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A.C.M., Levrard, M., 2004. A longterm numerical solution for the insolation quantities of the Earth. Astronom. Astrophys.
 428, 261-285. https://doi.org/10.1051/0004-6361:20041335.
- Laurin, J., Růžek, B., Giorgioni, M., 2017. Orbital Signals in Carbon Isotopes: Phase Distortion
 as a Signature of the Carbon Cycle. Paleoceanography 32, 1236–1255.
- 536 http://dx.doi.org/10.1002/2017PA003143.
- Little, C.T.S., Benton, M.J., 1995. Early Jurassic mass extinction: a global long-term event.
- 538 Geology 23, 495–498.
- 539 http://dx.doi.org/10.1130/0091-7613(1995)023<0495:EJMEAG>2.3.CO;2.
- Littler, K., Hesselbo, S.P., Jenkyns, H.C., 2010. A carbon-isotope perturbation at the
 Pliensbachian–Toarcian boundary: evidence from the Lias Group, NE England.
 Geological Magazine 147, 181–192. https://doi.org/10.1017/S0016756809990458.
- 543 Kemp, D.B., 2016. Optimizing significance testing of astronomical forcing in
 544 cyclostratigraphy. Paleoceanography 31, https://doi.org/10.1002/2016PA002963.

- Mann, M.E., Lees, J.M., 1996. Robust estimation of background noise and signal detection in
 climatic time series. Climatic Change 33, 409–445.
 http://dx.doi.org/10.1007/BF00142586.
- 548 Martinez, M., Deconinck, J.-F., Pellenard, P., Riquier, L., Company, M., Reboulet, S., Moiroud,
- 549 M., 2015. Astrochronology of the Valanginian-Hauterivian stages (Early Cretaceous):
- 550 chronological relationships between the Paraná–Etendeka large igneous province, the
- 551 Weissert and the Faraoni events. Global Planet. Change 131, 158–173. 552 https://doi.org/10.1016/j.gloplacha.2015.06.001.
- Martinez, M., Kotov, S., De Vleeschouwer, D., Pas, D., Pälike, H., 2016. Testing the impact of
 stratigraphic uncertainty on spectral analyses of sedimentary series. Clim. Past 12,
 1765–1783. https://doi.org/10.5194/cp-12-1765-2016.
- Martinez, M., Krencker, F.N., Mattioli E., Bodin, S., 2017. Orbital chronology of the
 Pliensbachian–Toarcian transition from the Central High Atlas Basin (Morocco).
 Newsletters on Stratigraphy 50, 47–69. https://doi.org/10.1127/nos/2016/0311.
- 559 Mattioli, E., Pittet, B., 2004. Spatial and temporal distribution of calcareous nannofossils along
- 560 a proximal-distal transect in the Lower Jurassic of the Umbria-Marche Basin (central
- 561 Italy). Palaeogeogr. Palaeoclim. Palaeoecol. 205, 295–316.
 562 https://doi.org/10.1016/j.palaeo.2003.12.013.
- Meyers, S.R., Sageman, B.B., 2007. Quantification of deep-time orbital forcing by average
 spectral misfit. Am. J. Sci. 307, 773–792. http://dx.doi.org/10.2475/05.2007.01.
- Meyers, S.R., 2012. Seeing red in cyclic stratigraphy: Spectral noise estimation for
 astrochronology. Paleoceanography 27, PA 3228.
 http://dx.doi.org/10.1029/2012PA002307.
- Meyers, S.R., 2014. Astrochron: An R Package for Astrochronology. [Available at <u>http://cran.r-</u>
 project.org/package=astrochron.]

570 Michard, A., 1976. Eléments de géologie Marocaine. Notes et Mémoires du Service Géologique
571 252, 408.

- Moulin, M., Fluteau, F., Courtillot, V., Marsh. J., Delpech, G., Quidelleur, X., and Gérard, M., 572 573 2017. Eruptive history of the Karoo lava flows and their impact on early Jurassic change. environmental J. Geophys. Res. Solid Earth. 122. 738–772. 574 http://dx.doi.org/10.1029/2011JB008210. 575
- Mutterlose, J., Ruffell, A., 1999. Milankovtich-scale palaeoclimate changes in pale–dark
 bedding rhythms from the Early Cretaceous (Hauterivian and Barremian) of eastern
 England and northern Germany. Palaeogeogr. Palaeoclim. Palaeoecol. 154, 133–160.
 https://doi.org/10.1016/S0031-0182(99)00107-8.
- Percival, L.M.E., Witt, M.L.I., Mather, T.A., Hermoso, M., Jenkyns, H.C., Hesselbo, S.P., AlSuwaidi, A.H., Storm, M.S., Xu, W., Ruhl, M., 2015. Globally enhanced mercury
 deposition during the end-Pliensbachian extinction and Toarcian OAE: a link to the
 Karoo–Ferrar Large Igneous Province. Earth Planet. Sci. Lett. 428, 267–280.
 https://doi.org/10.1016/j.epsl.2015.06.064.
- Pittet, B., Suan, G., Lenoir, F., Duarte, L.V., Mattioli E., 2014. Carbon isotope evidence for
 sedimentary discontinuities in the lower Toarcian of the Lusitanian Basin (Portugal):
 Sea level change at the onset of the Oceanic Anoxic Event. Sediment. Geol. 303, 1–14.
 https://doi.org/10.1016/j.sedgeo.2014.01.001.
- Ruebsam, W., Münzberger, P. and Schwark, L., 2014. Chronology of the Early Toarcian
 environmental crisis in the Lorraine Sub-Basin (NE Paris Basin). Earth Planet. Sci. Lett.
 404, 273–282. https://doi.org/10.1016/j.epsl.2014.08.005.
- Ruhl, M., Hesselbo, S.P., Hinnov, L., Jenkyns, H.C., Xu, W., Riding, J.B., Storm, M., Minisini,
 D., Ullmann, C.V., Leng, M.J., 2016. Astronomical constraints on the duration of the

- Early Jurassic Pliensbachian Stage and global climatic fluctuations. Earth Planet. Sci.
 Lett. 455, 149–165. https://doi.org/10.1016/j.epsl.2016.08.038.
- Sell, B., Ovtcharova, M., Guex, J., Bartolini, A., Jourdan, F., Spangenberg, J.E., Vicente, J., 596 597 Schaltegger, U., 2014. Evaluating the temporal link between the Karoo LIP and climatic-biologic events of the Toarcian Stage with high-precision U-Pb 598 geochronology. Earth Planet. Sci. 408. 48-56. 599 Lett. https://doi.org/10.1016/j.epsl.2014.10.008. 600
- Studer, M., Du Dresnay, R., 1980. Déformations synsédimentaires en compression pendant le
 Lias supérieur et le Dogger, au Tizi n'Irhil (Haut Atlas central de Midelt, Maroc). Bull.
 Soc. Geol. Fr. 22, 391–397. http://dx.doi.org/10.2113/gssgfbull.S7-XXII.3.391.
- Suan, G., Mattioli, E., Pittet, B., Mailliot, S. and Lécuyer, C., 2008a. Evidence for major
 environmental perturbation prior to and during the Toarcian (Early Jurassic) oceanic
 anoxic event from the Lusitanian Basin, Portugal. Paleoceanography. 23, PA1202.
 http://dx.doi.org/10.1029/2007PA001459.
- 608 Suan, G., Pittet, B., Bour, I., Mattioli, E., Duarte, L. and Mailliot, S., 2008b. Duration of the Early Toarcian carbon isotope excursion deduced from spectral analysis: Consequence 609 610 for its possible causes. Earth Planet. Sci. Lett. 267, 666–679. https://doi.org/10.1016/j.epsl.2007.12.017. 611
- Suan, G., van de Schootbrugge, B., Adatte, T., Fiebig, J., Oschmann, W., 2015. Calibrating the
 magnitude of the Toarcian carbon cycle perturbation. Paleoceanography. 30,
 PA002758. http://dx.doi.org/10.1002/2014PA002758.
- Svensen, H., Planke, S., Chevallier, L., Malthe-Sorenssen, A., Corfu, F., Jamtveit, B., 2007.
 Hydrothermal venting of greenhouse gases triggering Early Jurassic global warming.
- 617 Earth Planet. Sci. Lett. 256, 554–566. <u>https://doi.org/10.1016/j.epsl.2007.02.013</u>.

- Taner, M.T., 2003. Attributes Revisited, Technical Publication. Rock Solid Images, Inc.,
 Houston, TX (URL: rocksolidimages.com/pdf/attrib_revisited.htm.).
- Thibault, N., Rhul, M., Ullmann, C.V., Korte, C., Kemp, D.B., Gröcke, D.R., Hesselbo, S.P.,
 2018. The wider context of the Lower Jurassic Toarcian oceanic anoxic event in
 Yorkshire coastal outcrops, UK. Proceedings of the Geologists' Association, in press.
 https://doi.org/10.1016/j.pgeola.2017.10.007.
- Vaughan, S., Bailey, R.J., Smith, D.G., 2015. Cyclostratigraphy: data filtering as a source of
 spurious spectral peaks. Geological Society, London, Special Publications 404, 151156. <u>https://doi.org/10.1144/SP404.11</u>.
- Weedon, G.P., 2003. Time-Series Analysis and Cyclostratigraphy: Examining stratigraphic
 records of environmental cycles. Cambridge University Press, Cambridge.
- 629 Waltham. D., 2015. Milankovitch Period Uncertainties and Their Impact on Cyclostratigraphy.
- 631 Wilmsen, M. and Neuweiler, F., 2008. Biosedimentology of the Early Jurassic post-extinction

J. Sediment. Res. 85, 990–998. http://dx.doi.org/10.2110/jsr.2015.66.

632 carbonate depositional system, central High Atlas rift basin, Morocco. Sedimentology

633 55, 773–807. http://dx.doi.org/10.1111/j.1365-3091.2007.00921.x.

Wotzlaw, J.-F., Guex, J., Bartolini, A., Gallet, Y., Krystyn, L., McRoberts, C.A., Taylor, D.,
Schoene, B., Schaltegger, U., 2014. Towards accurate numerical calibration of the Late
Triassic: high-precision U–Pb geochronology constraints on the duration of the
Rhaetian. Geology. 42, 571–574. http://dx.doi.org/10.1130/G35612.1.

638





Fig. 1. Geological setting of the Issouka section. A. Palaeogeographic map of the western Tethys during the Early Jurassic. The red dash rectangle shows the limit of the second map. Label "1" indicates the location of the Peniche section. Label "2" indicates the location of Yorkshire area. B. Geographical map of Morocco and western Algeria showing the main geological provinces and the location of the Issouka section within the Middle Atlas. Label "3" indicates the location of the Foum Tillicht section (maps from Bodin et al., 2010).



Fig. 2. The different facies in the Issouka section. A. Limestone of the late Pliensbachian with
fauna (belemnites, sponges and brachiopods). B. The Lower Toarcian, characterized by grey
marls and limestone intercalations.



Fig. 3. Raw data, spectrograms and filters of the 100-kyr eccentricity band of the carbon isotope (δ^{13} C) and magnetic susceptibility (MS) signals. **A.** Raw δ^{13} C signal (in black) with long-term trend (in red). **B.** Standardised δ^{13} C signal (in grey) with the filter of the 100-kyr eccentricity (in orange). **C.** Spectrogram of the δ^{13} C signal performed with 15-m-width windows. **D.** Raw CaCO₃ signal (in black) with long-term trend (in red). **E.** Raw MS signal (in black) with long-term trend (in red). **F.** Standardised MS signal (in grey) with the filter of the 100-kyr eccentricity (in orange). **G.** Spectrogram of the MS signal performed with 15-m-width windows.



Fig. 4. Cross-plot of the magnetic susceptibility (MS) versus calcium carbonate content (CaCO₃). The full red line is the best-fit linear regression, while dash lines are the 95% confidence intervals of the linear regression. r indicates the coefficient correlation, and p is the p-value of the correlation.



Fig. 5. 2π -MTM spectra and ASM results of the δ^{13} C series. **A. and B.** Spectra of intervals 1 and 2 with confidence levels calculated with the Mann and Lees (1996) method (ML96) with a Tukey's endpoint rule (Meyers, 2014). **C. and D.** Spectra of intervals 1 and 2 with confidence levels calculated with the LOWSPEC method (Meyers, 2012). **E. and F.** H0 significance levels from the ASM applied on intervals 1 and 2. The red dashed line indicates the sedimentation rate for which the sedimentary frequencies fit the best with the orbital frequencies.



Fig. 6. 2π -MTM spectra of the magnetic susceptibility. **A. and B.** Spectra of intervals 1 and 2 with confidence levels calculated with the Mann and Lees (1996) method (ML96) with a Tukey's endpoint rule (Meyers, 2014). **C. and D.** Spectra of intervals 1 and 2 with confidence levels calculated with the LOWSPEC method (Meyers, 2012).



676

Fig. 7. 2π -MTM spectra of the (**A**.) calibrated δ^{13} C series, and (**B**.) calibrated MS series. The main periods are labeled in kyr. The theoretical period of the corresponding astronomical cycle is displayed in brackets. Abbreviations: E: 400-kyr eccentricity; e: 100-kyr eccentricity; O: obliquity.





683 Fig. 8.

Fig. 8. Comparison between the direct filters of the eccentricity on the calibrated δ^{13} C signal 689 and the amplitude modulation of the precession in the calibrated δ^{13} C. A. Filter of the 405-kyr 690 eccentricity (in red) and the 100-kyr eccentricity (in orange) of the calibrated δ^{13} C signal (in 691 grey). B. Filter of the precession (in blue) and its amplitude modulation (in red) of the calibrate 692 δ^{13} C signal (in grey). C. Filter of the 100-kyr cycle (in orange) and the 405-kyr cycle (in red) 693 obtained from the amplitude modulation of the precession cycle (in grey). In dashed brown: 694 Filter of the 405-kyr cycle obtained from the 100-kyr band of the amplitude modulation of the 695 696 precession. D. Filter of the 405-kyr cycle obtained from the amplitude modulation of the direct filter of the 100-kyr cycle (in grey). **E.** Spectrum of the calibrated δ^{13} C series, with the filters 697 used for the 405-kyr cycle (in red; frequency cut: 0.005833 cycles/kyr; roll-off rate: 10^{36}), the 698 100-kyr cycle (in orange; frequency cuts: 0.006667 - 0.01250 cycles/kyr; roll-off rate: 10^{36}) 699 700 and the precession cycle (in blue; frequency cuts: 0.03583 – 0.05417 cycles/kyr; roll-off rate: 10^{36}). **F.** 2π -MTM spectrum of the amplitude modulation of the precession cycles. The red and 701 orange curves represent the filter of the 405-kyr and 100-kyr eccentricity, respectively. The 702 703 405-kyr cycle is filtered with a Taner low-pass filter (frequency cut: 0.004833 cycles/kyr; rolloff rate: 10³⁶). The 100-kyr cycle is filtered with a Taner band-pass filter (frequency cuts: 704 0.006250 - 0.01558 cycles/kyr; roll-off rate: 10^{36}). G. 2π -MTM spectrum of the amplitude 705 modulation of the 100-kyr cycle filtered from the amplitude modulation of the precession 706 707 cycles. The 405-kyr cycle (in red) is filtered using a Taner low-pass filter (frequency cut: 0.005417 cycles/kyr; roll-off rate: 10^{36}). **H.** 2π -MTM spectrum of the amplitude modulation of 708 709 the 100-kyr cycle filtered from the calibrated series. The 405-kyr cycle (in red) is filtered using a Taner low-pass filter (frequency cut: 0.004583 cycles/kyr; roll-off rate: 10³⁶). Note that 710 spectra of panels F to H are obtained with a padding factor of 2. 711



Fig. 9. Correlations of astronomical time scales in between the Central High Atlas Basin

- 715 (Martinez et al., 2017), the Middle Atlas Basin (present study), the Peniche section (Suan et
- al., 2008b; Huang and Hesselbo, 2014) and the Yorkshire area (Thibault et al., 2018).