

Special issue “Akatsuki at Venus: The First Year of Scientific Operation”

Masato Nakamura, Dmitri Titov, Kevin Mcgouldrick, Pierre Drossart,
Jean-Loup Bertaux, Huixin Liu

► **To cite this version:**

Masato Nakamura, Dmitri Titov, Kevin Mcgouldrick, Pierre Drossart, Jean-Loup Bertaux, et al.. Special issue “Akatsuki at Venus: The First Year of Scientific Operation”. Earth Planets and Space, Springer/Terra Scientific Publishing Company, 2018, 70 (1), pp.art. 144. 10.1186/s40623-018-0916-3 . insu-01877518

HAL Id: insu-01877518

<https://hal-insu.archives-ouvertes.fr/insu-01877518>

Submitted on 8 Feb 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L’archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d’enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PREFACE

Open Access



Special issue “Akatsuki at Venus: The First Year of Scientific Operation”

Masato Nakamura^{1*}, Dmitri Titov², Kevin McGouldrick³, Pierre Drossart⁴, Jean-Loup Bertaux^{5,6} and Huixin Liu⁷

The JAXA Venus explorer, which had been developed in Japan since 2001, was launched in 2010. The spacecraft was named “Akatsuki” after the Japanese word meaning dawn. Akatsuki was inserted into the Venus orbit and began its observation program in 2015, after 5 years wandering around the sun due to the failure of orbit insertion in 2010. Nakamura et al. (2016) described the orbit insertion as well as some initial results from the IR1, UVI, and LIR cameras.

In the twentieth century, well before the start of Akatsuki mission, the former Soviet Union and the USA had explored Venus and elucidated basic characteristics such as atmospheric temperature, pressure, composition, wind speed, surface topography and the plasma environment. However, the physical processes leading to the present state were still unclear. Among them, the Akatsuki mission focused on the study of atmospheric dynamics as a major goal of Venus exploration in the early twenty-first century. The Venusian atmosphere encircles the planet from east to west at all latitudes at a speed much faster than the solid planet. This wind system, called the super-rotation, is one of the biggest mysteries of planetary meteorology and should have a major effect on Venus’ environment. Akatsuki is aimed at a comprehensive survey of the meteorological processes and a quantitative evaluation of the momentum transport sustaining the super-rotation. For this purpose, five cameras imaging the planet at different wavelengths are installed on the spacecraft to visualize atmospheric motions at different altitudes. Combined with a radio occultation instrument that probes the vertical structure, this suite

of instruments can monitor the dynamics of the whole atmosphere in three dimensions.

The second attempt of the orbit insertion using attitude control thrusters in 2015 was perfectly successful. The first scientific discovery was made just 3 h after the arrival: the longwave infrared camera captured a planetary-scale atmospheric gravity wave that was stationary with respect to the surface topography. Starting with this discovery, many other scientific results have been accomplished.

This issue presents initial results of the Akatsuki mission. Iwagami et al. (2018) reported the initial results from Akatsuki/1- μm camera (IR1). More than 600 day-side and 150 night-side images have been obtained since the beginning of regular operation on April 2016. The night-side images are less numerous due to limitations related to the light scattered from the bright day-side. Satoh et al. (2017) reported the performance of Akatsuki/2- μm camera (IR2), and the results obtained with IR2 camera from December 2015 through November 2016. A total of 3091 images of Venus (1420 dayside images at 2.02 μm and 1671 night-side images at 1.735, 2.26, and 2.32 μm) were acquired in this period. Additionally, 159 images, including those of stars for calibration and dark images for the evaluation of dark levels, were obtained. Yamazaki et al. (2018) described the initial results from Akatsuki/Ultraviolet Imager (UVI) at 283 nm and 365 nm. The UV images provide the spatial distribution of SO₂ and the unknown absorber at the cloud tops and characterize the cloud-top morphologies and haze properties. Nominal sequential images with 2-h intervals are used to understand the dynamics of the Venusian atmosphere by deriving the wind velocity field from measured motion vectors at the cloud tops, as well as the mass transportation of UV absorbers. Fukuhara et al. (2017) described the calibration of the Akatsuki/longwave infrared camera (LIR). The brightness

*Correspondence: mnakamur@stp.isas.jaxa.jp

¹ Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, Sagami-hara, Japan

Full list of author information is available at the end of the article

temperature derived from LIR images contained an unexpected bias related not to natural phenomena but to a thermal condition of the instrument. Deep-space images were acquired at different baffle temperatures, and a reference table for eliminating the bias from images was prepared. Takahashi et al. (2018) described the lightning search using the Akatsuki/lightning and airglow camera (LAC). LAC was designed to observe the light curve of possible flashes at a sufficiently high sampling rate to discriminate lightning from other sources and thereby perform a more definitive search for optical emissions. It was confirmed that the operational high voltage was achieved and that the triggering system functions correctly. LAC lightning search observations are planned to continue for several years.

Imamura et al. (2017) reported the initial results of the radio occultation experiments revealing the Venus atmosphere structure. The physical quantities retrieved include the pressure, the temperature, the H₂SO₄ vapor density, and the ionospheric electron density and their variations.

Archiving of the results of these observations is important. Ogohara et al. (2017) gave an overview of the data products. The level-2 images include calibrated radiances and geometry information. The level-3 data are global-grid data in a regular longitude–latitude coordinate system. The method of correcting the boresight pointing of each camera by fitting an ellipse to the observed Venusian limb is also described.

Three numerical modelling and theoretical works are included in this special issue. Horinouchi et al. (2018) studied the cloud-top wind field using images taken by the Ultraviolet Imager (UVI) at the wavelengths of 365 nm (unidentified absorber) and 283 nm (SO₂). These two wavelengths yield slightly different wind velocities, suggesting a difference in the altitude of the cloud features. The local-time dependence and an asymmetry with respect to the equator were observed. The geographic distribution of the zonal wind reported previously (Bertaux et al. 2016) was not seen in the data. McGouldrick (2017) studied the effects of variation in the coalescence efficiency of the Venus cloud particles on the structure of cloud using a one-dimensional cloud model. Specifically, they explored the consequence of allowing the coalescence efficiency of supercooled sulfuric acid in the upper clouds to tend to zero. The most significant result is the appearance of thick clouds of small particles near the transition between upper and middle clouds. Limaye et al. (2018) summarized the characteristics of Venus clouds seen in the multi-wavelength images taken by Akatsuki. The images reveal new and puzzling morphology of the complex cloud cover. The cloud morphologies provide some clues to the processes occurring in

the atmosphere and are thus a key diagnostic tool when quantitative dynamical analysis is not feasible due to insufficient information.

In this special issue, you will find the history of the spacecraft development, the design of the observation instruments, the data processing procedure and the initial scientific results. We hope this special issue will familiarize readers with the outline of the JAXA Akatsuki mission that opened up a new era of Venus exploration. Akatsuki follow-up Venus missions are planned by some space agencies and are discussed by Glaze et al. (2018).

Authors' contributions

All authors read and approved the final manuscript.

Author details

¹ Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, Sagami-hara, Japan. ² ESA/ESTEC, Keplerlaan 1, 2200 AG Noordwijk, The Netherlands. ³ Laboratory for Atmospheric and Space Physics, University of Colorado Boulder, 3665 Discovery Dr., Boulder, CO 80303, USA. ⁴ LESIA, Observatoire de Paris, Université PSL, CNRS, Sorbonne Université, Univ. Paris Diderot, Sorbonne Paris Cité, 5 place Jules Janssen, 92195 Meudon, France. ⁵ LATMOS/UVSQ, Guyancourt, France. ⁶ IKI/RAS, Moscow, Russia. ⁷ Department of Earth and Planetary Science, Faculty of Science, Kyushu University, Fukuoka, Japan.

Competing interests

The authors declare they don't have any competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 14 August 2018 Accepted: 29 August 2018

Published online: 10 September 2018

References

- Bertaux JL, Khatuntsev IV, Hauchecorne A, Markiewicz WJ, Marcq E, Lebonnois S, Patsaeva M, Turin A, Fedorova A (2016) Influence of Venus topography on the zonal wind and UV albedo at cloud top level: the role of stationary gravity waves. *J Geophys Res Planets* 121(6):1087–1101. <https://doi.org/10.1002/2015JE004958>
- Fukuohara T, Taguchi M, Imamura T, Hayashitani A, Yamada T, Futaguchi M, Kouyama T, Sato TM, Takamura M, Iwagami N, Nakamura M, Suzuki M, Ueno M, Hashimoto GL, Sato M, Takagi S, Yamazaki A, Yamada M, Murakami S-y, Yamamoto Y, Ogohara K, Ando H, Sugiyama K-i, Kashimura H, Ohtsuki S, Ishii N, Abe T, Satoh T, Hirose C, Hirata N (2017) Absolute calibration of brightness temperature of the Venus disk observed by the Longwave Infrared Camera onboard Akatsuki. *Earth Planets Space* 69:141. <https://doi.org/10.1186/s40623-017-0727-y>
- Glaze LS, Wilson CF, Zasova LV, Nakamura M, Limaye S (2018) Future of Venus research and exploration. *Space Sci Rev* 214:89. <https://doi.org/10.1007/s11214-018-0528-z>
- Horinouchi T, Kouyama T, Lee YJ, Murakami S, Ogohara K, Takagi M, Imamura T, Nakajima K, Peralta J, Yamazaki A, Yamada M, Watanabe S (2018) Mean winds at the cloud top of Venus obtained from two-wavelength UV imaging by Akatsuki. *Earth Planets Space* 70:10. <https://doi.org/10.1186/s40623-017-0775-3>
- Imamura T, Ando H, Tellmann S, Pätzold M, Häusler B, Atsushi Yamazaki TM, Sato KN, Futaana Y, Oschlisniok J, Sanjay Limaye RK, Choudhary YM, Takeuchi H, Hirose C, Ichikawa T, Toda T, Tomiki A, Abe T, Yamamoto Z-i, Noda H, Iwata T, Murakami S-y, Satoh T, Fukuohara T, Ogohara K, Sugiyama K-i, Kashimura H, Ohtsuki S, Takagi S, Yamamoto Y, Naru Hirata GL, Hashimoto MY, Suzuki M, Ishii N, Tomoko Hayashiyama YJ, Lee MN (2017) Initial

- performance of the radio occultation experiment in the Venus orbiter mission Akatsuki. *Earth Planets Space* 69:137. <https://doi.org/10.1186/s40623-017-0722-3>
- Iwagami N, Sakanoi T, Hashimoto GL, Sawai K, Ohtsuki S, Takagi S, Uemizu K, Ueno M, Kameda S, Murakami S-y, Nakamura M, Ishii N, Abe T, Satoh T, Imamura T, Hirose C, Suzuki M, Hirata N, Yamazaki A, Sato TM, Yamada M, Yamamoto Y, Fukuhara T, Ogohara K, Ando H, Sugiyama K-i, Kashimura H, Kouyama T (2018) Initial products of Akatsuki 1- μ m camera. *Earth Planets Space* 70:6. <https://doi.org/10.1186/s40623-017-0773-5>
- Limaye SS, Watanabe S, Yamazaki A, Yamada M, Satoh T, Sato TM, Nakamura M, Taguchi M, Fukuhara T, Imamura T, Kouyama T, Lee YJ, Horinouchi T, Peralta J, Iwagami N, Hashimoto GL, Takagi S, Ohtsuki S, Murakami S-y, Yamamoto Y, Ogohara K, Ando H, Sugiyama K-i, Ishii N, Abe T, Hirose C, Suzuki M, Hirata N, Young EF, Ocampo AC (2018) Venus looks different from day to night across wavelengths: morphology from Akatsuki multispectral images. *Earth Planets Space* 70:24. <https://doi.org/10.1186/s40623-018-0789-5>
- McGouldrick K (2017) Effects of variation in coagulation and photochemistry parameters on the particle size distributions in the Venus clouds. *Earth Planets Space* 69:161. <https://doi.org/10.1186/s40623-017-0744-x>
- Nakamura M, Imamura T, Ishii N, Abe T, Kawakatsu Y, Hirose C, Satoh T, Suzuki M, Ueno M, Yamazaki A, Iwagami N, Watanabe S, Taguchi M, Fukuhara T, Takahashi Y, Yamada M, Imai M, Ohtsuki S, Uemizu K, Hashimoto GL, Takagi M, Matsuda Y, Ogohara K, Sato N, Kasaba Y, Kouyama T, Hirata N, Nakamura R, Yamamoto Y, Horinouchi T, Yamamoto M, Hayashi Y-Y, Kashimura H, Sugiyama K, Sakanoi T, Ando H, Murakami S, Sato TM, Takagi S, Nakajima K, Peralta J, Lee YJ, Nakatsuka J, Ichikawa T, Inoue K, Toda T, Toyota H, Tachikawa S, Narita S, Hayashiyama T, Hasegawa A, Kamata Y (2016) AKATSUKI returns to Venus. *Earth Planets Space* 68:75. <https://doi.org/10.1186/s40623-016-0457-6>
- Ogohara K, Takagi M, Murakami S, Horinouchi T, Yamada M, Kouyama T, Hashimoto GL, Imamura T, Yamamoto Y, Kashimura H, Hirata N, Sato N, Yamazaki A, Satoh T, Iwagami N, Taguchi M, Watanabe S, Sato TM, Ohtsuki S, Fukuhara T, Futaguchi M, Sakanoi T, Kameda S, Sugiyama K, Ando H, Lee YJ, Nakamura M, Suzuki M, Hirose C, Ishii N, Abe T (2017) Overview of Akatsuki data products: definition of data levels, method and accuracy of geometric correction. *Earth Planets Space* 69:167. <https://doi.org/10.1186/s40623-017-0749-5>
- Satoh T, Sato TM, Nakamura M, Kasaba Y, Ueno M, Suzuki M, Hashimoto GL, Horinouchi T, Imamura T, Yamazaki A, Enomoto T, Sakurai Y, Takami K, Sawai K, Nakakushi T, Abe T, Ishii N, Hirose C, Hirata N, Yamada M, Murakami S-y, Yamamoto Y, Fukuhara T, Ogohara K, Ando H, Sugiyama K-i, Kashimura H, Ohtsuki S (2017) Performance of Akatsuki/IR2 in Venus orbit: the first year. *Earth Planets Space* 69:154. <https://doi.org/10.1186/s40623-017-0736-x>
- Takahashi Y, Sato M, Imai M, Lorenz R, Yair Y, Aplin K, Fischer G, Nakamura M, Ishii N, Abe T, Satoh T, Imamura T, Hirose C, Suzuki M, Hashimoto GL, Hirata N, Yamazaki A, Sato TM, Yamada M, Murakami S-y, Yamamoto Y, Fukuhara T, Ogohara K, Ando H, Sugiyama K-i, Kashimura H, Ohtsuki S (2018) Initiation of a lightning search using the lightning and airglow camera onboard the Venus orbiter Akatsuki. *Earth Planets Space* 70:88. <https://doi.org/10.1186/s40623-018-0836-2>
- Yamazaki A, Manabu Yamada YJ, Lee SW, Horinouchi T, Murakami S-y, Kouyama T, Ogohara K, Imamura T, Sato TM, Yamamoto Y, Fukuhara T, Ando H, Sugiyama K-i, Takagi S, Kashimura H, Ohtsuki S, Hirata N, Hashimoto GL, Suzuki M, Hirose C, Ueno M, Satoh T, Abe T, Ishii N, Nakamura M (2018) Ultraviolet imager on Venus orbiter Akatsuki and its initial results. *Earth Planets Space* 2018(70):23. <https://doi.org/10.1186/s40623-017-0772-6>

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)
