



HAL
open science

Modeling the impact of an X-class solar flare on the exosphere of Mercury

Elisabeth Werner, François Leblanc, Jean-Yves Chaufray, Ronan Modolo

► **To cite this version:**

Elisabeth Werner, François Leblanc, Jean-Yves Chaufray, Ronan Modolo. Modeling the impact of an X-class solar flare on the exosphere of Mercury. EPSC-DPS Joint Meeting 2019, Sep 2019, Geneva, Switzerland. pp. EPSC-DPS2019-48. insu-02161010

HAL Id: insu-02161010

<https://hal-insu.archives-ouvertes.fr/insu-02161010>

Submitted on 21 Jun 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Modeling the impact of an X-class solar flare on the exosphere of Mercury

Anita Linnéa Elisabeth Werner (1), François Leblanc (1), Jean-Yves Chaufray (2) and Ronan Modolo (2)
(1) LATMOS/IPSL, Sorbonne Université, UVSQ, CNRS, Paris, France
(2) LATMOS/IPSL, UVSQ Université Paris-Saclay, Sorbonne Université, CNRS, Guyancourt, France

Abstract

The weak magnetic field, tenuous atmosphere and close distance from the Sun makes Mercury more susceptible to solar transients than other terrestrial planets. For this reason short-term variations in Mercury's sodium exosphere have been associated with the passage of coronal mass ejections (CMEs) [5, 2, 4] and solar flares [1]. Helium ion enhancements have also been recently observed as the direct result of a CME arrival [6].

In this first work we simulate the response of Mercury's exosphere to an X-class solar flare. The momentaneous increase in the extreme ultraviolet (EUV) photon flux during a solar flare functions well as a proxy for modeling the response of the exosphere to extreme variations in the external plasma conditions. This test will allow us to extend such short-term time variations to model the impact of a CME on the precipitation of sodium and other species in Mercury's coupled exosphere-magnetosphere system.

We employ the Exospheric Global Model (EGM) [3], a 3-D parallelized Monte Carlo model, to model Mercury's exosphere. In this model, 10^7 test-particles with statistically representative weights are ejected at the start of the simulation and then followed as they move in the exosphere on ballistic trajectories. These particles can escape from the exosphere, stick to the surface or be re-ejected through one of four processes: thermal desorption (TD), photo-stimulated desorption (PSD), solar wind sputtering and meteoroid vaporization.

We investigate the flare-induced effects on the release of several exospheric species, including sodium and helium, through PSD, TD and photo-ionization. In addition, we will present simulation results with a test-particle model capable of tracing

the trajectories of newly created ions as they leave the exosphere and enter the magnetosphere.

References

- [1] Killen, R. M., Cremonese, G., Lammer, H., Orsini, S., Potter, A. E., Sprague, A. L. Wurz, P., Khodachenko, M. L., Lichtenegger, H. I. M., Milillo, A. and Mura, A.: Processes that promote and deplete the exosphere of Mercury, *Space Science Reviews*, Vol. 132, pp. 433-509, 2007.
- [2] Leblanc, F., Doressoundiram, A., Schneider, N., Masetti, S., Wedlund, M., López Ariste, A., Barbieri, C., Mangano, V. and Cremonese, G.: Short-term variations of Mercury's Na exosphere observed with very high spectral resolution, *Geophysical Research Letters*, Vol. 36, eid. L07201, 2009.
- [3] Leblanc, F., Oza, A. V., Leclercq, L., Schmidt, C., Cassidy, T., Modolo, R., Chaufray, J-Y. and Johnson, R. E.: On the orbital variability of Ganymede's atmosphere, *Icarus*, Vol. 293, pp. 185-198, 2017.
- [4] Orsini, S., Mangano, V., Milillo, A., Plainaki, C., Mura, A., Raines, J. M., De Angelis, E., Rispoli, R., Lazzarotto, F. and Aronica, A.: Mercury sodium exospheric emission as a proxy for solar perturbations transit, *Nature Scientific Reports*, Vol. 8, eid. 928, 2018.
- [5] Potter, A. and Morgan, T.: Evidence for magnetospheric effects on the sodium atmosphere of Mercury, *Science*, Vol. 248, pp. 835-838, 1990.
- [6] Raines, J. M., Wallace, K. L., Sarantos, M., Jasinski, J. M., Tracy, P. J., Dewey, R. M., Weberg, M. J. and Slavin, J. A.: First in-situ observations of exospheric response to CME impact at Mercury, *Mercury: Current and Future Science of the Innermost Planet*, 1–3 May 2018, Columbia, Maryland, 2018.