

Parallel bands of seismicity at the Mid-Atlantic Ridge, 12-14°N

Javier Escartin, Deborah Smith, Mathilde Cannat

► **To cite this version:**

Javier Escartin, Deborah Smith, Mathilde Cannat. Parallel bands of seismicity at the Mid-Atlantic Ridge, 12-14°N. *Geophysical Research Letters*, American Geophysical Union, 2003, 30 (12), pp.22. 10.1029/2003GL017226 . insu-02177675

HAL Id: insu-02177675

<https://hal-insu.archives-ouvertes.fr/insu-02177675>

Submitted on 9 Jul 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Parallel bands of seismicity at the Mid-Atlantic Ridge, 12–14°N

Javier Escartín,¹ Deborah K. Smith,² and Mathilde Cannat¹

Received 28 February 2003; revised 28 April 2003; accepted 6 May 2003; published 20 June 2003.

[1] Seismicity distribution along the Mid-Atlantic Ridge (MAR), monitored with the Autonomous Underwater Hydrophone (AUH) array, shows a band of seismicity ~70 km west of the axis between ~11.9°N and 14.2°N. Available focal mechanisms at the off-axis seismic band show extension directions consistent with the accommodation of the North-American (NA) and South-American (SA) relative plate motion. Axial seismicity shows a gap at ~14°N, coinciding with the 14°N hotspot-like region, and the focal mechanisms near this region are consistent with spreading of the African (AF) and the NA or SA plates. We speculate that the triple junction between the AF, NA and SA plates is close to the 14°N seismic gap. The westward continuation of the NA-SA plate boundary is unconstrained, but could initiate at ~13°N, at the south end of the off-axis seismic band, or be diffuse between ~12°N and ~15°N. *INDEX TERMS:* 3040 Marine Geology and Geophysics: Plate tectonics (8150, 8155, 8157, 8158); 7230 Seismology: Seismicity and seismotectonics; 7220 Seismology: Oceanic crust; 8150 Tectonophysics: Plate boundary—general (3040); 8164 Tectonophysics: Stresses—crust and lithosphere. **Citation:** Escartín, J., D. K. Smith, and M. Cannat, Parallel bands of seismicity at the Mid-Atlantic Ridge, 12–14°N, *Geophys. Res. Lett.*, 30(12), 1620, doi:10.1029/2003GL017226, 2003.

1. Introduction

[2] Understanding of the distribution of seismicity along the northern MAR has improved by the detection of low-magnitude events ($M > 2.5-3$) with the North-Atlantic AUH array [Smith *et al.*, 2002, 2003]. These data show an irregular distribution of earthquakes along the ridge, with zones of continuous seismicity and seismic gaps [Smith *et al.*, 2003]. As in the case of 14°N, many gaps coincide with shallow seafloor near segment centers associated with high magma supply, focused melt supply, or elevated mantle temperatures. These seismic gaps may therefore mark ridge sections with a thin lithosphere where seismicity is subdued or of low magnitude. Most of the events (>80%) occur within ~20 km of the axis or along fracture zones, while off-axis seismicity is scattered [Smith *et al.*, 2003]. The one exception is the ~12–14°N area where an axis-parallel band of seismicity is observed ~70 km west of the axis, extending ~100 km along-axis (Figure 1).

[3] Several processes can cause the intraplate seismicity between ~12–14°N, including off-axis magmatism, an

incipient westward ridge jump, a recent eastward ridge jump, or the presence in the area of the triple junction between the AF, NA, and SA plates. In this paper we study the nature of seismic activity from both the AUH and teleseismic events, and the direction of extension inferred from the teleseismic focal mechanisms. We compare these results with seafloor morphology from multibeam bathymetry acquired recently (Figure 2). Directions of extension from focal mechanisms are compared with predictions of relative plate motions, and the results are used to propose a possible location and configuration of the NA-SA-AF triple junction in the area (Figure 3).

2. Plate Dynamics and the North-South American Plate Boundary

[4] The Central Equatorial Atlantic accommodates the relative motion between the NA and SA plates, but the location and nature of this boundary is undetermined, as it is not clearly marked by seismicity or prominent tectonic features (Figure 1). The position of the triple junction between the NA, SA and AF plates is also ill defined, and has been placed between ~10°N and ~20°N along the MAR [e.g., Minster and Jordan, 1978; Roest and Collette, 1986; Müller and Smith, 1993]. The nature of the triple junction is unknown, and some models propose a zone of diffuse deformation over a portion of the ridge, or continuous position changes due to the unstable configuration of this triple junction [e.g., Roest and Collette, 1986]. While numerous plate models exist, the comparison of the distribution and nature of seismicity with predictions from plate models derived from Global Positioning System (GPS) data is more adequate than using models derived from magnetics or fracture zone orientations (see Figure 1, Table 5 and discussion in Sella *et al.* [2002]); In this paper we use the recent REVEL model, based on 7-year global GPS observations [Sella *et al.*, 2002]. At ~15°N this model predicts NA-SA plate separation at ~1.4 km/my along ~170°, while the predicted NA-AF or SA-AF plate separation is ~23.5 km/my along ~92–95° (Figure 3).

3. Bathymetry and Tectonic Structure

[5] The Mid Atlantic Ridge between the Marathon and the Fifteen-Twenty Fracture zones (MFZ and FTFZ) shallows by >1.5 km towards the segment centered at ~14°N. Geochemical, geological and geophysical data indicate that this segment is magmatically robust: Shallow topography and thick crust at its center; presence of a «hot spot» geochemical anomaly; regular faulting (Figure 2); no known peridotite or gabbro outcrops; no irregular faults and striated detachments, such as found on the ridge flanks to the north or south of the 14°N segment [e.g., Escartín and Cannat, 1999]. Bathymetry in the area south of 14°N

¹Laboratoire de Géosciences Marines (CNRS UMR 7097), Institut de Physique du Globe de Paris, France.

²Dept. of Geology and Geophysics, Woods Hole Oceanographic Institution, USA.




Figure 1. Location map and distribution of seismicity around the study area. Seismic events are from the USGS NEIC catalog (1973–2002, open circles) and the AUH array deployed at the Northern Atlantic (julian day 52/1999–245/2001, solid circles; data available at <http://autochart.pmel.noaa.gov:1776/autochart/GetPosit.html>). NA, SA and AF are the North-, South American and African plates. Diamonds are rotation poles for the NA-SA plate pair; REVEL model, *Sella et al.* [2002] (annotated R) and corresponding error ellipse); *Dixon and Mao* [1997] (D); *Roest and Collette* [1986] (RC); NUVEL-1 model, *DeMets et al.* [1990] (N); *Müller and Smith* [1993] (M); *Larson et al.* [1997] (L). Bathymetry from *Smith and Sandwell* [1997].

shows a normal subsidence of the oceanic crust over the off-axis seismicity band, with no subsidence anomaly or relict rift valley left (Figure 2). These features would be expected after a recent eastward ridge jump of a now dying or fossil ridge axis [e.g., *Mammerickx and Sandwell*, 1986; *Freed et al.*, 1995] from the off-axis seismicity band to the present-day axis. A large increase in depth (~ 500 m) is observed at the east edge of the off-axis seismic zone, corresponding to the western limit of the rift shoulders.

[6] The multibeam data show that the seafloor off-axis is characterized by short normal faults with irregular trends, locally oblique, and by irregular massifs that are not delimited by faults. These structures may correspond to tectonically elevated massifs where the fault escarpments have been degraded by mass wasting, or to later volcanic edifices that may cover the pre-existing topography. The deeps between abyssal hills are flat-bottomed sediment ponds that appear undeformed. The seafloor both north and south of the 14°N segment is similar, based on the irregular faulting, the occurrence of detachments (Figure 2), and the peridotite and gabbro outcrops.

4. Seismicity

[7] Numerous seismic events identified in the AUH array data are located along a well-defined, ridge-parallel band of seismicity, ~ 35 km wide and ~ 110 km long ~ 70 km west of the axis. This off-axis zone of seismicity is well constrained, as the error bars associated with the event locations are small (the calculated rms for errors in event location throughout the study area is 0.1° in latitude and 0.03° in longitude, see *Fox et al.* [2001] and *Smith et al.* [2003]). The southern limit corresponds to the MFZ, which may be locked, or deforming aseismically west of the $\sim 45^\circ\text{W}$ ridge-transform intersection. North of 14°N , the number of off-

axis events seem to decrease and approach the ridge axis. The ridge-axis seismicity show a marked seismic gap at ~ 14.1 – 14.3°N , separating a large number of events that concentrate at $\sim 14.5^\circ\text{N}$ (in proximity to the Logachev hydrothermal site), and a less abundant concentration of events south of 14°N (Figure 2).

[8] While the average acoustic magnitude of events is similar within the three seismic areas (off-axis, and on-axis north and south of the 14°N seismic gap), there are important variations in the rate of events among the areas. The off-axis band of seismicity shows 41 AUH events/100 km, similar to the rate of ~ 51 events/100 km at the ridge axis zone at the same latitude range (Figure 2). In contrast, the northern section of the axis shows the highest number of events (167/100 km), including several seismic swarms associated with teleseismic events (Figure 2), while the




Figure 2. Parallel bands of seismicity and bathymetry between the Marathon and Fifteen-Twenty fracture zones. Solid and open circles correspond to AUH and teleseismic events, and arrows to the direction of extension (T-axis) derived from focal mechanisms (Harvard CMT catalog). Shaded multibeam bathymetry for the $\sim 15^\circ\text{N}$ area [*Escartín and Cannat*, 1999; *Fujiwara et al.*, 2003] is complemented with data from the AT-4-4 (D. Smith, chief Scientist) and the JR63 cruises (C. J. MacLeod and J. Escartín, chief Scientists).




Figure 3. Proposed configuration of the NA-SA-AF plate system at the study area. The T-axes directions from focal mechanisms (thick arrows) are compared with the predicted spreading directions for the NA-SA-AF plate system (thin arrows) from the REVEL model [Sella *et al.*, 2002], and the distribution of AUH events (solid circles). The continuous line indicates the ridge axis and transform faults, and the dashed line the proposed additional plate boundaries. The off-axis seismic band may accommodate NA-SA plate separation, while the zone between the two seismic bands may be part of the SA plate (triple junction, TJ, located at $\sim 14.3^\circ\text{N}$), or correspond to microplate (mp) associated with the triple junction (TJ extending from $\sim 12.8^\circ\text{N}$ to $\sim 14.3^\circ\text{N}$).

seismic gap shows only a few events. This overall pattern of distribution of AUH seismicity is also observed in the teleseismic events, although it is less defined due to the lesser number of recorded events. Differences in the epicenter location between the Harvard CMT and the NEIC catalogs, or the NEIC and AUH-recorded events, be >50 km due to the less accurate locations of the teleseismically determined events with respect to AUH determined events, although no systematic shift in the locations of the same event given by the NEIC, AUH, and Harvard CMT catalogs is observed. A cluster of ~ 10 teleseismic events at the north of the off-axis band may indicate an increase in the intensity of seismicity towards the north of this band.

[9] Available focal mechanisms for some of the teleseismic events (Harvard CMT catalog) show a clear change

in stress orientation between the present-day ridge axis and the off-axis seismic band. All but one event at or near the ridge axis show focal mechanisms consistent with extension sub-parallel to spreading. One event at 14°N shows a T-axis highly oblique to spreading ($\sim 20^\circ$), but it is poorly located as the corresponding AUH and NEIC derived event locations are placed at the off-axis seismicity band (Figure 2). In contrast, the three focal mechanisms at the off-axis seismic band show T-axis directions at $\sim 150\text{--}165^\circ$, $<15^\circ$ from the $\sim 170^\circ$ direction of relative NA-SA spreading predicted by the REVEL model [Sella *et al.*, 2002]. Most transform fault events correspond to strike-slip mechanisms (Figure 3).

5. Discussion

[10] Possible explanations for the off-axis seismic band include a recent eastward ridge jump, a westward ridge jump taking place at the present time, and the complex stress and tectonic regime that may be expected from the presence of a NA-SA-AF triple junction in the area.

[11] *Eastward ridge jump.* Ridge jumps are common along the MAR [e.g., Müller *et al.*, 1998; Allerton *et al.*, 2000], and leave behind fossil rift valleys and anomalous seafloor subsidence [e.g., Mammerickx and Sandwell, 1986; Freed *et al.*, 1995]. The apparently normal subsidence of the seafloor (Figure 2) seems to be inconsistent with this ridge-jump model, and no magnetic data are available to constrain the spreading history of the area.

[12] *Westward ridge jump and incipient rifting.* The axis-parallel distribution of AUH-derived seismicity (Figure 2) may mark the development of a new ridge segment off-axis due to an active ridge jump. The lack of structures consistent with the development of an incipient rift valley, and the presence of sediment ponds, indicate that such a ridge jump would have initiated very recently, without substantial reshaping of the seafloor. Irregular structures several hundreds of meters high that seem to cover fault-controlled abyssal hills may correspond to recent volcanic edifices associated with off-axis magmatism, but sampling and direct observations are required to determine their nature. High-density magnetic data across the area would constrain the spreading history, and magnetic anomalies may be associated with the off-axis seismic band if it indicates off-axis magmatism. This model of a very recent ridge jump cannot be ruled out based on the existing data, but is not consistent with the direction of extension given by focal mechanisms that parallel the NA-SA separation direction, and that are highly oblique to both NA-AF and SA-AF plate separation at the ridge axis.

[13] *North-, South-American and African triple junction.* The orientations of the T-axes from focal mechanisms of teleseisms are consistent with the presence of a NA-SA boundary in close proximity to the ridge axis, and therefore placing the NA-SA-AF triple junction in the $12^\circ\text{--}14^\circ\text{N}$ area (Figures 2 and 3). All but one of the axial events show approximately east-west extension, consistent both with the NA-AF or SA-AF plate separation. The extension directions from events at the off-axis seismic band ($155\text{--}165^\circ$) suggest that this region may correspond to the NA-SA plate boundary, with an extension direction of $\sim 170^\circ$. However, the bathymetry does not show structures that may be associated with this plate boundary (i.e., normal faults trending $\sim 80^\circ$, or strike

slip faults trending $\sim 170^\circ$). Pre-existing normal faults that are parallel to the ridge axis trend could be re-activated mainly as strike-slip faults (events 3 and 4 in Figure 3), although normal faulting is also observed (event 5 in Figure 3).

[14] The link of the off-axis seismic zone with the axis is not well defined, but the scattered seismicity may place it near the $\sim 14^\circ\text{N}$ seismic gap (Figure 2). If the oceanic lithosphere between the seismic bands corresponds to the SA plate, the AF-NA-SA triple junction may be located at $\sim 14.3^\circ\text{N}$, and the MAR between ~ 12.9 and $\sim 14.2^\circ\text{N}$ must correspond to the AF-SA plate boundary (Figure 3). There may be a feedback between the position of the triple junction and the thin axial lithosphere at 14°N , as observed at other triple junctions along slow-spreading ridges [e.g., Ligi *et al.*, 1997; Georgen and Lin, 2002], but the mechanisms of these interactions are not fully constrained. Alternatively, the lithosphere between the seismic bands may be a $\sim 70 \times 110$ km microplate associated with the triple junction. The reduced seismicity at the north may be explained by the inferred thin lithosphere at $\sim 14^\circ\text{N}$, while the south boundary at the MFZ may be locked or deforming aseismically. In both models the triple junction induces an important rotation of stresses, with extension along $\sim 90^\circ$ at the axis and along $\sim 150^\circ$ – 170° 70-km off-axis at the seismic band.

[15] We cannot provide constraints on the position of the NA-SA plate boundary extending west towards the Caribbean subduction zone (Figure 1). If this boundary were continuous, we expect it to initiate at the southern end of the off-axis seismicity band, coinciding approximately with the trace of the MFZ (Figure 3). Alternatively, the plate boundary may be diffuse along the corridor defined by the FTFZ and MFZ ($\sim 12.5^\circ\text{N}$ and $\sim 15.3^\circ$).

[16] **Acknowledgments.** We thank T. Fujiwara for providing multi-beam bathymetry of the area, and D. J. Fornari and S. Bazin for suggestions. The paper was improved thanks to comments by an anonymous reviewer, K. C. Macdonald and the Editor A. Zollo. JE was partially supported by a DORSALES grant (INSU-CNRS, France). DKS was funded by NSF grant OCE-0136808. IGP contribution number 1916 and WHOI contribution number 10919.

References

Allerton, S., J. Escartin, and R. C. Searle, Evidence for asymmetric accretion at a slow-spreading ridge segment, *Geology*, 28, 179–182, 2000.
 Dixon, T. H., and A. Mao, A GPS estimate of relative motion between North and South America, *Geophys. Res. Lett.*, 24(5), 535–538, 1997.

DeMets, C., R. G. Gordon, D. F. Argus, and S. Stein, Current plate motions, *Geophys. J. Int.*, 101, 425–478, 1990.
 Escartin, J., and M. Cannat, Ultramafic exposures and the gravity signature of the lithosphere near the Fifteen-Twenty Fracture Zone (Mid-Atlantic Ridge, 14° – 16.5°N), *Earth Planet. Sci. Lett.*, 171, 411–424, 1999.
 Fox, C. G., H. Matsumoto, and T.-K. A. Lan, Monitoring Pacific Ocean seismicity from an autonomous hydrophone array, *Geophys. Res. Lett.*, 106, 4183–4206, 2001.
 Freed, A. M., J. Lin, P. Shaw, and H. J. Melosh, Long term survival of the axial valley morphology at abandoned slow-spreading centers, *Geology*, 23, 971–974, 1995.
 Fujiwara, T., J. Lin, P. B. Matsumoto, B. Tucholke, and J. F. Casey, Crustal evolution of the Mid-Atlantic Ridge near the Fifteen-Twenty Fracture Zone in the last 5 Ma, *Geochem. Geophys. Geosys.*, 4, 10.1029/2002GC000364, 2003.
 Georgen, J. E., and J. Lin, Three-dimensional passive flow and temperature structure beneath ridge-ridge-ridge triple junctions, *Earth Planet. Sci. Lett.*, 204, 115–132, 2002.
 Larson, K. M., J. T. Freymueller, and S. Philipson, Global plate velocities from the Global Positioning System, *J. Geophys. Res.*, 102, 9961–9981, 1997.
 Ligi, M., E. Bonatti, G. Bortoluzzi, G. Carrara, P. Fabretti, D. Penitenti, D. Gilod, A. A. Peyve, S. Skolotnev, and N. Turko, Death and transfiguration of a triple junction in the South Atlantic, *Science*, 276, 243–245, 1997.
 Mammerickx, J., and D. Sandwell, Rifting of old oceanic lithosphere, *Geophys. Res. Lett.*, 91, 1975–1988, 1986.
 Minster, J. B., and T. H. Jordan, Present-day plate motions, *Geophys. Res. Lett.*, 83, 5331–5354, 1978.
 Müller, R. D., and W. H. F. Smith, Deformation of oceanic crust between the North American and South American plates, *J. Geophys. Res.*, 98(B5), 8275–8291, 1993.
 Müller, R. D., W. R. Roest, and J.-Y. Royer, Asymmetric sea-floor spreading caused by ridge-plume interactions, *Nature*, 396, 456–459, 1998.
 Roest, W. R., and B. J. Collette, The Fifteen Twenty Fracture Zone and the North American-South American plate boundary, *J. Geol. Soc. London*, 143, 833–843, 1986.
 Sella, G. F., T. H. Dixon, and A. Mao, REVEL: A model for recent plate velocities from space geodesy, *Geophys. Res. Lett.*, 107, 10.1029/2000JB000033, 2002.
 Smith, D. K., M. Tolstoy, C. G. Fox, D. R. Bohnenstiehl, H. Matsumoto, and M. Fowler, Hydroacoustic monitoring of seismicity at the slow-spreading Mid-Atlantic Ridge, *Geophys. Res. Lett.*, 29, 10.1029/2001GL013912, 2002.
 Smith, D. K., J. Escartin, M. Cannat, M. Tolstoy, C. G. Fox, D. Bohnenstiehl, and S. Bazin, Spatial and temporal distribution of seismicity along the northern Mid-Atlantic Ridge (15° – 35°N), *Geophys. Res. Lett.*, 108, 10.1029/2002JB001964, 2003.
 Smith, W. H., and D. T. Sandwell, Global seafloor topography from satellite altimetry and ship depth soundings, *Science*, 227, 1956–1962, 1997.

J. Escartin and M. Cannat, Laboratoire de Géosciences Marines (CNRS UMR 7097) Institut de Physique du Globe de Paris, France.

D. K. Smith, Dept. of Geology and Geophysics Woods Hole Oceanographic Institution, USA.