



**HAL**  
open science

## **The LOFAR Tied-Array All-Sky Survey (LOTAAS): Characterization of 20 pulsar discoveries and their single-pulse behavior**

Daniele Michilli, C. Bassa, S. Cooper, J.W.T. Hessels, V.I. Kondratiev, S. Sanidas, B.W. Stappers, C.M. Tan, J. van Leeuwen, Ismaël Cognard, et al.

### ► **To cite this version:**

Daniele Michilli, C. Bassa, S. Cooper, J.W.T. Hessels, V.I. Kondratiev, et al.. The LOFAR Tied-Array All-Sky Survey (LOTAAS): Characterization of 20 pulsar discoveries and their single-pulse behavior. *Monthly Notices of the Royal Astronomical Society*, 2020, 491 (1), pp.725-739. 10.1093/mnras/stz2997 . insu-02407092

**HAL Id: insu-02407092**

**<https://insu.hal.science/insu-02407092>**

Submitted on 12 Dec 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# The LOFAR tied-array all-sky survey (LOTAAS): Characterization of 20 pulsar discoveries and their single-pulse behaviour

D. Michilli<sup>1,2,3,4\*</sup>, C. Bassa<sup>2</sup>, S. Cooper<sup>5</sup>, J. W. T. Hessels<sup>1,2</sup>, V. I. Kondratiev<sup>2,6</sup>, S. Sanidas<sup>1,5</sup>, B. W. Stappers<sup>5</sup>, C. M. Tan<sup>1,5</sup>, J. van Leeuwen<sup>1,2</sup>, I. Cognard<sup>7,8</sup>, J.-M. Grießmeier<sup>7,8</sup>, A. G. Lyne<sup>5</sup>, J. P. W. Verbiest<sup>9,10</sup> and P. Weltevrede<sup>1,5</sup>

<sup>1</sup>Anton Pannekoek Institute for Astronomy, University of Amsterdam, Science Park 904, NL 1098 XH Amsterdam, The Netherlands

<sup>2</sup>ASTRON, Netherlands Institute for Radio Astronomy, Oude Hoogeveensedijk 4, NL 7991 PD Dwingeloo, The Netherlands

<sup>3</sup>Department of Physics, McGill University, 3600 University Str, Montréal, QC H3A 2T8, Canada

<sup>4</sup>McGill Space Institute, McGill University, 3550 University Str, Montréal, QC H3A 2A7, Canada

<sup>5</sup>Jodrell Bank Centre for Astrophysics, School of Physics and Astronomy, The University of Manchester, M13 9PL Manchester, UK

<sup>6</sup>Astro Space Centre, Lebedev Physical Institute, Russian Academy of Sciences, Profsoyuznaya Str 84/32, 117997 Moscow, Russia

<sup>7</sup>LPC2E - Université d'Orléans / CNRS, F-45071 Orléans cedex 2, France

<sup>8</sup>Station de Radioastronomie de Nançay, Observatoire de Paris, PSL Research University, CNRS, Univ. Orléans, OSUC, F-18330 Nançay, France

<sup>9</sup>Fakultät für Physik, Universität Bielefeld, Postfach 100131, D-33501 Bielefeld, Germany

<sup>10</sup>Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany

Accepted 2019 October 17. Received 2019 October 15; in original form 2019 May 2

## ABSTRACT

We are using the LOw-Frequency ARray (LOFAR) to perform the LOFAR Tied-Array All-Sky (LOTAAS) survey for pulsars and fast transients. Here, we present the astrometric and rotational parameters of 20 pulsars discovered as part of LOTAAS. These pulsars have regularly been observed with LOFAR at 149 MHz and the Lovell telescope at 1532 MHz, supplemented by some observations with the Lovell telescope at 334 MHz and the Nançay Radio Telescope at 1484 MHz. Timing models are calculated for the 20 pulsars, some of which are among the slowest spinning pulsars known. PSR J1236 – 0159 rotates with a period  $P \sim 3.6$  s, while five additional pulsars show  $P > 2$  s. Also, the spin-down rates  $\dot{P}$  are, on average, low, with PSR J0815 + 4611 showing  $\dot{P} \sim 4 \times 10^{-18}$ . Some of the pulse profiles, generically single-peaked, present complex shapes evolving with frequency. Multifrequency flux measurements show that these pulsars have generically relatively steep spectra but exceptions are present, with values ranging between  $\sim -4$  and  $-1$ . Among the pulsar sample, a large fraction shows large single-pulse variability, with four pulsars being undetectable more than 15 per cent of the time and one tentatively classified as a Rotating Radio Transient. Two single-peaked pulsars show drifting sub-pulses.

**Key words:** methods: observational – ephemerides – pulsars: general.

## 1 INTRODUCTION

Radio pulsars are rotating neutron stars where a small fraction of the spin-down energy powers beamed radio emission that can cross our line of sight on every rotation resulting in an observable pulsed signal (Pacini 1967; Gold 1968). Since their discovery by Hewish et al. (1968), pulsars have provided a great wealth of scientific discoveries largely because of their use as uniquely precise astronomical clocks (Manchester 2017). This has motivated ongoing pulsar surveys at a wide range of radio frequencies, which to date have found close to 3000 sources (ATNF catalogue;

Manchester et al. 2005).<sup>1</sup> In order to take advantage of these pulsar discoveries, it is necessary to construct a pulsar timing model by measuring the pulses' times-of-arrival (TOAs). This model describes the rotational and astrometric properties of the pulsar and the propagation of the pulses through the interstellar medium (ISM: Edwards, Hobbs & Manchester 2006). The pulsar spin and spin-down rate are indicative of the pulsar evolutionary history. Assuming a simplified model (i.e. dipole braking with constant magnetic field), pulsar parameters can be estimated from these, such as the characteristic age and magnetic field (Goldreich & Julian 1969). For this reason, a scatter-plot of pulsar periods  $P$

\* E-mail: danielemichilli@gmail.com

<sup>1</sup><http://www.atnf.csiro.au/people/pulsar/psrcat>

and period derivatives  $\dot{P}$  (the so-called  $P-\dot{P}$  diagram) provides valuable information on the properties of the pulsar population as a whole. TOAs for a given pulsar can be obtained directly from its single pulses. However, the low signal-to-noise ratio (S/N) and erratic shapes of individual pulses result in larger uncertainties. Therefore, hundreds of rotational periods are usually added together to increase the S/N and form a stable average pulse profile. Pulse profiles are correlated with noiseless templates in order to produce TOAs. A detailed description of this ‘timing’ procedure is provided by Edwards et al. (2006).

All pulsars manifest some level of pulse-to-pulse variation in flux and pulse shape. In some cases, the emission switches between bistable states, and these sources are classified as mode changing pulsars (Wang, Manchester & Johnston 2007). In other cases, the single pulses form patterns called drifting sub-pulses (Taylor, Manchester & Huguenin 1975). Pulsars that are undetected for one or multiple rotational periods are classified as nullers (Backer 1970; here sufficient sensitivity is needed to distinguish between nulling and weak pulses). The nulling fraction can vary from a small fraction of the rotations to nearly 100 per cent (e.g. Wang et al. 2007). In the latter case, sources are often discovered through their single pulses and termed Rotating Radio Transients (RRATs; McLaughlin et al. 2006). These are pulsars whose emission is detected over single rotations separated by large periods of apparent inactivity ( $\gtrsim 1$  min and up to hours).

Radio waves propagating through the cold plasma present in the ISM undergo various propagation effects (e.g. Rickett 1990), which are usually more evident for lower frequency waves ( $\lesssim 300$  MHz). These effects are highly relevant both in searching for new pulsars and in measuring precise TOAs. Dispersion is the frequency dependence of the wave group velocity. It is quantified by the dispersion measure (DM) and scales as  $\nu^{-2}$ , where  $\nu$  is the observing frequency. Diffractive scintillation is the phase perturbation of the waves induced by smaller scale inhomogeneities in the ISM. It creates intensity modulations of the signal both in time and frequency. Diffractive scintillation is typically averaged out by wide-band observations ( $> 10$  MHz) at low frequencies. Refractive scintillation is the angular broadening of the radio signal due to larger scale inhomogeneities in the ISM. It typically manifests in the time domain as an exponential scattering tail in the pulse profiles, which scales roughly as  $\nu^{-4}$ . Scattering can strongly limit the detectability of pulsars at low frequencies because it can wash out the pulsed signal.

The majority of current pulsar surveys are being carried out at frequencies above 300 MHz, where the sky background brightness is lower and the aforementioned radio propagation effects are less severe (Lorimer & Kramer 2004; Stovall, Lorimer & Lynch 2013). However, low-frequency observations ( $\lesssim 300$  MHz) present some practical advantages as well and they can probe the pulsar population in a way that complements the view from higher frequencies. First, the telescope’s field of view is typically larger at lower frequencies, allowing a larger survey speed for all-sky surveys (Stappers et al. 2011). Secondly, most pulsars are brighter at lower frequencies (Bilous et al. 2016). The flux density  $S$  of radio pulsars is usually described by a power law of the observing frequency  $\nu$  whose exponent  $\alpha$  is called the spectral index ( $S \propto \nu^\alpha$ ). If a pulsar has a spectrum steeper than the sky background ( $\alpha \sim -2.55$ ; Mozdzen et al. 2017), it can potentially be detected more easily at lower frequencies – as long as scattering is modest. The average spectral index of pulsars is  $\alpha = -1.4$ , with a standard deviation of 1 (Bates, Lorimer & Verbiest 2013).

Here, we report the timing models and other properties of 20 radio pulsars discovered using the LOw-Frequency ARray (LOFAR; Stappers et al. 2011; van Haarlem et al. 2013), a sensitive radio interferometer that operates at low radio frequencies. We are using this telescope in the frequency range 119–151 MHz to perform the LOFAR Tied-Array All-Sky Survey (LOTAAS; Coenen et al. 2014)<sup>2</sup> for pulsars and fast transients in the Northern sky. A detailed description of the survey is presented in Sanidas et al. (2019). Among the LOTAAS discoveries presented in this paper, PSR J0815 + 4611 has been first presented by Jelić et al. (2015), who identified a steep spectrum, unresolved, and polarized point source in continuum images of the 3C 196 field observed by the LOFAR Epoch of Reionization project (Candi2, Ger de Bruyn, private communication; Yatawatta et al. 2013). Pulsations were then discovered using a targeted LOFAR beam-formed observation (DDT2.004, PI: Hessels) and subsequent search over a range of trial DMs (V. Kondratiev, private communication; Jelić et al. 2015). PSR J1404 + 1159 was discovered by Chandler (2003) and blindly re-detected by LOTAAS. It did not have a timing model at the time of the LOTAAS re-discovery and we detected it at a very different DM than the value given by Chandler (2003), so there was initially some ambiguity about whether it was indeed the same source (Sanidas et al. 2019). Brinkman et al. (2018) recently presented a timing model for the source compatible with the one we obtain. PSR J0302 + 2252 was first reported by Tyul’bashev et al. (2016); PSRs J0122 + 1416, J1635 + 2332, and J2051 + 1248 by Tyul’bashev et al. (2017); and PSRs J0139 + 3336, J1404 + 1159, and J1848 + 1516 by Tyul’bashev, Tyul’bashev & Malofeev (2018). These sources have been blindly detected by LOTAAS around the same time, and we present their timing models for the first time.

Pulsars discovered by LOTAAS are regularly monitored using multiple telescopes; these subsequent timing observations are described in Section 2. The timing models obtained for these pulsars are presented in Section 3 and the characteristics of the pulse profiles are described in Section 4. Flux densities and spectral indices are analysed in Section 5. Individual sources presenting interesting variations within single observations are further described in Section 6. Finally, conclusions are drawn in Section 7.

## 2 OBSERVATIONS

As discussed in greater detail by Sanidas et al. (2019), the LOTAAS survey is performed using the LOFAR ‘Superterp’, a part of the telescope where six stations of antennas are closely spaced. After a promising candidate is found, its rough sky position is re-observed using the full LOFAR core (up to 24 stations) for confirmation and refined localization. Because of the longer baselines of the full core, the localization improves to roughly arcminute precision. The increased sensitivity of the full core compared to the Superterp also means that significantly shorter integrations, typically 15 min, can be used to achieve an S/N sufficient to obtain an adequately precise TOA. The resulting discoveries are added to the LOFAR timing campaign, where selected pulsars are observed monthly using the full LOFAR core. All the pulsars presented here have been observed for a span of at least 1 yr; the total set of observations used in this study is reported in Table 1. Typically, each pulsar is observed for 10 min in the timing campaign. However, due to their weak or sporadic signals, PSRs J0139 + 3336, J0518 + 5125, J1848 +

<sup>2</sup><http://www.astron.nl/lotaas>

**Table 1.** Number of single detections of pulsars in each frequency band. Non-detections are indicated with a dash. Pulsars not observed at a certain frequency are highlighted with an ‘X’. The two last columns indicate the total span ranged by the observations and the pulsar names reported by Sanidas et al. (2019), before a timing model was available.

PSR	LOFAR 149 MHz	Lovell 334 MHz	NRT 1484 MHz	Lovell 1532 MHz	Span months	Name in Sanidas et al. (2019)
J0115 + 6325	17	X	–	41	19	J0115 + 63
J0122 + 1416	14	–	–	–	17	J0121 + 14
J0139 + 3336	9	–	X	31	15	J0139 + 33
J0302 + 2252	23	1	X	20	24	J0302 + 22
J0349 + 2340	20	–	X	–	22	J0349 + 23
J0518 + 5125	18	–	X	–	21	J0518 + 51
J0742 + 4334	17	–	X	–	24	J0742 + 43
J0815 + 4611	31	1	X	20	33	J0815 + 4611
J0857 + 3349	9	1	X	–	17	J0857 + 33
J1226 + 0005	12	1	X	–	24	J1226 + 00
J1236 – 0159	13	1	X	50	25	J1235 – 02
J1343 + 6634	24	–	–	–	23	J1344 + 66
J1404 + 1159	14	1	11	38	19	J1404 + 11
J1635 + 2332	15	1	X	33	16	J1635 + 23
J1735 + 6320	25	1	X	50	29	J1735 + 63
J1848 + 1516	21	1	X	52	31	J1848 + 15
J1849 + 2559	21	1	X	–	24	J1849 + 25
J1933 + 5335	11	X	X	X	15	J1933 + 53
J2051 + 1248	24	–	X	40	25	J2051 + 12
J2329 + 4743	15	1	X	31	15	J2329 + 47

1516, and J1236 – 0159, were observed for 15–20 min per epoch. During the timing campaign, pulsars are coherently de-dispersed at the best DM value resulting from the confirmation observation in order to correct for the intrachannel smearing (Hankins & Rickett 1975). The LOFAR PULsar Pipeline (PULP), an automatic pipeline described by Stappers et al. (2011) and Kondratiev et al. (2016), processes the data from the telescope using PSRCHIVE (Hotan, van Straten & Manchester 2004; van Straten, Demorest & Osłowski 2012)<sup>3</sup> and DSPSR (van Straten & Bailes 2011)<sup>4</sup> to produce an *archive* file. The archives are data cubes containing the signal folded at the approximate pulsar spin period determined from the initial confirmation observation as a function of phase, polarization, frequency, and time. Full Stokes information is recorded, the 78 MHz of available bandwidth is divided into 400 channels, the pulse phase is divided into 1024 bins, and the time resolution is usually 5 s. Only for PSRs J0139 + 3336 and J1848 + 1516 did we store single-pulse-resolved, total intensity archives in order to study their variability over short time-scales.

All the pulsars except for PSR J1933 + 5335 were observed with the 76-m Lovell Telescope at an observing frequency of 1532 MHz with a bandwidth of 384 MHz (Bassa et al. 2016). Each pulsar was first observed between 4–5 times in a span of 10 d, with single observation lasting between 40 and 60 min. If a pulsar was detected, a regular timing campaign began with an average observing cadence of two weeks and observing lengths between 6 and 60 min depending on the detected S/N. The data were processed with the digital filter-bank backend (dfb), which incoherently de-disperses and produces a data cube containing 1024 frequency channels, 10 s-long sub-integrations and 1024 phase bins. For the intermittent source PSR J0139 + 3336, the Apollo backend was used to store single-pulse resolved data. These data have a time resolution of 256  $\mu$ s and 672 frequency channels of 0.5 MHz.

A sample of 18 pulsars were also observed in one occasion with a bandwidth of 64 MHz centred at 334 MHz with the Lovell Telescope. Each pulsar was observed for a duration of 30 min. The data were also processed using the dfb with 512 channels, 10 s-long sub-integrations, and 512 phase bins.

Four of the pulsars reported have also been observed with the Nançay radio telescope (NRT) using the NUPPI backend with a bandwidth of 512 MHz centred around 1484 MHz. The data, which are coherently de-dispersed, have a frequency resolution of 4 MHz, sub-integrations with a duration of 15 s and 2048 phase bins. Typical observation durations were between  $\sim$ 10 and 40 min. While PSRs J0115 + 6325 and J0122 + 1416 have been observed two times each, PSR J1343 + 6634 has been observed nine times without detecting the source.

### 3 TIMING MODELS

LOFAR observations supplemented by Lovell TOAs, when available, have been used to construct the timing models presented here. For each pulsar, initial timing parameters were obtained using PRESTO (Ransom 2001)<sup>5</sup> to maximize the S/N of the pulse profile in the confirmation observation. This resulted in approximate values for the period and DM of the sources. The period derivative was initially set to zero. The position determined by maximizing the pulse profile S/N of the beam grid of the confirmation observation (Sanidas et al. 2019) was used as a starting point in the timing model.

LOFAR TOAs were obtained using standard pulsar timing methods. The PAZ utility from the PSRCHIVE package and CLEAN.PY from COAST GUARD (Lazarus et al. 2016)<sup>6</sup> were used to automatically remove radio frequency interference (RFI) present in the observations. After a visual inspection of the data to remove additional

<sup>3</sup><http://psrchive.sourceforge.net>

<sup>4</sup><http://dspsr.sourceforge.net>

<sup>5</sup><https://www.cv.nrao.edu/~sransom/presto>

<sup>6</sup>[https://github.com/plazar/coast\\_guard](https://github.com/plazar/coast_guard)



**Table 2.** Parameters of the timing models for the 20 pulsars presented. The different columns report the source name, Right Ascension and Declination referenced to the J2000 frame, the reference epoch of the parameters, the measured values of spin period, and its time derivative and DM. Values in parentheses are  $1\sigma$  uncertainties on the last digit.

PSR (J2000)	RA (J2000)	Dec. (J2000)	Epoch (MJD)	$P$ (s)	$\dot{P}$ ( $10^{-15}$ )	DM ( $\text{pc cm}^{-3}$ )
J0115 + 6325	01:15:45.87(1)	+63:25:50.8(1)	57876	0.521455427473(5)	1.599(3)	65.069(4)
J0122 + 1416	01:22:01.31(3)	+14:16:17(1)	57876	1.38899395038(2)	3.803(1)	17.693(3)
J0139 + 3336	01:39:57.23(4)	+33:36:59.7(9)	57901	1.2479609557(1)	2.064(8)	21.23(1)
J0302 + 2252	03:02:31.990(4)	+22:52:12.1(2)	57811	1.207164839778(2)	0.0825(1)	18.9922(6)
J0349 + 2340	03:49:57.38(4)	+23:40:53(2)	57813	2.42077097760(4)	1.0995(7)	62.962(5)
J0518 + 5125	05:18:26.145(8)	+51:25:58.7(2)	57829	0.912511685262(8)	0.191(1)	39.244(2)
J0742 + 4334	07:42:42.13(1)	+43:34:02.2(3)	57998	0.606190680037(6)	0.371(2)	36.255(4)
J0815 + 4611	08:15:59.4623(8)	+46:11:53.24(2)	57662	0.4342422517307(2)	0.0039(1)	11.2738(3)
J0857 + 3349	08:57:07.07(2)	+33:49:17.0(6)	57876	0.242961077060(4)	0.24(1)	23.998(3)
J1226 + 0005	12:26:14.4(2)	+00:05:44(7)	57785	2.28507617026(8)	2.474(2)	18.50(1)
J1236 – 0159	12:36:02.5(6)	–01:59:10(20)	57803	3.5975735967(2)	5.103(2)	19.08(3)
J1343 + 6634	13:43:59.26(2)	+66:34:25.05(9)	57785	1.39410378554(1)	1.0882(7)	30.031(3)
J1404 + 1159	14:04:36.987(8)	+11:59:16.0(2)	57820	2.65043929892(4)	1.3768(6)	18.499(4)
J1635 + 2332	16:35:05.36(1)	+23:32:23.2(3)	57935	1.20869424580(2)	0.863(5)	37.568(6)
J1735 + 6320	17:35:06.562(4)	+63:20:00.06(3)	57760	0.510718135408(1)	0.3226(4)	41.853(1)
J1848 + 1516	18:48:56.13(2)	+15:16:44.1(4)	57655	2.23376977466(5)	1.6813(8)	77.436(9)
J1849 + 2559	18:49:47.555(1)	+25:59:57.66(2)	57785	0.5192634055906(6)	0.1798(3)	75.0016(4)
J1933 + 5335	19:33:01.1(1)	+53:35:43(1)	57546	2.052574490(4)	1.26(2)	33.54(3)
J2051 + 1248	20:51:29.66(2)	+12:48:21.5(6)	57811	0.55316745256(2)	0.019(6)	43.45(1)
J2329 + 4743	23:29:31.548(7)	+47:43:39.73(6)	57950	0.728408609085(4)	0.016(2)	44.012(2)

RFI or corrupted observations, sub-integrations and channels were summed to obtain a single-pulse profile for each observation. Since PSRs J0139 + 3336 and J1848 + 1516 show only very sporadic radio pulses, specific periods where the sources were active have been manually extracted for these pulsars. An analytic pulse profile template was generated for each pulsar by fitting the profile having the highest S/N with von Mises functions using PAAS from PSRCHIVE. For each pulsar, PAT has been used to cross-correlate the observed profiles with this analytic template in order to obtain one TOA per observation. LOFAR TOAs are referenced to the topocentric position of the centre of the LBA CS002 station. For the 11 pulsars detected by the Lovell telescope at 1532 MHz, a single-pulse profile and a TOA were generated from each observation following Hobbs et al. (2004).

The TOAs obtained for each pulsar were fitted with TEMPO2 (Hobbs, Edwards & Manchester 2006)<sup>7</sup> using the initial timing model. Most of the pulsars that were detected with the Lovell telescope at 1532 MHz were observed at high cadence, allowing us to resolve any phase ambiguities and get an initial coherent timing model. However, since TEMPO2 can only fit TOAs with known integer rotation counts in between (Freire & Ridolfi 2018), for pulsars only detectable at LOFAR the cadence of our observations meant that in some cases the phase ambiguities could not be resolved. We therefore used a brute-force algorithm in these cases. This algorithm fitted a set of initial spin periods around the value from the discovery observation, producing a series of plots with the relative residuals. In this way, assuming that the other parameters have a much smaller impact on the fit, it is possible to identify a more precise spin period. Using this simple yet effective technique, a timing model could be obtained for all the pulsars in the sample.

In order to obtain more precise values of the pulsar DMs, LOFAR observations were split into two frequency sub-bands and a TOA was calculated for each one. The same templates were used for the


two sub-bands; the good precision of the obtained timing models (see next) justifies this choice. In the analysis, we did not include possible DM or profile variations over time that are sometimes detected at low frequencies (e.g. Michilli et al. 2018; Donner et al. 2019). The low frequency of LOFAR observations and the availability of the 1532 MHz observations with Lovell for most of the pulsars allowed DM uncertainties  $\lesssim 0.01 \text{ pc cm}^{-3}$  (see Table 2). In all cases, the TOAs from the LOFAR and Lovell telescopes were well described by the same timing model after fitting an arbitrary jump in phase between the two instruments to account for a possible phase offset due to different cable length and differences in the reference pulse phase of the template profile. Both the observatory clocks are referenced to the GPS time system.

The timing models, obtained using the Solar system ephemeris model DE405 (Standish 1998) and performing an unweighted fit, are reported in Table 2. Some of the pulsars presented here are among those with the slowest periods ever measured (Manchester et al. 2005). It is interesting to note that PSR J0250 + 5854, the slowest radio pulsar ever found, was also discovered by LOTAAS (Tan et al. 2018; see also Sanidas et al. 2019 for a discussion of LOTAAS sensitivity to slow pulsars). The pulsars in the sample presented here also have, on average, low values of  $\dot{P}$ .

In Fig. 1, the new pulsar discoveries are plotted on the  $P-\dot{P}$  diagram together with known normal, non-recycled radio pulsars (from the ATNF catalogue; Manchester et al. 2005). The relatively high  $P$  and low  $\dot{P}$  of the sample imply that the new pulsars are on average closer to the death line than the majority of the pulsar population. It is unclear whether this is due to survey observational selection effects or if it is a real effect, e.g. due to older pulsars having on average steeper radio spectra. This will be further investigated in a future study using the full sample of LOTAAS discoveries.

Physical quantities were derived from the timing model parameters using standard assumptions. The characteristic age, dipole magnetic field strength, and spin-down energy of the pulsars (Lorimer & Kramer 2004) are reported in Table 3. Also reported in the table are the pulsar distances derived from their DMs using

<sup>7</sup><https://bitbucket.org/psrsoft/tempo2>



**Figure 1.**  $P-\dot{P}$  diagram of the new pulsar discoveries represented with the black dots. The grey dots represent non-recycled, ‘slow’ radio pulsars (ATNF catalogue; Manchester et al. 2005). The dotted lines represent the indicated characteristic ages and magnetic fields, while the dashed line represents the ‘classical’ death line below which dipolar rotators are not expected to emit radio signals (Ruderman & Sutherland 1975).

both the NE2001 (Cordes & Lazio 2002) and the YMW16 (Yao et al. 2017) models for the free electron density distribution in the Milky Way. The latter model implies a maximum expected Galactic contribution lower than the value measured for three pulsars (PSRs J1343 + 6634, J1635 + 2332, and J1735 + 6320), indicating possible improvements needed in the model (see the discussion in Sanidas et al. 2019, which includes a larger pulsar sample). Given the relatively long rotation period and short observing time span of the pulsars presented here, it was not possible to obtain reliable proper motion values since they did not affect the residuals significantly.

The timing residuals, i.e. the difference between observed and model-predicted TOAs, are shown in Fig. 2. The long period and sometimes irregular emission of some of the sources (see discussion in Section 6) imply that the integrated pulse profile for some of the observations might be formed by too few single pulses to stabilize. This would contribute to the scatter in the TOAs. However, the timing precision achieved is on average higher than what it is often obtained on similar time spans, and a few pulsars are particularly good timers compared to typical slow pulsars (e.g. Hobbs et al. 2004; the residuals of PSRs J0302 + 2252, J0815 + 4611, and J1849 + 2559 have a root mean square lower than 200  $\mu$ s). This relatively high precision could be due to the narrowness of the peaks in the pulse profiles (Section 4) and the low impact of timing noise for these pulsars (e.g. Hobbs et al. 2004).

#### 4 PULSE PROFILES

We obtained a refined pulse profile from each observation by applying the timing models presented in Table 2. For each pulsar and observing frequency, the profiles from all observations have been added together to form a global profile; these are presented in Fig.

3. The profiles are stored in the EPN Database of Pulsar Profiles,<sup>8</sup> where they can be accessed online. Since the flux of PSR J0139 + 3336 is highly variable, rotations where the pulsar was active have been manually selected before forming the integrated profile.

Most pulse profiles are single peaked, with the exceptions of PSRs J0302+2252 and J2329+4743, which are double peaked. In addition, the single peaks of PSRs J0815+4611, J1236-0159, J1635+2332, J1848+1516, and J2051 + 1248 have complex shapes, while the rest are well fitted by a single von Mises function. The features of the profiles can be seen in detail in the online version stored in the EPN data base.

Full widths at 20 per cent ( $W_{20}$ ) and 50 per cent ( $W_{50}$ ) of the profile peak were calculated. In order to improve the precision of the calculated pulse limits, the number of phase bins in the profiles was increased by a factor of 10 000 with a linear interpolation. Full widths were obtained as phase differences between the first and last intersects of the profile with a line at 20 per cent and 50 per cent of the peak, respectively. Only the phase bins containing the pulse were selected in order to avoid spurious noise peaks to bias the result for low-S/N profiles. The pulsar duty cycles  $\delta$  were obtained by calculating the ratio between the width  $W$  and the pulsar period. The results are summarized in Table 4.

The obtained duty cycles are usually below 10 per cent. The only exceptions are PSRs J1848+1516 and, most notably, J2051 + 1248, whose peak occupies more than a quarter of the pulse profile at 149 MHz. For most of the pulsars, the duty cycle decreases with increasing observing frequency. This is a common behaviour explained, for example, by the radius-to-frequency-mapping model (Cordes 1975; Ruderman & Sutherland 1975). However, a few pulsars have wider peaks at higher frequencies. This could be due to the appearance of additional components in the higher frequency profile, e.g. in PSRs J1236 – 0159, J1635 + 2332, and J1848 + 1516, similar to the exceptional cases reported by Pilia et al. (2016).

The pulse profile of PSR J1848 + 1516 is remarkably different at different observing frequencies. The phase of the main peak at 149 MHz corresponds to the phase of a secondary component in the 1532-MHz profile. Instead, no features are present in the 149-MHz profile coincident with the main peak in the 1532-MHz profile, although it could become the secondary component shifted at an earlier phase. In addition, the different components overlap in the 1532-MHz profile, even though this could be due to the higher noise with respect the 149-MHz profile. Finally, the 334-MHz profile presents only one narrow component clearly detected, coincident with the secondary component in the 149-MHz profile. However, only  $\sim 25$  of the 816 pulsar rotations summed to obtain the pulse profile at 334-MHz contained detectable pulses.

None of the pulse profiles are heavily scattered. The main peaks of some pulsars, such as PSRs J0115 + 6325, J0139 + 3336, J1236 – 0159, J1635 + 2332, and J1735 + 6320, show a tail at 149 MHz that might be consistent with a scattered component. However, we did not attempt to model the eventual scattering because of its weak effect.

#### 5 FLUX DENSITIES AND SPECTRAL INDICES

Mean flux densities have been calculated using the following version of the radiometer equation, obtained by expanding equations 7.1 and A1.21 in Lorimer & Kramer (2004), after normalizing the pulse

<sup>8</sup><http://www.epta.eu.org/epndb/>

**Table 3.** Quantities derived from the timing parameters presented in Table 2 assuming dipole braking with constant magnetic fields, short initial periods, and a moment of inertia  $I = 10^{45}$  g cm<sup>2</sup>.  $gl$  and  $gb$  are the Galactic coordinates,  $t_c$  is the characteristic age,  $B$  is the surface magnetic field,  $\dot{E}$  is the spin-down energy, and  $d$  the distance of the pulsars.

PSR	$gl$ (deg)	$gb$ (deg)	$\log t_c$ (yr)	$\log B$ (G)	$\log \dot{E}$ (erg s <sup>-1</sup> )	$d^a$ (kpc)	$d^b$ (kpc)
J0115 + 6325	125.65	0.69	6.7	12.0	32.6	2.2	1.9
J0122 + 1416	134.03	-47.94	6.8	12.4	31.7	0.8	1.6
J0139 + 3336	134.38	-28.17	7.0	12.2	31.6	1.0	1.5
J0302 + 2252	158.44	-30.82	8.4	11.5	30.3	0.7	1.0
J0349 + 2340	167.43	-23.38	7.5	12.2	30.5	3.3	3.7
J0518 + 5125	158.26	7.90	7.9	11.6	31.0	1.4	1.3
J0742 + 4334	175.54	27.21	7.4	11.7	31.8	1.3	1.6
J0815 + 4611	173.63	33.45	9.3	10.6	30.3	0.4	0.4
J0857 + 3349	190.14	39.74	7.2	11.4	32.8	0.9	1.6
J1226 + 0005	289.28	62.30	7.2	12.4	30.9	0.9	1.9
J1236 - 0159	295.06	60.65	7.0	12.6	30.6	0.9	2.0
J1343 + 6634	114.90	49.73	7.3	12.1	31.2	1.8	< 13.8
J1404 + 1159	355.08	67.11	7.5	12.3	30.5	1.4	2.2
J1635 + 2332	42.00	39.75	7.3	12.0	31.3	4.8	< 15.7
J1735 + 6320	92.72	32.55	7.4	11.6	32.0	3.2	< 19.1
J1848 + 1516	46.33	7.44	7.3	12.3	30.8	3.3	3.5
J1849 + 2559	56.24	11.87	7.7	11.5	31.7	3.9	6.1
J1933 + 5335	85.59	15.76	7.4	12.2	30.8	2.2	2.5
J2051 + 1248	59.36	-19.45	8.7	11.0	30.7	2.5	4.1
J2329 + 4743	108.96	-12.91	8.9	11.0	30.2	2.2	2.4

<sup>a</sup> Value based on the NE2001 electron density model (Cordes & Lazio 2002).

<sup>b</sup> Value based on the YMW16 electron density model (Yao, Manchester & Wang 2017).

profiles:


$$S_{\text{mean}} = \frac{T_{\text{sky}} + T_{\text{rec}}}{G\sqrt{2t\Delta f}} \sqrt{\frac{\max(p)}{n \max(p) - \sum_1^n p_i}} \sum_1^n p_i,$$

where  $p$  is the signal amplitude in a phase bin,  $n$  is the total number of phase bins in the pulse profile,  $T_{\text{sky}}$  is the sky temperature,  $T_{\text{rec}}$  is the receiver noise temperature,  $G$  is the telescope gain,  $t$  is the integration time, and  $\Delta f$  is the effective bandwidth free from RFI. Pulse width and period have been expressed in units of phase bins so that  $P = n$ . For Lovell telescope observations, we assumed a gain  $G = 1 \text{ K Jy}^{-1}$  and a system temperature  $T_{\text{sys}} = 25 \text{ K}$  for the 1532-MHz receiver (Bassa et al. 2016) and  $T_{\text{sys}} = 50 \text{ K}$  for the 334-MHz receiver. The typical RFI environment at the Lovell telescope is estimated to leave  $\sim 60$  per cent and 75 per cent of clean data at 334 and 1532 MHz, respectively. For the NRT telescope, we used a gain  $G = 1.6 \text{ K Jy}^{-1}$  and a system temperature  $T_{\text{sys}} = 35 \text{ K}$  (Theureau et al. 2005), while the RFI environment is estimated to leave  $\sim 50$  per cent of clean data. Since the LOFAR gain depends on source elevation, number of functional antennas, and observing frequency, LOFAR observations were calibrated following the procedure described by Noutsos et al. (2015) and Kondratiev et al. (2016). The sky temperature  $T_{\text{sky}}$  was calculated for each source and frequency using the 408-MHz sky map by Haslam et al. (1982) and a spectral index  $\alpha \sim -2.55$  (Mozdzen et al. 2017). Observations too contaminated by RFI or with pointing positions too inaccurate that were refined at a later stage were excluded from the analysis. LOFAR observations were split into two frequency sub-bands to increase the constraint on the spectral index. This was not possible for PSRs J1933 + 5335 because the source was too weak and the full LOFAR bandwidth was thus used to calculate the flux of this pulsar. We decided to exclude PSR J0139 + 3336 from the analysis because its nulling fraction is too extreme to obtain reliable flux values in the duration of our observations. PSR

J2051 + 1248 was barely visible at 334 MHz and the signal was too weak to calculate a flux density.

For each pulsar, all Lovell and NRT observations from the same receiver have been added together using PSRADD from PSRCHIVE, which weights the data by the observation duration and RFI level. The mean flux density was then calculated from the resulting integrated profile. This is expected to effectively remove the effect of diffractive scintillation observed at higher frequencies, which caused the flux to vary up to five times the average value for different observations of the same pulsar. It was not possible to follow the same method for LOFAR observations since the number of active antennas varied between observations. However, since the resulting array sensitivity is not expected to be significantly affected and the RFI level and observation duration are approximately constant, we obtained the mean flux density by averaging the flux density measured in different observations. The resulting values of flux densities are reported in Table 5 and shown in Fig. 4. We calculated a spectral index or upper limits in case of non-detections. Due to the small number of measurements and large uncertainties, all the flux values have been fitted with a single power law. However, a spectral turnover is sometimes observed in pulsars at LOFAR frequencies (e.g. Bilous et al. 2016) and this could be the case of, e.g. PSRs J1236 - 0159 and J1848 + 1516. The values of the spectral indices obtained vary significantly (Table 5), but the spectra of most pulsars are steeper than the average pulsar population ( $\alpha \approx -1.4$ ; Bates et al. 2013). In the case of a non-detection with one of the telescopes, upper limits were derived by assuming that an  $S/N > 5$  is needed to confidently detect a source.

For all the telescopes, the random error on flux densities for specific pulsars and frequencies was obtained as the standard error if more than nine observations were available. The systematic error due to uncertain estimates of the telescope gain, temperature, and average RFI environment is estimated to be 50 per cent of the flux




**Figure 2.** Timing residuals of the models presented in Table 2. Different symbols represent different observing frequencies, with ‘+’ for LOFAR lower band ( $\sim 130$  MHz), ‘x’ for LOFAR higher band ( $\sim 170$  MHz) and the dots for Lovell at 1532 MHz.

density value for LOFAR measurements (Kondratiev et al. 2016) and 25 per cent for NRT and Lovell. The resulting uncertainties on the flux density were estimated using the largest of the two errors. In addition to these estimated uncertainties, however, there

are potentially significant errors on the flux density values that are not accounted for. Most important, the observations were acquired before timing models were available and thus approximate source positions were used. The offset of the refined position from the






**Figure 2.** *Continued from previous page..*

beam centre implies actual flux densities somewhat higher than those reported, and consequently steeper spectra. The uncertainties on the initial positions were of the order of arcminutes, as discussed in Section 2. Given the full width at half-maximum (FWHM) of LOFAR beams ( $\sim 3.5$  arcmin) the offset from the centre of telescope

is important for some pulsars. For comparison, the FWHM of Lovell beams are  $\sim 40$  and  $9$  arcmin at  $334$  and  $1532$  MHz, respectively. Correcting for the complex beam shape of LOFAR (e.g. Obrocka, Stappers & Wilkinson 2015) is difficult and the simple approach used by Sanidas et al. (2019) to model the beam as a  $\text{sinc}^2$  function



**Figure 3.** Cumulative pulse profiles of the pulsars presented here. Pulse peaks are all normalized to the same height and full rotational phase windows are shown. The profiles at different frequencies have been aligned by applying the timing models presented in Table 2 and then rotated to show the main peak at the centre for the 149-MHz profile.

**Table 4.** Characteristics of the pulse profiles shown in Fig. 3.  $W$  is the width at a fraction of the peak intensity and  $\delta$  is the duty cycle. The subscripts indicate the percentage of the peak intensity that the value refers to. Uncertainties are  $\sim 1$  ms on the width values and  $\sim 0.1$  per cent on the duty cycle values.

PSR	$W_{20}$ (ms)			$\delta_{20}$ (per cent)			$W_{50}$ (ms)			$\delta_{50}$ (per cent)		
	149 MHz	334 MHz	1532 MHz	149 MHz	334 MHz	1532 MHz	149 MHz	334 MHz	1532 MHz	149 MHz	334 MHz	1532 MHz
J0115 + 6325	35	–	28	6.7	–	5.3	22	–	19	4.3	–	3.7
J0122 + 1416	48	–	–	3.4	–	–	32	–	–	2.3	–	–
J0139 + 3336	32	–	22	2.6	–	1.8	15	–	14	1.2	–	1.2
J0302 + 2252	79	66	55	6.6	5.4	4.6	70	59	47	5.8	4.9	3.9
J0349 + 2340	71	–	–	2.9	–	–	50	–	–	2.0	–	–
J0518 + 5125	51	–	–	5.5	–	–	33	–	–	3.7	–	–
J0742 + 4334	44	–	–	7.3	–	–	30	–	–	5.0	–	–
J0815 + 4611	21	17	16	4.8	3.8	3.7	15	12	9	3.4	2.8	2.2
J0857 + 3349	14	14	–	5.6	5.6	–	9	5	–	3.5	2.1	–
J1226 + 0005	69	47	–	3.0	2.1	–	48	32	–	2.1	1.4	–
J1236 – 0159	150	110	176	4.2	3.1	4.9	94	96	90	2.6	2.7	2.5
J1343 + 6634	91	–	–	6.5	–	–	57	–	–	4.1	–	–
J1404 + 1159	81	65	55	3.0	2.5	2.1	58	45	36	2.2	1.7	1.4
J1635 + 2332	38	57	53	3.2	4.7	4.4	14	32	46	1.1	2.6	3.8
J1735 + 6320	15	16	9	3.0	3.0	1.7	9	10	6	1.7	2.0	1.1
J1848 + 1516	208	106	261	9.3	4.7	11.7	40	20	146	1.8	0.9	6.5
J1849 + 2559	11	17	–	2.1	3.2	–	6	5	–	1.3	1.0	–
J1933 + 5335	67	–	–	3.3	–	–	53	–	–	2.6	–	–
J2051 + 1248	147	–	87	26.5	–	15.7	87	–	49	15.7	–	8.8
J2329 + 4743	55	44	48	7.5	6.0	6.7	10	6	33	1.4	0.8	4.5

**Table 5.** Mean flux densities  $S$  measured at different frequencies (indicated in units of MHz as subscripts) and the inferred spectral indices  $\alpha$ . Flux density values have been fitted with a single spectral index. The last column reports the offset between the centre of the telescope beams and the position refined with timing models.


PSR	$S_{129}$ (mJy)	$S_{168}$ (mJy)	$S_{334}$ (mJy)	$S_{1484}$ (mJy)	$S_{1532}$ (mJy)	$\alpha$	offset (arcmin)
J0115 + 6325	16(8)	10(5)	–	<0.1	0.05(1)	–2.38(6)	2.3
J0122 + 1416	9(4)	4(2)	<1.6	<0.1	<0.2	–3(3)	0.6
J0302 + 2252	20(10)	18(9)	5(1)	–	0.7(2)	–1.43(6)	1.6
J0349 + 2340	3(2)	1.1(6)	<1.6	–	<0.2	–4(3)	0.8
J0518 + 5125	4(2)	2(1)	<2.0	–	<0.2	–3(3)	1.9
J0742 + 4334	4(2)	3(2)	<1.6	–	<0.2	–1(3)	0.0
J0815 + 4611	18(9)	16(8)	2.5(6)	–	0.36(9)	–1.5(2)	0.4
J0857 + 3349	4(2)	2(1)	0.4(1)	–	<0.2	–2.30(8)	0.9
J1226 + 0005	20(10)	8(4)	0.5(1)	–	<0.2	–4.05(2)	2.0
J1236 – 0159	7(3)	6(3)	3.0(8)	–	0.05(1)	–2.3(4)	3.9
J1343 + 6634	14(7)	7(4)	<1.5	<0.1	<0.2	–3(3)	1.4
J1404 + 1159	30(20)	18(9)	2.7(7)	0.28(7)	0.09(2)	–2.1(2)	1.5
J1635 + 2332	9(4)	2(1)	0.4(1)	–	0.025(6)	–2.1(3)	0.8
J1735 + 6320	2(1)	1.8(9)	0.6(1)	–	0.06(1)	–1.52(8)	0.6
J1848 + 1516	6(3)	9(5)	1.5(4)	–	0.12(3)	–1.7(2)	0.4
J1849 + 2559	7(4)	4(2)	0.5(1)	–	< 0.2	–2.9(2)	1.1
J1933 + 5335		1.0(5) <sup>a</sup>	–	–	–	–	3.0
J2051 + 1248	50(30)	40(20)	<1.8	–	0.05(1)	–2.9(2)	1.1
J2329 + 4743	3(2)	3(1)	0.5(1)	–	0.10(2)	–1.4(2)	0.8

<sup>a</sup>Value referenced to a central frequency of 149 MHz.

proved insufficient for this study. Moreover, ionospheric effects are expected to cause a jitter of LOFAR beams up to  $\sim 1$  arcmin. Therefore, we do not attempt to correct for the positional offset and caution the reader that the values reported for LOFAR fluxes and spectral indices are indicative and should not be used for detailed studies. We report the offset between the beam centre and the refined position of the pulsars in the last column of Table 5.


NRT observations of PSR J1404 + 1159 were independently calibrated by regularly observing known calibration sources. This method could not be applied to Lovell data because calibration

sources were not observed nor to LOFAR data because of the dependence of the telescope gain on the source position. The average flux density obtained for PSR J1404 + 1159 with NRT through calibration sources was 0.7(2) mJy,  $\sim 1.5\sigma$  away from the value of 0.28(7) mJy obtained through the radiometer equation and reported in Table 5. We could not identify a clear reason for this discrepancy and it could originate from the systematic errors described above. In addition, Brinkman et al. (2018) report a flux density for the source of 4.3(9) mJy at 327 MHz and 0.027(5) mJy at 1400 MHz using Arecibo. While the first value is compatible with our measurement



**Figure 4.** Mean flux densities (the dots) and fitted power-law spectra (the lines) reported in Table 5 for the different pulsars and observing frequencies. The triangles represent upper limits considering a fiducial value for detection of  $S/N > 5$ , with the filled triangles for Lovell observations and the empty triangles for NRT observations. The shadowed regions represent  $1\sigma$  uncertainties on the spectral indices referenced to 149 MHz.





**Figure 5.** Phase-resolved flux density variations over time for six pulsars observed with LOFAR at 149 MHz. The grey scale is normalized independently for each plot. Pulsar names are indicated on the individual panels. Each time bin contains a single pulsar rotation for PSRs J0139 + 3336 and J1848 + 1516; time bins are 5 s for the rest of the sources. There are 1024 phase bins over the full phase, 10 per cent of the full rotation is shown here. The horizontal white stripes indicate data that have been excised to remove RFI.

at 334 MHz, the second is lower than our estimates at both 1484 and 1532 MHz. However, the flux density of the pulsar is highly variable at 1.4 GHz and we measure values between  $\sim 0.04$  and 1.1 mJy in individual NRT observations calibrated with known sources.

We checked the TIFR GMRT Sky Survey (Intema et al. 2017) source catalogue around the position of the brightest pulsars at 149 MHz in our sample, but we did not find any counterpart.

## 6 INDIVIDUAL SOURCES

Here, we discuss six pulsars of the sample that show sporadic emission or interesting single-pulse behaviour. The flux densities of these sources as a function of rotational phase and time are shown in Fig. 5 for LOFAR observations. Two pulsars show drifting sub-pulses and five are nullers, with PSR J0139 + 3336 tentatively classified as a RRAT (extreme nuller). We calculated the nulling fractions of these pulsars following the procedure of Wang et al. (2007). All the observations, with the exception of those relative to PSRs J0139 + 3336 and J1848 + 1516, are averaged every 5 s and single pulses are not stored.

After excluding PSR J0139 + 3336, four pulsars in our sample show nulling fractions  $> 15$  per cent. Therefore, the percentage of nulling pulsars in our sample is more than double the percentage in the total pulsar population, where nulling pulsars are  $\lesssim 10$  per cent (Yang, Han & Wang 2014). Since the characteristic age of the pulsars in our sample is on average larger than the rest of the population, this could support the evidence found by Ritchings (1976) and Wang et al. (2007) that the nulling fraction is related to the pulsar characteristic age, or it could be due to the long dwell time of LOTAAS observations (1 h each). The nulling fractions found in our sample (between 15 and 50 per cent) is large with respect to the rest of the nulling pulsars but not unheard of (e.g. Biggs 1992; Wang et al. 2007).


### PSR J0139 + 3336

The source shows the behaviour of a RRAT, with only sporadic single pulses detected. We stored single-pulse resolved data at both 149 and 1532 MHz. Pulses are visible in single pulsar rotations separated by minutes. An example of a few bright pulses is reported in Fig. 6.

Using LOFAR observations, we selected pulses having an S/N  $> 10$  at the phase of the main peak in the integrated pulse profile. This threshold was chosen to select pulses clearly separated from the noise distribution. A total of 30 pulses were detected in LOFAR observations above this threshold. An average of  $\sim 3$  pulses per observation was detected, implying a rate of one pulse every  $\sim 5$  min. Given the small number of detected events, the rate of pulses among different observations was roughly compatible with a Poisson distribution.

The same analysis was repeated for Lovell observations at 1532 MHz. The S/N of the brightest pulses is similar in the two cases. Also, the rate is similar, with a pulse detected every  $\sim 5$  min with an S/N  $> 10$  and a distribution roughly consistent with a Poisson distribution. The lack of a robust estimate of the source spectral index prevents a more detailed comparison of the pulses at the two frequencies.

A peak in the integrated pulse profile was detected in all of LOFAR and most of Lovell observations containing pulsar rotations where emission could be visually identified. After excluding these single rotations, the integrated pulse profile is indistinguishable from noise.



**Figure 6.** Flux density (in arbitrary units) as a function of phase in single rotations of PSR J0139 + 3336. The two observations (LOFAR at 149 MHz, top, and Lovell at 1532 MHz, bottom) are not simultaneous. The same number of rotations are shown for the two observations.

### PSR J0302 + 2252

The flux of this nulling pulsar is highly variable on short time-scales for both the peaks in the profile (Fig. 5). Unfortunately, we did not store single pulses for this source; rather, the flux is averaged every 5 s (about four rotational periods). Therefore, it is impossible

to assess the flux variability over single rotations. The degree and time-scale of variation are similar for the two peaks, with a nulling fraction  $\sim 15$  per cent. However, the flux density of the two peaks in single sub-integrations is not obviously correlated.

### PSR J1226 + 0005

A null emission lasting  $\sim 40$  s can be seen for this pulsar in Fig. 5. around 200 s after the start of the observation. Longer nulls are detected as well, with the pulsar being detected for only the first  $\sim 2$  min of one 10-min observation. The average nulling fraction for the pulsar is  $\sim 50$  per cent.

Fig. 5. also reveals drifting sub-pulses for PSR J1226 + 0005. With no individual pulses being recorded (Fig. 5. shows 5 s, or  $\sim 2.2$  pulse period averages), it is hard to quantify this further. Nevertheless, the drift-rate appears to be variable with the drift-rate being lower as seen at the top of Fig. 5. (i.e. the drift bands are steeper) compared to what it is at  $\sim 120$  s into the observation. In addition, the emission appears to wander slightly in pulse phase (e.g. the emission is slightly late  $\sim 120$  s into the observation shown in Fig. 5.). Individual pulse observations might reveal if the observed variability is related to discrete mode changes, or if the effect is smoother.

### PSR J1343 + 6634

This pulsar shows a nulling fraction  $\sim 35$  per cent. The source switches between detectable and non-detectable states on a time-scale of a few tens of seconds. This behaviour is consistent throughout the different observations.

### PSR J1404 + 1159

During the preparation of this manuscript, the source has also been studied by Brinkman et al. (2018) at 327 and 1400 MHz using the Arecibo telescope. The parameters that they present are in agreement with our measurements. We also detect the bright drifting sub-pulses forming the main peak. The sub-pulses are clearly visible in the 5-s long sub-integrations visible in Fig. 5. The relatively high S/N of the pulsar and detection of the drifting sub-pulses over multiple frequencies could allow detailed studies of the drifting evolution with frequency (e.g. Hassall et al. 2013).

### PSR J1848 + 1516

The source switches between detectable and non-detectable states every few tens of rotations, with an average nulling fraction of  $\sim 50$  per cent. While the pulsar is relatively active in some observations (as shown in Fig. 5), it is undetected in several 15-min observations. Sporadically, a second peak appears trailing the main one, becoming the brightest in three observations. Only on a very few occasions, a third peak leading the main one has been detected for a few rotations.

## 7 CONCLUSIONS

We have presented the properties of 20 radio pulsars discovered by the LOFAR telescope as part of the LOTAAS survey. Since their discovery, the sources have been regularly observed at multiple frequencies using LOFAR, Lovell, and NRT telescopes. This allowed us to calculate the astrometric and rotational parameters

of the pulsars. They have, on average, longer periods and lower spin-down rates than the majority of the pulsar population. This places the pulsars closer to the death line than the average of the global pulsar population. It is unclear whether this is a real effect or a selection bias and this will be explored in a subsequent paper using a larger LOTAAS sample. Integrated pulse profiles were calculated at different frequencies using the obtained timing models. They are mostly single-peaked and show frequency evolution with a complex behaviour in some cases. Values of mean flux densities at the different observing frequencies have been calculated. Even keeping in mind the systematic errors present, the resulting spectra are steeper than average for most pulsars. Five out of the 20 pulsars in the sample are undetectable for more than 15 per cent of time, with PSR J0139 + 3336 tentatively classified as a RRAT. Two of the pulsars show drifting sub-pulses.

## ACKNOWLEDGEMENTS

We thank Anne M. Archibald for her help with the analysis. DM and JWTH acknowledge funding from the European Research Council under the European Union's Seventh Framework Programme (FP/2007-2013) / ERC Starting Grant agreement no. 337062 (DRAGNET). JWTH also acknowledges funding from an NWO Vidi fellowship. DM is a Banting fellow. JvL acknowledges funding from the European Research Council under the European Union's Seventh Framework Programme (FP/2007-2013) / ERC Grant Agreement no. 617199 (ALERT), and from Vici research programme 'ARGO' with project number 639.043.815, financed by the Netherlands Organisation for Scientific Research (NWO). This paper is based in part on data obtained with the International LOFAR Telescope (ILT). LOFAR (van Haarlem et al. 2013) is the Low Frequency Array designed and constructed by ASTRON. It has facilities in several countries. These are owned by various parties (each with their own funding sources) and are collectively operated by the ILT foundation under a joint scientific policy. The data for this project were taken as part of the LOFAR long-term proposals 'Pulsar Timing with LOFAR' (PI: Verbiest), under proposal codes LC0\_011, DDT0003, LC1\_027, LC2\_010, LT3\_001, LC4\_004, LT5\_003, LPR6\_002, LPR7\_001, LPR8\_001, LC9\_041, and LT10\_004, and 'Additional Timing Observations on LOTAAS Pulsar Discoveries' (PI: Tan), under proposal codes LC8\_035, LC9\_021, and LT10\_015. Pulsar research at Jodrell Bank and access to the Lovell Telescope is supported by a Consolidated Grant from the UK's Science and Technology Facilities Council. The Nançay Radio Observatory is operated by the Paris Observatory, associated with the French Centre National de la Recherche Scientifique (CNRS) and Université d'Orléans.

## REFERENCES

- Backer D. C., 1970, *Nature*, 228, 42
- Bassa C. G. et al., 2016, *MNRAS*, 456, 2196
- Bates S. D., Lorimer D. R., Verbiest J. P. W., 2013, *MNRAS*, 431, 1352
- Biggs J. D., 1992, *ApJ*, 394, 574
- Bilous A. V. et al., 2016, *A&A*, 591, A134
- Brinkman C., Freire P. C. C., Rankin J., Stovall K., 2018, *MNRAS*, 474, 2012
- Chandler A. M., 2003, PhD thesis, California Institute of Technology
- Coenen T. et al., 2014, *A&A*, 570, A60
- Cordes J. M., 1975, *ApJ*, 195, 193
- Cordes J. M., Lazio T. J. W., 2002, arXiv Astrophysics e-prints (astro-ph/0207156)
- Donner J. Y. et al., 2019, *A&A*, 624, A22

- Edwards R. T., Hobbs G. B., Manchester R. N., 2006, *MNRAS*, 372, 1549
- Freire P. C. C., Ridolfi A., 2018, *MNRAS*, 476, 4794
- Gold T., 1968, *Nature*, 218, 731
- Goldreich P., Julian W. H., 1969, *ApJ*, 157, 869
- Hankins T. H., Rickett B. J., 1975, in Alder B., Fernbach S., and Rotenberg M., eds, Academic, Vol. 14, Methods in Computational Physics, Radio astronomy. New York, p. 55
- Haslam C. G. T., Salter C. J., Stoffel H., Wilson W. E., 1982, *A&AS*, 47, 1
- Hassall T. E. et al., 2013, *A&A*, 552, A61
- Hewish A., Bell S. J., Pilkington J. D. H., Scott P. F., Collins R. A., 1968, *Nature*, 217, 709
- Hobbs G., Lyne A. G., Kramer M., Martin C. E., Jordan C., 2004, *MNRAS*, 353, 1311
- Hobbs G. B., Edwards R. T., Manchester R. N., 2006, *MNRAS*, 369, 655
- Hotan A. W., van Straten W., Manchester R. N., 2004, *Publ. Astron. Soc. Aust.*, 21, 302
- Intema H. T., Jagannathan P., Mooley K. P., Frail D. A., 2017, *A&A*, 598, A78
- Jelić V. et al., 2015, *A&A*, 583, A137
- Kondratiev V. I. et al., 2016, *A&A*, 585, A128
- Lazarus P., Karuppusamy R., Graikou E., Caballero R. N., Champion D. J., Lee K. J., Verbiest J. P. W., Kramer M., 2016, *MNRAS*, 458, 868
- Lorimer D. R., Kramer M., 2004, Handbook of Pulsar Astronomy. Cambridge Univ. Press, Cambridge
- Manchester R. N., 2017, in Journal of Physics Conference Series, Vol. 932, Presented at Physics of Neutron Stars - 2017, 50 years later. Saint Petersburg, p. 012002,
- Manchester R. N., Hobbs G. B., Teoh A., Hobbs M., 2005, *AJ*, 129, 1993
- McLaughlin M. A. et al., 2006, *Nature*, 439, 817
- Michilli D. et al., 2018, *MNRAS*, 476, 2704
- Mozdzen T. J., Bowman J. D., Monsalve R. A., Rogers A. E. E., 2017, *MNRAS*, 464, 4995
- Noutsos A. et al., 2015, *A&A*, 576, A62
- Obrocka M., Stappers B., Wilkinson P., 2015, *A&A*, 579, A69
- Pacini F., 1967, *Nature*, 216, 567
- Pilia M. et al., 2016, *A&A*, 586, A92
- Ransom S. M., 2001, PhD thesis, Harvard University
- Rickett B. J., 1990, *ARA&A*, 28, 561
- Ritchings R. T., 1976, *MNRAS*, 176, 249
- Ruderman M. A., Sutherland P. G., 1975, *ApJ*, 196, 51
- Sanidas S. et al., 2019, *A&A*, 626, A104
- Standish E. M., 1998, JPL Planetary and Lunar Ephemerides, DE405/LE405, Jet Propulsion Laboratory, Interoffice Memorandum, F-98\_048
- Stappers B. W. et al., 2011, *A&A*, 530, A80
- Stovall K., Lorimer D. R., Lynch R. S., 2013, *Class. Quantum Gravity*, 30, 224003
- Tan C. M. et al., 2018, *ApJ*, 866, 54
- Taylor J. H., Manchester R. N., Huguenin G. R., 1975, *ApJ*, 195, 513
- Theureau G. et al., 2005, *A&A*, 430, 373
- Tyul'bashev S. A., Tyul'bashev V. S., Oreshko V. V., Logvinenko S. V., 2016, *Astron. Rep.*, 60, 220
- Tyul'bashev S. A. et al., 2017, *Astron. Rep.*, 61, 848
- Tyul'bashev S. A., Tyul'bashev V. S., Malofeev V. M., 2018, *A&A*, 618, A70
- van Haarlem M. P. et al., 2013, *A&A*, 556, A2
- van Straten W., Bailes M., 2011, *Publ. Astron. Soc. Aust.*, 28, 1
- van Straten W., Demorest P., Osłowski S., 2012, *Astron. Res. Technol.*, 9, 237
- Wang N., Manchester R. N., Johnston S., 2007, *MNRAS*, 377, 1383
- Yang A., Han J., Wang N., 2014, *Sci. China Phys. Mech. Astron.*, 57, 1600
- Yao J. M., Manchester R. N., Wang N., 2017, *ApJ*, 835, 29
- Yatawatta S. et al., 2013, *A&A*, 550, A136

This paper has been typeset from a  $\text{\TeX}/\text{\LaTeX}$  file prepared by the author.