Geochemical characteristics of basalts related to incipient oceanization: The example from the Alpine-Tethys OCTs

Méderic Amann, Marc Ulrich, Gianreto Manatschal, Eric Pelt, Marie-eva Epin, Julia Autin, Daniel Sauter

► To cite this version:

Méderic Amann, Marc Ulrich, Gianreto Manatschal, Eric Pelt, Marie-eva Epin, et al.. Geochemical characteristics of basalts related to incipient oceanization: The example from the Alpine-Tethys OCTs. Terra Nova, 2019, 32 (1), pp.75-88. 10.1111/ter.12438 . insu-02456734

HAL Id: insu-02456734

<https://hal-insu.archives-ouvertes.fr/insu-02456734>

Submitted on 30 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Geochemical characteristics of basalts related to incipient oceanisation: the example from the Alpine-Tethys OCTs

Amann M.*¹, Ulrich M.¹, Manatschal G.¹, Pelt E.,² Epin M-E.¹, Autin J.¹, and Sauter D.¹

¹Institut de Physique du Globe de Strasbourg (IPGS)-EOST, CNRS-UMR 7516, Université de Strasbourg, 5, rue René Descartes, 67084 Strasbourg, France

²Laboratoire d'Hydrologie et de Géochimie de Strasbourg (LHyGeS), CNRS-UMR 7517, Université de Strasbourg, 1, rue Blessig, 67084 Strasbourg, France

Abstract

Basalts exposed in the Platta and Tasna nappes (SE Switzerland) derive from the Alpine-Tethys Ocean-Continent Transitions (OCT) and overlie SubContinental Lithospheric Mantle (SCLM). We show that the trace element signatures of these basalts differ from Mid-Ocean Ridge Basalts (MORB). Two types of basalts occur in the OCT: a type-1 showing a “garnet signature” that can be modeled by the partial melting of the SCLM in the spinel stability field and a type-2 characterized by an enrichment in incompatible elements that can be explained by the mixing between garnet-pyroxenite-derived melts and the melting of either a depleted MORB mantle or a refertilized SCLM. Based on geological and geochemical observations we propose that the basalts from the Alpine-Tethys OCTs result from a poly-phase magmatic system that carry an inherited SCLM signature. These basalts should therefore be referred to as OCT-basalts rather than as MOR-basalts.

Keywords: Ocean-Continent Transition, Alpine-Tethys, basalt, inheritance, subcontinental mantle, OCT-basalt.

22 **1. Introduction**

23 The geochemical composition of mid-ocean ridge basalts (MORB) is due to the partial melting of a
24 depleted mantle source (DMM). One remaining key question is whether basalts from magma-poor ocean-
25 continent transitions (OCTs) are formed by similar melting processes as MORB or if they reflect more complex
26 mantle processes associated with lithospheric thinning of a sub-continental lithospheric mantle (SCLM).
27 Basaltic samples derived from present-day magma-poor OCTs are rare and limited to few samples drilled at
28 the Iberia-Newfoundland (Chazot et al., 2005; Cornen et al., 1999; Robertson, 2007), the South China Sea (Sun
29 et al., 2016) or the Diamantine OCT (Beslier et al., 2004; Chatin et al., 1998). In contrast, Alpine-Apennine
30 ophiolites expose remnants of the OCTs of the ancient magma-poor Jurassic Alpine Tethys, which provide
31 information not only on the basalts but also on the underlying mantle (Picazo et al., 2016). Previous studies on
32 basalts derived from the fossil Alpine-Tethys OCT show that they are characterized by a wide range of trace
33 element compositions, from depleted basalts (Bill et al., 2000; Desmurs et al., 2002; Durand-Delga et al., 1997;
34 Kramer et al., 2003; Montanini et al., 2008; Piccardo, 2008; Sacconi et al., 2008) to enriched basalts (Frisch et
35 al., 1994; Renna et al., 2018; Steinmann and Stille, 1999; Vannucci et al., 1993) leading to various
36 interpretations concerning the parental sources of these basalts. Over the last decade, a considerable data set
37 has been acquired on the mantle rocks from Alpine-Tethys OCTs (Müntener et al., 2010; Picazo et al., 2016;
38 Piccardo, 2016; Rampone et al., 2008). This new knowledge, combined with the geological observations,
39 enable to better discriminate between the various interpretations and to understand the processes involved
40 in magma-production at deeper parts of the lithosphere during mantle exhumation and lithospheric breakup.

41 In this paper, we present a compilation of Alpine-Tethys basalt chemistry including new data from the
42 Platta and Tasna nappes (SE Switzerland, Fig. 1a). The lack of a major Alpine metamorphic overprint, the
43 excellent exposure and the understanding of the mantle evolution prior and during breakup of these two
44 nappes allowed to propose a tectono-magmatic model that may explain the peculiar geochemistry of the
45 basalts of the Alpine-Tethys OCT.

46 **2. Geological setting**

47 **2.1. Remnants of OCTs in the Alps**

48 The Tasna and Platta nappes represent well preserved remnants of a fossil OCT of the ancient European
49 and Adriatic conjugate rifted margins (Manatschal et al., 2006; Manatschal and Nievergelt, 1997) that were
50 related to the opening of the Jurassic Liguro-Piemonte/Alpine Tethys ocean (Fig. 1b). This ocean probably was
51 a narrow basin (<600km-wide) that developed as a result of an ultra-slow spreading activity (Marroni and
52 Pandolfi, 2007; Li et al., 2013). The Tasna nappe is part of the northeastern European OCT preserving a wedge
53 of continental crust that is laterally replaced by exhumed subcontinental mantle and grades into proto-oceanic
54 crust with first magmatic additions (Manatschal et al., 2006). The Penninic Platta nappe is subdivided into two
55 Alpine tectonic units, the Upper and the Lower Platta units (Supplementary material S1). The Upper Platta
56 nappe is bounded at its top by the Lower Austroalpine Err nappe (part of the Adria distal continental margin)
57 and overlies Middle and Lower Penninic nappes (remnant of the European distal margin; Manatschal and
58 Nievergelt, 1997). Both the Platta and Tasna nappes, as well as the majority of Alpine-Apennine ophiolites,
59 consist of highly serpentinized peridotites intruded by large gabbroic bodies and overlain locally by massive
60 basaltic lava flows, continent-derived extensional allochthons and post-rift sediments (supplementary material
61 S2, Desmurs, 2002; Desmurs et al., 2001; Peters et al., 2008).

62 **2.2. Nature of the exhumed mantle**

63 Two main types of mantle have been distinguished all around the Alpine-Tethys and more particularly
64 in the Tasna and Platta nappes (Fig. 2a, Picazo et al., 2016). The first type identified in the Tasna and Upper
65 Platta units consists of spinel lherzolites containing abundant garnet-bearing pyroxenites (Desmurs, 2002;
66 Müntener et al., 2010; Peters et al., 2008). Major and trace element compositions in clinopyroxene
67 demonstrate the fertile character of these mantle rocks (Picazo et al., 2016). This mantle, referred to as the
68 inherited subcontinental mantle, is similar to the mantle exhumed in the proximal parts of the present-day
69 Iberia-Newfoundland OCTs (Müntener and Manatschal, 2006).

The second type of mantle composing the lower Platta unit consists of spinel lherzolites with minor harzburgites and dunites and shows only few occurrences of pyroxenites (Desmurs, 2002; Müntener et al., 2010, 2004; Peters et al., 2008). This peridotite is pervasively impregnated by ascending tholeiitic melts (Müntener et al., 2010, 2004) that pre-date the Middle Jurassic mantle exhumation (Piccardo, 2016; Rampone et al., 1995). This mantle type has also been described from the distal part of the Iberia margin (Müntener and Manatschal, 2006) and is considered to be the refertilized part of exhumed subcontinental mantle (see type 2 mantle in Picazo et al. 2016).

3. Basalts from the Platta and Tasna OCTs

3.1. Petrology of extrusive magmatic rocks and basaltic intrusions

Extrusive basaltic rocks exposed in the Platta and Tasna nappes (supplementary material S3) occur as hyaloclastites, pillow breccias, pillow lavas and massive lava flows that can reach up to 300 m thickness (Fig. 2a-b). Basalts lie on exhumed mantle and are covered by Upper Jurassic to Lower Cretaceous marine sediments (Fig. 2c). Basalts exhibit intersertal to porphyric textures and contain plagioclase phenocrysts embedded in a fine-grained matrix of plagioclase, chlorite, epidote, pumpellyite and calcite (Fig. 2d). Hyaloclastite breccias consist of fragments of lava flows, which probably derive from the dislocation of former volcanic deposits. A Jurassic age (167-161 Ma) is based on the dating of radiolarian cherts that are either interlayered or overlie the basalts (Bill et al., 2001) and U-Pb on gabbros that are genetically linked to the basalts (Schaltegger et al., 2002).

A younger magmatic event consisting of sills of 10 to 20 m thickness (Fig. 2e) has been found in the Platta nappe. These sills intrude limestones and shales of Lower Cretaceous age (Peters et al., 2008), representing the youngest pre-Alpine magmatic event identified so far in the Alpine OCT. The sills are characterized by fine-grained textures at the rims and coarse-grained at the center. They are composed of plagioclase, chlorite, pumpellyite and epidote (Fig. 2f).

3.2. Major and trace element composition of basalts

A comprehensive description of analytical methods for chemical analysis are given in supplementary material S4. Major element content of our samples is similar to those of other basalts from the Platta nappe (Desmurs et al., 2002; Frisch et al., 1994). Considering both their mineralogical and geochemical compositions, they range between tholeiite and alkali basalt in nature (SiO_2 : 45-50 wt.%; Al_2O_3 : 15-19 wt.%; $\text{Na}_2\text{O}+\text{K}_2\text{O} < 5$ wt.%) and display relatively low Mg# (< 0.65) showing that they represent differentiated melts rather than primitive liquids (Table 1).

On the basis of trace element compositions, two different types of Jurassic basalts can be distinguished (Fig. 3). Type-1 is depleted in incompatible elements, shows a positive Zr anomaly, a low $(\text{La}/\text{Sm})_N$ and is comparable to “normal” MORB. Type-1 shows a large fractionation of HREE over MREE, $(\text{Sm}/\text{Yb})_N=0.82\text{-}1.76$, and a widespread Nb/La ratio (0.55 to 3.25). Samples characterized by medium to high HREE over MREE and a low Nb/La ratio are volumetrically predominant in the Platta-Tasna nappes as well as in the entire Alpine Tethys margins (Fig. 5). The type-2 is enriched in incompatible elements (>10 times PUM concentrations) and high field strength elements (HFSE, Nb, Ta, Zr, Hf; Fig. 3). Type-2 basalts have high $(\text{La}/\text{Sm})_N > 1.3$ and high $(\text{Sm}/\text{Yb})_N > 1.5$ ratios, and are similar to present-day enriched MORB (E-MORB; Gale et al., 2013). Although of a different apparent age, trace element patterns of Cretaceous sills (type-3) are similar to those of Jurassic E-MORB. This geochemical diversity is not restricted to Platta and Tasna basalts but have been described in several ophiolites of the Alpine-Tethys domain (e.g. Bill et al., 2000 and Supplementary material S6).

4. Origin of OCT-basalts

Basalts from this study are representative of basalts from the Alpine-Tethys (Fig. 4). The Cr vs. Y diagram shows that all these basalts result from 5% to 15% partial melting followed by various degrees of fractional crystallization (Fig. 4). However, their variable trace element patterns (Fig. 3), illustrate that the Platta-Tasna basalts (types-1, 2 and 3) as well as other basalts from the Alpine Tethys may not derived from the melting of a depleted mantle source but rather may reflect the melting of another mantle source.

4.1. The depleted “N-MORB”-like basalts (type-1)

Type-1 basalts depletion in LREE makes them to a first order comparable to N-MORB (Erreur ! Source du renvoi introuvable.a). However, most of these basalts as well as most of the Alpine-Tethys basalts substantially differ from present-day MORB by lower Nb/La ratios and higher $(\text{Sm}/\text{Yb})_N$ ratios (Erreur ! Source du renvoi introuvable. and supplementary material S5). High $(\text{Sm}/\text{Yb})_N$ ratios are classically used as a proxy to identify the presence of a garnet-bearing component in the MORB source (Hellebrand et al., 2002; Montanini et al., 2008; Saccani, 2015; Sobolev et al., 2007; Stracke et al., 1999). The melting of a DMM-like source in the spinel stability field hardly produces basalts with $(\text{Sm}/\text{Yb})_N > 1.2$, even at low degrees of melting, while most of the Alpine basalts display $(\text{Sm}/\text{Yb})_N > 1.2$ (see supplementary material S5). Such high $(\text{Sm}/\text{Yb})_N$ ratios are easily simulated by melting a depleted mantle source in the garnet stability. However, both modeling approaches fail to reproduce the low Nb/La ratio that characterize most of the Alpine basalts (Erreur ! Source du renvoi introuvable.b). Since Nb is more incompatible than La ($D_{\text{Nb}} < D_{\text{La}}$), Nb/La ratios tend to decrease in both the residual mantle and the extracted melts with increasing melt degree. Thus, basalts with $\text{Nb}/\text{La} < 0.7$ must be derived from the melting of a mantle source more refractory than the asthenospheric DMM.

The inherited SCLM can hardly be considered as a potential source for type-1 basalts since it is too fertile (mantle type 1a from Picazo et al., 2016) or too depleted (mantle type 1b from Picazo et al., 2016). Similarly, the oceanic mantle (type 3 from Picazo et al., 2016) is too depleted and occurs too oceanward from the location where basalts erupted. However, the refertilized mantle section (mantle type 2; Picazo et al., 2016) may be a plausible source. The geochemical composition of this refertilized mantle is calculated following the modelling approach of Müntener et al. (2010). The starting mantle composition is the PUM, which has undergone 4% partial melting in the garnet (4%) stability field, followed by 4% in the spinel stability field. This residual mantle is then refertilized by ascending melts (up to 12% in volume) deriving from the melting of an asthenospheric depleted source (DMM) in the spinel stability field. Fig. 5a, b shows that the melting of such a refertilized source is able to produce melts similar to type-1 basalts with low Nb/La and high $(\text{Sm}/\text{Yb})_N$. The calculated melting degree of the refertilized mantle is rather low ($< 15\%$), which is consistent with previous estimates

made on Alpine basalts (Desmurs et al., 2002). This model is very similar to the one proposed previously by Meyzen et al., (2003) to explain the peculiar REE signature of some MORB from the northern magma-poor section of the ultraslow-spreading Southwest Indian Ridge. More importantly, this model is consistent with geological observations made by previous studies (Müntener et al., 2010; Picazo et al. 2016).

4.2. The formation of “E-MORB”-like basalts (types-2 and 3)

Aside from type-1 basalts, a few percent of extrusive basalts can be ascribed as enriched-basalts (E-MORB, types-2 and 3). Additional occurrences of E-MORBs have been reported previously in other fossil OCT in the Alps (External Ligurides and Corsica; Montanini et al., 2008; Renna et al., 2018; Sacconi et al., 2008). The enriched nature of these basalts was attributed to the simultaneous melting of the Jurassic ascending asthenosphere (i.e. the DMM) and garnet-pyroxenites. This hypothesis was tested by applying a similar modeling approach to that of Montanini et al., (2008): variable amounts of garnet-pyroxenite derived melts were mixed with (i) aggregated melts of a depleted spinel peridotite source (red mixing lines in **Erreur ! Source du renvoi introuvable.**c, d) and (ii) type-1 basalts extracted from the refertilized SCLM (blue mixing lines in **Erreur ! Source du renvoi introuvable.**c, d). Three different partial melting degrees are considered for the formation of depleted melts ($F=0.05$, 0.1 and 0.15), while a high partial melting degree of 0.25 is considered for the pyroxenite melt (Montanini et al., 2008). Other modeling features (partition coefficients, pyroxenite source composition, etc.) are given in **Erreur ! Source du renvoi introuvable.** caption. Results highlight that E-MORB melts may derive from the mixing between 80-90% of depleted MORB and 10-20% partial melts of garnet-pyroxenite, these results being in good agreement with previous calculations of Sobolev et al., (2007) and Montanini et al., (2008).

4.3. Geodynamic constraints and conclusions

Ophiolites from the Alpine-Tethys domain have provided strong evidence for processes of interaction between inherited SCLM and the ascending asthenosphere (Bodinier et al., 1991; Montanini et al., 2008; Müntener et al., 2004; Müntener and Piccardo, 2003; Piccardo et al., 2004b, 2007; Tribuzio, 2004), which is

different from basalts formed at steady state, mature MORs. Almost all extrusive basalts from the Alpine-Tethys OCT overlie an exhumed, refertilized SCLM (Picazo et al., 2016). Only a few overlie the inherited SCLM in Central Alps (Upper Platta, Malenco, Totalp, Picazo et al., 2016). We therefore suggest here that the specific geochemistry of basalts from the Alpine-Tethys OCT is related to their emplacement over the exhumed, refertilized SCLM. The upward melt infiltration coming from the partial melting of the DMM in the garnet stability field creates the widespread refertilized SCLM (type 2 mantle of Picazo et al. 2016). This melting episode is enhanced by a near-adiabatic decompression regime (McKenzie and Bickle, 1988) during the very first stages of lithospheric thinning and asthenospheric uplift (Piccardo et al., 2004a, Fig. 5a). Melt percolation occurred during lithospheric/crustal thinning and was controlled by tectonic processes and the thermal structure of the extending lithosphere, both of which are not well constrained. Kaczmarek and Müntener (2010) proposed, using the example of the Lanzo shear zone, that percolation occurred in the footwall of a major extensional shear zone that acted as a permeability barrier. The thermal advection related to both asthenospheric upwelling and melt percolation is likely able to increase the thermal gradient in order to reach the solidus of the refertilized SCLM at shallower depths in the spinel stability field (Piccardo et al., 2004a, 2004b). Integrating this inherited history of the SCLM may explain the peculiar geochemical signature characterizing most of basalts from the Alpine-Tethys OCTs that were erupted on exhumed SCLM in the distal part of the hyperextended margin (i.e. type-1 basalts, Fig. 5b). In contrast, the enriched basalts may correspond to low degree melts of the DMM with small amounts of garnet-pyroxenite, both before and after the onset of seafloor spreading (Fig. 5b-c). This study supports the idea that basalts erupting in the Alpine-Tethys OCT have long-lived, poly-phase and variable mantle sources. The peculiar trace element geochemistry of basalts and the well-established geodynamic setting of OCT in the Alps with complex mantle inheritance enable to explain the differences between OCT and MOR basalts. Therefore, basalts formed along OCTs should be referred to as OCT-basalts rather than MOR-basalts.

5. Acknowledgements

We greatly thank Thibault Wiedemann, Nicolas Mattioni, Jean Zielinski and René Boutin for their help during laboratory works. We also thank John Moine for the realization of thin sections. This work is a part of M. Amann Ph.D thesis and has benefited of the financial support provided by the French Ministry of Research and INSU-CNRS through the SYSTER program. None of the authors of the above manuscript has declared any conflict of interest.

6. References

- Anders, E., Grevesse, N., 1989. Abundances of the elements: Meteoritic and solar. *Geochim. Cosmochim. Acta*, **53**, 197–214. doi.org/10.1016/0016-7037(89)90286-X
- Beccaluva, L., Ohnenstetter, D., Ohnenstetter, M., Venturelli, G., 1977. The trace element geochemistry of Corsican ophiolites. *Contrib. Mineral. Petrol.*, **64**, 11–31.
- Beslier, M.-O., Royer, J.-Y., Girardeau, J., Hill, P.J., Boeuf, E., Buchanan, C., Chatin, F., Jacovetti, G., Moreau, A., Munsch, M., 2004. Une large transition continent-océan en pied de marge sud-ouest australienne: premiers résultats de la campagne MARGAU/MD110. *Bull. Soc. Geol. Française*, **175**, 629–641.
- Bill, M., Nägler, T.F., Masson, H., 2000. Major, minor, trace element, Sm-Nd and Sr isotope compositions of mafic rocks from the earliest oceanic crust of the Alpine Tethys. *Schweizerische Mineralogische und Petrographische Mitteilungen*, **80-2**, 131-145.
- Bill, M., O'Dogherty, L., Guex, J., Baumgartner, P.O., Masson, H., 2001. Radiolarite ages in Alpine-Mediterranean ophiolites: Constraints on the oceanic spreading and the Tethys-Atlantic connection. *Geol. Soc. Am. Bull.*, **113**, 129–143.

211 Bodinier, J., Guiraud, M., Dupuy, C., Dostal, J., 1986. Geochemistry of basic dikes in the Lanzo massif
 212 (Western Alps): petrogenetic and geodynamic implications. *Tectonophysics*, **128**, 77–95.

213 Bodinier, J.-L., Menzies, M.A., Thirlwall, M.F., 1991. Continental to oceanic mantle transition—REE and Sr-Nd
 214 isotopic geochemistry of the Lanzo lherzolite massif. *J. Petrol.*, **2**, 191–210.

215 Chalot-Prat, F., Ganne, J., Lombard, A., 2003. No significant element transfer from the oceanic plate to the
 216 mantle wedge during subduction and exhumation of the Tethys lithosphere (Western Alps). *Lithos*, **69**, 69–
 217 103.

218 Chatin, F., Robert, U., Montigny, R., Whitechurch, H., 1998. La zone Diamantine (océan Indien oriental),
 219 témoin de la séparation entre l’Australie et l’Antarctique: arguments pétrologiques et géochimiques.
 220 Comptes Rendus Académie des Sciences, *Earth Planet. Sci.*, **326**, 839–845.

221 Chauvel, C., Bureau, S., Poggi, C., 2011. Comprehensive Chemical and Isotopic Analyses of Basalt and
 222 Sediment Reference Materials. *Geostand. Geoanalytical Res.*, **35**, 125–143. doi.org/10.1111/j.1751-
 223 908X.2010.00086.x.

224 Chazot, G., Charpentier, S., Kornprobst, J., Vannucci, R., Luais, B., 2005. Lithospheric Mantle Evolution during
 225 Continental Break-Up: The West Iberia Non-Volcanic Passive Margin. *J. Petrol.*, **46**, 2527–2568.
 226 doi.org/10.1093/petrology/egi064.

227 Cornen, G., Girardeau, J., Monnier, C., 1999. Basalts, underplated gabbros and pyroxenites record the rifting
 228 process of the West Iberian margin. *Mineral. Petrol.*, **67**, 111–142.

229 Desmurs, L., 2002. Mantle evolution and magmatism in an evolving ocean-continent transition: The Platta
 230 nappe, eastern Switzerland. Diss., Naturwissenschaften, *ETH Zürich*, **14429**, **2002**.

231 Desmurs, L., Manatschal, G., Bernoulli, D., 2001. The Steinmann trinity revisited: mantle exhumation and
 232 magmatism along an ocean-continent transition: the Platta nappe, eastern Switzerland. *Geol. Soc. Lond.*
 233 *Spec. Publ.*, **187**, 235–266.

234 Desmurs, L., Müntener, O., Manatschal, G., 2002. Onset of magmatic accretion within a magma-poor rifted
 235 margin: a case study from the Platta ocean-continent transition, eastern Switzerland. *Contrib. Mineral.*
 236 *Petrol.*, **144**, 365–382. doi.org/10.1007/s00410-002-0403-4

237 Durand-Delga, M., Peybernès, B., Rossi, P., 1997. Arguments en faveur de la position, au Jurassique, des
 238 ophiolites de Balagne (Haute-Corse, France) au voisinage de la marge continentale européenne. *Académie*
 239 *Sci.*, **325**, 973–981.

240 Dürr, S., Ring, U., Frisch, W., 1993. Geochemistry and geodynamic significance of North Penninic ophiolites
 241 from the Central Alps. *Schweiz. Mineral. Petrogr. Mitteilungen*, **73**, 407–419.

242 Ferrara, G., Innocenti, F., Ricci, C.A., Serri, G., 1976. Ocean-floor affinity of basalts from north Apennine
 243 ophiolites: geochemical evidence. *Chem. Geol.*, **17**, 101–111.

244 Frisch, W., Ring, U., Dürr, S., Borchert, S., Biehler, D., 1994. The Arosa Zone and Platta Nappe ophiolites
 245 (Eastern Swiss Alps): geochemical characteristics and their meaning for the evolution of the Penninic Ocean.
 246 *Jahrb Beol BA*, **137**, 19–23.

247 Gale, A., Dalton, C.A., Langmuir, C.H., Su, Y., Schilling, J.-G., 2013. The mean composition of ocean ridge
 248 basalts: MEAN MORB. *Geochem. Geophys. Geosystems*, **14**, 489–518. doi.org/10.1029/2012GC004334

249 Hart, S.R., Dunn, T., 1993. Experimental cpx/melt partitioning of 24 trace elements. *Contrib. Mineral. Petrol.*,
 250 **113**, 1–8.

251 Hellebrand, E., Snow, J.E., Hoppe, P., Hofmann, A.W., 2002. Garnet-field melting and late-stage refertilization
 252 in ‘residual’ abyssal peridotites from the Central Indian Ridge. *J. Petrol.*, **43**, 2305–2338.

253 Herzberg, C., 2006. Petrology and thermal structure of the Hawaiian plume from Mauna Kea volcano. *Nature*,
 254 **444**, 605.

255 Hirschmann, M.M., Stolper, E.M., 1996. A possible role for garnet pyroxenite in the origin of the “garnet
 256 signature” in MORB. *Contrib. Mineral. Petrol.*, **124**, 185–208.

257 Ionov, D.A., Bodinier, J., Mukasa, S.B., Zanetti, A., 2002. Mechanisms and sources of mantle metasomatism:
 258 major and trace element compositions of peridotite xenoliths from Spitsbergen in the context of numerical
 259 modelling. *J. Petrol.*, **43**, 2219–2259.

260 Jochum, K.P., Brueckner, S.M., Nohl, U., Stoll, B., Weis, U., 2008. Geostandards and Geoanalytical Research
 261 bibliographic review 2007. *Geostand. Geoanalytical Res.*, **32**, 509–514.

262 Johnson, K., Dick, H.J., Shimizu, N., 1990. Melting in the oceanic upper mantle: an ion microprobe study of
 263 diopsides in abyssal peridotites. *J. Geophys. Res. Solid Earth*, **95**, 2661–2678.

264 Johnson, K.T.M., 1998. Experimental determination of partition coefficients for rare earth and high-field-
 265 strength elements between clinopyroxene, garnet, and basaltic melt at high pressures. *Contrib. Mineral.*
 266 *Petrol.*, **133**, 60–68. doi:10.1007/s004100050437

267 Kogiso, T., Tatsumi, Y., Nakano, S., 1997. Trace element transport during dehydration processes in the
 268 subducted oceanic crust: 1. Experiments and implications for the origin of ocean island basalts. *Earth Planet.*
 269 *Sci. Lett.*, **148**, 193–205.

270 Kramer, J., Abart, R., Müntener, O., Schmid, S.M., Stern, W.B., 2003. Geochemistry of metabasalts from
 271 ophiolitic and adjacent distal continental margin units: Evidence from the Monte Rosa region (Swiss and
 272 Italian Alps). *Swiss Bull. Mineral. Petrol.*, **83**, 217–240.

273 Lewis, A.D., Smewing, J.D., 1980. The Montgenevre ophiolite (Hautes Alpes, France): Meta—morphism and
 274 trace-element geochemistry of the volcanic sequence. *Chem. Geol.*, **28**, 291–306.

275 Li, X.-H., Faure, M., Lin, W., and Manatschal, G., 2013. New isotopic constraints on age and magma genesis of
 276 an embryonic oceanic crust: The Chenaillet Ophiolite in the Western Alps. *Lithos*, **160-161**, 283-291,
 277 doi.org/10.1016/j.lithos.2012.12.016

278 Liberi, F., Morten, L., Piluso, E., 2006. Geodynamic significance of ophiolites within the Calabrian Arc. *Isl. Arc*,
 279 **15**, 26–43.

280 Manatschal, G., Engström, A., Desmurs, L., Schaltegger, U., Cosca, M., Müntener, O., Bernoulli, D., 2006.
 281 What is the tectono-metamorphic evolution of continental break-up: The example of the Tasna Ocean–
 282 Continent Transition. *J. Struct. Geol.*, **28**, 1849–1869. doi.org/10.1016/j.jsg.2006.07.014

283 Manatschal, G., Müntener, O., 2009. A type sequence across an ancient magma-poor ocean–continent
 284 transition: the example of the western Alpine Tethys ophiolites. *Tectonophysics*, **473**, 4–19.
 285 doi.org/10.1016/j.tecto.2008.07.021

286 Manatschal, G., Nievergelt, P., 1997. A continent-ocean transition recorded in the Err and Platta nappes
 287 (Eastern Switzerland). *Eclogae Geol. Helvetiae*, **90**, 3–28.

288 Marroni, M., Molli, G., Montanini, A., Tribuzio, R., 1998. The association of continental crust rocks with
 289 ophiolites in the Northern Apennines (Italy): implications for the continent-ocean transition in the Western
 290 Tethys. *Tectonophysics*, **292**, 43–66.

291 Marroni, M., Pandolfi, L., 2007. The architecture of an incipient oceanic basin: a tentative reconstruction of
 292 the Jurassic Liguria-Piemonte basin along the Northern Apennines–Alpine Corsica transect. *Int. J. Earth Sci.*,
 293 **96**, 1059–1078. doi.org/10.1007/s00531-006-0163-x

294 Mazzeo, F.C., De Vita, P., Aulinas, M., Arienzo, I., Cirillo, G., Iovine, R.S., Sparice, D., 2016. New constraints on
 295 the origin of the ophiolitic rocks within sin-orogenic turbiditic sequences at Cilento region (southern Italy).
 296 *Geol. Acta Int. Earth Sci. J.*, **14**(3), 299–314.

297 McDonough, W.F., Sun, S.-S., 1995. The composition of the Earth. *Chem. Geol.*, **120**, 223–253.

298 McKenzie, D., Bickle, M. J. 1988. The volume and composition of melt generated by extension of the
 299 lithosphere. *J. Petrol.* **29**, 625–679.

300 Meyzen, C.M., Toplis, M.J., Humler, E., Ludden, J.N., Mével, C., 2003. A discontinuity in mantle composition
 301 beneath the southwest Indian ridge. *Nature*, **421**, 731–733. doi.org/10.1038/nature01424

302 Montanini, A., Tribuzio, R., Thirlwall, M., 2012. Garnet clinopyroxenite layers from the mantle sequences of
303 the Northern Apennine ophiolites (Italy): Evidence for recycling of crustal material. *Earth Planet. Sci. Lett.*,
304 **351**, 171–181. doi.org/10.1016/j.epsl.2012.07.033

305 Montanini, A., Tribuzio, R., Vernia, L., 2008. Petrogenesis of basalts and gabbros from an ancient
306 continent?ocean transition (External Liguride ophiolites, Northern Italy). *Lithos*, **101**, 453–479.
307 doi.org/10.1016/j.lithos.2007.09.007

308 Mugnier, J.-L., Cannic, S., Lapierre, H., 2008. The tholeiites of the Valaisan domain (Versoyen, western Alps): a
309 Carboniferous magma emplaced in a small oceanic basin. *Bull. Société Géologique Fr.*, **179**, 357–368.

310 Müntener, O., Manatschal, G., 2006. High degrees of melt extraction recorded by spinel harzburgite of the
311 Newfoundland margin: The role of inheritance and consequences for the evolution of the southern North
312 Atlantic. *Earth Planet. Sci. Lett.*, **252**, 437–452. doi.org/10.1016/j.epsl.2006.10.009

313 Müntener, O., Manatschal, G., Desmurs, L., Pettke, T., 2010. Plagioclase Peridotites in Ocean-Continent
314 Transitions: Refertilized Mantle Domains Generated by Melt Stagnation in the Shallow Mantle Lithosphere. *J.*
315 *Petrol.*, **51**, 255–294. doi.org/10.1093/petrology/egp087

316 Müntener, O., Pettke, T., Desmurs, L., Meier, M., Schaltegger, U., 2004. Refertilization of mantle peridotite in
317 embryonic ocean basins: trace element and Nd isotopic evidence and implications for crust–mantle
318 relationships. *Earth Planet. Sci. Lett.*, **221**, 293–308. doi.org/10.1016/S0012-821X(04)00073-1

319 Müntener, O., Piccardo, G.B., 2003. Melt migration in ophiolitic peridotites: the message from Alpine-
320 Apennine peridotites and implications for embryonic ocean basins. *Spec. Publ. Geol. Soc. Lond.*, **218**, 69–90.

321 Ottonello, G., Ernst, W.G., Joron, J.L., 1984. Rare earth and 3d transition element geochemistry of peridotitic
322 rocks: I. Peridotites from the Western Alps. *J. Petrol.*, **25**, 343–372.

323 Padoa, E., Saccani, E., Durand-Delga, M., 2001. Structural and geochemical data on the Rio Magno Unit:
 324 Evidence for a new 'Apenninic' ophiolitic unit in Alpine Corsica and its geodynamic implications. *Terra Nova*,
 325 **13**, 135–142.

326 Peters, T., Dietrich, V.J., Office fédéral de la topographie, 2008. Geologischer Atlas der Schweiz Blätt 1256
 327 Bivio.

328 Picazo, S., Müntener, O., Manatschal, G., Bauville, A., Karner, G., Johnson, C., 2016. Mapping the nature of
 329 mantle domains in Western and Central Europe based on clinopyroxene and spinel chemistry: Evidence for
 330 mantle modification during an extensional cycle. *Lithos*, **266–267**, 233–263.
 331 doi.org/10.1016/j.lithos.2016.08.029

332 Piccardo, G.B., 2016. Evolution of the lithospheric mantle during passive rifting: Inferences from the Alpine–
 333 Apennine orogenic peridotites. *Gondwana Res.*, **39**, 230–249. doi.org/10.1016/j.gr.2016.03.001

334 Piccardo, G.B., 2008. The Jurassic Ligurian Tethys, a fossil ultraslow spreading ocean: the mantle perspective.
 335 *Geol. Soc. Lond. Spec. Publ.*, **293**, 11–34. doi.org/10.1144/SP293.2

336 Piccardo, G.B., Müntener, O., Zanetti, A., 2004a. Alpine–Apennine ophiolitic peridotites: new concepts on
 337 their composition and evolution. *Ophioliti*, **29**, 63–74.

338 Piccardo, G.B., Müntener, O., Zanetti, A., Pettke, T., 2004b. Ophiolitic peridotites of the Alpine–Apennine
 339 system: mantle processes and geodynamic relevance. *Int. Geol. Rev.*, **46**, 1119–1159.

340 Piccardo, G.B., Padovano, M., Guarnieri, L., 2014. The Ligurian Tethys: Mantle processes and geodynamics.
 341 *Earth-Sci. Rev.*, **138**, 409–434. doi.org/10.1016/j.earscirev.2014.07.002

342 Piccardo, G.B., Zanetti, A., Müntener, O., 2007. Melt/peridotite interaction in the Southern Lanzo peridotite:
 343 Field, textural and geochemical evidence. *Lithos*, **94**, 181–209. doi.org/10.1016/j.lithos.2006.07.002

344 Pognante, U., Rösli, U., Toscani, L., 1985. Petrology of ultramafic and mafic rocks from the Lanzo peridotite
 345 body (Western Alps). *Lithos*, **18**, 201–214.

346 Rampone, E., Hofmann, A., Piccardo, G., Vannucci, R., Bottazzi, P., Ottolini, L., 1995. Petrology, mineral and
 347 isotope geochemistry of the External Liguride peridotites (Northern Apennines, Italy). *J. Petrol.*, **36**, 81–105.

348 Rampone, E., Hofmann, A.W., 2012. A global overview of isotopic heterogeneities in the oceanic mantle.
 349 *Lithos*, **148**, 247–261. doi.org/10.1016/j.lithos.2012.06.018

350 Rampone, E., Hofmann, A.W., Raczek, I., 1998. Isotopic contrasts within the Internal Liguride ophiolite (N.
 351 Italy): the lack of a genetic mantle–crust link. *Earth Planet. Sci. Lett.*, **163**, 175–189.

352 Rampone, E., Piccardo, G.B., Hofmann, A.W., 2008. Multi-stage melt–rock interaction in the Mt. Maggiore
 353 (Corsica, France) ophiolitic peridotites: microstructural and geochemical evidence. *Contrib. Mineral. Petrol.*,
 354 **156**, 453–475.

355 Renna, M.R., Tribuzio, R., Sanfilippo, A., Thirlwall, M., 2018. Role of melting process and melt–rock reaction
 356 in the formation of Jurassic MORB-type basalts (Alpine ophiolites). *Contrib. Mineral. Petrol.*, **173**, 31.

357 Robertson, A., 2007. Evidence of continental breakup from the Newfoundland rifted margin (Ocean Drilling
 358 Program Leg 210): Lower Cretaceous seafloor formed by exhumation of subcontinental mantle lithosphere,
 359 and the transition to seafloor spreading. Presented at the Proceedings of the Ocean Drilling Program,
 360 Scientific Results, pp 1–69.

361 Rudnick, R.L., Barth, M., Horn, I., McDonough, W.F., 2000. Rutile-Bearing Refractory Eclogites: Missing Link
 362 Between Continents and Depleted Mantle. *Science*, **287**, 278–281. doi.org/10.1126/science.287.5451.278

363 Saccani, E., 2015. Continental Margin Ophiolites of Neotethys: Remnants of Ancient Ocean–Continent
 364 Transition Zone (OCTZ) Lithosphere and Their Geochemistry, Mantle Sources and Melt Evolution Patterns.
 365 *Episodes*, **38**. doi.org/10.18814/epiugs/2015/v38i4/82418

366 Saccani, E., Principi, G., Garfagnoli, F., Menna, F., 2008. Corsica ophiolites: geochemistry and petrogenesis of
 367 basaltic and metabasaltic rocks. *Ophioliti*, **33**, 207.

368 Schaltegger, U., Desmurs, L., Manatschal, G., Müntener, O., Meier, M., Frank, M., Bernoulli, D., 2002. The
 369 transition from rifting to sea-floor spreading within a magma-poor rifted margin: field and isotopic
 370 constraints. *Terra Nova*, **14**, 156–162.

371 Sobolev, A.V., Hofmann, A.W., Kuzmin, D.V., Yaxley, G.M., Arndt, N.T., Chung, S.-L., Danyushevsky, L.V.,
 372 Elliott, T., Frey, F.A., Garcia, M.O., others, 2007. The amount of recycled crust in sources of mantle-derived
 373 melts. *Science*, **316**, 412–417.

374 Steinmann, M., Stille, P., 1999. Geochemical evidence for the nature of the crust beneath the eastern North
 375 Penninic basin of the Mesozoic Tethys ocean. *Geol. Rundsch.*, **87**, 633–643. doi.org/10.1007/s005310050236

376 Stracke, A., Salters, V.J., Sims, K.W., 1999. Assessing the presence of garnet-pyroxenite in the mantle sources
 377 of basalts through combined hafnium-neodymium-thorium isotope systematics. *Geochem. Geophys.*
 378 *Geosystems*, **1**, 12

379 Sun, Z., Stock, J., Jian, Z., McIntosh, K., Alvarez-Zarikian, C., Klaus, A., 2016. Expedition 367/368 scientific
 380 prospectus: South China Sea rifted margin. *Int. Ocean Discov. Program*.

381 Tribuzio, R., 2004. Origin of the Gabbro-Peridotite Association from the Northern Apennine Ophiolites (Italy).
 382 *J. Petrol.*, **45**, 1109–1124. doi.org/10.1093/petrology/egh006

383 Vannucci, R., Rampone, E., Piccardo, G.B., Ottolini, L., Bottazzi, P., 1993. Ophiolitic magmatism in the Ligurian
 384 Tethys: an ion microprobe study of basaltic clinopyroxenes. *Contrib. Mineral. Petrol.*, **115**, 123–137.

385 Venturelli, G., Capedri, S., Thorpe, R., Potts, P., 1979. Rare-earth and other element distribution in some
 386 ophiolitic metabasalts of Corsica, Western Mediterranean. *Chem. Geol.*, **24**, 339–353.

387 Venturelli, G., Thorpe, R.S., Potts, P.J., 1981. Rare earth and trace element characteristics of ophiolitic
 388 metabasalts from the Alpine-Apennine belt. *Earth Planet. Sci. Lett.*, **53**, 109–123.

389 Workman, R.K., Hart, S.R., 2005. Major and trace element composition of the depleted MORB mantle
 390 (DMM). *Earth Planet. Sci. Lett.*, **231**, 53–72.

391

392 7. List of figures and tables

393

394 Fig. 1: (a) Location of major ophiolite massifs in the Alps-Apennine-Corsica (redrawn after Rampone and
395 Hofmann, (2012); Schaltegger et al., (2002). [1]: Tertiary basins; [2]: European Units; [3]: Penninic Units; [4]:
396 Ophiolitic Units; [5]: Adriatic Units. (b) Schematic paleogeographic reconstruction of the Alpine-Tethys during
397 the Late Jurassic (redrawn after Picazo et al., 2016). The reconstruction is based on structural (Manatschal and
398 Müntener, 2009) as well as mantle geochemical (Picazo et al., 2016; Rampone and Hofmann, 2012)
399 considerations. Ophiolite with (*) highlight ambiguous paleo-location. CE: Cecina Valley, CH: Chenaillet, EL:
400 External Ligurides, ET: Erro-Tobbio, GE: Gets Nappe, IL: Internal Ligurides, LA: Lanzo, LP: Lower Platta, MA:
401 Malenco, MM: Monte Maggiore, MV: Mont-Viso, UP: Upper Platta, TA: Tasna, TO: Totalp, ZS: Zermatt-Saas.
402 Br: Briançonnais, Sa: Sardinia.

403

404 Fig. 2: Macroscopic field and microscopic observations of basalts. (a) Panoramic view of the Err-Platta Ocean-
405 Continent Transition zone illustrating the position of the inherited and refertilized (infiltrated) subcontinental
406 mantle. UP: Upper Platta; LP: Lower Platta; SCLM: subcontinental lithospheric mantle. (b) Platta outcrop with
407 gabbro body and refertilized mantle, overlain by basalt flows and radiolarian chert. Note that the topographic
408 surface is controlled by both Alpine thrusts and detachment faults. (c) Radiolarian chert sealing a basalt flow
409 (inverse serie). (d) Photomicrograph of Jurassic extrusive basalt showing plagioclase intersertal texture with
410 occurrence of little epidote. (e) Early Cretaceous basaltic sill intruding post-rift sediments. (f) Photomicrograph
411 of Cretaceous basaltic sill showing subhedral granular albitite (doleritic texture). Ep: epidote, Pl: plagioclase.

412

413 Table 1: Major and trace element compositions of basaltic rocks from the Tasna and Platta nappes. Mg# =
414 $Mg/(Mg+Fe)$ (calculated in molar proportions).

415

416 Fig. 3: Whole-rock incompatible trace element (normalized to the primitive mantle, PM, McDonough and
417 Sun, 1995) and rare-earth element compositions (normalized to C1 chondrite, Anders and Grevesse, 1989) of
418 basalts from the Platta-Tasna nappes. (a) Type-1 depleted Jurassic extrusive basalts, (b) Type-2 enriched
419 Jurassic extrusive basalts, (c) Type-3 Cretaceous basaltic sills. Dashed black lines N-MORB and E-MORB are
420 from Gale et al., (2013) and other basalts from Platta are from Desmurs et al., (2002).

421

422 Fig. 4: Cr (ppm) versus Y (ppm) of whole-rock Platta-Tasna basalts. The partial melting trend starts from the
423 mean DMM (Workman and Hart, 2005). The arrows indicate the calculated trends of the liquids for fractional
424 crystallization of clinopyroxene + plagioclase (cpx + plg) assemblages starting from the relatively primitive
425 basaltic sample 13-04 (residual liquid fraction F up to 0.5). Mineral mass fractions are 0.2cpx+0.8plg. Partition
426 coefficients sources for Y and Cr are from Ionov et al., (2002) and from Hart and Dunn, (1993) respectively.
427 Note that the starting mantle composition does not dramatically change the partial melting models nor the
428 crystal fractionation trends. 1 : Gale et al., 2013, 2 : Other Platta basalts are from Desmurs et al., (2002) and
429 Frisch et al., (1994). Alpine-Apennine basalts (little blue circles) are the overall Alpine-Tethys domain. Internal
430 Ligurides (Ferrara et al., 1976; Ottonello et al., 1984; Rampone et al., 1998; Venturelli et al., 1981), External
431 Ligurides (Marroni et al., 1998; Montanini et al., 2008; Vannucci et al., 1993; Venturelli et al., 1981), Corsica
432 (Beccaluva et al., 1977; Durand-Delga et al., 1997; Padoa et al., 2001; Renna et al., 2018; Saccani et al., 2008;
433 Venturelli et al., 1981, 1979), Western Alps (Bodinier et al., 1986; Chalot-Prat et al., 2003; Lewis and Smewing,
434 1980; Pognante et al., 1985), Central Alps (Desmurs et al., 2002; Frisch et al., 1994), Tuscany (Ferrara et al.,
435 1976; Mazzeo et al., 2016), Calabre (Liberi et al., 2006) and inferred Valais domain (Bill et al., 2000; Dürr et al.,
436 1993; Kramer et al., 2003; Mugnier et al., 2008; Steinmann and Stille, 1999).

437

Fig. 5: (a) Nb/La vs. La showing the compositions of Alpine basalts. The modelling consists of the calculation of Nb/La ratio of aggregated non-modal fractional liquids that derive from the partial melting of the refertilized SCLM. The chemistry of the refertilized mantle source is calculated following the study of Müntener et al. (2010) : a fertile peridotite (primitive upper mantle, PUM; McDonough and Sun, 1995) experienced 4% partial melting in the garnet stability field (A) followed by 4% of additional melting in the spinel stability field (B). The residue is refertilized by adding a 4% fractional melt derived from a spinel DMM source (1% increment, C). A volume of 1%, 3%, 6%, 9% and a maximum of 12% of melts stored in the peridotite is considered (Müntener et al., 2010). The red curves correspond to melts extracted from the calculated refertilized mantle in the spinel stability field at F up to 20%. Partition coefficients are from Johnson et al., (1998), starting mineralogy and modes are from Johnson et al., (1990). (b) (Sm/Yb)_N vs. Yb_N of alpine basalts. Red curves are calculated in the same way as in Fig. 5a. (c) Nb/La vs. La of E-MORB alpine basalts, assuming that they derive from the mixing of aggregated non-modal fractional melts (F=5, 10, 15%) from a spinel DMM source (in red) or a refertilized SCLM (8% refertilization, in blue) with garnet pyroxenite derived melts (F=25%; Montanini et al., 2008). The garnet pyroxenite source derives from high degrees of batch melting (F=40%; Montanini et al., 2012) of a recycled oceanic crust (i.e. an eclogite; Sobolev et al., 2007). The eclogite composition (C_{eclogite}) is calculated assuming that eclogite corresponds to basaltic, N-MORB type crust ($C_{\text{N-MORB}}$) that experienced element mobility during subduction prior being recycled into the mantle: $C_{\text{eclogite}} = C_{\text{N-MORB}} - \text{retention factors}$ where retention factors = $1 - \text{mobility coefficients}$ (Kogiso et al., 1997). The calculated eclogite has a high Nb/La ratio (~2, not shown here), consistent with values calculated by Rudnick et al., (2000). Gt-pyroxenite mineralogy: 60% cpx, 30% opx, 10% gt (Hirschmann and Stolper, 1996); melting modes: cpx=gt=50%. Eclogite mineralogy: 50% gt, 50% cpx. Numbers in italic represent the proportion of mantle derived melts in the mixing. (d) (Sm/Yb)_N vs. Yb_N of alpine basalts. Red and blue curves are calculated in the same way as in Fig. 5c. N-star: N-MORB average $\pm 2\sigma$ (grey area); Present-day MORBs and N-MORB average are from Gale et al. (2013).

Fig. 5 : Schematic representation of the geodynamic model leading to the OCT-basalts. (a) Melt percolation. Near adiabatic upwelling of the asthenosphere (DMM) during this early rifting stage. Onset of asthenosphere

partial melting in the garnet stability field (pink circles), and initial percolation of melts towards the subcontinental mantle (light brown area) during the stretching of the lithosphere. Porous-flow melt percolation induces (i) the refertilization of Permian (?) depleted SCLM, (ii) the formation of plagioclase-bearing peridotite at shallower levels, and (iii) the formation of a weakened mantle wedge for the lithospheric breakup (Piccardo, 2016). The distal subcontinental mantle is melt-impregnated (up to 12%) and refertilized. Major crustal/lithospheric thinning is followed by mantle exhumation and onset of sea-floor exposure of the melt-modified peridotites. (b) Creation of OCT-basalts. Thermal advection (Piccardo et al., 2014) enhanced by porous-flow melt percolation moves the solidus upwards allowing the melting of the refertilized subcontinental mantle in the spinel stability field (light blue circles) forming the type-1 basalts (light blue pipes and volcanoes). Mixing of melts from the partial melting of the DMM (dark blue circles) and garnet-pyroxenite (black dashes, orange circles) creates type-2 basalts (red pipes and volcanoes). (c) Formation of type-3 off-axis basalts (yellow volcanoes) by the same process as type-2. The mixing between the two sources is located at the two edges of the upwelling asthenosphere and is probably related to thermal readjustments. Active spreading ridge followed by MOR-basalt formation caused by the partial melting of the DMM under spinel (shallow) stability field.

8. Supplementary material

Supplementary material S1: (a) Location of the Platta and the Tasna nappes in the Alpine realm. (b) Tectonic units and paleogeographic domains close to Platta and Tasna Nappes. (c) Geological map of Platta modified after Schaltegger et al., (2002). with sample locations. Cretaceous magmatic sills (red) have been magnified for clearance.

Supplementary material S2: Schematic section representing the lithologies and their stratigraphic position in the Platta nappe. Note that the shear-zone in mantle and the sulfide deposit are truncated by the detachment fault. Redrawn after Peters et al., (2008) and own fieldwork observations.

489 Supplementary material S3: Locations and mineralogical assemblage of studied magmatic samples.
490 Coordinates grid is WGS84 UTM 32N, decimal degrees. Fds (Ab): feldspar (albite), Cal: calcite, Chl: chlorite,
491 Cpx: clinopyroxene, Ep: epidote. Crosses indicate relative mineral proportions. Note that samples 13-03 and
492 13-04 are from the Tasna nappe.

493 Supplementary material S4: Analytical method

494 Major and trace elements were measured using the analytical method of Chauvel et al., (2011) at
495 Laboratoire d'Hydrologie et de Géochimie de Strasbourg (LHyGeS, University of Strasbourg, France). For major
496 elements, ~50 mg of sample powder was precisely weighted and dissolved in closed Savilex® beakers using a
497 mixture of 0.3 mL HF 24N and 0.85 mL HNO₃ 14N in a hot plate at 90°C for 5 days. 20 ml of boric acid (H₃BO₃
498 at 20 g.l⁻¹) was added to the solutions after cooling to neutralize excess HF and were further diluted with 250
499 ml of Milli-Q® water. The solutions were then stored in a refrigerator for at least 2 days to ensure complete
500 neutralization of HF by H₃BO₃ and analyzed within a week using an ICP-AES (Thermo Scientific ICAP 6500). For
501 trace elements, 100 mg of sample powder were dissolved in a HF-HClO₄ mixture (5:1) in closed teflon beaker
502 on a hot plate at 140°C for 7 days. The mixture was then completely evaporated at 150°C, and the residue was
503 taken up in concentrated HNO₃. After another complete evaporation, the residue is finally diluted in 40 mL of
504 HNO₃ 7N. The international standard PM-S (gabbro) was analysed repeatedly as unknown to ensure the validity
505 of our measurements. Results show that PM-S concentrations are consistent with the preferred reference
506 values from GeoREM (Jochum et al., 2008). Differences are less than 0.6% for SiO₂ and less than 2% for all
507 other major oxides except for Na₂O, K₂O and P₂O₅ which are present in very low concentrations. In addition,
508 sample 24-04 has been analyzed twice as a duplicate and results show no relevant differences in trace
509 elements between the two aliquots. It is noticed that the anomalous composition of one sample (20-03), which
510 is characterized by very low SiO₂ (36.92 wt.%), high CaO (15.33 wt.%) and high LOI (9.33 wt.%), is explained by
511 the amount of calcite veins and is not used in the discussion.

512

Supplementary material S5: (a) $(La/Sm)_N$ vs. $(Sm/Yb)_N$ of Alpine basalts compared to present-day MORBs (Gale et al., 2013). Sample position reflects their shape of REE patterns, which depends on melting processes and/or the source composition. Most of present-day MORBs (N-MORBs) are explained by moderate to high degrees of partial melting (Gale et al., 2013) of a depleted mantle source in the spinel stability field (Workman and Hart, 2005) as represented by the small graph in the lower left corner. In contrast, most Alpine basalts (including basalts from the Platta and Tasna nappes) are characterized by high $(Sm/Yb)_N$, suggesting the involvement of garnet in the mantle source. Samples that plot in the upper right part of the graph (i.e. E-MORB) can be related to low degrees of melting of a garnet depleted mantle source although they more likely derive from the melting of an heterogeneous, enriched source. N-star: N-MORB average $\pm 2\sigma$ (grey area); E-star: E-MORB average (Gale et al., 2013). PM: partial melting, sp: spinel, gt: garnet. (b) Nb/La vs. $(Sm/Yb)_N$ of Alpine basalts compared to present-day MORBs. The calculated aggregated non-modal fractional melts derived from a spinel lherzolite (in red) and a garnet lherzolite (in green) are shown. The mantle source chemistry is the DMM from Workman and Hart, 2005, and results using the enriched (E-) and depleted (D-)DMM are also shown to not have a narrow vision of the heterogeneous geochemistry of the DMM. Numbers in italic indicate the degree of partial melting. Partition coefficients are from Johnson et al., (1998), starting mineralogy and modes are from Johnson et al., (1990).

(a) Melt percolation

(b) OCT-basalts

(c) MOR-basalts

Refertilization process

- Porous melt flow percolation
- Partial melting of DMM

JURASSIC CRETACEOUS

Tonschiefer and Palombini

Aptychus limestone

- phyllite, schist
- alkali-gabbro
- calc-schist and calcarenite
- limestone
- Mn-deposit
- hyaloclastite flow/pillow breccia
- radiolarian cherts
- basalt flow and pillow basalt
- sulfide deposit
- opihicalcite
- rodingite dike
- gabbro with sigmoidal magmatic pocket
- mantle-derived rocks and garnet-pyroxenite layers

Detachment fault
(gouge, breccia with gabbro and serp. clasts)

Shear-zone
(rodingitized basaltic dikes)