High temperature in-situ study of radiative properties of basaltic dry magmas

Jonas Biren, Leire del Campo, Lionel Cosson, Hao Li, Aneta Slodczyk, Joan Andujar

► To cite this version:

Jonas Biren, Leire del Campo, Lionel Cosson, Hao Li, Aneta Slodczyk, et al.. High temperature in-situ study of radiative properties of basaltic dry magmas. EGU General Assembly 2020, May 2020, Online, France. 10.5194/egusphere-egu2020-546 . hal-03554445

HAL Id: hal-03554445

<https://hal.science/hal-03554445>

Submitted on 3 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

High temperature in-situ study of radiative properties of basaltic dry magmas

Jonas Biren¹, Leire del Campo², Lionel Cosson², Hao Li¹, Aneta Slodczyk^{1,2}, and Joan Andujar¹

¹ISTO, UMR 7327, Université d'Orléans-CNRS-BRGM, 45071 Orléans, France (jonas.biren@cnrs-orleans.fr)

²CEMHTI, UPR 3079, CNRS-Université Orléans, 45071 Orléans, France

Temperature is a key parameter controlling the evolution of lava flows. The hazardous behavior of eruptions prevents direct measurements of hot magmatic bodies. Hence, the temperature of magma is mostly retrieved by using non-contact methods (ground-based or satellite-based thermal cameras) based on measuring the infrared (IR) emission flux (E) of the body [1]. These well-established techniques are however subjected to important errors, ± 100 °C, related to surrounding environment [2], large temperature gradients of cooling lavas [3], constant changes in composition and texture and especially an apparent lack of radiative emission properties during the lava emplacement. Despite that reducing the uncertainties of environmental and thermal gradients when measuring E is ultimately challenging, our study aimed to minimizing the uncertainty in one of the critical hitherto poorly known oversimplified parameters [3,4,5] namely spectral emissivity. Therefore, we performed optical measurements at relevant magmatic temperatures (up to 1200 °C) of representative basaltic dry magmas (MORB, alkaline, calc-alkaline). Emissivity has been systematically determined over a wide spectral ($400\text{-}15000\text{ cm}^{-1}$) and thermal range (from room up to 1200 °C) using a non-contact in situ IR emissivity apparatus [6]. SEM, EMPA and Raman spectroscopy techniques were also used in order to characterize and understand the complex radiative behavior of these natural magmatic compositions. Emissivity varies accordingly with temperature and wavenumber but our results also show that small changes in bulk-rock composition produce drastic changes in emissivity at given T , with iron content and its oxidation state being the main agents controlling this parameter. Appropriate emissivity values will then help to refine current field or (space) satellite IR monitoring data (i.e. Holuhraun 2014-2015, Iceland; [3]) and to implement the thermo-rheological models of lava flows [7] as to support hazard assessment and risk mitigation.

References: [1] Kolzenburg et al. 2017. Bull. Volc. 79:45. [2] Ball and Pinkerton 2006. J. Geophys. Res., 111. [3] Aufaristama et al. 2018. Remote Sens. 10, 151 [4] Harris, A. 2013: Cambridge University press. 728. [5] Rogic et al. 2019 Remote Sens. 2019, 11, 662 [6] De Sousa Meneses et al. 2015. Infrared Physics & Technology 69. [7] Ramsey et al. 2019. Annals of Geophysics, 62, 2.

Keywords: Emissivity, temperature, vibrational spectroscopy, remote sensing, basalt