Journal Pre-proof

Taxonomic additions to the peleciniid wasps (Hymenoptera, Peleciniidae) of the 'mid-Cretaceous' Burmese amber

Corentin Jouault, Valerie Ngô-Muller, Jean-Marc Pouillon, André Nel

PII: S0195-6671(20)30191-9

DOI: <https://doi.org/10.1016/j.cretres.2020.104505>

Reference: YCRES 104505

To appear in: *Cretaceous Research*

Received Date: 14 December 2019

Revised Date: 9 March 2020

Accepted Date: 8 May 2020

Please cite this article as: Jouault, C., Ngô-Muller, V., Pouillon, J.-M., Nel, A., Taxonomic additions to the peleciniid wasps (Hymenoptera, Peleciniidae) of the 'mid-Cretaceous' Burmese amber, *Cretaceous Research*, <https://doi.org/10.1016/j.cretres.2020.104505>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Elsevier Ltd. All rights reserved.

Taxonomic additions to the peleciniid wasps (Hymenoptera, Peleciniidae) of the ‘mid-Cretaceous’
Burmese amber

Corentin Jouault^{a,*}, Valerie Ngô-Muller^{b,d}, Jean-Marc Pouillon^c, André Nel^d

^a *Uni., CNRS, Géosciences Rennes, UMR 6118, F-35000 Rennes, France*

^b *UFR Sciences du Vivant, Université Paris Diderot, Université de Paris, F-75013 Paris, France*

^c *Nivolas Vermelle, France*

^d *Institut de Systématique, Évolution, Biodiversité, ISYEB - UMR 7205 – CNRS, MNHN, UPMC, EPHE, Muséum national d’Histoire naturelle, Sorbonne Universités, F-75005 Paris, France*

* Corresponding author: jouaultc0@gmail.com

ABSTRACT.

A new species of peleciniid wasp, *Eopelecinus inopinatus* sp. nov., is described and figured from Cenomanian Burmese amber based on a single female specimen. Additionally, the examination of a new, well preserved specimen of *Zoropelecinus periosus* Guo, Shih & Ren 2016, allows to emend the diagnosis of this species. These discoveries confirm that Peleciniidae were clearly highly diverse during the Cretaceous. We provide a list of *Eopelecinus* species with their respective metasomal ratio.

Key words.

New species, Burmese amber, *Zoropelecinus*, emended diagnosis, *Eopelecinus*

1. Introduction.

While the extant parasitoid wasp superfamily Proctotrupoidea is modest in diversity compared to the mega-diverse Ichneumonoidea, the former group currently comprises no less than 99 species during the Cretaceous (<http://fossilworks.org>). This superfamily comprises several unplaced genera (viz. †*Amitchellia*, †*Hemerocynipites*, †*Paraulacus*, †*Sinocynipites*) and the families Austroniidae, Heloridae, Jurapriidae, Mesoserphidae, Peleserphidae, Peradeniidae, Proctotrupidae, Roproniidae, and Pelecinidae.

Pelecinidae have a fossil record dating back to the Jurassic, with four species described from China and Mongolia: *Archaeopelecinus jinzhouensis* Shih, Liu & Ren 2009, *Archaeopelecinus tebei* Shih, Liu & Ren 2009, *Cathaypelecinus daohugouensis* Shih, Liu & Ren 2009, *Praescopinus excellens* Rasnitsyn 2008 (Shih et al., 2009; Rasnitsyn, 2008). The family was most diverse during the Cretaceous with more than 48 described species. Following this relative diversity, the Pelecinidae gradually decreased in taxa number and currently comprise only three described species in the sole extant genus *Pelecinus* Latreille 1800, distributed in North, Central, and South America (Johnson and Musetti, 1999). Pelecinid wasps are easily distinguishable from all other Hymenoptera in having a long, thin, pluri-segmented metasoma with elongated segments (conspicuous on female specimens). Females are glossy wasps and possess a very long (up to 7 cm) abdomen, used to lay eggs directly on scarab larvae buried in the soil (Mason, 1984; Johnson and Musetti, 1999). Extinct pelecinids are mostly documented from rock imprints or compression fossils, while six species only have been described from amber. Fossil taxa are known from the Eocene Baltic amber (Brues, 1933; Engel, 2002), Eocene Sakhalin amber (Kozlov, 1974; Johnson, 1998) Eocene of Montana (Greenwalt and Engel, 2014), Turonian New Jersey amber (Engel and Grimaldi, 2006), Cenomanian Burmese amber (Engel et al., 2013; Guo et al., 2016a) and Cretaceous of Russia and China (Zhang and Rasnitsyn 2004; Rasnitsyn, 2008; Feng et al., 2010; Shih et al., 2010; Guo et al., 2016b). Recently, Engel et al. (2013) synonymized Iscopiniinae with Pelecinidae, this synonymization being accepted by Guo et al. (2016a,b) and by us. With the growing number of studies on Burmese amber and re-

cent publications describing new families and genera (Ross, 2019), discoveries of new peleciniid taxa seem inevitable.

In this paper, we report a new species, *Eopelecinus inopinatus* sp. nov., based on one inclusion from the amber collection of the Geological Department and Museum (IGR) of the University of Rennes, France. We also propose an emended diagnosis of *Zoropelecinus periosus* Guo et al., 2016a.

2. Material and methods.

The two amber pieces containing the specimens studied herein derive from the deposits of Noiye Bum in the Hukawng Valley (26° 29' N, 96° 35' E), Kachin State, northern Myanmar (see detailed map in Grimaldi and Ross, 2017: fig. 2). Radiometric data established an Early Cenomanian age (98.79 ± 0.62 Ma) for Kachin amber, based on zircons from volcanic clasts found within the amber-bearing sediments (Shi et al., 2012). Some ammonites found in the amber-bearing bed and within amber corroborates a late Albian–early Cenomanian age (Cruickshank and Ko, 2003; Yu et al., 2019).

The holotype of *Eopelecinus inopinatus* sp. nov. is embedded in a piece of dark orange amber. It contains a nearly complete individual with some metasomal segments missing. The new *Zoropelecinus* is complete and embedded in a piece of clear yellow amber. The holotype of *Eopelecinus inopinatus* sp. nov. is housed in the amber collection of the Geological Department and Museum (IGR) of the University of Rennes, France, under the collection number IGR.BU-007, while the new specimen of *Zoropelecinus* (Pouillon number 6) is deposited in the Musée of Colmar, France. The specimens were examined with a Leica MZ APO stereomicroscope and a Nikon SMZ25. Photographs were made using a Canon EOS 5D mark II mounted on a Leica MZ APO stereomicroscope and a Nikon D800. All images are digitally stacked photomicrographic composites of several individual focal planes, which were obtained using HeliconFocus 6.7 and Nikon Capture

76 NXD. The figures were composed with Adobe Illustrator CC2018 and Photoshop CC2018 soft-
77 wares.

78 Acronyms for measurements and indices are listed below:

79 HL: maximum head length measured laterally from anteriormost point of head capsule to posteri-
80 ormost point of vertex; HW: maximum head width measured at eyes level (eyes included); HH:
81 maximum head height measured at midlength; AL: antennal length; length of all antennomeres
82 combined; ED: maximum diameter of eye as measured in lateral view of head to show full surface
83 of eye; PW: pronotum width in dorsal view; ML: mesosomal length measured in lateral view from
84 anterior surface of pronotum to posterior extension of propodeum; WH: mesosoma height; in side
85 view, maximum height measured from lowermost point of mesopleuron (in front of middle coxa) to
86 dorsal edge of mesosoma; MIL: maximal length of first metasomal segment; MIH: maximal height
87 of first metasomal segment; MIIL: maximal length of second metasomal segment; MIIH: maximal
88 height of second metasomal segment; CI: cephalic index ($HH/HW \times 100$); AI: antennal index
89 ($AL/HW \times 100$); OI: ocular index ($ED/HL \times 100$).

91 **3. Systematic paleontology**

92 Order Hymenoptera Linnaeus, 1758

93 Suborder Apocrita Gerstaecker, 1867

94 Superfamily Proctotrupoidea Latreille, 1802

95 Family Pelecinidae Haliday, 1840

96 Genus *Zoropelecinus* Engel & Grimaldi, 2013 (in Engel et al., 2013)

97 Type species. *Zoropelecinus zigrasi* Engel & Grimaldi, 2013, other species *Zoropelecinus periosus*
98 Guo, Shih & Ren 2016.

100 *Zoropelecinus periosus* Guo, Shih & Ren 2016

101 Fig. 1

102 *New material.* MHNE.2019.X.1 (female, 06, coll. Jean-Marc Pouillon), deposited in the Musée of
103 Colmar, France.

104 *Locality and horizon.* Noiye Bum Hill, Hukawng Valley, Kachin State, Myanmar; lower Cenomani-
105 an, Upper Cretaceous.

106 *Remark.* This fossil is extremely similar to the holotype of *Zoropelecinus periosus*, in the body
107 shape and proportion (Fig. 1A), wing venation and even in the typical pattern of coloration of legs
108 and metasomal segments, which are basally clear and distally darkened (see Guo et al., 2016a: fig.
109 4). We can thus safely attribute this new specimen to the same species. Guo et al. (2016a: 89) had
110 some doubt about the exact number of antennomeres (indicated 14?). *Zoropelecinus zigrasi* has 14
111 antennomeres in females. Engel et al. (2013: 4) indicated the ‘combined lengths of scape and pedi-
112 cel as slightly greater than length of first flagellomere’ in this last species. Our fossil clearly shows
113 14 antennomeres (see Fig. 1B), with a scape twice as long as pedicel and a pedicel slightly shorter
114 than first flagellomere. It thus shows that *Zoropelecinus periosus* differs from *Zoropelecinus zigrasi*
115 in the antennal structure. Nevertheless, the number of antennomeres is variable among species of
116 some other peleciniid genera, thus this difference cannot support a generic separation between the
117 two species of *Zoropelecinus*.

118 We emend the diagnosis of *Zoropelecinus periosus* as follows: antenna 14-segmented with scape
119 twice as long as pedicel, short pedicel slightly shorter than first flagellomere.

120

121 Genus *Eopelecinus* Zhang et al., 2002

122 Type species: *Eopelecinus vicinus* Zhang et al. 2002. Other species: see Tab. 1.

123 ***Eopelecinus inopinatus*** sp. nov.

124 Figs. 2-3

125 *Material.* Holotype IGR.BU-007, nearly complete and well preserved specimen in a rectangular
126 piece of amber measuring $5 \times 4 \times 3$ mm.

127 *Locality and horizon.* Noije Bum Hill, Hukawng Valley, Kachin State, Myanmar; lower Cenomani-
128 an, Upper Cretaceous.

129 *Etymology.* From the Latin word *inopinatus*, meaning “unexpected”. The epithet refers to the rather
130 unexpected nature of the discovery of *Eopelecinus* in Burmese amber.

131 *Diagnosis.* *Eopelecinus inopinatus* sp. nov. differs from all known *Eopelecinus* species in having a
132 unique metasomal segment ratio (for the three preserved segments): 1.0; 1.25; 0.85, the pronotal
133 and mesonotal lateral surfaces areolate; and the propodeum coarsely and regularly reticulate. It also
134 differs from other *Eopelecinus* known from compression fossil in having its relatively smaller size
135 and its 14-segmented antennae.

136 *Description.* Female; male unknown.

137 Head (Fig. 2A, D). Hypognathous, ovoid (CI: 88), wider than high, widest above mid-length in full
138 face view; mandibles dentate, oriented ventrally; eyes large, nearly oval; vertex with 3 ocelli form-
139 ing an equilateral triangle. Antennae 14-segmented (Fig. 2D), filiform; scape 0.16 mm long x 0.06
140 mm wide, pedicel 0.8 x 0.06 mm, F1 0.08 x 0.04 mm, F2 0.08 x 0.05 mm, F3 0.08 x 0.05 mm, F4
141 0.10 x 0.05 mm, F5 0.07 x 0.05 mm, F6 0.08 x 0.05 mm, F7 0.07 x 0.05 mm, F8 0.08 x 0.05 mm,
142 F9 0.08 x 0.05 mm, F10 0.08 x 0.05 mm, F11 0.07 x 0.06 mm, F12 0.14 x 0.05mm; scape narrow
143 basally, widened apically, 2.6x longer than wide, longer than the others segments; pedicel globular,
144 narrower basally, about as long as wide; first flagellomere narrower basally than pedicel apex,
145 gradually increasing in width apically, twice as long as wide; second flagellomere less than twice
146 as long as wide, gradually increasing in width apically; remaining flagellomere similar to second
147 one except tenth through apical one cylindrical slightly long than wide; apical flagellomere longer
148 than wide, rounded apically. Mesosoma (Fig. 2A, C) 0.9 mm long, rectangular, convex in lateral
149 view; pronotum extremely short; mesonotum nearly straight; notauli not clearly visible, metanotal
150 suture conspicuous; mesonotum and propodeum separated by a deep constriction; pronotum and
151 mesonotum with lateral surfaces areolate; propodeum coarsely and regularly reticulate (Fig. 3). Fore
152 and mid legs short. Hind leg with larger coxa; trochanter small; femur swollen medially; tibia nar-

153 row basally, gradually broadened apically and slightly longer than femur; basitarsomere 0.19 mm
154 long and more than twice as short as tibia; second tarsomere 0.07 mm long and more than twice as
155 short as basal tarsomere; third 0.06 mm long, slightly shorter than second; fourth 0.04 mm long,
156 shortest and half as long as third; fifth slightly clavate and sub-equal to second tarsomere. Tarsal
157 claws unarmed.

158 Fore wing 1.3 mm long, with broad costal area; pterostigma short and broadest in its apical third,
159 with posterior margin arched. Hind wing not visible.

160 Metasoma elongated and pluri-segmented (Fig. 2A, B). First metasomal segment 0.75 mm long,
161 elongated, trapezoidal and broadest slightly before its mid-length; tergite and sternite of first meta-
162 somal segment convex laterally and dorso-ventrally, slightly shorter than mesosoma and slightly
163 broader basally than apically, without conspicuous sculpture; second and third metasomal segments
164 tubiform, slender and arched medially.

165 *Color pattern.* Colors black to dark brown (as preserved in amber). Tibiae, tarsi and basal 2/3rd of
166 metasomal segments 2-3-4 (assumed) lighter than the rest of the body; apical third of metasomal
167 segments 2-3-4 (assumed) with a dorsal darker area.

168

169 *Measurements (in mm):* HL 0.4; HW 0.57; HH 0.5; AL 1.22; ED 0.25; PW 0.53; ML 0.9; WH 0.45;
170 MIL 0.75; MIH 0.34; MIIL 1; MIIH 0.12; MIIIL 0.88; MIIIH 0.12 ; CI 88; AI 214; OI 62.

171

172 4. Discussion

173 Although only partially preserved, *Eopelecinus inopinatus* sp. nov. possesses a fore wing
174 only with veins C and R present; the pterostigma short; the pronotum short medially; the pro-
175 podeum reticulate, and the metasoma with one basal segment wide and remaining tubular. Among
176 *Eopelecinus* species, *E. inopinatus* sp. nov. can be easily distinguished by the metasomal ratio (see
177 differential ratio for all species in tab. 1) is one of the major criteria used in the species diagnoses

178 (Zhang et al., 2002; Zhang and Rasnitsyn, 2004; Zhang, 2005; Duan and Cheng, 2006; Liu et al.,
179 2011).

180 In addition to these data (Tabl. 1), *Eopelecinus inopinatus* sp. nov. differs from *E. giganteus*,
181 *E. mecometasomatus*, *E. mesomicrus*, *E. pusillus* and *E. laiyangicus* in its smaller size (about three
182 times as short as *E. giganteus*), different metasomal proportions, and 14-segmented antennae. It dif-
183 fers from *E. leptaleus* in its smaller head (0.4 mm vs. 1 mm) and mesosoma (0.9 mm vs. 2.5 mm),
184 and in antennae with 14 segments vs. 15-segments. *Eopelecinus inopinatus* sp. nov. differs from *E.*
185 *hodoiporus* in its metasomal proportions, a smaller size and a thinner mesosoma. It also differs from
186 *E. eucallus* in its metasomal proportions, mesonotal lateral surfaces, and propodeum coarsely and
187 regularly reticulate. *Eopelecinus inopinatus* sp. nov. differs from *E. minutus*, *E. fragilis* and *E. tu-*
188 *midus* in its larger size and higher metasomal proportions; and from *E. yuanjiawaensis*, *E. similis*,
189 *E. shangyuanensis*, *E. vicinus*, *E. huangi*, *E. exquisitus*, *E. scorpioideus*, and *E. rudis* its smaller
190 size and metasomal proportions (Zhang et al., 2002; Zhang and Rasnitsyn, 2004; Zhang, 2005;
191 Duan and Cheng, 2006; Liu et al., 2011). *Eopelecinus inopinatus* sp. nov. only shares with *E.*
192 *mecometasomatus* Zhang 2005 a second metasomal segment longer than the first one. This charac-
193 teristic suggests a close phylogenetic relation between these two species.

194 The current fossil record of peleciniid wasps suggests an Asian origin for the family, more
195 precisely around China, and a later spreading to Europe through Central and Eastern Asia. Peleci-
196 nids are only known today in the Americas, having probably migrated from Eurasia to North Amer-
197 ica through one of the 'routes' that existed between these regions (Beringia) present from the Late
198 Cretaceous till the Paleocene, as this wasp family is present in the Eocene North America and Baltic
199 amber in the Eocene (Sanmartin et al., 2001; Shih et al., 2009; Brikiatis, 2014). Shih et al. (2010)
200 highlight this hypothesis by comparing the distribution of peleciniid wasps during geological times.
201 The recent studies of Zhang et al. (2018) and Westerweel et al. (2019) suggest an island endemism
202 for the Burmese amber biota. These new data have to be taken into consideration to explain possible
203 speciation and therefore the establishment of a unique Burmese fauna. We can assume that *Eo-*

pelecinus inopinatus sp. nov. comes from a radiation of the genus *Eopelecinus* from the Cretaceous Asian terrane.

According to a phylogenetic analysis based on wing venation, Shih et al. (2009) proposed *Allopelecinus* as the sister lineage of *Eopelecinus*. The description of *Eopelecinus inopinatus* sp. nov. as the youngest representative of the genus seems to corroborate this result. *Eopelecinus inopinatus* sp. nov. possesses the typical wing venation of the genus and allows us to say that the reduced wing venation of the *Eopelecinus* species was maintained at least until the Cenomanian.

5. Conclusion

As aforementioned in the introduction, only six species of Peleciniidae were previously described in amber (Brues, 1933; Engel and Grimaldi, 2006; Engel and Grimaldi, 2013; Guo et al., 2016a). *Eopelecinus inopinatus* sp. nov. is the first record of this genus in amber, which is otherwise known as imprints from n species from China and Russia. This also highlights the underestimated diversity of peleciniid wasps in Burmese amber and a possible origin from Asian biota.

Acknowledgements

We are grateful to Vincent Perrichot for access to the imaging facilities at the University of Rennes, France (IGR); and to the editor (Dr. Eduardo Koutsoukos) and the reviewers (Prof. Alexandr Rasnitsyn and an anonymous reviewer) for insightful comments on the manuscript.

Bibliography:

- Brikiatis, L., 2014. The De Geer, Thulean and Beringia routes: key concepts for understanding early Cenozoic biogeography. *Journal of Biogeography* 41, 1036–1054. <https://doi.org/10.1111/jbi.12310>
- Brues, C.T., 1933. The parasitic Hymenoptera of the Baltic amber. Part 1. *Bernstein Forschungen* 3, 4–178.

- 228 Cruickshank, R.D., Ko, K., 2003. Geology of an amber locality in the Hukawng Valley, northern
229 Myanmar. *Journal of Asian Earth Sciences* 21, 441–455. [https://doi.org/10.1016/S1367-](https://doi.org/10.1016/S1367-9120(02)00044-5)
230 [9120\(02\)00044-5](https://doi.org/10.1016/S1367-9120(02)00044-5)
- 231 Duan, Y., Cheng, S.L., 2006. A new species of Pelecinidae (Insecta, Hymenoptera, Proctotrupoidea)
232 from the Lower Cretaceous Jiufotang Formation of western Liaoning. *Acta Palaeontologica*
233 *Sinica* 45, 393–398
- 234 Engel, M.S., 2002. The fossil pelecinid *Pelecinopteron tubuliforme* Brues in Baltic amber (Hyme-
235 noptera: Pelecinidae). *Journal of Hymenoptera Research* 11, 5–11.
- 236 Engel, M.S., Grimaldi, D.A., 2006. A diminutive pelecinid wasp in Cretaceous amber from New
237 Jersey (Hymenoptera: Pelecinidae). *Northeastern Naturalist* 13, 291–297.
238 [https://doi.org/10.1656/1092-6194\(2006\)13\[291:ADPWIC\]2.0.CO;2](https://doi.org/10.1656/1092-6194(2006)13[291:ADPWIC]2.0.CO;2)
- 239 Engel, M.S., Grimaldi, D.A., Ortega-Blanco, J., 2013. *Zoropelecinus zigrasi*, a pelecinid wasp in
240 mid-Cretaceous amber from Myanmar (Hymenoptera: Pelecinidae). *Novitates Paleoentomologi-*
241 *cae* 4, 1–10. <https://doi.org/10.17161/np.v0i4.4571>
- 242 Feng, H., Shih, C.K., Ren, D., Liu, C.X., 2010. New male pelecinid wasps (Hymenoptera: Pelecini-
243 dae) from the Yixian formation of western Liaoning (China). *Geologica Carpathica* 61, 463–468.
244 <https://doi.org/10.2478/v10096-010-0028-x>
- 245 Greenwalt D.E., Engel, M.S., 2014. A diminutive pelecinid wasp from the Eocene Kishenehn For-
246 mation of northwestern Montana (Hymenoptera: Pelecinidae). *Novitates Paleoentomologicae* 8,
247 1–9. <https://doi.org/10.17161/np.v0i8.4750>
- 248 Grimaldi, D. & Ross, A., 2017. Extraordinary lagerstätten in amber, with particular reference to the
249 Cretaceous of Burma. In: Fraser, N.C. & Sues, H.-D. (Eds.), *Terrestrial conservation lagerstätten:*
250 *Windows into the evolution of life on land*. Dunedin Academic Press, Edinburgh, pp. 287–342.
- 251 Guo, L.C., Shih, C.K., Li, L.F., Ren, D., 2016a. New pelecinid wasps (Hymenoptera: Pelecinidae)
252 from Upper Cretaceous Myanmar amber. *Cretaceous Research* 67, 84–90.
253 <https://doi.org/10.1016/j.cretres.2016.01.009>

- 254 Guo, L.C., Shih, C.K., Li, L.F., Ren, D., 2016b. New peleciniid wasps (Hymenoptera: Peleciniidae)
255 from the Yixian Formation of western Liaoning, China. *Cretaceous Research* 61, 151–160.
256 <https://doi.org/10.1016/j.cretres.2016.01.009>
- 257 Johnson, N.F., 1998. The fossil peleciniids *Pelecinopteron* Brues and *Isocopinus* Kozlov (Hymenop-
258 tera: Peleciniidae). *Proceedings of the Entomological Society of Washington* 100, 1–6.
- 259 Johnson, N.F., Musetti, L., 1999. Revision of the proctotrupoid genus *Pelecinus* Latreille
260 (Hymenoptera: Peleciniidae). *Journal of Natural History* 33, 1513–1543.
- 261 Kozlov, M.A., 1974. An early Cretaceous ichneumon of the family Peleciniidae (Hymenoptera,
262 Pelecinoidea). *Paleontological Journal* 8, 136–138
- 263 Liu, C.X., Gao, T.P., Shih C.K., Ren, D., 2011. New peleciniid wasps (Hymenoptera: Proctotrupoi-
264 dea: Peleciniidae) from the Yixian Formation of western Liaoning, China. *Acta Geologica Sinica*
265 85, 749–757. <https://doi.org/10.1111/j.1755-6724.2011.00480.x>
- 266 Mason, W.R.M., 1984. Structure and movement of the abdomen of female *Pelecinus polyturator*
267 (Hymenoptera: Peleciniidae). *The Canadian Entomologist* 116, 419–426.
- 268 Rasnitsyn, A.P., 2008. Hymenopterous insects (Insecta: Vespida) in the Upper Jurassic deposits of
269 Shar Teg, SW Mongolia. *Russian Entomological Journal* 17, 299–310.
- 270 Ross, A.J., 2019. Burmese (Myanmar) amber checklist and bibliography 2018. *Palaeoentomology* 2,
271 22–84. <http://dx.doi.org/10.11646/palaeoentomology.2.1.5>
- 272 Sanmartin, I., Enghoff, H., Ronquist, F., 2001. Patterns of animal dispersal, vicariance and diversi-
273 fication in the Holarctic. *Biological Journal of the Linnean Society* 73, 345–390.
274 <https://doi.org/10.1006/bijl.2001.0542>
- 275 Shih, C.K., Liu, C.X., Ren, D., 2009. The earliest fossil record of peleciniid wasps (Insecta: Hyme-
276 noptera: Proctotrupoidea: Peleciniidae) from Inner Mongolia, China. *Annals of the Entomologi-
277 cal Society of America* 102, 20–38. <https://doi.org/10.1603/008.102.0103>

- 278 Shih, C.K., Feng, H., Liu, C.X., Zhao, Y.Y., Ren, D., 2010. Morphology, phylogeny, evolution, and
279 dispersal of peleciniid wasps (Hymenoptera: Proctotrupoidea: Peleciniidae) over 165 million years.
280 *Annals of the Entomological Society of America* 102, 20–38. <https://doi.org/10.1603/AN09043>
- 281 Shi, G.H., Grimaldi, D.A., Harlow, G.E., Wang, J., Wang, J., Yang, M., et al., 2012. Age constraint
282 on Burmese amber based on U-Pb dating of zircons. *Cretaceous Research* 37, 155–163.
283 <https://doi.org/10.1016/j.cretres.2012.03.014>
- 284 Yu, T., Kelly, R., Mu, L., Ross, A., Kennedy, J., Broly, P., Xia, F., Zhang, H., Wang, B., Dilcher,
285 D., 2019. An ammonite trapped in Burmese amber. *Proceedings of the National Academy of*
286 *Sciences of the USA*, 116, 11345–11350. <https://doi.org/10.1073/pnas.1821292116>
- 287 Westerweel, J., Roperch, P., Licht, A., Dupont-Nivet, G., Win, Z., Poblete, F., Ruffet, G., Swe,
288 H.H., Thi, M.K., Aung, D.W., 2019. Burma Terrane part of the Trans-Tethyan arc during colli-
289 sion with India according to palaeomagnetic data. *Nature Geoscience* 12, 863–868.
290 <https://doi.org/10.1038/s41561-019-0443-2>
- 291 Zhang, J.F., 2005. Eight new species of the genus *Eopelecinus* (Hymenoptera: Proctotrupoidea:
292 Peleciniidae) from the Laiyang Formation, Shandong Province, China. *Paleontological Journal* 39,
293 417–427.
- 294 Zhang, H.C., Rasnitsyn, A.P., Junfeng, Z., 2002. Peleciniid wasps (Insecta: Hymenoptera: Procto-
295 trupoidea) from the Yixian Formation of western Liaoning, China. *Cretaceous Research* 23:87–
296 98. <https://doi.org/10.1006/cres.2002.0301>
- 297 Zhang, H.C., Rasnitsyn, A.P., 2004. Peleciniid wasps (Insecta: Hymenoptera: Proctotrupoidea) from
298 the Cretaceous of Russia and Mongolia. *Cretaceous Research* 25, 807–825.
299 <https://doi.org/10.1016/j.cretres.2004.08.002>
- 300 Zhang, Q., Rasnitsyn, A.P., Wang, B., Zhang, H., 2018. Hymenoptera (wasps, bees and ants) in
301 mid-Cretaceous Burmese amber: A review of the fauna. *Proceedings of the Geologists' Associa-*
302 *tion* 129, 736–747. <https://doi.org/10.1016/j.pgeola.2018.06.004>

304 **Table. 1.** List of world species of *Eopelecinus* spp. with data on their metasomal ratio (X: refers to
305 incomplete data).

306 **Fig. 1.** *Zoropelecinus periosus* Guo, Shih & Ren 2016, MHNE.2019.X.1, collection J.-M. Poullion,
307 Musée de Colmar. A, habitus in lateral view. B, antenna. Scale bars: 1 mm.

308 **Fig. 2.** *Eopelecinus inopinatus* sp. nov., holotype IGR.BU-007. A, habitus in dorso-lateral view. B,
309 detailed view of mesosoma. C, detailed view of left antenna. Scale bars: 0.5 mm.

310 **Fig. 3.** Line drawing of *Eopelecinus inopinatus* sp. nov., holotype IGR.BU-007. Habitus in lateral
311 view. Scale bars: 0.5 mm.

	Metasomal ratio	Distribution	Age	Reference
<i>Eopelecinus inopinatus</i> sp. nov.	1; 1.25; 0.85; X; X; X	Myanmar	Cenomanian, Cretaceous	This study
<i>Eopelecinus eucallus</i> Zhang 2005	1; 0.6; 0.7; 0.7; 0.6; 0.5	China	Aptian, Cretaceous	Zhang, 2005
<i>Eopelecinus exquisitus</i> Zhang & Rasnitsyn 2004	1; 1; 0.9; 0.6; 0.4; 0.4	Russia	Aptian, Cretaceous	Zhang and Rasnitsyn, 2004
<i>Eopelecinus fragilis</i> Zhang & Rasnitsyn 2004	1; 0.6; 0.6; 0.5; 0.4; 0.3	Mongolia	Valanginian, Cretaceous	Zhang and Rasnitsyn, 2004
<i>Eopelecinus giganteus</i> Zhang 2005	1; 1; 0.9; 1; 0.9; 0.3	China	Aptian, Cretaceous	Zhang, 2005
<i>Eopelecinus hodoiporus</i> Zhang 2005	1; 0.9; 0.8; 0.7; 0.6; 0.5	China	Aptian, Cretaceous	Zhang, 2005
<i>Eopelecinus huangi</i> Liu et al. 2011	1; 0.9; 1; 1; 1; 0.4	China	Aptian, Cretaceous	Liu et al., 2011
<i>Eopelecinus laiyangicus</i> Zhang 2005	1; 0.5; 0.5; 0.4; 0.4; 0.3	China	Aptian, Cretaceous	Zhang, 2005
<i>Eopelecinus leptaleus</i> Zhang 2005	1; 0.8; 0.7; 0.6; 0.4; 0.3	China	Aptian, Cretaceous	Zhang, 2005
<i>Eopelecinus mecometasomatus</i> Zhang 2005	1; 1.1; 1.1; 1; 0.9; 0.6	China	Aptian, Cretaceous	Zhang, 2005
<i>Eopelecinus mesomicrus</i> Zhang 2005	1; 0.5; 0.6; 0.4; 0.5; 0.4	China	Aptian, Cretaceous	Zhang, 2005
<i>Eopelecinus minutus</i> Zhang & Rasnitsyn 2004	1; 0.4; 0.6; 0.5; 0.5; 0.4	Mongolia	Valanginian, Cretaceous	Zhang and Rasnitsyn, 2004
<i>Eopelecinus pusillus</i> Zhang 2005	1; 0.7; 0.7; 0.4; 0.6; 0.5	China	Aptian, Cretaceous	Zhang, 2005
<i>Eopelecinus rudis</i> Zhang & Rasnitsyn 2004	NA	Russia	Aptian, Cretaceous	Zhang and Rasnitsyn, 2004
<i>Eopelecinus scorpioideus</i> Zhang & Rasnitsyn 2004	1; 0.8; 0.8; 0.5; 0.4; 0.3	Russia	Aptian, Cretaceous	Zhang and Rasnitsyn, 2004
<i>Eopelecinus shangyuanensis</i> Zhang et al. 2002	1; 0.4; X; X; X; X	China	Aptian, Cretaceous	Zhang et al., 2002
<i>Eopelecinus similis</i> Zhang et al. 2002	1; 0.8; 0.7; 0.7; 0.6; 0.6	China	Aptian, Cretaceous	Zhang et al., 2002
<i>Eopelecinus tumidus</i> Liu et al. 2011	1; 0.45; 0.7; 0.7; 0.45; 0.45	China	Aptian, Cretaceous	Liu et al., 2011
<i>Eopelecinus vicinus</i> Zhang et al. 2002	1; 0.8; 0.7; 0.6; 0.5; 0.5	China	Aptian, Cretaceous	Zhang et al., 2002
<i>Eopelecinus yuanjiawaensis</i> Duan & Cheng 2006	1; 0.5; 1; 0.9; 0.8; X	China	Aptian, Cretaceous	Duan and Cheng, 2006

Journal Pre-proof

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.