



HAL
open science

Recent tectonic and volcanic evolution of the southern termination of the Main Ethiopian Rift

Zara Franceschini, Raffaello Cioni, Giacomo Corti, Federico Sani, Stéphane Scaillet, Ilaria Isola, Francesco Mazzarini, Asfaw Erbello, Ameha Muluneh, Sascha Brune

► **To cite this version:**

Zara Franceschini, Raffaello Cioni, Giacomo Corti, Federico Sani, Stéphane Scaillet, et al.. Recent tectonic and volcanic evolution of the southern termination of the Main Ethiopian Rift. 21st EGU General Assembly, Apr 2019, Vienne, Austria. insu-02949256

HAL Id: insu-02949256

<https://hal-insu.archives-ouvertes.fr/insu-02949256>

Submitted on 3 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NoDerivatives | 4.0 International License



Recent tectonic and volcanic evolution of the southern termination of the Main Ethiopian Rift

Zara Franceschini (1), Raffaello Cioni (1), Giacomo Corti (2), Federico Sani (1), Stephane Scaillet (3), Ilaria Isola (4), Francesco Mazzarini (4), Asfaw Erbello (5), Ameha Muluneh (6), Sascha Brune (7,8)

(1) Dipartimento di Scienze della Terra, Università degli Studi di Firenze, Florence, Italy, (2) Istituto di Geoscienze e Georisorse, Consiglio Nazionale delle Ricerche, Florence, Italy, (3) Institut des Sciences de la Terre d'Orléans (ISTO), Orléans, France, (4) Istituto Nazionale di Geofisica e Vulcanologia, Sezione di Pisa, Pisa, Italy, (5) School of Applied Natural Sciences, Adama Science and Technology University, Adama, Ethiopia, (6) School of Earth Sciences, Addis Ababa University, Addis Ababa, Ethiopia, (7) German Research Centre for Geosciences GFZ, Potsdam, Germany, (8) Institute of Earth and Environmental Science, University of Potsdam, Germany

The Main Ethiopian Rift, in (East Africa), is a classic example where to analyse the interplay between tectonics and magmatic activity during rifting. This continental rift records indeed all different stages of rift evolution and is characterised by significant volcanic activity spanning from the Oligocene to present. However, despite recent improvements in the knowledge of single volcanic centres thanks to recent scientific projects (e.g., RiftVolc), volcanic activity in many rift sectors still remains poorly documented. In this work we provide new constraints on the timing, evolution and characteristics of the poorly documented volcanic activity at the southern termination of the Main Ethiopian Rift. We present new fieldvolcanological, petrographic, geochemical and geochronological data, together with the results of a statistical analysis of the distribution of volcanic vents from remote sensing, together with petrographic, geochemical and geochronological analyses on the collected samples, for the Mega and the Dilo-Dukana volcanic fields, near the Kenya-Ethiopia border. In line with previous works, our data delineate the occurrence of two distinct groups of rocks: 1) a group of subalkaline basalts, observed only in the Dilo-Dukana volcanic field, forming a Pliocenic lava basement during Pliocene, faulted during an important rifting phase; 2) a group of alkaline basalts, characterising the Quaternary activity in both areas and sampled from Quaternary volcanic vents or from their associated lava flows, that are associated to several, mainly monogenetic volcanic centres. The products of this activity lay in discordance above the lava platform and seal the rift-related faults. Ar/Ar dating allowed to constrain the large basal lava plateau to ~ 3.7 Ma, whereas the youngest volcanic activity characterizing the two areas dates back to 134 ka for Dilo-Dukana and Holocene age for Mega volcanic field. Interestingly, no direct relations are observed between the Pliocene, roughly N-S major boundary faults of the Ririba Rift and the NE-SW structures feeding the Quaternary volcanic activity, indicating a non-correlation between recent volcanism and rifting in the area. Conversely, our analysis suggests a major influence exerted by deep, inherited structures on the distribution of the recent volcanic vents under a complex, local stress field resulting from a major episode of reorganization of extensional structures in the region, recording the history of related to rift propagation and abandonment.