Source rocks in foreland basins: a preferential context for the development of natural hydraulic fractures

Alain Zanella, Peter R Cobbold, Nuno Rodrigues, H Loseth, Marc Jolivet, F Gouttefangeas, D Chew

► To cite this version:

Alain Zanella, Peter R Cobbold, Nuno Rodrigues, H Loseth, Marc Jolivet, et al.. Source rocks in foreland basins: a preferential context for the development of natural hydraulic fractures. AAPG Bulletin, 2021, 105 (4), pp.647-668. 10.1306/08122018162 . insu-03010282

HAL Id: insu-03010282

<https://hal-insu.archives-ouvertes.fr/insu-03010282>

Submitted on 17 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Source rocks in foreland basins: a preferential context for the development of natural hydraulic fractures.

A. Zanella, P.R. Cobbold, N. Rodrigues, H. Loseth, M. Jolivet, F. Gouttefangeas, and D. Chew

AAPG Bulletin published online 28 August 2020
doi: 10.1306/08122018162

Disclaimer: The AAPG Bulletin Ahead of Print program provides readers with the earliest possible access to articles that have been peer-reviewed and accepted for publication. These articles have not been copyedited and are posted “as is,” and do not reflect AAPG editorial changes. Once the accepted manuscript appears in the Ahead of Print area, it will be prepared for print and online publication, which includes copyediting, typesetting, proofreading, and author review. ***This process will likely lead to differences between the accepted manuscript and the final, printed version.*** Manuscripts will remain in the Ahead of Print area until the final, typeset articles are printed. Supplemental material intended, and accepted, for publication is not posted until publication of the final, typeset article.

Cite as: Zanella, A., P.R. Cobbold, N. Rodrigues, H. Loseth, M. Jolivet, F. Gouttefangeas, and D. Chew, **Source rocks in foreland basins: a preferential context for the development of natural hydraulic fractures.**, (*in press; preliminary version published online Ahead of Print 28 August 2020*): AAPG Bulletin, doi: 10.1306/08122018162.

Source rocks in foreland basins: a preferential context for the development of natural hydraulic fractures.

A. Zanella^{a*}, P.R. Cobbold^b, N. Rodrigues^{b,c}, H. Loseth^c, M. Jolivet^b, F. Gouttefangeas^d, D. Chew^e

^a*Le Mans Université, L.P.G-UMR 6112, avenue Olivier Messiaen, 72085 Le Mans, Cedex 9, France*

^b*Géosciences Rennes, UMR 6118, Université de Rennes 1, 35042 Rennes, France*

^c*Statoil Research Centre, 7005 Trondheim, Norway*

^d*Centre de Microscopie Electronique à Balayage et Microanalyse, Université de Rennes 1, 35042 Rennes, France*

^e*Department of Geology, Trinity College Dublin, Dublin 2, Ireland*

*Corresponding author: alain.zanella@univ-lemans.fr

KEYWORDS

beef veins; natural hydraulic fractures; petroleum source rocks; fluid overpressure; foreland basins

ABSTRACT

Bedding-parallel veins of fibrous calcite (also called BPV or ‘beef’) occur in many sedimentary basins, especially those containing low-permeability strata with organic source material for petroleum. The formation of such veins is often linked with fluid overpressure in these source rocks. In this review, we demonstrate that beef veins are most commonly present in foreland basins worldwide or in basins that recorded a compressive tectonic period. The formation of beef veins is related to two main phases: (1) the initiation of bedding-parallel fracture and (2) the infilling of the fracture.

Previous structural studies have shown that formation of beef veins occurred during a period of compressive stress activity. This is especially the case for the Wessex Basin (UK) and the Neuquén Basin (Argentina). Here we provide more observations for other basins: the Cordillera Oriental (Colombia), the Paris Basin (France), the northern Pyrenees (France), the Uinta Basin (US), the Tian Shan Mountains (central Asia) and the Appalachian Mountains (US). In the Paris Basin, beef vein formation is dated at 155 Ma (U/Pb calcite method) and is coeval with the compressional deformation in the eastern part of the basin.

Because of the timing of generation for such veins and even if the theory and the experiments of fracturing demonstrate that bedding-parallel fractures can be generated only with a distributed fluid overpressure, the formation of beef veins seems to be a consequence of both fluid overpressures and a compressional tectonic stress.

1. Introduction

Many of the major mountain ranges around the world have resulted from recent horizontal shortening and vertical thickening (Figure 1). Others have resulted from rifting, strike-slip faulting or uplift associated with magmatism. In contrast, most older mountain ranges have tended to disappear by active erosion of their sharp topographic profiles. For recent shortening and thickening, the main causes are (1) subduction beneath active continental margins, especially in the Atlantic and Pacific oceans, or (2) continental collision, for example between Europe and Africa or central Asia and India (Figure 1). In such tectonic settings, there has been progressive development of adjacent foreland basin systems as a result of thrusting and local sedimentation and in many cases, they contain petroleum systems.

Because they are commonly hydrocarbon-bearing, foreland basins are major targets for the study of source rocks and reservoirs. In the last twenty years, organic-rich source rocks have been extensively studied because of their hydrocarbon potential. Within these sedimentary units, many fractures occur and some of them can affect the permeability as well as the cap capability of the source rocks. This is especially the case for the natural hydraulic fractures, such as bedding-parallel veins (also called BPV or 'beef'). The first examples of beef to be described were likely those in the Wessex Basin of Southern England (Buckland & De la Beche, 1835). The veins were easily observable along coastal cliffs and provided useful material for building walls and roads. More generally, bedding-parallel veins are common worldwide in sedimentary basins, especially those containing hydrocarbons, indicating that the host rock has reached maturity (Cobbold et al., 2013; Gale et al., 2014). The formation of such veins is often linked to fluid overpressure during hydrocarbon generation (Rodrigues et al., 2009; Zanella et al., 2015a). Beef veins therefore consist of natural hydraulic fractures, infilled by a fibrous mineral, such as calcite, gypsum or quartz (Cobbold et al. 2013). Other mechanisms are also involved, but seem to be more minor, such as, for example, the force of crystallization (e.g. Taber, 1916; Means & Li, 2001; Gratier et al., 2012). Recent studies have shown that many bedding-parallel veins formed during horizontal shortening, as in the Wessex Basin, southern England, (Zanella et al., 2015b), in the Bristol Channel Basin, England and Wales, (Meng et al., 2017), or in the Neuquén Basin of western Argentina (Rodrigues et al., 2009; Ukar et al., 2017; Ukar et al. 2018).

This paper presents a worldwide review of bedding-parallel veins (natural hydraulic fractures) within foreland basins and discusses the development of such natural hydraulic fracturing processes within this particular tectonic basin setting. We also present new observations and data to complete those of previous studies, particularly with respect to four localities: (1) the Northern Pyrenees, France; (2) the Uinta Basin, USA; (3) the Tian Shan Basin, China and (4) the Appalachian Basin, USA.

2. Formation of beef veins

The simplest calcite beef veins, which form in tectonically quiescent basins, are typically bedding-parallel (horizontal), of regular thickness and contain calcite fibers, which have formed almost vertically and antitaxially, separating the two boundaries of the fracture. The best evidence for epitaxial separation comes from the presence of flattened fossils (for example, ammonites), which remain in the central part of the vein, while counterparts of them are visible at the two outer boundaries, but displaced along the directions of the calcite fibers, which may be somewhat oblique (Rodrigues et al., 2009). Other forms of calcite beef are cone-in-cone structures, which likely form as a result of shear faulting during epitaxial growth of the calcite veins (e.g. Cobbold et al. 2013).

Two successive phases characterize the formation of a beef vein: (1) the initiation of the fracture and (2) the opening of the fracture. For the initiation of the fracture, the model can explain the generation of horizontal hydraulic fractures without external tectonic stresses (e.g. compressive stress) (Cobbold et al. 2007, Mourgues & Cobbold, 2003). However, much of our knowledge of the mechanisms involved in such processes come from experimental models developed during the early 21st century. In the early iterations of these experimental models, the pore fluids were injected from the outside, at measured rates and pressures (Cobbold & Castro, 1999; Mourgues & Cobbold, 2003). Thus, it became clear that vertical gradients of overpressure counteract the weight of the granular material. When noncohesive materials were also compressed horizontally, the pore fluid gradient facilitated detachments for thrust faults and caused these to become more nearly horizontal (Cobbold et al., 2001). When the material was cohesive, the vertical pressure gradient generated horizontal tensile fractures, which opened progressively (Cobbold & Rodrigues, 2007). Moreover, when the models were also compressed horizontally, the resulting stresses facilitated the formation of horizontal tensile fractures. Because the early experimental models simulated the origin of the fluid overpressure as an injector-type system, they were not adapted for the study of the origin and mechanisms involved in the development of such fluid overpressures.

Thus, to be able to study the origin of fluid overpressures and the parameters involved in the process of natural hydraulic fracturing within source rocks, later experiments have used transition phases (solid to liquid), to produce a fluid within a closed experimental setup. From solid organic particles (such as beeswax microspheres), models were able to generate a fluid by chemical compaction-like mechanisms, which occur during burial of organic-rich source rock (Fig. 2; Lemrabott & Cobbold, 2010; Zanella et al., 2014a). In such models, the wax was able to melt, on heating the model from below. The resulting decrease in underlying support allowed solid particles to move downwards, causing compaction of the underlying framework and an increase in pore fluid

pressure, by a mechanism of load transfer (Zanella et al. 2014a). In other models, which were saturated with water but not subject to horizontal shortening (Figure 2), the overpressure developed soon after the temperature had reached the melting point of beeswax, at the base of the model. This overpressure became great enough to cause horizontal hydraulic fracturing of the wax-rich layer. Molten wax then migrated vertically and horizontally through the pore space and filled the opening generated by hydraulic fractures. In similar models, which were also subject to horizontal shortening, the hydraulic fractures developed more strongly, becoming wider and longer. Furthermore, many of the fractures varied laterally in thickness and orientation, as a result of local folding and thrust faulting.

For the filling of the natural hydraulic fractures, beef veins appear to have incrementally grown by successive phases of crystallization (e.g. Taber, 1918; Ramsay, 1980; Rodrigues et al., 2009). Thus, fibers appear to have grown incrementally during the displacement of the edges of the veins by a crack-seal mechanism (Ramsay, 1980) or more continuously (Taber 1918; Durney & Ramsey, 1973; Means & Li, 2001). The mechanisms involved in the formation of beef veins are complex and several of them have been postulated by previous authors. The opening of the vein implies a key mechanism already describe by several authors: the force of crystallization (e.g. Baker et al., 2006; Bons, 2001; Bons et al., 2012; Keulen et al., 2001; Mean & Li, 2001). Indeed, several experiments have demonstrated the great importance of the force of crystallization in the beef vein opening (Bons & Jessel, 1997; Hilgers et al., 2001; Nolle et al., 2005, 2009) Thus, the tectonic stress, the pore fluid pressure (fluid overpressures) and the force of crystallization appear to be the major mechanisms (e.g. Sibson, 2003; Shearman et al., 1972; Stoneley, 1983; Taber, 1916; Gratier et al., 2012).

In many basins worldwide, the formation of beef veins seems to be linked to the generation of hydrocarbons, as showing by scanning electron microscopy for calcite beef of the Wessex Basin (Zanella et al., 2015b), the Neuquén Basin (Rodrigues et al., 2009; Ukar et al. 2017, 2018), the Magellan Basin (Zanella et al., 2014b) and the Paris Basin. All of these examples show that beef veins contain inclusions of organic matter (liquid, solid or both) within or next to the calcite crystals, indicating that the organic matter was diffusing, while the calcite crystals were growing. Other examples of calcite beef, e.g. in the Lourdes area, the Uinta Basin or the Appalachians, may either include or be adjacent to veins of organic material. Investigating the relationship between the organic matter and the hydraulic fracturing, Zanella et al. (2014a) showed with their experiments that the maturity degree is a key parameter to explain the increase of pore fluid pressure within source rocks and that a horizontal compressive stress can favor the development of horizontal hydraulic fractures.

3. Worldwide distribution of beef veins within foreland basins

3.1. Foreland basins

A global topographic map (Figure 1) illustrates the occurrence of significant present-day mountain ranges, especially along continental margins such as the Andes or within zones of continental collision (such as central Asia or the Alps). Many recent foreland basins have formed next to these current mountain ranges. Additional evidence for compressional reactivation of such mountains comes from stress maps on a global scale, which show directions of horizontal compressional stress (e.g. Heidbach et al., 2009). Although many such mountain ranges have formed recently and expose Mesozoic or younger rocks, others expose older rocks (Paleozoic or even Precambrian). In many cases, however, older mountain belts have been subject to recent localized reactivation, such as the Appalachian Mountains in the eastern US and Canada, but also parts of the mountains of Scotland, Norway, Brazil, northern Africa or Australia. Indeed, such reactivation explains why the mountains still exist today, despite relatively active erosion.

Foreland basins (such as those next to the Andes) tend to have relatively low and flat upper surfaces, but deep basements, which have subsided, as a result of (1) the weight of nearby mountains, (2) the propagation of outward-verging thrust faults, (3) the local accumulation of sediments derived from erosion of the mountains, (4) the accumulation of evaporites or limestones within local depressions and (5) the accumulation of volcanic and plutonic rocks propagated out from the mountain belt, especially in subduction zone settings.

Petroleum source rocks (commonly of Mesozoic age, but also of Paleozoic age) often have formed within many of these foreland basins, in part because the surface depressions yield sufficient accommodation space, but also because of suitable climatic and environmental conditions. Progressive subsidence of foreland basins leads to heating and maturation of source rocks at depth. Fluid overpressure tends to be common within such source rocks, as a result of the heating, maturation, increases in volume and chemical compaction (under the weight of the overlying rock column and the effect of horizontal tectonic compression). Such heating is mainly associated with burial but may also be due to magmatic intrusions, especially in subduction zone systems (Zanella et al. 2015a).

Finally, thrust detachments are common in foreland basins, especially within overpressured source rocks, where seepage forces counteract the weight of gravity and sedimentary strata providing an anisotropic mechanical response (Cobbold et al., 2001).

3.2. Calcite beef within foreland basins (Mesozoic host rocks)

Calcite beef is common within petroleum source rocks, especially those which have reached maturity (Table. 1). In the last few years, several authors have described examples of beef within foreland basins, especially within Mesozoic rocks of the Wessex Basin (UK), the Neuquén Basin

(Argentina), the Magellan Basin (Chile and Argentina) and the Paris Basin (France). Other examples have been described worldwide (e.g. Cobbold et al., 2013; Gale et al. 2014), although not necessarily in such detail.

Where a basin has been subject to compressional tectonics, it may also contain synchronous calcite beef. Typically, the veins will then be variable in thickness and orientation (Figs. 3, 4 and 5). A vein may vary in thickness across small amplitude folds becoming thicker within synclines and thinner over anticlines. Other variations in thickness may occur across reverse faults (see the examples from the Wessex Basin; Figure 3). In the Magellan Basin, Zanella et al. (2014b) show that beef vein occurrences are more numerous closed to main thrust faults. Finally, some veins may have formed obliquely to bedding, (i.e. nearly horizontally) after the bedding has rotated, as a result of compressional deformation. These structures are similar to some of those described from the experimental modeling with fluid overpressures (Zanella et al., 2014a).

3.2.1. Wessex Basin, southern England

The Wessex Basin covers much of southern England (Figure 3A) and consists of Mesozoic and Cenozoic rocks that young in general towards the east, but is subject to folding and faulting, especially near the southern basin margin (Wight-Bray and Purbeck faults). A north-south section across the Purbeck fault (Figure 3A) and its restoration, published by Underhill & Stoneley (1998), show that much of the slip has been of Late Cretaceous to Cenozoic age. The Wessex Basin also has several oilfields, which produce hydrocarbons from Jurassic source rocks (Underhill & Stoneley, 1998).

Within the Liassic source rocks, calcite beef are common and have drawn attention for many years from geologists (e.g. Lang et al., 1923; Richardson, 1923; Buckland & De la Beche, 1835; Marshall, 1982), who at first attributed them to early diagenesis of the sediments. However, later work revealed epitaxial growth of calcite fibers (e.g. Cobbold & Rodrigues, 2007), after splitting of the rock.

On studying other calcite veins in the Purbeck Formation (latest Jurassic or Early Cretaceous) at Lulworth Cove (above the Purbeck fault), Zanella et al. (2015b) discovered that many of them are localized very close to thrust faults (Figure 3). These veins show sigmoidal fibers (Fig. 3B & C) as well as variations in thickness around folds (Figure 3D). Such observations argue for a vein development synchronous with a horizontal shortening period. Indeed, recent uranium-lead dating of calcite (Chew et al., work in progress) has produced Albian ages (about 107 Ma) for such compressional beef veins, both at Lulworth Cove and also at Charmouth (in the Shales-with-Beef Formation of Liassic age). Thus, the beef veins in the Wessex Basin formed during the onset of inversion tectonics.

3.2.2. Neuquén Basin, western Argentina

The Neuquén Basin covers an area of western Argentina, in the Andean foreland (Figure 4A). Its western margin is in the sub-Andes and it contains many north-northwest-trending folds. An eroded anticline exposes large areas of the Vaca Muerta Formation (Figure 4A), which is an Upper Jurassic organic-rich shale and a world-class source rock for oil. A geological section (Figure 4B, after Vera et al., 2014), through the city of Chos Malal (eastern side of the cross-section on Figure 4B), illustrates uplifted blocks of Palaeozoic basement, as well as thrust faults, many of which have detached near the base of the Vaca Muerta Formation.

Also, in the Vaca Muerta Formation are many examples of calcite beef. Although some of these veins are as much as 30 cm thick and are laterally continuous over distances of hundreds of meters (e.g. Cobbold et al., 2013, fig. 3A; Gale et al., 2014), others show local variations in thickness across minor folds and reverse faults (Figure 4A), indicating that they formed during horizontal compression and shortening. By analyses of the domal structures from beef veins, Ukar et al. (2017) concluded that beef veins, in the area of Loncopué, developed during the Late Cretaceous. This conclusion is consistent with Rodrigues (2008) and Weger et al. (2018), which concluded by isotopes analysis that beef veins growth occurred at depth related to high temperatures (120°C to 185°C). Beef veins also occur elsewhere in the Neuquén Province (Cobbold et al., 2013) within shales of the Los Molles Formation (Lower Jurassic) or Agrio Formation (Lower Cretaceous). Finally, beef veins also occur in the Vaca Muerta Formation further north in Mendoza Province (Zanella et al., 2015a).

In summary, there is good evidence in the Neuquén Basin for recent overpressure, thrust detachments and multiple veins of compressional calcite beef, all within source-rock shale. Much of the heating was probably related to foreland subsidence, but some may have come from magmatic intrusions, which are abundant, especially in Mendoza Province (Zanella et al., 2015a).

3.2.3. Edges of the Cordillera Oriental, central Colombia

The Cordillera Oriental is an Andean mountain belt in eastern part of Colombia (Fig. 5A, after Mora et al., 2015). Within it, folds and thrusts trend north-northeast, affecting mainly Cretaceous rocks. The geological section shows uplift of Paleozoic to Precambrian basement between reverse faults, as well as major folds within Cretaceous sedimentary rocks, some of which are source rocks for petroleum (Figure 5A). On the western side of the Cordillera, folds and thrusts verge towards the west (Figure 5A) and Cretaceous shales contain layers of calcite beef (white to yellow), which vary in thickness across folds (Figure 5C). At the eastern edge of the Cordillera, in the Macanal Formation (Lower Cretaceous), some beef contains thin layers of yellow pyrite, which have folded (Figure 5B, bottom right), while calcite has grown antitaxially above and below them, filling synclines more than anticlines, indicating a synchronous development with a horizontal shortening.

In summary, there is good evidence, at both edges of the Cordillera Oriental, for recent overpressure, thrust detachments and multiple veins of compressional calcite beef veins, all within organic-rich mature shale source rocks.

3.2.4. Paris Basin, northcentral France

The Paris Basin occupies a large area of northcentral France (Figure 6A) and consists mainly of Triassic, Jurassic and Cretaceous marine strata, but also some Cenozoic strata (mainly lacustrine) in its central part (Guillocheau et al., 2000). A cross-section (Figure 6A) shows that the basin has undergone some uplift and erosion, especially on its eastern margin. In the past, there have been some descriptions of calcite beef or cone-in-cone structures, mostly within Jurassic shales of the Schistes Carton Formation (lower Toarcian, about 185 Ma) on the eastern basin margin (e.g. Denaeyer, 1943, 1947), but also within Callovian shales at the northern edge (Voisin, 1999). The Schistes Carton have been subject to much exploration as potential source rocks for oil, and this has revealed the presence of fluid overpressure at several localities. Cobbold et al. (2015) investigated a ditch through the Schistes Carton Formation at Gélaucourt, near Nancy at the eastern edge of the basin (Fig. 6B, left). The ditch (Figure 6B, left) contains many beef veins, several cm thick and several meters long (Fig. 6B, right). These veins had been identified previously by Denaeyer (1947). They consist mainly of calcite fibers, almost perpendicular to bedding, but they also contain some inclusions of hydrocarbons, which are visible in hand specimens, but especially by scanning electron microscopy (Figure 7B; Cobbold et al., 2015). Recently, by uranium-lead dating of calcite, we have determined the age of formation of this calcite beef to be 155 Ma (Figure 7A). This is about 30 m.y. younger than the stratigraphic age of the host rock (about 185 Ma). Indeed, the age of the calcite almost coincides with the age of onset of compressional deformation on the eastern edge of the Paris Basin (see Guillocheau et al., 2000).

4. New results: field data for calcite beef in other compressional basins

4.1. Lourdes, northern Pyrenees, France (locality 22)

At the northern edge of the Pyrenees, a major thrust fault zone separates this mountain range to the south from the Aquitaine Basin to the north. Geological maps of Lourdes (e.g. BRGM, Carte Géologique Détaillée de la France, 1:50 000, sheet XVI-46, 1970) show multiple folds and faults within Mesozoic strata, especially near the town of Lourdes (Choukroune, 1969). Indeed, an oblique view of the Pic du Béout hills, to the east of Lourdes (Figure 8A) shows repetitions of resistant white Aptian-Albian limestones, which dip 30° to 40° towards the south. Between the limestones are darker and softer layers of upper Aptian shale, which have acted as thrust detachments and exhibit some

cleavage and down-dip lineations. The plains surrounding the hills consist mainly of unconformably overlying Upper Cretaceous flysch. Some of the upper Aptian shale contains layers of fibrous calcite beef (Figure 8B), especially near the village of Aspin-en-Lavedan (Figure 8A). Many of the veins are almost parallel to bedding, whereas younger ones are less steeply dipping and therefore somewhat oblique to bedding. The dipping of the sediments is the result of the compressional deformation from the Late Cretaceous to present day. Because the compression is active since these geological times, beef veins in this area are synchronous with this deformation. Thus, they probably formed once the bedding had rotated and partly as a result of compressional deformation. Our analyses, by scanning electron microscopy (Figure 8C), have revealed steep fractures across a flat-lying vein of calcite beef, the fractures containing much more carbon (orange color) than pure calcite.

It also happens that the Aquitaine Basin is hydrocarbon prone, especially beneath low-permeability upper Aptian shale (Biteau & Canérot, 2007). Thus, there is evidence, around Lourdes, for synchronicity of (1) Late Cretaceous or Cenozoic compressional deformation, (2) generation and accumulation of organic-rich fluids and (3) formation of calcite beef.

4.2. Uinta Basin, Utah (locality 12)

While investigating the presence of bitumen veins in the Uinta Basin, we discovered veins of calcite beef (up to 3 cm thick), which are common within Eocene shales of the Green River Formation, which are also source rocks for petroleum with several major oil fields. The Uinta Basin is a typical intermontane basin (Fig. 9A). At its northern edge, the basin abuts a major thrust fault, which has uplifted Paleozoic basement (part of the Uinta Mountains). At the northern margin of the basin, overpressure occurs within hydrocarbons of the Altamont-Bluebell oil field (Dubiel, 2003). Elsewhere, exposures of Green River shales (for example, within open mines) contain visible beef, which consist mainly of calcite with dominantly vertical fibers (Fig. 9B). However, in some places the veins consist not only of calcite, but also of solid hydrocarbons (gilsonite; Fig. 9C). Some veins also contain bitumen between fibers of calcite. Thus, there is evidence for synchronicity of compressional tectonics, maturation of source rocks and growth of beef veins.

4.3. Tian Shan Mountains, central Asia (localities 30 to 32)

In central Asia, the Tian Shan Mountains (up to 7439 m high) separate the Junggar Basin to the north from the Tarim Basin to the south, and the Fergana Basin to the west (Figure 10A). The Junggar Basin has a long history of development (Jolivet et al., 2010; 2013; Jolivet, 2015), from the Paleozoic onwards, and experienced significant Cenozoic deformation as a result of tectonic reactivation of the Tian Shan intracontinental range (Figure 10B). Our recent fieldwork has shown that calcite beef (several cm thick) and cone-in-cone (Figure 10C) occur frequently within mid-Jurassic strata (mainly

the Xishanyao and Totounhe Formations), which crop out in the Junggar Basin on its southern margin (Wusu and Totoun localities) or eastern edge (Kalameili region), along a series of thrust faults and folds, related to the development of the Tian Shan and Altai ranges, respectively. Inside the Tian Shan range itself, cone-in-cone structures occur at Nileke (Figure 10D), at the eastern tip of the intra-mountain Yili Basin within the Middle Jurassic Totounhe Formation (Figures 1 and 10). This area has been subject to large-scale thrusting, during Neogene growth of the northern Tian Shan subrange. Finally, to the west in the intramountain Issik-Kul Basin (Kyrgyzstan), bedding-parallel beef with vertical fibers (Figure 10E, F) occur in Jurassic strata along the southern edge of the basin, which has been subject to Cenozoic compressional deformation in the Terzkey range (Figures 1 and 10). The proximity between our beef occurrences and major thrust faults suggests that beef and the Cenozoic reactivation of the basin are synchronous.

At all of these localities, the beef or cone-in-cone occur within organic-rich fine-grained alluvial plain deposits (Heilbronn, 2014). In the Junggar and Yili basins, they are also close to coal layers, which are several metres thick. At some localities, especially in the Yili and Issik-Kul basins, the calcite beef is close to iron-rich sandy layers of probable diagenetic origin. At Issik-Kul, strong uranium enrichment of the Jurassic series (Kaji Sai mine) containing the beef again indicates post-sedimentary fluid circulation. In the Junggar Basin, the main source rocks for oil are the upper Permian, Upper Triassic and Middle Jurassic detrital series (Jiao et al., 2007). The occurrence of calcite beef within the Middle Jurassic series and the systematic association between beef and organic-rich siltstone or coal layers suggests a link between hydrocarbon source rocks and calcite beef.

In the southern Junggar Basin (locality 8, Figure 1), Jiao et al. (2007) (Fig. 2) described and illustrated thin bedding-parallel veins of calcite within the upper Permian Lucaogou Formation near Urumqi. These authors did not refer to fibrous calcite (beef) or cone-in-cone, but they showed many examples of solid hydrocarbons within cavities, the Lucaogou Formation being a mature petroleum source rock.

4.4. Appalachian Mountains, United States (locality 2, Fig. 1).

In the eastern US (Figure. 11A), the Appalachian Mountains trend northeast-southwest and consist mainly of Paleozoic strata, folds and thrusts (Gilman & Metzger, 1967; Evans, 1995; Tobin et al., 1996). At the southwestern end of the mountains, a major unconformity underlies Cretaceous strata, which form a broad anticline (Figure 11A). This provides good evidence for Late Cretaceous or Cenozoic reactivation of the mountain belt. Indeed, even today, the belt is subject to earthquakes, resulting from compressive stress (Heidbach et al., 2009), which appears to derive from ridge-push of the Atlantic spreading center. Veins of calcite beef and bitumen (e.g. Figure 11B) occur within the Devonian Marcellus Shale (Gale et al., 2014; Aydin & Engelder, 2014). Some of the veins are undulating and have additional lenses within synclines (Gale et al., 2014, their fig. 9E). Even today,

the Marcellus Shale is a source rock for oil and locally reaches overpressure (Aydin & Engelder, 2014), possibly as a result of long-term and recent burial. Furthermore, the layers of Marcellus Shale have acted as detachments for thrust faults in the Appalachians (Aydin & Engelder, 2014). Thus, the Appalachians, like other areas, provide evidence for synchronicity of (1) compressional deformation, (2) generation of organic-rich fluids and (3) formation of calcite beef. A possible problem in the Appalachians is to date these features, which may have started long ago, but still be occurring today.

5. Discussion

Because beef veins seem to be proxies for natural hydraulic fracturing in rocks, especially within source rocks for petroleum, the studies of such fractures are key to the understanding of such geological processes and for the migration and interactions between fluids and rocks. Their formation depends on 2 main phases: (1) the generation of the fracture and (2) the filling of the fracture.

For the initiation of the natural hydraulic fractures, and thus the initiation of beef veins, the theory and the experiments of fracturing demonstrate that horizontal (or bedding-parallel) fractures can be generated, due to a distributed fluid overpressure (Cobbold & Rodrigues, 2007; Mourgues et al. 2011; our review; Zanella et al. 2014a). Nevertheless, previous reviews (Cobbold et al. 2003; Gale et al. 2014) and more recent and local studies (e.g. Rodrigues et al. 2007; Zanella et al. 2014a, 2014b, 2015a, 2015b; Weng et al. 2017; Ukar et al. 2017; Ukar et al. 2018) have shown that the natural hydraulic fracturing often occurred within sedimentary basins which experienced a compressive tectonic history. Moreover, according to several previous studies, the timing of development of beef veins was synchronous with a compressive period in the basin. This is, in particular, well-illustrated for the Neuquén Basin (Rodrigues et al. 2009; Zanella et al. 2015a; Ukar et al. 2017; Ukar et al. 2018), the Wessex Basin (Zanella et al. 2015b), the Bristol Channel (Weng et al. 2017) and the Magallanes Basin (Zanella et al., 2014). In our study, we demonstrate that this observation is also true for the development of beef veins in the Paris Basin (beef veins dated at 155 Ma), in the northern Pyrenees, in the Uinta Basin, in the Tian Shan Mountains and in the Appalachian Mountains. In view of all of these observations and conclusions, we ask some questions: even if in theory the compressive tectonic stress is not necessary to develop bedding-parallel natural hydraulic fractures, is, in nature, this stress crucial for the development of such fractures? Is the fluid overpressure, generated by hydrocarbons, enough to induce hydraulic fracturing of shales?

We thus propose that compressional tectonic stress is one of the key parameters in the development of bedding-parallel veins in shales. This could have major consequences for the understanding of fluid migration in sedimentary rocks, because of the historical complexity of such geological processes. Concerning the generation of hydrocarbons, even if tectonic activity has a big role, the maturation of organic matter, which leads to the development of distributed fluid

overpressures and then to natural hydraulic fracturing, is still the main parameter. Indeed, as already demonstrated, during the maturation of the source rock a part of the solid framework (the organic matter) will be transform into fluid (hydrocarbons), implying a collapse and a load transfer responsible of the increasing of the pore fluid pressure (Zanella et al. 2014a). These fractures are always within or near source rocks for petroleum (Cobbold et al. 2013; Gale et al. 2014). All of our examples respect the previous observations. So, as already suggested by previous authors (Ukar et al. 2017, 2018; Zanella et al. 2014b; Zanella et al. 2015a; Zanella et al. 2015b), the link between the presence of beef veins and organic matter is strong. Moreover, the degree of maturity of the source rock is a key parameter for the development of beef veins and other tectonic structures, such as detachments, as demonstrated in the Magellan Basin (Zanella et al. 2014b). In this basin, Zanella et al. (2014b) also demonstrated that there is a link between beef vein composition and the degree of maturity of the source rock. Thus, we infer that this natural fluid generation process is the main driver for inducing fluid overpressure in mature shales, but needs to be assisted by another force, such as the compressive tectonic stress, to be able to induce natural fracturing of the host rocks.

Concerning the filling of the fracture and thus the cementation and growth of the bedding-parallel fractures, other mechanisms have to be involved to precipitate minerals. Indeed, the opening of the fractures is in mode 1 and is facilitated by a horizontal compressive stress. Nevertheless, the force of crystallization also participates to pushing outward the vein walls. Currently, which of these two processes plays the dominant role is not yet known.

6. Conclusions

Beef veins (BPV) are common in or near source rocks for petroleum. The study of such geological evidences can help to understand the mechanisms involved in their formation: (1) the generation of the fracture and (2) the opening of the fracture, as well as the migration of fluids in sedimentary basins. Studies of beef vein occurrences around the world have led us to conclude that it is especially common within foreland basins. Here we have reviewed examples (or described new ones) from the Wessex Basin (UK), the Neuquén Basin (Argentina), the Cordillera Oriental (Colombia), the Uinta Basin (USA), the Paris Basin (France), the northern Pyrenees (France), the Tian Shan Mountains (central Asia) and the Appalachian Mountains (USA). However, we have discovered similar beef within other localities of foreland basins (some of which are visible in Fig. 1).

In this review, we demonstrate that the development of beef veins occurs worldwide within or near source rocks for petroleum and during a period of hydrocarbon generation. It is now becoming clearer that the maturation of the organic matter can lead to fluid overpressures. Beef veins (and more generally the natural hydraulic fractures) can therefore be used as proxy to determine very quickly if a

source rock was or has been mature. On a global scale, many foreland basins contain source rocks, near active or ancient mountain belts, the latter of which may have been reactivated by recent tectonic stress. Many such basins contain enough organic material to have acted as source rocks for petroleum systems, especially where recent burial has generated sufficiently high temperatures. In some examples (such as the Neuquén Basin of Argentina), next to subduction-zone systems, magmatic intrusions or extrusions have added heat and facilitated the maturation of the source rocks, also during compressional tectonic activity.

The timing of beef vein generation in foreland basins is always coeval with shortening periods, due to compressive tectonic stress. Thus, even if the theory and the experiments of fracturing demonstrate that bedding-parallel fractures can be generated only with a distributed fluid overpressure, the beef veins formation seems to require an external tectonic stress to develop in nature. The filling of the fractures is likely related to force of crystallization, during the compressive period and facilitates the vertical opening of veins.

Acknowledgements

During the years 2005 to 2012, Ian West of Southampton University introduced us to some of the best beef localities in the Wessex Basin, along the coast of southern England (West, 2015). We thank Tony Boassen (Statoil Research Centre, Trondheim), for help with scanning electron microscopy of calcite beef from the Wessex Basin and Neuquén. We also thank Andres Mora of Ecopetrol for showing Peter Cobbold some beef veins at the eastern edge of the Cordillera Oriental, Colombia (Fig. 5). Fieldwork in central Asia was funded by the DARIUS group. Elsewhere, much of the fieldwork and experimental work, as well as PhD studies, were funded by Statoil (Norway). DC acknowledges support from Science Foundation Ireland under Grant Number 15/IACA/3365.

References

- Aydin, M.G., Engelder, T., 2014. Revisiting the Hubbert-Rubey pore pressure model for overthrust faulting: Inferences from bedding-parallel detachment surfaces within Middle Devonian gas shale, the Appalachian Basin, USA. *J. Struct. Geol.* 69, 519-537.
- Barker, S.L.L., Cox, S.F., Eggins, S.M., Gagan, M.K., 2006. Microchemical evidences for episodic growth of antitaxial veins during fracture-controlled fluid flow. *Earth and Planetary Science Letters* 250, 331-344.
- Biteau, J.-J., Canérot, J., 2007. La Chaîne des Pyrénées et ses avant-pays d'Aquitaine et de l'Ebre: systèmes pétroliers et gisements d'hydrocarbures. *Géologues* 155, 29-41.

475 Bons, P., 2001. Development of crystal morphology during unitaxial growth in a progressively
476 widening vein: I. The numerical model. *Journal of Structural Geology*, 23(6-7): 865-872.

477 Bons, P.D., Elburg, M.A. and Gomez-Rivas, E., 2012. A review of the formation of tectonic veins and
478 their microstructures. *Journal of Structural Geology*, 43: 33-62.

479 Bons, P.D., Jessell, M.W., 1997. Experimental simulation of the formation of fibrous veins by localised
480 dissolution-precipitation creep. *Mineralogical Magazine* 61, 53-63.

481 Bradbury, K.K., Davis, C.R., Shervais, J.W., Janecke, S.U., Evans, J.P., 2015. Composition, alteration,
482 and texture of fault-related rocks from Safod core and surface outcrop analogs: evidence for
483 deformation processes and fluid-rock interactions. *Pure and Applied Geophysics* 172, 1053–1078.

484 Buckland, W., De la Beche, H.T., 1835. On the geology of the neighbourhood of Weymouth and the
485 adjacent parts of the coast of Dorset. *Trans. Geol. Soc. Lond.*, s2 4, 1-46.

486 Chatellier, J.-Y., 2013. How to maximize the use of TMAX, compositional ratios and stable carbon
487 isotopes to accurately infer the maturity of a shale. *Tight Oil Canada*, 2013.

488 Choukroune, P., 1969. Sur la presence, le style et l'âge des tectoniques superposées dans le Crétacé
489 nord-pyrénéen de la région de Lourdes (Hautes-Pyrénées). *Bulletin du Bureau de Recherches*
490 *Géologiques et Minières* Section I, No. 2, 11-20.

491 Cobbold, P.R., 2013. Geological evidence for fluid overpressure in mature source rocks within
492 foreland basins of the Americas. *American Association of Petroleum Geologists, International*
493 *Conference and Exhibition, Cartagena, Colombia, 8-11 September. Search and Discovery Article*
494 *#30291*.

495 Cobbold, P.R., Castro, L., 1999. Fluid pressure and effective stress in sandbox models.
496 *Tectonophysics* 301, 1-19.

497 Cobbold, P.R., Durand, S., Mourgues, R., 2001. Sandbox modelling of thrust wedges with fluid-
498 assisted detachments. *Tectonophysics* 334, 245-258.

499 Cobbold, P.R., Rodrigues, N., 2007. Seepage forces, important factors in the formation of horizontal
500 hydraulic fractures and bedding-parallel fibrous veins ("beef" and "cone-in-cone"). *Geofluids* 7,
501 313-332.

502 Cobbold, P.R., Zanella, A., Ruffet, G., Rodrigues, N., Loseth, H., 2013. Bedding-parallel fibrous veins
503 (beef and cone-in-cone): Worldwide occurrence and possible significance in terms of fluid
504 overpressure, hydrocarbon generation and mineralization. *Mar. Petrol. Geol.* 43, 1-20.

505 Cobbold, P., Zanella, A., Fourdan, B., Néraudeau, D., Gouttefangeas, F., 2015. Natural hydraulic
506 fractures in the Eastern Paris Basin. *Bull. Inf. Géol. Bassin Paris* 52 (2), 23-30.

507 David, L., 1952. Présence de la structure "beef" et "cone-in-cone" dans le Crétacé de l'Est-
508 Constantinois (Algérie). *Comptes Rendus Sommaires de la Société Géologique de France* 3, 51-
509 52.

- Denaeyer, M.-E., 1943. Les cone-in-cone de la France métropolitaine et d'outre-mer. *Bull. Soc. Fr. Minéral.* 66 (1-6), 173-221.
- Denaeyer, M.-E., 1947. Les gisements de cone-in-cone de France et de Grande-Bretagne. Première Partie. *Bulletin de la Société Belge de Géologie, Paléontol. Hydrol.* 56, 21-46.
- Dubiel, R.F., 2003. Geology, depositional models, and oil and gas assessment of the Green River total petroleum system, Uinta-Piceance Province, Eastern Utah and Western Colorado. In: *Petroleum systems and geologic assessment of oil and gas in the Uinta-Piceance Province, Western Colorado*. U.S. Geological Survey, Digital Data Series DDS-69-B, 1-41.
- Evans, M.A., 1995. Fluid inclusions in veins from the Middle Devonian shales: A record of deformation conditions and fluid evolution in the Appalachian Plateau. *Geol. Soc. Am. Bull.* 107 (3), 327-339.
- Fischer, M.P., Higuera-Diaz, I.C., Evans, M.A., Perry, E.C., Lefticariu, L., 2009. Fracture-controlled paleohydrology in a map-scale detachment fold: Insights from the analysis of fluid inclusions in calcite and quartz veins. *J. Struct. Geol.* 31, 1490-1510.
- Fitz-Diaz, E., Hudleston, P., Siebenaller, L., Kirschner, D., Camprubi, A., Tolson, G., Puig, T.P., 2011. Insights into fluid flow and water-rock interaction during deformation of carbonate sequences in the Mexican fold-thrust belt. *J. Struct. Geol.* 33, 1237-1253.
- Gale, J.F.W., Laubach, S.E., Olson, J.E., Eichhubl, P., Fall, A., 2014. Natural fractures in shale: a review and new observations. *Am. Assoc. Petrol. Geol. Bull.* 98 (11), 2165-2216.
- Gely, J-P., Hanot, F., 2014. Le bassin parisien : un nouveau regard sur la géologie. In: *Bulletin d'information des géologues du bassin de Paris*, 9, 229 pp.
- Gilman, R.A., Metzger, W.J., 1967. Cone-in-cone concretions from western New York. *J. Sed. Petrol.* 37, 87-95.
- Gressier, J.B., Mourgues, R., Bodet, L., Matthieu, J.Y., Galland, O., Cobbold, P.R., 2010. Control of pore fluid pressure on depth of emplacement of magmatic sills: An experimental approach. *Tectonophysics* 489, 1-13.
- Guillocheau, F., Robin, C., Allemand, P., Bourquin, S., Brault, N., Dromart, G., Friedenberg, R., Garcia, J.-P., Gaullier, J.-M., Gaumet, F., Grosdoy, B., Hanot, F., Le Strait, P., Mettraux, M., Nalpas, T., Prijac, C., Rigollet, C., Serrano, O., Grandjean, G., 2000. Meso-Cenozoic geodynamic evolution of the Paris Basin: 3D stratigraphic constraints. *Geodinamica Acta* 13, 189-246.
- Heidbach, O., Tingay, M., Bartyh, A., Reinecker, J., Kurfeth, B., Müller, B., 2009. World stress map, 2nd Edition. GFZ German Research Centre for Geosciences.
- Heilbronn, G., 2014. Evolution paléogéographique et paléotopographique du Tian Shan Chinois au Mésozoïque. PhD thesis, Univ. Rennes, 267 pp.
- Hilgers, C., Koehn, D., Bons, P.D. and Urai, J.L., 2001. Development of crystal morphology during unitaxial growth in a progressively widening vein: II. Numerical simulations of the evolution of antitaxial fibrous veins. *Journal of Structural Geology*, 23(6-7): 873-885.

547 Hillier, R.D., Cosgrove, J.W., 2002. Core and seismic observations of overpressure-related
548 deformation within Eocene sediments of the Outer Moray Firth, UKCS. *Petroleum Geoscience* 8,
549 141-149.

550 Jiao, Y.Q., Wu, L.Q., He, M.C., Mason, R., Wang, M.F., Xu, Z.C., 2007. Occurrence, thermal
551 evolution and primary migration processes derived from studies of organic matter in the
552 Lucaogou source rock at the southern margin of the Junggar Basin, NW China. *Science in China,*
553 *Series D - Earth Sciences* 50, 114-123.

554 Jolivet, M., Dominguez, S., Charreau, J., Chen, Y., Li, Y., Wang, Q., 2010. Mesozoic and Cenozoic
555 tectonic history of the central Chinese Tian Shan: Reactivated tectonic structures and active
556 deformation. *Tectonics* 29, 1-30.

557 Jolivet, M., Heilbronn, G., Robin, C., Barrier, L., Bourquin, S., Guo, Zh., Jia, Y., Guerit, L., Yang, W.,
558 Fu, B., 2013. Reconstructing the Late Palaeozoic – Mesozoic topographic evolution of the
559 Chinese Tian Shan: available data and remaining uncertainties. *Advances in Geosciences* 37, 7-
560 18.

561 Jolivet, M., 2015. Mesozoic tectonic and topographic evolution of Central Asia and Tibet: a
562 preliminary synthesis. *In: Brunet, M.-F., McCann, T. & Sobel, E.R. (eds) Geological evolution of*
563 *Central Asian Basins and the Western Tien Shan Range. Geol. Soc. London Sp. Publ., 427,*
564 <http://doi.org/10.1144/SP427.2>

565 Kemp, J., 2014. The Big One: Russia's Bazhenov shale.
566 <http://www.reuters.com/article/2014/07/18/us-russia-shale-kemp-idU>.

567 Kershaw, S., Guo, L. 2016. Beef and cone-in-cone calcite fibrous cements associated with the end-
568 Permian and end-Triassic mass extinctions: Reassessment of processes of formation. *J.*
569 *Palaeogeog.* 5 (1), 28-42.

570 Keulen, N. T., den Brok, S.W.J., Spiers, C.J., 2001. Force of crystallisation of gypsum during
571 hydration of synthetic anhydrite rock. 13th DRT conference, Deformation Mechanisms,
572 Rheology, and Tectonics, Noordwijkerhout, The Netherlands.

573 Kowal-Linka, M., 2010. Origin of cone-in-cone calcite veins during calcitization of dolomites and
574 their subsequent diagenesis: A case study from the Gogolin Formation (Middle Triassic), SW
575 Poland. *Sed. Geol.* 224, 54-64.

576 Kozłowski, E.E., Cruz, C.E., Sylwan, C., 1998. Geología estructural de la zona de Chos Malal, Cuenca
577 Neuquina, Argentina. *In: XIII Congreso Geológico Argentino y III Congreso de Exploración de*
578 *Hidrocarburos, Buenos Aires, Actas 1, 15-26.*

579 Labaume, P., Sheppard, S.M.F., Moretti, I., 2001. Fluid flow in cataclastic thrust fault zones in
580 sandstones, Sub-Andean Zone, southern Bolivia. *Tectonophysics* 340, 141-172.

581 Lamb, S., 2004. Devil in the mountain. A search for the origin of the Andes. Princeton University
582 Press 335 pp.

583 Lang, W.D., Spath, L.F., Richardson, W.A., 1923. Shales-With-'Beef', a sequence in the Lower Lias of
584 the Dorset Coast. Q. J. Geol. Soc. Lond. 79, 47-99.

585 Le Breton, E., Cobbold, P.R., Zanella, A., 2013. Cenozoic reactivation of the Great Glen Fault,
586 Scotland: additional evidence and possible causes. J. Geol. Soc. Lond. 170, 403-415, doi:
587 10.1144/jgs2012-067.

588 Lee, G.W., 1920. The Mesozoic rocks of Applecross, Raasay and North-East Skye. H.M. Geological
589 Survey, Scotland, 93 pp.

590 Lemrabott, A., Cobbold, P.R., 2010. Physical modeling of fluid overpressure and compaction during
591 hydrocarbon generation in source rock of low permeability. Search and Discovery, Article
592 #40518.

593 Leythaeuser, D., Littke, R., Radke, M., Schaefer, R.G., 1988. Geochemical effects of petroleum
594 migration and expulsion from Toarcian source rocks in the Hils Syncline area, NW-Germany. In:
595 Mattavelli, L. & Novelli, L. (Eds.), Advances in Organic Geochemistry 1987. Organic
596 Geochemistry 13, 489-502.

597 Lilloc-Olsen, T., Bang, N.A., 2012. DNO and Tawke in Kurdistan: How an Iraq oil giant has emerged.
598 Oil & Gas Journal 1-19. [http://www.ogj.com/articles/print/vol-110/issue-2/exploration-](http://www.ogj.com/articles/print/vol-110/issue-2/exploration-developmet/dno-and-tawke-in...)
599 [developmet/dno-and-tawke-in...](http://www.ogj.com/articles/print/vol-110/issue-2/exploration-developmet/dno-and-tawke-in...)

600 Macsotay, O., Erlich, R.N., Peraza, T., 2003. Sedimentary structures of the La Luna, Navay and
601 Querecual Formations, Upper Cretaceous of Venezuela. Palaios 18, 334-348.

602 Mahboubi, A., Moussavi-Harami, R., Collins, R.B., Muhling, J.R., 2010. Petrography and
603 geochemical signatures in cracks filling calcite sequences in septarian concretions, Sanganeh
604 Formation, Kopet-Dagh Basin, NE Iran. J. Appl. Sci. 10, 526-534.

605 Maillot, H., Bonte, A., 1983. Cone-in-cone texture from Deep Sea Drilling Project Leg 71, Site 511,
606 Falkland Plateau, South Atlantic Ocean. Initial Reports of the Deep Sea Drilling Project 71(1),
607 345-349.

608 Mangenot, X., Gasparrini, M., Rouchon, V., Bonifacie, M., 2018. Basin-scale thermal and fluid flow
609 histories revealed by carbonate clumped isotopes ($\Delta 47$)—Middle Jurassic carbonates of the Paris
610 Basin depocentre. *Sedimentology*, 65(1), 123-150.

611 Marques, F.O., Nogueira, F.C.C., Bezerra, F.H.R., Castro, D.L. de, 2014. The Araripe Basin in NE
612 Brazil: An intracontinental graben inverted to a high-standing horst. Tectonophysics 630, 251–
613 264.

614 Marshall, J.D., 1982. Isotopic composition of displacive fibrous calcite veins; reversals in pore-water
615 composition trends during burial diagenesis. J. Sediment. Petrol. 52, 615-630.

616 Martinius, A.W., Hegner, J., Kaas, I., Bejarano, C., Mathieu, X., Mjoes, R., 2012. Sedimentology and
617 depositional model for the Early Miocene Oficina Formation in the Petrocedeno Field (Orinoco
618 heavy-oil belt, Venezuela). Mar. Petrol. Geol. 35, 354-380.

619 Means, W. and Li, T., 2001. A laboratory simulation of fibrous veins: some first observations. Journal

of Structural Geology, 23(6-7): 857-863.

Mora, A., Blanco, V., Naranjo, J., Sanchez, N., Ketcham, R.A., Rubiano, J., Stockli, D.F., Quintero, I., Nemcok, M., Horton, B.K., Davila, H., 2013. On the lag time between internal strain and basement involved thrust induced exhumation: The case of the Colombian Eastern Cordillera. *J. Struct. Geol.* 52, 96-118.

Mora, A., Casallas, W., Ketcham, R.A., Gomez, D., Parra, M., Namson, J., Stockli, D., Almendral, A., Robles, W., Ghorbal, B., 2015. Kinematic restoration of contractional basement structures using thermokinematic models. *Am. Assoc. Petrol. Geol. Bull.* 99 (8), 1575–1598.

Mourgues, R., Cobbold, P.R., 2003. Some tectonic consequences of fluid overpressures and seepage forces as demonstrated by sandbox modeling. *Tectonophysics* 376, 75-97.

Nollet, S., Urai, J.L., Bons, P.D., Hilgers, C., 2005. Numerical simulations of polycrystal growth in veins. *Journal of Structural Geology* 27, 217-230.

Nollet, S., Koerner, T., Kramm, U., Hilgers, C., 2009. Precipitation of fracture fillings and cements in the Buntsandstein (NW Germany). *Geofluids* 9, 373-385.

Nomura, S.F., Sawakuchi, A.O., Bello, R.M.S., Méndez-Duque, J., Fuzikawa, K., Giannini, P.C.F., Dantas, M.S.S., 2014. Paleotemperatures and paleofluids recorded in fluid inclusions from calcite veins from the northern flank of the Ponta Grossa dyke swarm: Implications for hydrocarbon generation and migration in the Paraná Basin. *Mar. Petrol. Geol.* 52, 107-124.

Parnell, J., Blamey, N.J.F., Costanzo, A., Feely, M., Boyce, A.J., 2014. Preservation of Mesoproterozoic age deep burial fluid signatures, NW Scotland. *Mar. Petrol. Geol.* 55, 275-281.

Pearson, P.N., Nicholas, C.J., Singano, J.M., Bown, P.R., Coxall, H.K., van Dongen, B.E., Huber, B.T., Karega, A., Lees, J.A., MacLeod, K., McMillan, I.K., Pancost, R.D., Pearson, M., Msaky, E., 2006. Further Paleogene and Cretaceous sediment cores from the Kilwa area of coastal Tanzania: Tanzania Drilling Project Sites 6–10. *J. Afr. Earth Sci.* 45, 279-317.

Richardson, W.A., 1923. Petrology of the Shales-with-“Beef.” *Q. J. Geol. Soc. Lond.* 79, 88-99.

Rodrigues, N., Cobbold, P.R., Loseth, H., Ruffet, G., 2009. Widespread bedding-parallel veins of fibrous calcite (“beef”) in a mature source rock (Vaca Muerta Fm, Neuquén Basin, Argentina): evidence for overpressure and horizontal compression. *J. Geol. Soc. Lond.* 166, 695-709, doi 10.1144/0016-76492008-111.

Rybak-Ostrowska, B., Konon, A., Nejbert, K., Kozłowski, A., 2014. Bedding-parallel calcite veins in the Holy Cross Mountains Fold Belt, central Poland. *Geological Quarterly* 58 (1), 99–116.

Silva, A.L. Da, 2003. Estratigrafia física e deformação do sistema lacustre carbonático (Aptiano-Albiano) da Bacia do Araripe em afloramentos selecionados. Dissertação de Mestrado, Univ. Fed. Pernambuco, Pós-graduação em Geociências, 118 pp.

Smith, A.P., Fischer, M.P., Evans, M.A., 2014. On the homogeneity of fluids forming bedding-parallel veins. *Geofluids* 14, 45-57.

656 Suchy, V., Dobes, P., Filip, J., Stejskal, M., Zeman, A., 2002. Conditions for veining in the Barrandian
657 Basin (Lower Palaeozoic), Czech Republic: evidence from fluid inclusion and apatite fission
658 track analysis. *Tectonophysics* 348, 25-50.

659 Tarney, J., Schreiber, B.C., 1976. Cone-in-cone and beef-in-shale textures from DSDP site 330,
660 Falkland Plateau, South Atlantic. In: Barker, P.F., Dalziel, I.W.D. et al. (Eds), *Initial Reports of*
661 *the Deep Sea Drilling Project*. U.S. Government Printing Office, Washington, D.C. 36, 865-870.

662 Tobin, K.J., Walker, K.R., Steinhauß, D.M., Mora, C.I., 1996. Fibrous calcite from the Ordovician of
663 Tennessee: preservation of marine oxygen isotopic composition and its implications.
664 *Sedimentology* 43, 235-251.

665 Trabuco-Alexandre, J., Dirkx, J., Veld, H., Klaver, G., De Boer, P.L., 2012. Toarcian black shales in
666 the Dutch Central Graben: Record of energetic, variable depositional conditions during an
667 oceanic anoxic event. *J. Sed. Res.* 82, 104-120.

668 Ukar, E., Lopez, R.G., Gale, J.F., Laubach, S. E., Mancada, R., 2017. New type of kinematic indicator
669 in bed-parallel veins, Late Jurassic–Early Cretaceous Vaca Muerta Formation, Argentina: EW
670 shortening during Late Cretaceous vein opening. *Journal of Structural Geology*, 104, 31-47.

671 Ukar, E., López, R., Hryb, D., Gale, J.F., Mancada, R., Fall, A., Brisson, I., Hernandez-Bilbao, E.,
672 Weger, R., Marchal, D., Zanella, A., Lanusse, I., 2018. Natural fractures in the Vaca Muerta
673 Formation: from core and outcrop analog observations to subsurface models. *AAPG Memoir*,
674 120, (submitted).

675 Underhill, J.R., Stoneley, R., 1998. Introduction to the development, evolution and petroleum geology
676 of the Wessex Basin. In: *Geol. Soc., London, Sp. Publ.* 133, 1-18.

677 Vera, E.A.R., Folguera, A., Valcarce, G.Z., Bottesi, G., Ramos, V. A., 2014. Structure and
678 development of the Andean system between 36 and 39 S. *Journal of Geodynamics*, 73, 34-52.

679 Voisin, L., 1999. Le “beef” de Chaumiau (08). *Bull. d’Inform. Géol. Bassin de Paris* 36 (2), 13-16.

680 Volk, H., Horsfield, B., Mann, U., Suchy, V., 2002. Variability of petroleum inclusions in vein, fossil
681 and vug cements - a geochemical study in the Barrandian Basin (Lower Palaeozoic, Czech
682 Republic). *Organic Geochemistry* 33, 1319-1341.

683 Watts, N.L., 1978. Displacive calcite: Evidence from recent and ancient calcretes. *Geology* 6, 699-
684 703.

685 Weger, R.J., Murray, S.T., McNeill, D.J., Swart, P.K., Eberli, G.P., Rodriguez Blanco, L., Tenaglia,
686 M., Rueda, L., 2018. Paleo thermometry and distribution of calcite beef in the Vaca Muerta
687 Formation, Neuquén Basin, Argentina. *AAPG Bulletin*, (submitted).

688 West, I.M., 2015. Geology of the Wessex coast of southern England – the World Heritage Jurassic
689 Coast – and more. Internet page: <http://www.southampton.ac.uk/~imw/index.htm>.

690 Woodland, B.G., 1964. The nature and origin of cone-in-cone structure. *Fieldiana: Geology* 13 (4),
691 187-305.

692 Zanella, A., Cobbold, P.R., Le Carlier de Veslud, C., 2014a. Physical modelling of chemical

compaction, overpressure development, hydraulic fracturing and thrust detachments in organic-rich source rock. *Mar. Petrol. Geol.* 55, 262-274.
<http://dx.doi.org/10.1016/j.marpetgeo.2013.12.017>.

Zanella, A., Cobbold, P.R., Rojas, L., 2014b. Beef veins and thrust detachments in Early Cretaceous source rocks, foothills of Magallanes-Austral Basin, southern Chile and Argentina: structural evidence for fluid overpressure during hydrocarbon maturation. *Mar. Petrol. Geol.* 55, 250-261.
<http://dx.doi.org/10.1016/j.marpetgeo.2013.10.006>.

Zanella, A., Cobbold, P.R., Ruffet, G., Leanza, H.A., 2015a. Geological evidence for fluid overpressure, hydraulic fracturing and strong heating during maturation and migration of hydrocarbons in Mesozoic rocks of the northern Neuquén Basin, Mendoza Province, Argentina. *J. S. Am. Earth Sci.* 62, 229-242.

Zanella, A., Cobbold, P.R., Boassen, T., 2015b. Natural hydraulic fractures in the Wessex Basin, SW England: widespread distribution, composition and history. *Mar. Petrol. Geol.* 68, 438-448.

Zhang, B., Yin, C.Y., Gu, Z.D., Zhang, J.J., Yan, S.Y., Wang, Y., 2015. New indicators from bedding-parallel beef veins for the fault valve mechanism. *Science China: Earth Sci.*, doi: 10.1007/s11430-015-5086-6.

Figure captions

Table. 1. Global distribution of compressional basins, where calcite beef or cone-in-cone occur. For localities (numbers at left), see Figure 1. Not all references are to previous descriptions of calcite beef.

Figure 1. Map showing distribution of calcite beef (bedding-parallel veins), either within Mesozoic or Cenozoic sedimentary rocks (light triangles) or within Paleozoic sedimentary rocks (dark triangles). The numbers next to the triangles refer to the localities in Table 1.

Figure 2. A. Physical model of horizontal hydraulic fracturing with no deformation (after Zanella et al., 2014a); B. Physical model of horizontal hydraulic fracturing with shortening (after Zanella et al., 2014a); C. Cross-section of a 3-D physical model showing the different styles of deformation due to the propagation of a detachment linked to overpressure development (after Zanella et al., 2014a).

Figure 3. A. Geological map and cross-section of the Wessex Basin, southwestern England (after Zanella et al. 2015B, modified from Underhill et al. 1998). B. folded and faulted beef veins, the thickness of which is again variable, especially across the main reverse fault (center). C. Locally folded beef, the thickness of which is variable (thicker in syncline, thinner in anticline). D. Steeply dipping but curved calcite fibers, which comprise a layer of beef. (scales: pen: 14cm; coin diameter:

2.4 cm).

Figure 4. A. Calcite beef in the Neuquén Basin, Argentina (white locality 4, Figure 1) (after Rodrigues et al., 2009). The Landsat image (left) shows sub-Andean folds, trending north-northwest. Field observations (bottom right) show that beef is common in the Vaca Muerta Formation at various scales. B. Geological section through the fold and thrust belt of the Neuquén Basin (located with the red line in Figure 4A.) (after Vera et al., 2014).

Figure 5. A. Geological map and cross-section, showing the main structures of the eastern Cordillera (after Mora et al., 2013, 2015). B. Calcite beef (bedding-parallel) veins (white to yellow), which vary in thickness across folds (Cobbold, 2013). C. On the western side of the Cordillera, near Villeta, folds and thrusts verge westward and Cretaceous shales contain layers of calcite beef. D. At the eastern edge of the Cordillera, in the Macanal Formation (Lower Cretaceous) near Villavicencio, some beef contains thin layers of yellow pyrite, which are folded, while calcite has grown epitaxially above and below them, filling in synclines more than anticlines. (scales: coin diameter: 2.2 cm).

Figure 6. A. Geological map of the Paris Basin (after Gely and Hanot, 2014; Mangenot et al. 2018). The basin has an elliptical shape with a long axis trending approximately northeast-southwest. The line of section is indicated in red. B. Gélaucourt, near Nancy, where ditches have exposed Liassic shales of the Schistes Carton Formation (lower Toarcian, Denaeyer, 1943; ca 185 Ma.), which contain abundant veins of fibrous calcite beef (hammer length: 33 cm). C. Calcite beef from the Toarcian at Gélaucourt (coin diameter: 2.3 cm).

Figure 7. A. U-Pb Tera-Wasserburg calcite lower intercept age of 155 ± 19 Ma (Oxfordian) for the formation of the calcite beef (Chew et al., work in progress). B&C. Scanning electron microscopy (SEM) analyses of calcite beef (bedding-parallel veins) from the Liassic “Schistes Carton” near Gélaucourt in the Paris Basin (see Fig. 6). Scanning electron microscopy (SEM) has yielded significant quantities of calcium (B) and carbon (C), which are typical of calcite. However, the amount of carbon is locally greater (pink, top right), due to inclusions of hydrocarbons within the calcite crystals (Cobbold et al., 2015).

Figure 8. Calcite beef near Lourdes, at the northern edge of the Pyrenees (white locality 22, Figure 1). A. Google Earth oblique view of the Pic du Béout hills (1530 m high, near the southeast end of Lourdes city), shows repetitions of white, thick, resistant layers of Aptian-Albian limestones, which dip at about 30° to 40° to the south forming scarps. The white line represents the main thrust fault. B. Road outcrop showing fibrous calcite beef (bedding-parallel veins) in upper Aptian shales (coin diameter: 2.3 cm). C. Scanning electron microscopy and the repartition of the carbon, the calcium and

the oxygen in the beef vein.

Figure 9. Calcite beef and bitumen in the Uinta Basin, Utah (white locality 13, Fig. 1). A. A simplified geological section (top, north-northeast to southwest, after Dubiel, 2003) shows the asymmetric structure of the basin. B. Outcrop, discovered in 2009, showing gently dipping beds of the Green River Formation (grey) and numerous veins, either of pure fibrous calcite (orange, left), or of fibrous calcite (coin diameter: 2.4 cm). C. Beef veins with gilsonite (whitish and grey; beneath hammer) (hammer length: 33 cm).

Figure 10. Calcite beef around the Tian Shan Mountains, central Asia (localities 30 to 32, Fig. 1). A: Topographic and tectonic map of the Tian Shan and Junggar region (modified from Jolivet et al., 2013). B: Geological cross-section (approximately north-south) of the Junggar Basin (red line, A). C: Cone-in-cone structures in the Xishanyao Formation (Middle Jurassic) at Wusu (top) (lens cover diameter: 5.2 cm). D: Cone-in-cone structures in the Totounhe Formation (Middle Jurassic) at Nileke (top). E and F: bedding-parallel calcite beef in Jurassic strata at Kaji Sai (Issik Kul Basin, top) (pen: 14 cm).

Figure 11. Calcite beef and bitumen in the Appalachian Mountains, US (locality 2, Figure 1). A. Simplified geological map showing the Paleozoic thrust belt (red), trending northeast-southwest. At its southwestern end, a major unconformity marks the base of Cretaceous strata (green), which nevertheless form a broad anticline, plunging southwest. Red dot refers to Fig. 11B. B. Veins of calcite beef (bedding-parallel veins) and bitumen occurring within the Marcellus Shale (Devonian) along Route 250, Highland County, Virginia (38°19'34.31°N; 79°26'32.29°W, south-southeast of Pittsburgh, Pennsylvania) (coin diameter: 2.4 cm).

- Areas of outcropping Vaca Muerta Fm
- Studied localities within Vaca Muerta Fm
- River
- + Anticline
- Thrust fault
- Strike-slip fault
- AFTB Agrio fold-and-thrust belt
- CMFTB Chos-Malal fold-and-thrust belt

Ca

C

<i>No. Location</i>	<i>References</i>
A. Mesozoic or Cenozoic host rocks	
1 Magallanes Basin (Chile, Tierra del Fuego), Rio Jackson Fm (Early Cretaceous)	Zanella et al., 2014a
2 Magallanes-Austral Basin (Chubut, Argentina), Rio Mayer Fm (Early Cretaceous)	Zanella et al., 2014a
3 Falkland Plateau, Maurice Ewing Bank (Late Jurassic to Early Cretaceous)	Tarney & Schreiber, 1976; Maillot & Bonte, 1983
4 Neuquén Basin, Argentina, Los Molles, Vaca Muerta, Agrio Fms (Jurassic to Cretaceous)	Fig. 4; Rodrigues et al., 2009; Cobbold et al., 2013; Gale et al., 2014; Ukar et al. 2017; Zanella et al., 2015
5 Sub-Andean Zone, southern Bolivia (Tertiary strata)	Labaume et al., 2001; Lamb, 2004
6 Araripe Basin, NE Brazil (Early Cretaceous)	Silva, 2003; Marques et al., 2014
7 Eastern Cordillera, Colombia, (Early Cretaceous source rock)	Fig. 5; Cobbold et al., 2013; Mora et al., 2013; Mora et al., 2015
8 Northern Venezuela, La Luna Fm (Late Cretaceous)	Macsoy et al., 2003
9 Northern Venezuela, Oficina Fm (Early Miocene)	Martinius et al., 2012
10 Central Mexico Fold-and-Thrust Belt, (Cretaceous)	Fitz-Diaz et al., 2011
11 Sierra Madre Oriental, NE Mexico (Jurassic-Cretaceous)	Fischer et al., 2009; Smith et al., 2014
12 SW California, USA, Franciscan Complex (Late Jurassic to Cretaceous)	Bradbury et al., 2015
13 Uinta Basin, Utah, Green River Fm (Eocene)	Fig. 9; Woodland, 1964; Dubiel, 2003
14 Texas, Haynesville Shale (Jurassic)	Gale et al., 2014
15 Outer Hebrides (Eigg, Raasay, Skye), Scotland, UK (Jurassic)	Lee, 1920; Marshall, 1982; Parnell et al., 2014
16 Eathie, Great Glen, NE Scotland (Jurassic)	Le Breton et al., 2013
17 Alba Field, Outer Moray Firth, United Kingdom (Eocene)	Hillier & Cosgrove, 2002
18 Lavernock Point, South Wales (Triassic)	Kershaw & Guo, 2016
19 Wessex Basin, UK (Liassic to Mid-Cretaceous)	Fig. 3; Buckland & De la Beche, 1835; Richardson, 1923; Marshall, 1982; Underhill & Stoneley, 1998; Cobbold & Rodrigues, 2007; Zanella et al., 2015b; Kershaw & Guo, 2016
20 Dutch Central Graben (Toarcian)	Trabucho-Alexandre et al., 2012
21 Eastern and Northern Paris Basin, France (Triassic, Liassic)	Figs. 6 & 7; Denayer, 1943, 1947; Voisin, 1999; Cobbold et al., 2015
22 Lourdes, North-Central Pyrenees, France (Aptian-Albian)	Fig. 8; Choukroune, 1969; Biteau & Canérot, 2007
23 Hils Syncline, NW Germany (Toarcian)	Leythaeuser et al., 1988
24 West Siberia Basin, Russia, Bazhenov Shale (Tithonian-Berriasian)	Kemp, 2014; Fjellanger et al., 2015
25 Algeria-Tunisia (Cretaceous)	David, 1952
26 Kalahari Desert, South Africa and Botswana (Quaternary)	Watts, 1978
27 Kilwa, coastal Tanzania (Cretaceous, Paleogene)	Pearson et al., 2006
28 Tawke Field, Kurdistan, NW Iraq, Sargelu Fm (Jurassic)	Lilloe-Olsen & Bang, 2012
29 Kopet-Dagh Basin, NE Iran, Sanganeh Fm (Late Cretaceous)	Mahboubi et al., 2010
30 Junggar Basin, China, Xishanyao Fm. (Middle Jurassic)	Fig. 10; Jolivet et al., 2010; Heilbronn, 2014
31 Yili Basin, China, Totounhe Fm. (Middle Jurassic)	Fig. 10
32 Issyk-Kul Basin, Kyrgyzstan (Jurassic)	Fig. 10
33 Sichuan Basin, China, Jialingjiang Fm (Triassic)	Zhang et al., 2015

B. Palaeozoic or Precambrian host rocks

1	Parana Basin, SE Brazil, Teresina Fm (Permian)	Nomura et al., 2014
2	Appalachian Mountains, USA, Marcellus Fm (Devonian)	Fig. 11; Gilman & Metzger, 1967; Evans, 1995; Tobin et al., 1996; Gale et al., 2014; Aydin & Engelder, 2014
3	Appalachian Mountains, Quebec, Canada, Utica Shale Fm (Ordovician)	Séjourné et al., 2005; Chatellier, 2013
4	Barrandian Basin, Czech Republic (Lower Palaeozoic)	Suchy et al., 2002; Volk et al., 2002
5	Holy Cross Mountains, Poland (Devonian, Triassic)	Kowal-Linka, 2010; Rybak-Ostrowska et al., 2014
6	Junggar Basin, China, Lucaogou Fm (Upper Permian)	Jiao et al., 2007
7	Kalahari Desert, South Africa and Botswana, (Silurian-Devonian)	Watts, 1978
8	Australia, New South Wales, Murrumbidgee Fm (Devonian)	Barker et al., 2006