



HAL
open science

Electrokinetic and magnetic fields generated by flow through a fractured zone: A Sensitivity study for La Fournaise Volcano

Pierre M Adler, Jean-Louis Le Mouel, Jacques Zlotnicki

► **To cite this version:**

Pierre M Adler, Jean-Louis Le Mouel, Jacques Zlotnicki. Electrokinetic and magnetic fields generated by flow through a fractured zone: A Sensitivity study for La Fournaise Volcano. *Geophysical Research Letters*, 1999, 26 (6), pp.795-798. 10.1029/1999GL900095 . insu-03039723

HAL Id: insu-03039723

<https://hal-insu.archives-ouvertes.fr/insu-03039723>

Submitted on 4 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Electrokinetic and magnetic fields generated by flow through a fractured zone: a sensitivity study for La Fournaise volcano

Pierre M. Adler, Jean-Louis Le Mouel

Institut de Physique du Globe de Paris, Paris, France

Jacques Zlotnicki

Laboratoire de Géophysique d'Orléans, CNRS, Orléans, France

Abstract. A number of electric and magnetic signals have been observed on La Fournaise volcano, attributed to electrokinetic effects. A simple model is proposed here to check this hypothesis. The volcano is idealized as a 2d heterogeneous structure composed of a fractured zone located between two porous zones through which meteoritic waters flow downwards; flow occurs predominantly in the fractured zone and induces electromagnetic fields. Correct orders of magnitude are obtained for the measured surface fields. The importance of the heterogeneous character of the underground medium is demonstrated; local measurements of various quantities are recommended.

1. Introduction

It has been proposed that anomalous electric and magnetic fields observed before earthquakes and volcanic activity could be generated by the electrokinetic effect induced by water flow resulting from internal stresses, thermal buoyancy effects and meteoritic waters. The order of magnitude and the sign of the magnetic field were correctly obtained for the Matsushiro earthquake swarm [Mizutani and Ishido, 1976]. Self-potentials and magnetic measurements at La Fournaise volcano have been analyzed and interpreted in the same way [Zlotnicki and Le Mouel, 1990; Michel and Zlotnicki, 1998].

These two series of large-scale observations were substantiated by local measurements made on plugs in order to obtain dzeta-potentials and streaming potential coefficients of various minerals and rocks as functions of physico-chemical parameters such as salt concentration, pH and temperature [Ishido and Mizutani, 1981; Martin et al., 1982; Morgan et al., 1989; Antraygues and Aubert, 1993; Hovhannissian, 1999; Lorne et al., 1998]. These data confirmed that the electrokinetic effect could generate the observed large-scale fields, and is more plausible for volcanoes than previously suggested thermo- and piezo-magnetism mechanisms.

However, there has been little quantitative modeling of a precise physical situation, although Fitterman [1978,1979] analyzed the surface electric potential anomaly induced by a buried pressure source in a layered half-space and the surface

magnetic field in the vicinity of a vertical fault (*cf* also the recent review by Johnston, 1997).

The major purpose of the present note is to provide a sensitivity analysis of the electrokinetic effect for La Fournaise volcano. This paper is organized as follows. An idealized model of the system is proposed and the governing equations are presented as well as the method of resolution and the range of parameters. The last section is devoted to a presentation and discussion of the results.

2. General

La Fournaise volcano (Réunion Island), 2630 m high, is one of the most active basaltic volcanoes in the world with one eruption every 18 months. It is characterized by heavy rainfalls reaching 6 m per year and large anomalies in the electric potential, which may be of the order of 2 V [Michel and Zlotnicki, 1998]. A large fracture zone, about 500 m wide, cuts across the whole volcano. As mentioned by Zlotnicki and Le Mouel, [1990], this zone is likely to play an important part in the water circulation through the massif.

For this study, we adopt the simplified structure shown in Figure 1 which presents a cross-section of the volcano perpendicular to the fracture zone; the topography is simplified so that the cross-section is of a constant height $L \sim 2$ km, of the order of the height of the volcano; the top corresponds roughly to the top of the volcano and the bottom to sea level. This cross-section consists of three zones; the two external zones are assumed to be a porous medium with identical properties surrounding a fractured zone of width $2h \approx 500$ m with different properties. For simplicity, the structure is assumed to be translationally invariant along the z -axis which is perpendicular to the plane of the figure.

Meteoritic water is assumed to flow through this structure. The upper and lower boundaries of the porous medium ($x=0$ and L , $|y| \geq h$) are assumed to be impermeable; hence, water flows through the central zone and drags along water contained in the porous medium. We investigate the influence of this flow on the electric potential and the magnetic field which can be measured at the volcano surface ($x = 0$).

When an electrolyte flows through a charged medium, electric and convective phenomena are coupled [de Groot and Mazur, 1969; Nourbehecht, 1963]. The seepage velocity u and the current density I are linearly related to the driving forces which are the gradients of the pressure p and of the

Copyright 1999 by the American Geophysical Union.

Paper number 1999GL900095.
0094-8276/99/1999GL900095\$05.00




Figure 1. Model schematic of La Fournaise volcano. V_0 and J_0 are defined in (5). The x - and y -axes are vertical and horizontal, respectively.

electric potential ψ

$$I_i = -\sigma_i \nabla \psi_i - \alpha_i \nabla p_i \quad (1a)$$

$$u_i = -\alpha_i \nabla \psi_i - \frac{K_i}{\mu} \nabla p_i \quad (1b)$$

where the subscript i stands for the porous medium ($i = p$) or for the fractured medium ($i = f$). σ_i is the electrical conductivity of the medium i , K_i its permeability and α_i the electroosmotic coupling coefficient. μ is the viscosity of the fluid.

Conservation of mass and current requires that u_i and I_i satisfy the continuity equations

$$\nabla \cdot I_i = 0, \quad \nabla \cdot u_i = 0, \quad \text{for } i = p, f \quad (2)$$

This system has to be supplemented by boundary conditions. At $|y| = h$, $x \geq 0$, pressure, potential and normal fluxes are all continuous,

$$p_p = p_f, \quad \psi_p = \psi_f, \quad n \cdot I_p = n \cdot I_f, \quad n \cdot u_p = n \cdot u_f \quad (3)$$

where n denotes the unit normal to the discontinuity.

At $x = 0$ and L , for $|y| \geq h$, no flux crosses the surface. For $|y| \leq h$, the local seepage velocity V_0 and local current density J_0 are assumed not to vary with y . To summarize,

$$|y| \geq h, \quad x = 0 \quad \text{and} \quad L : u_{px} = 0, \quad I_{px} = 0 \quad (4)$$

$$|y| \leq h, \quad x = 0 \quad \text{and} \quad L : u_{fx} = V_0, \quad I_{fx} = J_0 \quad (5)$$

Other boundary conditions could have been chosen, such as $\psi = \text{constant}$ at $x = L$; however, measurements are performed at $x \sim 0$, and are insensitive to the conditions at $x = L$.

3. Solution and range of parameters

Because of the symmetry of the model with respect to the x -axis, the general solution for $y > 0$ can be written as

$$p_p = \sum_{n=0}^{\infty} a_n \cos\left(\frac{n\pi x}{L}\right) \exp\left(-\frac{n\pi y}{L}\right) \quad (6)$$

$$\psi_p = \sum_{n=0}^{\infty} c_n \cos\left(\frac{n\pi x}{L}\right) \exp\left(-\frac{n\pi y}{L}\right) \quad (7)$$

$$p_f = \Delta x + \sum_{n=0}^{\infty} b_n \cos\left(\frac{n\pi x}{L}\right) \cosh\left(\frac{n\pi y}{L}\right) \quad (8)$$

$$\psi_f = \Gamma x + \sum_{n=0}^{\infty} d_n \cos\left(\frac{n\pi x}{L}\right) \cosh\left(\frac{n\pi y}{L}\right) \quad (9)$$

The constants Δ and Γ , which are the macroscopic gradients along the x -direction in the fractured zone, are related to the local fluxes V_0 and J_0 by

$$V_0 = -\alpha_f \Gamma - \frac{K_f}{\mu} \Delta \quad (10)$$

$$J_0 = -\sigma_f \Gamma - \alpha_f \Delta \quad (11)$$

These relations can be easily inverted in order to obtain Δ and Γ as functions of V_0 and J_0 .

The form of this solution automatically satisfies the boundary conditions (4) and (5) at the upper and lower boundaries. The four unknown coefficients a_n , b_n , c_n and d_n are easily obtained by expressing the boundary conditions (3) at $|y| = h$. However, the series are often slowly convergent and some care has to be taken to obtain precise results.

Once these coefficients are obtained, the magnetic field $B(R)$ at point M (cf Figure 1) is obtained by using the Biot-Savart law

$$B(R) = \frac{\mu_0}{4\pi} \iiint_{x' \geq 0} I(R') \times \frac{R - R'}{|R - R'|^3} dR' \quad (12)$$

where μ_0 is the magnetic permittivity ($= 4\pi \cdot 10^{-7}$). The integration is performed over the whole medium (i.e., for $x' \geq 0$); x' is the abscissa of the point M' . $B(R)$ has a single horizontal component B_z . The observation point M is usually located slightly above the ground level (here, calculations are made for 1 m). The integration over z' can be performed analytically because of the translational symmetry along the z -axis; the domain can also be reduced to the quarter plane $x \geq 0$, $y \geq 0$, $z = 0$ because of the symmetry with respect to the x -axis.

Table 1. Range of studied parameters. All quantities are in SI units: σ (S), α ($\text{m}^2 \text{s}^{-1} \text{V}^{-1}$), K (m^2). Data are for $J_0 = 0$.

Case	σ_p	α_p	K_p	σ_f	α_f	K_f
c1	0.01	$3 \cdot 10^{-9}$	10^{-12}	0.1	$3 \cdot 10^{-9}$	10^{-10}
c2	0.1	$3 \cdot 10^{-9}$	10^{-10}	0.1	$3 \cdot 10^{-9}$	10^{-10}
c3	0.001	$3 \cdot 10^{-9}$	10^{-12}	0.01	$3 \cdot 10^{-9}$	10^{-10}
c4	0.01	10^{-8}	10^{-12}	0.1	10^{-8}	10^{-10}
c5	0.01	$3 \cdot 10^{-9}$	10^{-11}	0.1	$3 \cdot 10^{-9}$	10^{-9}




Figure 2. The electromagnetic fields: the current densities (a and b) in the central upper part of the volcano; the potential fields ψ (c and d) in the right half of the volcano. Data are for: case c2 (a and c); case c1 (b and d). Grey levels and arrow lengths are not consistent from one plot to another. The current densities in a and b at $x = y = 200\text{m}$ are of the order of 10^{-20} and 10^{-4} A/m², respectively. The potential differences in c and d between the dark and the white levels are equal to 2.7 and 1.5 Volts, respectively.

The many parameters which govern this situation can be divided into several groups. The first group is the geometry of the layers; since it is imposed by the geometry of the volcano, it is kept fixed

$$h = 260\text{m}, \quad L = 2000\text{m} \quad (13)$$

The second group includes the ingoing (and thus outgoing) fluxes. Usually, J_0 is taken equal to zero. An order of magnitude of the seepage velocity is estimated by assuming that when water is provided to the system, it falls under its own weight in the fractured zone

$$V_0 = \rho g \frac{K_f}{\mu} \quad (14)$$

The third group of parameters consists of the local properties K_z , α_z and σ_z . For a given medium and fluid, they can be either measured or calculated given certain assumptions [Coelho *et al.*, 1996]. Some of their properties are not intrinsic and also depend on water salinity and other physicochemical conditions. Hence, all these quantities can vary independently, at least to a first approximation.

The studied parameters are summarized in Table 1. The problem stated in Section 2 is rendered difficult by the large number of unknowns. With present knowledge, it was found preferable to make a sensitivity study around a set of central values, given as case c1.

4. Results and discussion

Overall views of the electric fields are given in Figure 2 for the standard case c1 and the homogeneous case c2. For




Figure 3. The potential field (Volts) on the ground ($x=0$) as a function of the distance y (m) to the symmetry axis. Data are for: c1 (solid line), c2 (*), c3 (···), c4 (o), c5 (---). c5 is overprinted by c1.

the sake of clarity, only half of the central upper part is shown in Figure 2a and b. Because of the continuity of the fields at $|y| = h$, the fractured zone does not present any exceptional feature. These two representations were also used in order to check the solutions, their symmetries and the boundary conditions.

The electromagnetic fields predicted for ground level are displayed in Figures 3 and 4 for all the cases listed in Table 1. First of all, for case c1, it should be emphasized that the order of magnitude of ψ ($\approx -1\text{V}$) is consistent with the observations made by Michel and Zlotnicki [1998]. However, the amplitude of B_z ($\approx 1\text{nT}$) is too small by a factor 10 [Zlotnicki and Le Mouel, 1990], a feature which will be discussed again later. These fields present a remarkable angular point at $|y| = h$ which is the limit of the fractured zone; this is consistent with the boundary conditions (3) where the normal components of the fluxes are continuous, but not the derivatives of the fields. Note that ψ tends to zero far from the fractured zone. For $|y| < 1000\text{m}$, B_z does not depend




Figure 4. The magnetic field B_z (nT) at 1m above the ground as a function of the distance y (m) to the symmetry axis. Data are for: c1 (solid line), c2 (*), c3 (···), c4 (o), c5 (---). c3 and c5 are overprinted by c1.

on the size of the integration domain of (12) along the y -axis; computations were performed for sizes equal to 4000 and 10000 m without any significant influence.

The influence of the heterogeneous character of the medium is demonstrated in these Figures by the homogeneous situation $c2$ where the permeability of the porous medium is equal to that of the fractured zone. First, the kinks disappear from the curves as they should; second, ψ and B_z are reduced by a factor 10. Hence, as is well known, the heterogeneous character of the underground medium is an essential feature to produce large fields. The residual fields which remain in $c2$ are due to the no flux condition 4 at the surface of the porous medium.

The influence of the conductivities was analyzed in case $c3$ where they were both divided by a factor 10 (cf Table 1). This parameter may vary widely; it depends on the local structure of the media and of the chemical content of the water. It is seen in Figures 3 and 4 that ψ is multiplied by a factor 10, while B_z is superposed with case $c1$ and thus does not appear clearly in Figure 4. Local water measurements are necessary to fix this important parameter.

The influence of larger coupling factors α_i was also clearly demonstrated by case $c4$ where these two factors are multiplied by a factor 3 with respect to $c1$. ψ and B_z are both roughly multiplied by a factor 3. These orders of magnitude are the ones which fit the best the measured values of Zlotnicki and Le Mouel [1990]; note that the proposed variation is very small when compared with variations measured on plugs [Ishido and Mizutani, 1981; Hovhannissian, 1999]. Hence, this factor should also be measured on samples extracted at La Fournaise.

Stronger permeabilities of the medium were studied in case $c5$ where both permeabilities were multiplied by a factor 10 with respect to $c1$. u is modified accordingly and its influence on ψ and B_z is negligible.

Finally, the influence of an hypothesized outgoing current density J_0 was studied. It was shown that for $|J_0| \leq 10^{-7}$ A/m², the influence of this overall current is negligible on the external measurements ψ and B_z ; such a value is much larger than any likely atmospheric current density.

5. Concluding remarks

Magnetic and self-potential observations on the volcano [Zlotnicki and Le Mouel, 1990; Zlotnicki et al., 1993; Michel and Zlotnicki, 1998] consist of the time dependent differences between the intensities of the geomagnetic field measured at various points distant of a few km. Only variations are useful, since constant differences are generated by the magnetized rocks of the volcano. These variations have a typical amplitude of 5-10 nT and time constants ranging from a few minutes to weeks and years. They were conjectured to be due to the modulation of a fluid flow in the rift zone by forcing mechanisms such as volcano activity and associated changes in the stress field, atmospheric forcing (temperature, rain), and long term variations in the structure of the rift zone. This hypothesis is supported by the present steady-state analysis which yields the correct orders of magnitude.

This study also demonstrates the importance of the heterogeneous character of the underground medium. Moreover, it is crucial that properties such as conductivities and coupling coefficients are measured on plugs extracted from the volcano with the local water.

This study is preliminary; other boundary conditions could have been chosen which would modify the fields, but probably not the orders of magnitude which are obtained. Flow conditions could also have been varied, and storage effects considered. Thus, this study will be extended in a near future to a time-dependent analysis in a more realistic three-dimensional representation of the volcano.

Acknowledgments. This work was partly supported by a PNRN grant (CNRS/INSU).

References

- Antraygues P. and M. Aubert, Self potential generated by two-phase flow in a porous medium: experimental study and volcanological applications, *J. Geophys. Res.*, *B98*, 22,273-22,281, 1993.
- Coelho D., Shapiro M., Thovert J.-F. and P.M. Adler, Electro-osmotic phenomena in porous media, *J. Colloid Interf. Sci.*, *181*, 169-190, 1996.
- de Groot S.R. and P. Mazur, *Non-equilibrium thermodynamics*, 510 pages, North-Holland, Amsterdam, 1969.
- Fitterman D.V., Electrokinetic and magnetic anomalies associated with dilatant regions in a layered earth, *J. Geophys. Res.*, *83*, 5923-5928, 1978.
- Fitterman D.V., Calculations of self-potential anomalies near vertical contacts, *Geophys.*, *44*, 195-205, 1979.
- Hovhannissian G.M., IPGP, unpublished data, 1999.
- Ishido T. and H. Mizutani, Experimental and theoretical basis of electrokinetic phenomena in rock-water systems and its applications to geophysics, *J. Geophys. Res.*, *86*, 1763-1775, 1981.
- Johnston M.J.S., Review of electric and magnetic fields accompanying seismic and volcanic activity, *Surveys in Geophys.*, *18*, 441-475, 1997.
- Lorne B., Perrier F. and J.-P. Avouac, Streaming potential measurements, I: Properties of the electrical double layer from crushed rock samples, *J. Geophys. Res.*, in press.
- Martin R.J., Haupt R.W. and R.J. Greenfield, The effect of fluid flow on the magnetic field in low porosity crystalline rock, *Geophys. Res. Lett.*, *12*, 1301-1304, 1982.
- Michel S. and J. Zlotnicki, Self-potential and magnetic surveying of La Fournaise volcano (Réunion Island): Correlations with faulting, fluid circulation, and eruption, *J. Geophys. Res.*, *B103*, 17,845-17,857, 1998.
- Mizutani H. and T. Ishido, A new interpretation of magnetic field variation associated with the Matushiro earthquakes, *J. Geomag. Geoelectr.*, *28*, 179-188, 1976.
- Morgan F.D., Williams E.R. and T.R. Madden, Streaming potential properties of Westerly granite with applications, *J. Geophys. Res.*, *B94*, 12,449-12,461, 1989.
- Nourbehech B., *Irreversible thermodynamic effects in inhomogeneous media and their applications in certain geological problems*, Ph. D. thesis, Mass. Inst. Technol., Cambridge, MA, 1963.
- Zlotnicki J. and J.-L. Le Mouel, Possible electrokinetic origin of large magnetic variations at La Fournaise volcano, *Nature*, *343*, 633-635, 1990.
- Zlotnicki J., Le Mouel J.-L., Delmond J.-C., Pambrun C. and H. Delorme, Magnetic variations on Piton de la Fournaise volcano and volcanomagnetic signals associated with the November 6 and 30, 1987, eruptions, *J. Volcanol. Geotherm. Res.*, *56*, 281-296, 1993.
- P.M. Adler, J.-L. Le Mouel, Institut de Physique du Globe de Paris, 4, Place Jussieu, 75252-Paris Cedex 05, France, (e-mail: adler@io.cnusc.fr, lemouel@ipgp.jussieu.fr)
- J. Zlotnicki, Laboratoire de Géophysique d'Orléans, 3D Avenue de la Recherche Scientifique, 47071 Orléans Cedex 02, France, (e-mail: jacques.zlotnicki@cnrs-orleans.fr)

(Received October 29, 1998; revised December 14, 1998; accepted January 27, 1999.)