

Isotopic composition of water vapour in the Martian atmosphere: vertical profiles from ACS MIR on ExoMars TGO

Juan Alday, Patrick G. J. Irwin, Colin F. Wilson, Kevin Olsen, Lucio Baggio, Franck Montmessin, Oleg Korablev, Alexander Trokhimovskiy, Anna A. Fedorova, Denis A. Belyaev, et al.

► To cite this version:

Juan Alday, Patrick G. J. Irwin, Colin F. Wilson, Kevin Olsen, Lucio Baggio, et al.. Isotopic composition of water vapour in the Martian atmosphere: vertical profiles from ACS MIR on ExoMars TGO. Europlanet Science Congress 2020. EPSC 2020, Sep 2020, Online, Germany. pp.EPSC2020-590, 10.5194/epsc2020-590 . insu-03121180

HAL Id: insu-03121180

<https://hal-insu.archives-ouvertes.fr/insu-03121180>

Submitted on 26 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Isotopic composition of water vapour in the Martian atmosphere: vertical profiles from ACS MIR on ExoMars TGO

Juan Alday¹, Patrick G. J. Irwin¹, Colin F. Wilson¹, Kevin S. Olsen¹, Lucio Baggio², Franck Montmessin², Oleg Korablev³, Alexander Trokhimovskiy³, Anna A. Fedorova³, Denis A. Belyaev³, Alexey Shakun³, Andrew Patrakee³, and Alexey Grigoriev³

¹University of Oxford, Atmospheric, Oceanic and Planetary Physics, Atmospheric, Oceanic and Planetary Physics, Oxford, United Kingdom of Great Britain and Northern Ireland (juan.aldayparejo@physics.ox.ac.uk)

²LATMOS/CNRS, Paris, France.

³Space Research Institute (IKI), Moscow, Russia

Abstract

We report retrievals in the 2.53 – 2.79 μm spectral range using the Atmospheric Chemistry Suite onboard of ExoMars Trace Gas Orbiter, which allow the simultaneous measurement of vertical profiles of H and O isotope ratios in water vapour in the Martian atmosphere. The large coverage of ACS MIR solar occultations provides the possibility to analyse the spatial, seasonal and diurnal variations of these ratios.

1. Introduction

Isotopic ratios in C, O and H provide important clues to understand the history and evolution of volatiles on Mars. Enrichment of the D/H ratio by a factor of approximately 5.5 in atmospheric water vapour with respect to the Vienna Standard Mean Ocean Water [1] has often been understood as evidence of the substantial atmospheric escape into space.

On the other hand, oxygen isotope ratios in both water vapour and carbon dioxide do not show substantial enrichment of the heavier isotopes (e.g. [2],[3]). The lack of substantial fractionation therefore indicates the presence of a large reservoir of oxygen to exchange with the atmosphere [4].

In this work, we use solar occultation measurements made with the mid-infrared (MIR) channel of the Atmospheric Chemistry Suite (ACS), onboard ExoMars Trace Gas Orbiter (TGO) to measure the evolution of the vertical structure of the isotopic composition of water vapour (D/H, $^{18}\text{O}/^{16}\text{O}$ and $^{17}\text{O}/^{16}\text{O}$) at the terminator for a full Martian year.

2. Measurements

ACS consists of a set of three infrared spectrometers covering a total wavelength range from 0.7 to 17 μm . The MIR channel, used in this study, is a cross-dispersion echelle spectrometer dedicated to solar occultation measurements in the 2.3-4.2 μm (2300-4400 cm^{-1}) range, with the main objective of measuring high resolution spectra ($\lambda/\Delta\lambda \approx 30000$ -50000) in a wide instantaneous spectral range (width 0.15-0.3 μm). In order to cover the full spectral range, MIR is equipped with a steerable

secondary grating that allows the selection of different diffraction orders [5].

In this work, we analyse MIR spectra in secondary grating positions 4 and 5, which allow the simultaneous measurement of several diffraction orders covering a spectral range from 2.67 to 2.79 μm (3584-3745 cm^{-1}) and 2.53 to 2.67 μm (3745-3952 cm^{-1}), respectively. We select spectral windows in several diffraction orders, which show absorption lines of the four main oxygen isotopologues of water vapour (H_2^{16}O , H_2^{18}O , H_2^{17}O , HD^{16}O) and CO_2 (see Figure 1).




Figure 1: Example of one ACS MIR spectral window in diffraction order 224, measured at tangent heights of 19, 35 and 50 km. The black dots represent the measured spectra, and the green line shows the best fit to the data. The contribution from each species to the spectra is also shown, following the colours in the legend.

3. Radiative Transfer Analysis

The analysis of the spectra is performed using the NEMESIS code [6], which works under the optimal estimation framework. In particular, for each solar occultation, we retrieve simultaneous vertical profiles of pressure, temperature and volume mixing ratio of the four main water isotopologues.

The pressure and temperature profiles can be constrained from the CO_2 absorption lines under the assumption of hydrostatic equilibrium and a given CO_2 volume mixing ratio profile, which we obtain from the Mars Climate Database [7]. In the case of the water vapour isotopologues, the volume mixing ratios can be constrained from the depth of the corresponding absorption lines, which are observed up to 60 km approximately.

The retrieval scheme is applied to all ACS MIR observations made with secondary grating positions 4 and 5 for a full Martian year, covering a wide range of latitudes, seasons, and local time, and

enabling, for the first time, the study of the climatology of both the O and H isotopic ratios in Martian water vapour.

4. Summary and Conclusions

Vertical profiles of the isotope ratios in the four main water vapour isotopologues (D/H, $^{18}\text{O}/^{16}\text{O}$ and $^{17}\text{O}/^{16}\text{O}$) at the terminator are obtained using solar occultation measurements by the Atmospheric Chemistry Suite onboard ExoMars Trace Gas Orbiter.

The large coverage of ACS solar occultation measurements allows, for the first time, the analysis of the variations of the O and H isotopic composition of water vapour for a full Martian year.

References

- [1] Owen, T., Maillard, J. P., de Bergh, C., and Lutz, B. L.: Deuterium on Mars: The abundance of HDO and the values of D/H, *Science*, 240, 1767 LP - 1767, 1988.
- [2] Webster, C. R., Mahaffy, P. R., Flesch, G. J., Niles, P. B., Jones, J. H., Leshin, L. A., Atreya, S. K., Stern, J. C., Chrsitensen, L. E., Owen, T., Franz, H., Pepin, R. O., and Steele, A.: Isotope Ratios of H, C and O in CO₂ and H₂O of the Martian Atmosphere, *Science*, 341, 260 LP – 263, 2013.
- [3] Krasnopolsky, V.A., Maillard, J. P., Owen, T., Toth, R. A., Smith, M. D.: Oxygen and carbon isotope ratios in the martian atmosphere, *Icarus*, Vol. 192, Issue 2, 2007.
- [4] Jakosky, B. M.: Mars Volatile Evolution: Evidence from Stable Isotopes, *Icarus*, 94, 14-31, 1991.
- [5] Korablev et al.: The Atmospheric Chemistry Suite (ACS) of Three Spectrometers for the ExoMars 2016 Trace Gas Orbiter, *Space Science Reviews*, vol. 214, pag. 7, 2017.
- [6] Irwin, P., Teanby, N., de Kok, R., Fletcher, L., Howett, C., Tsang, C., Wilson, C., Calcutt, S., Nixon, C., Parrish, P.: The NEMESIS planetary atmosphere radiative transfer and retrieval tool, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 109, 1136-1150.
- [7] Forget, F., Hourdin, F., Fournier, R., Hourdin, C., Talagrand, O., Collins, M., Lewis, S. R., Read, P. L., and Huot, J-P: Improved general circulation models of the Martian atmosphere from the surface to above 80 km, *Journal of Geophysical Research: Planets*, 104, 24155-24175, 1999.