An intracontinental orogen exhumed by basement-slice imbrication in the Longmenshan Thrust Belt of the Eastern Tibetan Plateau

Zhenhua Xue, Wei Lin, Yang Chu, Michel Faure, Yan Chen, Wenbin Ji,
Huaning Qiu

► To cite this version:

Zhenhua Xue, Wei Lin, Yang Chu, Michel Faure, Yan Chen, et al.. An intracontinental orogen exhumed by basement-slice imbrication in the Longmenshan Thrust Belt of the Eastern Tibetan Plateau. Geological Society of America Bulletin, 2021, 10.1130/B35826.1 . insu-03186417

HAL Id: insu-03186417

<https://hal-insu.archives-ouvertes.fr/insu-03186417>

Submitted on 1 Apr 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **An intracontinental orogen exhumed by basement-slice-imbrication in the**
2 **Longmenshan Thrust Belt of the Eastern Tibetan Plateau**

3
4 **Zhenhua Xue^{1,2,3}, Wei Lin^{2,4}, Yang Chu^{2,4}, Michel Faure³, Yan Chen³, Wenbin Ji⁵,**
5 **Huaning Qiu⁶**

6 ¹ School of Earth Sciences, China University of Geosciences, Wuhan 430074, China

7 ² State Key Laboratory of Lithospheric Evolution, Institute of Geology and Geophysics,
8 Chinese Academy of Sciences, Beijing 100029, China

9 ³ Univ. Orléans, CNRS, BRGM, ISTO, UMR 7327, F-45071, Orléans, France

10 ⁴ University of Chinese Academy of Sciences, Beijing 100049, China

11 ⁵ Department of Geology, Northwest University, Xi'an 710069, China

12 ⁶ Key Laboratory of Tectonics and Petroleum Resources (China University of
13 Geosciences), Ministry of Education, Wuhan 430074, China

14 Corresponding author: Wei Lin (linwei@mail.iggcas.ac.cn)

15
16 **Key Points**

- 17 1. The double-vergent structure of the Longmenshan belt was established during the
18 late Triassic intracontinental underthrusting
19 2. Boundary faults of the NW-normal faulting and SE-thrusting facilitated the SE-ward
20 basement-slice imbrication
21 3. The episodic basement-slice imbrication exhumed the early Mesozoic metamorphic
22 rocks

23
24 **Abstract** The Longmenshan Thrust Belt (LMTB) in the Eastern Tibet resulted from a
25 Mesozoic orogeny and Cenozoic reworking. It is generally believed that the Cenozoic
26 tectonics along the LMTB are mostly inherited from the Mesozoic one. Reconstruct the
27 Mesozoic tectonics of the LMTB is therefore important for understanding its
28 evolutionary history. On the basis of detailed structural analysis, we recognized a Main
29 Central Boundary that divides the LMTB into a Southeastern Zone and a Northwestern
30 Zone. Both zones underwent a main D₁ event characterized by the D_{1E} top-to-the-SE
31 thrusting in the Southeastern Zone and the D_{1W} top-to-the-NW/N thrusting in the
32 Northwestern Zone. In the Southeastern Zone, a D₂ top-to-the-NW/N normal faulting

that cuts the D_{1E} structures is developed along the NW boundary of the basement complexes. Newly obtained and previous geochronological data indicate that the D_{1E} and D_{1W} event occurred synchronously at *ca.* 224–219 Ma, and the D₂ top-to-the-NW/N normal faulting was episodically activated at *ca.* 166–160 Ma, 141–120 Ma, 81–47 Ma and 27–25 Ma. Episodically and synchronously activated top-to-the-NW normal faulting and top-to-the-SE thrusting along the northwest and southeast boundary of the basement complexes, respectively, let us propose that the basement-slices were episodically imbricated to the SE during the late Jurassic–early Cretaceous and late Cretaceous–earliest Paleocene. The D₁ amphibolite-facies metamorphic rocks above the basement complexes recorded fast exhumation during the late Jurassic–early Cretaceous. We propose that the early Mesozoic northwestward basement underthrusting along a crustal “weak zone” was responsible for the D₁ double-vergent thrusting and amphibolite-facies metamorphism. Subsequent basement-slice imbrications reworked the LMTB and exhumed the amphibolite-facies rocks. Our results highlight the importance of basement underthrusting and imbrication in the formation and reworking of the intracontinental Longmenshan Thrust Belt in the Eastern Tibet.

1. Introduction

The Longmenshan Thrust Belt (LMTB; Figure 1a), at the eastern margin of the Tibetan Plateau, has attracted attentions of geologists due to the paradox between low convergence rate (Zhang et al., 2004; Shen et al., 2009) and extremely high topographic gradient (Clark and Royden, 2000; Royden et al., 2008) as well as intense seismic activities (e.g., 2008 Wenchuan Mw 7.9 earthquake, 2013 Lushan Mw 7.0 earthquake, and 2017 Jiuzhaigou Mw 7.0 earthquake). This paradox has spawned a vigorous debate regarding the roles of upper crustal faulting (Hubbard and Shaw, 2009) and lower crust channel flow (Royden et al., 2008) in building and maintaining such a high Tibetan Plateau margin. However, both models attribute shortening and crustal thickening of the LMTB to the late Cenozoic evolution, and underestimate the importance of widely recognized Mesozoic tectonics along the LMTB (Xu et al., 1992; Burchfiel et al., 1995; Worley and Wilson, 1996). Furthermore, it has been reached a consensus that the LMTB is mainly inherited from the Mesozoic tectonic framework (Burchfiel et al.,

1995; Meng et al., 2005). Elucidating the Mesozoic tectonics of the LMTB and its adjacent areas is therefore crucial to understand the tectonic evolution of the LMTB.

The prominent deformation recorded in the LMTB and adjacent areas is indicated by observations including: 1) the peripheral foreland basin formed due to the southeastward thrusting and loading of the LMTB (Chen et al., 1994; Li et al., 2003); 2) the early Jurassic rocks unconformably overlies extensively folded late Triassic rocks (BGMRSF, 1991; Xu et al., 1992; Burchfiel et al., 1995); 3) the pervasive orogen-parallel klippen and nappe belt sitting on the foot hills of the Longmen Mountains are believed related to the regional top-to-the-SE thrusting (Figure 2); 4) the consistent top-to-the-SE thrusting indicated by different indicators at various scales (Xu et al., 1992; Burchfiel et al., 1995; Chen et al., 1995; Chen and Wilson, 1996; Jia et al., 2006; Yan et al., 2011; Yan et al., 2018); 5) the pervasive upright folding in the adjacent Songpan–Ganzi Terrane was thought synchronous to the early Mesozoic deformation of the LMTB (Burchfiel et al., 1995; Chen et al., 1995; Harrowfield and Wilson, 2005). This prominent deformation event was constrained to *ca.* 234–200 Ma by basin tectonics, timing of the top-to-the-SE/SSE shearing and syntectonic metamorphism, which indicated an early Mesozoic orogeny of the LMTB (Chen et al., 1994; Li et al., 2003; Yan et al., 2011; Zheng et al., 2014; Xue et al., 2016; Zheng et al., 2016; Airaghi et al., 2018a; Airaghi et al., 2018b; Yan et al., 2018).

The early Mesozoic tectonic evolutionary mechanism accounting for the above observations, however, is still in dispute. Sengör (1984) and Luo (1991) suggested that the LMTB is an early Mesozoic intracontinental subduction zone accommodating multi-stage crustal shortening. Then, Xu et al. (1992) proposed a bivergent subduction model in which the Songpan–Ganzi Terrane sedimentary cover has been “pushed” to the southeast on top of the Yangtze Block due to the bivergent subduction along the Aymaqen and Jinshajiang Suture Zones. The current models suggested that the Songpan–Ganzi sedimentary cover were overthrust southwards on the Yangtze Block along a decollement in response to the closure of the Paleotethys (Mattaier et al., 1992; Calassou, 1994; Dirks et al., 1994; Chen and Wilson, 1996; Worley and Wilson, 1996; Roger et al., 2004; Harrowfield and Wilson, 2005; de Sigoyer et al., 2014; Yan et al., 2018).

Apart from the early Mesozoic top-to-the-SE/SSE thrusting along the LMTB, an increasing number of studies proposed a top-to-the-NW normal faulting that developed

98 along the NW boundary of the orogen-parallel and stripped Neoproterozoic complexes
99 (Figure 1b)(Xu et al., 2008; Zhou et al., 2008; Tian et al., 2016; Billerot et al., 2017;
100 Xue et al., 2017; Airaghi et al., 2018b). However, timing and significance of the top-
101 to-the-NW normal faulting are in dispute that have resulted in several different models,
102 such as: 1) mid-Jurassic to early Paleogene (173–160 Ma and 81–47 Ma) post orogenic
103 extension (comparing to the early Mesozoic orogeny)(Zhou et al., 2008), 2) early
104 Cretaceous (~120 Ma) crustal-scale wedge extrusion (Xu et al., 2008), 3) episodic
105 eastward basement-slice-imbrication during the late Jurassic–early Cretaceous and late
106 Cretaceous–Earliest Paleogene (Billerot et al., 2017; Xue et al., 2017), and, 4)
107 formation of a late Cretaceous–earliest Paleogene crustal scale duplex (Tian et al.,
108 2016). Based on the above compilation, most previous works have proposed that the
109 top-to-the-NW normal faulting occurred in the late Mesozoic.

110 Major questions therefore remain unsolved: 1) Considering previous early Mesozoic
111 tectonic models of the LMTB, whether did the significant shortening recorded in the
112 LMTB correspond to the imbricated thrust system during the Paleotethys' closure, or
113 to the intracontinental underthrusting along the Longmenshan belt? 2) What is the
114 tectonic significance of the top-to-the-NW normal faulting (mostly late Mesozoic) and
115 emplacement of the Neoproterozoic complexes? Therefore, an overall accurate
116 reconstruction of the Mesozoic tectonics of the LMTB and its adjacent areas, especially
117 its detailed structural analysis and timing, are of great importance.

118 This study presents detailed field investigations focusing on structural geometry and
119 kinematic reconstruction, and geochronological constraints including zircon U–Pb
120 dating of synkinematic plutonic rocks and $^{40}\text{Ar}/^{39}\text{Ar}$ dating on the synmetamorphic
121 micas of the LMTB. In conjunction with previous published geochronological and
122 metamorphic data, a new geodynamic model was proposed to interpret the Mesozoic
123 tectonic evolutionary history of the LMTB.

125 **2. Geological Setting**

126 At the eastern margin of the Tibetan Plateau, the LMTB separates the intensely
127 folded and metamorphosed (Greenschist facies grade) Songpan–Ganzi Terrane to the
128 NW, and the weakly/undeformed Sichuan Basin to the SE (Figure 1a). The triangle-
129 shaped Songpan–Ganzi Terrane is separated from the western Qinling–Dabie Orogenic

Belt to the north by the East Kunlun–Anymaqen Suture Zone, from the Qiangtang Block to the southwest by the Jinshajiang Suture Zone (Figure 1a).

2.1 Songpan–Ganzi Terrane

The Songpan–Ganzi Terrane mainly comprises ~5–15 km-thick intensely folded late Triassic turbidites (Calassou, 1994; Harrowfield and Wilson, 2005; Zhang et al., 2006c; Zhang et al., 2008; Yin, 2015). The provenance of the turbidites includes the Kunlun Mountains, North China Block, South China Block and southwest Qinling Orogenic Belt (Yin and Nie, 1993; Enkelmann et al., 2007; Weislogel et al., 2010). Voluminous felsic plutons intruded the Songpan–Ganzi Terrane at *ca.* 226–205 Ma, dominantly derived from partial melting of the Neoproterozoic basement and sedimentary cover of the Songpan–Ganzi Terrane, with negligible amounts of mantle material (Figure 1b)(Roger et al., 2004; Hu et al., 2005; Zhang et al., 2006b; Zhang et al., 2007; de Sigoyer et al., 2014). Both geophysical and geological data indicate a continental crust of Yangtze Block affinity underlying the eastern Songpan–Ganzi Terrane (Roger et al., 2004; Hu et al., 2005; Zhang et al., 2009; Weislogel et al., 2010; Zhang et al., 2010; Guo et al., 2013).

2.2 Sichuan Basin of the Yangtze Block

The Sichuan Basin, located in the Yangtze Block of northwest South China Block (Figure 1a), preserves a Paleoproterozoic to Neoproterozoic basement overlain by Neoproterozoic to Cenozoic sedimentary sequences. Silurian–Devonian abyssal deposits and radiolarian chert suggested a long period of rifting during the Silurian to Devonian ages along the western part of the Yangtze Block (Long, 1991; Luo, 1991; Chang, 2000; Li, 2009). This rift was revealed by high-resolution seismic reflection profile as well (Jia et al., 2006). During middle–late Triassic ages, coeval with the southeastward thrusting and loading of the LMTB, the western Sichuan Basin began subsiding flexurally and evolved into an epicontinental foreland basin (Figure 2). Thrust sheets were transported to the southeast synchronously and intensely folded at the end of the late Triassic (Chen et al., 1994; Chen and Wilson, 1996; Li et al., 2013c). The continuous sedimentation of the middle Jurassic–upper Cretaceous deposits implies a pronounced subsidence at the western Sichuan foreland basin during this period (Burchfiel et al., 1995; Chen et al., 1995). Episodic SE-ward thrusting and loading of the LMTB were indicated by the wedge-shaped megasequences of the western Sichuan foreland basin, namely, in late Triassic–early Jurassic, late Jurassic–

early Cretaceous and late Cretaceous–earliest Paleocene (Xu et al., 2008; Li et al., 2013c; Tian et al., 2016).

2.3 The LMTB

The NE-striking LMTB merges into the western Qinling Orogenic Belt to the northeast and is truncated by the Xianshuihe Fault to the southwest (Figure 1a).

2.3.1 Major faults of the LMTB

The regional tectonic architecture of the LMTB is defined by four major NE–SW striking faults, namely from the southeast to the northwest, the Anxian–Guanxian Fault (T_1 in Figure 1b), the Yingxiu–Beichuan Fault (T_2 in Figure 1b), the Wenchuan–Maoxian Fault (T_3 in Figure 1b), and the Qingchuan–Pingwu Fault (T_4 in Figure 1b).

The Anxian–Guanxian Fault dips to the northwest by $30\text{--}50^\circ$ and crosscuts the late Triassic–early Jurassic rocks. Approaching northwestwards, the Yingxiu–Beichuan Fault dipping $40\text{--}60^\circ$ to the NW separates the Sichuan foreland basin to the southeast from the LMTB hinterland (BGMRSF, 1991; Xu et al., 1992; Burchfiel et al., 1995).

The Wenchuan–Maoxian Fault dipping $30\text{--}50^\circ$ to the NW was episodically activated since *ca.* 220 Ma (Burchfiel et al., 1995; Yan et al., 2008; Zheng et al., 2016). Seismic reflection profile showed that the ENE-striking Qingchuan–Pingwu Fault dips to the NNW or SSE at variable angles (Wang et al., 2000; Jia et al., 2006; Lin et al., 2014).

The northern LMTB was dissected by the Qingchuan–Pingwu Fault and incorporated to northwestward- and southeastward-thrusting that resemble a positive flower structure (Jia et al., 2006). The Wenchuan–Maoxian Fault and Qingchuan–Pingwu Fault are characterized by mylonite overprinted by breccia and gouge (Chen and Wilson, 1996; Li et al., 2013a; Li et al., 2015).

2.3.2 Basement complexes in the LMTB

The basement complexes that distributes from the NE to the SW in the LMTB are composed of Neoproterozoic meta-granites overlain by a Neoproterozoic meta-sedimentary cover, such as the Danba (DB), Kanding (KD), Wulong (WL), Baoxing (BX), Pengguan (PG), Moutuo (MT), Xuelongbao (XLB) and Jiaoziding (JZD) complexes (Figure 1b). These basement complexes share similar lithology with the Yangtze Block basement (BGMRSF, 1991). Field observations, Anisotropy of Magnetic Susceptibility (AMS) and gravity modeling have suggested that these basement complexes were strongly deformed and imbricated southeastwards as

basement-slices (Ma et al., 1996; Xu et al., 2008; Tian et al., 2016; Airaghi et al., 2017b; Billerot et al., 2017; Xue et al., 2017).

2.3.3 Metamorphism in the LMTB

Two amphibolite facies metamorphic zones have been recognized in the meta-sedimentary cover in the Wenchuan–Xuelongbao area and the Danba area (Dirks et al., 1994; Huang et al., 2003a; Zhou et al., 2008; Airaghi et al., 2018a; Airaghi et al., 2018b). Petrological studies (garnet–amphibolite–biotite thermometer, pseudosection using Perplex modeling and chlorite–white mica–quartz thermal dynamic equilibrium) constrain the peak metamorphic conditions at $\sim 11 \pm 2$ kbar, ~ 620 °C in the Wenchuan–Xuelongbao area (Dirks et al., 1994; Airaghi et al., 2018a) and ~ 5.3 –8 kbar, 570–680 °C in the Danba area (Huang et al., 2003a; Weller et al., 2013). The Wenchuan–Xuelongbao metamorphic belt reached amphibolite facies conditions at *ca.* 210–180 Ma (Airaghi et al., 2018a; Airaghi et al., 2018c) and recorded fast exhumation at *ca.* 140–120 Ma (Arne et al., 1997; Airaghi et al., 2018c). The Danba amphibolite-facies metamorphic belt reached peak conditions at *ca.* 205–180 Ma (Huang et al., 2003a; Weller et al., 2013) and initiated enhanced rapid cooling at *ca.* 168–158 Ma (Huang et al., 2003a; Zhou et al., 2008).

Apart from these two amphibolite facies metamorphic zones, late Triassic blueschists occur as lenses along the Qingchuan–Pingwu Fault, with peak metamorphic pressure attained to 5.2–6.5 kbar (Wei, 1994; Xu et al., 2020). The blueschist then have been retrograded to greenschist in a nearly isothermal decompression way (Wei, 1994).

2.3.4 Tectonic events of the LMTB

The LMTB recorded at least three phases of deformation, including (1) D₁, the late Triassic to early Jurassic regional top-to-the-SE thrusting, (2) D₂, the top-to-the-NW/N normal faulting of poorly constrained timing and tectonic significance, and (3) D₃, the Cenozoic shortening reactivation (Xu et al., 1992; Burchfiel et al., 2008; Richardson et al., 2008; Xu et al., 2008; Li et al., 2013a; Li et al., 2015; Tian et al., 2016; Xue et al., 2017). The D₁ deformation is well identified by penetrative cleavages (S₁), mineral and stretching lineation (L₁) in the Wenchuan area, and the top-to-the-SE thrust sheets and nappes over the intensely folded late Triassic sequences in the western Sichuan foreland basin (Xu et al., 1992; Burchfiel et al., 1995; Worley and Wilson, 1996; Li et al., 2003). Most parts of the Songpan–Ganzi Terrane show slow cooling rate at *ca.* 1 °C/Ma since the early Mesozoic (Huang et al., 2003a; Huang et al., 2003b; Roger et al., 2011).

However, an enhanced cooling event (*ca.* 8°C/Ma) during the late Jurassic to Eocene was reported at the peripheries of the Songpan–Ganzi Terrane, especially in the Wenchuan and Danba areas (Yuan et al., 1991; Arne et al., 1997; Huang et al., 2003a). This rapid cooling event probably responded to the D₂ top-to-the-NW/N normal faulting. The belt characterized by the top-to-the-NW normal faulting developed along the northwest/north boundary of the basement complexes was entitled as the East Tibet Detachment (Wang et al., 2012b). The D₃ deformation was thought to have initiated since *ca.* 30–25 Ma that builds the present Longmen mountains of the eastern Tibetan Plateau (Clark and Royden, 2000; Kirby et al., 2002; Wang et al., 2012a).

3. Structural analysis of the LMTB

3.1 Structural subdivisions of the LMTB

The LMTB was interpreted as an allochthonous sheet thrusting southeastward upon the autochthonous western Sichuan Foreland Basin (Xu et al., 1992; Burchfiel et al., 1995; Worley and Wilson, 1996; Robert et al., 2010a). Based on structural investigations, we present revised structural subdivisions of the LMTB. Firstly, the LMTB is divided into a Southeastern Zone and a Northwestern Zone by a Main Central Boundary (see Section 3.2; Figure 3a). According to the lithology, deformation intensity, and metamorphic grade, the Southeastern Zone can be further subdivided into four subdivisions (Figure 3). They are, from the southeast to the northwest: i) the undeformed Autochthon that comprises the western part of the Sichuan Foreland Basin, ii) the Para-autochthon, transported southeastwards along the Anxian–Guanxian Fault and rests over the Autochthon, iii) the Allochthon that traveled southeastwards much further on top of the Para-autochthon, and iv) the Inner Metamorphic Core that recorded the highest metamorphic conditions (*i.e.*, amphibolite facies) among the Southeastern Zone. The four subdivisions in the Southeastern Zone are separated by the three major listric faults of the LMTB (Figure 3; see in Section 2.3.1).

i) The Autochthon, located to the southeast of the Anxian–Guanxian Fault, is mainly composed of Jurassic–Cretaceous strata and overlying Cenozoic sediments (Figures 3 and 4). The Jurassic–Cretaceous strata were deposited in circumstances of alluvial, fluvio-deltaic, and lacustrine fills, characterized by medium and coarse-grained sandstone with some conglomeratic layers at the base. Total thickness of the Jurassic–Cretaceous sequences is more than 5 km (BGMRSF, 1991). In the northern part of the

LMTB, close to Anxian and Jiangyou county, the early Jurassic and overlying Cretaceous to Cenozoic sequences are dipping to the southeast that unconformably rest on the intensively folded late Triassic rocks (Figures 2 and 4c–d)(BGMRSF, 1991).

ii) The Para-autochthon is sandwiched by the Anxian–Guanxian Fault and the Yingxiu–Beichuan Fault, with its half area covered by klippen and nappes from the Allochthon (Figure 3). The Late Triassic Xujiahe Formation dominated in this subdivision either passes up from the earliest late Triassic marine sandstone or unconformably overlies the middle Triassic limestone (Li et al., 2003; Meng et al., 2005). The Xujiahe Formation shows markable lateral variation in both thickness and sedimentary facies. Coarse-grained sediments, such as alluvial conglomerate bodies occur close to the Yingxiu–Beichuan Fault, but change into fine-grained lacustrine sediments toward to the basin interior.

iii) The Allochthon is mainly exposed between the Wenchuan–Maoxian Fault and the Yingxiu–Beichuan Fault, and also includes the klippen and nappes resting on top of the Para-autochthon (Figures 2 and 3a). Northwest to the Yingxiu–Beichuan Fault, this subdivision mainly consists of Silurian shale and greywacke with the bedding completely transposed into foliation. Greenschist-facies metamorphism is indicated by mineral assemblages of fine-grained muscovite + chlorite + quartz (Airaghi et al., 2017b; Xue et al., 2017). Metamorphic condition jumps of ~50 °C, 3–5 kbar southeastwards across the Yingxiu–Beichuan Fault suggest that the Allochthon are more deformed and metamorphosed in a deeper structural level than the Para-autochthon (Robert et al., 2010a; Airaghi et al., 2018b). Neoproterozoic complexes including the Pengguan and Jiaoziding complexes are distributed in this subdivision. Southeast to the Yingxiu–Beichuan Fault, klippen and nappes are mainly composed of Devonian to early Triassic thick-bedded limestone and marine sandstone (Figures 2, 3 and 4a–d).

iv) The Inner Metamorphic Core comprises phyllite and micaschist that are mostly metamorphosed Silurian sediments (Figures 2 and 3). Amphibolite-facies metamorphic zone that wraps the oval-shaped Xuelongbao (XLB in Figure 2) and stripped Moutuo (MT in Figure 2) complexes records the highest-grade metamorphism within the Southeastern Zone (Dirks et al., 1994; Airaghi et al., 2017a; Airaghi et al., 2018a). Peak P–T condition attained to $\sim 11 \pm 2$ kbar and ~ 620 °C at the southernmost margin of the Xuelongbao complex calculated from metamorphic assemblages of biotite + garnet +

294 staurolite + muscovite + kyanite (Dirks et al., 1994; Airaghi et al., 2018a). Based on
295 the enhanced metamorphic grade and deformation intensity, the Inner Metamorphic
296 Core are therefore more deformed and metamorphosed in a deeper structural level than
297 the Allochthon.

298 The Northwestern Zone includes eastern Songpan–Ganzi Terrane with
299 Neoproterozoic sequences outcropped in the north and Paleozoic sequences outcropped
300 in the south (Figures 2 and 3). The northern Northwestern Zone mainly consists of two
301 Neoproterozoic Groups, namely, 1) the volcanic Bikou Group defined by mafic lavas
302 within its lower part and ranges from mafic to intermediate–felsic within its upper
303 portion, and 2) the Hengdan Group tectonically resting on the Bikou Group is composed
304 of black, thin-bedded pyrite-bearing shales and interbedded thinly volcanoclastic
305 sandstone and siltstone (Yan et al., 2003; Druschke and Wang, 2006). The southern
306 Northwestern Zone includes Devonian Weixian Formation and the Triassic sequences.
307 The Devonian sequences consist mainly of quartziferous phyllite to schist and
308 interbedded quartzite. The overlying Triassic Xikang Group is dominated by thick and
309 highly repetitious flysch sediments. Although different rock units comprising the
310 northern and southern Northwestern Zone, it was metamorphosed under similar
311 greenschist facies conditions (Figures 1b and 2)(Xu et al., 1992; Liu et al., 2008)

313 3.2 The Main Central Boundary

314 The Main Central Boundary serves as a major tectonic boundary separating the
315 Northwestern Zone and the Southeastern Zone. It corresponds to the southern branch
316 of the Qingchuan Fault (T₄ in figure 2) in the north (BGMRSF, 1991), and to a vertical
317 cleavage zone within the Paleozoic rocks in the south (Figure 2). To the SE of the Main
318 Central Boundary, all structural planes in the Southeastern Zone, including foliations,
319 cleavages, and fold axial planes, are exclusively dipping to the NW. Various kinematic
320 indicators in the Southeastern Zone suggest a consistent top-to-the-SE/S thrusting. To
321 the NW of the Main Central Boundary, the Northwestern Zone is characterized by the
322 SSE-dipping planar structures and associated top-to-the-NW thrusting (Figure 3).

323 The Qingchuan–Pingwu Fault, as the northern segment of the Main Central
324 Boundary, is a major tectonic boundary. Seismic profiles, drill data (Jia et al., 2006)
325 and field observations indicate that the Qingchuan–Pingwu Fault is a positive flower
326 structure developed in a transpressional system. Microfabric study (Wang et al., 2000)

indicated that the Qingchuan–Pingwu Fault was originated from a deep-seated ductile shear zone during late Triassic and overprinted by Cenozoic brittle deformation at very least. It is the late Triassic deep-seated ductilely sheared Qingchuan–Pingwu Fault rather than the Cenozoic brittle one that separates the LMTB into the Southeastern Zone and Northwestern Zone.

The southern segment of the Main Central Boundary roughly distributes along the geological boundary between the Silurian and Devonian–Permian sequences (Figure 2). Macroscopically, this segment of Main Central Boundary shows predominant vertical cleavage and foliation within intensely deformed argillaceous limestone, graphitic pelite and thin-bedded limestone–quartz ribbon interlayers (Figures 3a and 5). Their bedding (S_0) can be distinguished by the compositional variations that is parallel to the cleavages (Figure 3a and 5b). Associated rootless folds imply significant shortening during the formation of the Main Central Boundary (Figures 3a and 5b–c). P-wave receive function and seismic reflection profiles revealed a Moho at ~36–40 km depth beneath the Sichuan Basin that gradually deepens northwestwards at ~63 km depth beneath the eastern Tibet (Guo et al., 2013). This transition occurs beneath the Wenchuan Shear Zone (corresponding to Inner Metamorphic Core) over a horizontal distance of fewer than 30 km (Robert et al., 2010b; Zhang et al., 2010; Guo et al., 2013). Geophysical profiles, therefore, support the statement that the southern Main Central Boundary, as the northwest boundary of the Inner Metamorphic Core, serves as a major tectonic boundary.

3.3 The main deformation (D_1)

The D_1 main deformation is defined by a NW–SE striking stretching lineation recognized in both the Southeastern and Northwestern zones which is separated by the Main Central Boundary (Figure 2). Kinematic indicators indicate a top-to-the-SE thrusting and a top-to-the-N/NW thrusting in the Southeastern and the Northwestern zones, respectively. Since there is no evidence of superposition between these two senses of shear, it is difficult to precisely define the relative timing of these opposite shearing events. Nevertheless, to clarify the presentation of the deformation events, in the following, we distinguish D_{1E} , the top-to-the-SE thrusting in the Southeastern Zone, and D_{1W} , the top-to-the-NW/N thrusting in the Northwestern Zone, respectively.

3.3.1 Top-to-the-SE thrusting in the Southeastern Zone (D_{1E})

The top-to-the-SE shearing and thrusting are widespread in the Southeastern Zone at variable scales, except that the Autochthon merely recorded long-wavelength fold and monoclines toward the interior of the Sichuan Basin (Figure 4). The D_{IE} top-to-the-SE thrusting is most developed in the Inner Metamorphic Core and generally attenuated southeastwards.

The Inner Metamorphic Core was ductilely deformed showing predominant NW-dipping foliation (S_1) at varied angles between 60–90°. Specifically, the foliation gradually steepens when approaching the Main Central Boundary, and eventually presents in a vertical attitude (Figures 2–4). This changing trend of foliation implies that the deformation intensified toward the Main Central Boundary. Foliations mostly consist of an alternation of biotite-white mica-rich and light-colored, pelitic interlayers (Figures 3a and 6a–b). Along the transect from Wenchuan county to Xuecheng town, an NW-plunging mineral and stretching lineation (L_1) defined by aligned newly formed biotite, white mica, elongated quartz aggregates, or iron-oxide pods are well developed (Figures 3a and 6a). Kinematic indicators document a consistent top-to-the-SE thrusting in the Inner Metamorphic Core. Such as asymmetric pyrite pressure shadow in mica-schist (Figures 3a and 6b) and sigma-shaped feldspar porphyroclasts in paragneiss (Figures 3a and 6c) both show top-to-the-SE thrusting. Photomicrographs (perpendicular to the foliation and parallel to the lineation) show that newly formed white mica oriented as compaction plane (S) and shear plane (C), both of which composes an S-C fabric documenting a consistent top-to-the-SE thrusting (Figures 3a and 6d).

In the Allochthon, the Neoproterozoic complexes such as the Pengguan and Jiaoziding complexes, mostly show brittle deformation characterized by subhorizontal joints. Inside the Pengguan complex, several NE–SW trending ductile shear zones dissected and thrust the granitic rocks to the southeast (Airaghi et al., 2017b; Xue et al., 2017). Upon the granitic rocks, the Neoproterozoic Huangshuihe Group recorded extensive shearing and sigmoid pelitic lens incorporated to the top-to-the-SE thrusting (Figures 3a and 6e). Structural planes including axial planar cleavages and spaced cleavages dip to the northwest at angles of 30–65° (Figures 2–4). Fine-grained white mica and chlorite are developed on these structural planes demonstrating a greenschist facies grade of deformation. Sigma-shaped sandy lens in the Silurian pelitic schist indicate a top-to-the-SE thrusting as well (Figures 3a and 6f).

392 As part of the Allochthon, most of the klippen are synclines overturned to the SE,
393 with SE-ward reverse fault which dissected the thick Devonian limestone at the rear of
394 the klippe (Figures 2 and 6g), and show a consistent top-to-the-SE thrusting (BGMRSF,
395 1991).

396 In the Para-autochthon, the late Triassic Xujiahe sandstone is intensively folded as
397 indicated by the NE-SW trending π -circle-projected axis of bedding (Figures 3b-d). In
398 the south part, multi π circle-Axis exist roughly trending NE-SW, implying multiphase
399 of shearing and thrusting. Folds are mostly overturned to the SE indicating a top-to-
400 the-SE thrusting (Figure 2). Axial planar cleavages are locally developed that gradually
401 intensified and more penetrative approaching northwestward to the Yingxiu–Beichuan
402 Fault. This cleavage-enhancing trend implies a northwestward increase in deformation
403 intensity. In mudstone–sandstone interlayered rocks, competence contrast facilitates
404 the formation of layer-parallel slip and shearing. Sigmoid lens as well as “cold” shear-
405 band fabrics consistently indicate a top-to-the-SE thrusting (Figures 3a and 6h).

406 In the Autochthon, lower Jurassic strata and overlying continuous Cretaceous–
407 Cenozoic sequences are mostly concordantly folded in a long-wavelength way. The
408 fact that early Jurassic rocks locally overturned to the southeast (North to the Tangbazi
409 town; Figure 2) therefore implies a post-Jurassic reactivation of the SE-directed
410 thrusting (Figures 2 and 4d).

411 3.3.2 Top-to-the-NW/N thrusting in the Northwestern Zone (D_{1W})

412 The structures and kinematics within the Northwestern Zone are different from those
413 within the Southeastern Zone. In the northern Northwestern Zone, S-dipping foliation
414 and associated downdip lineation are well developed. Lineations are defined by aligned
415 calcite, fine-grained white mica, and oxide pods (Figures 2, 3a and 7a). Along the
416 lineation, a consistent top-to-the-N/NW shearing and thrusting is determined by various
417 kinematic indicators, such as sigma-shaped pebbles in strongly sheared meta-
418 conglomerate (Figures 3a and 7b) and asymmetric pressure shadow in sandy–pelitic
419 schist. Photomicrographs show that quartz ribbon (Figures 3a and 7c) and sigma-shaped
420 feldspar porphyroclasts (Figures 3a and 7d) both indicate a top-to-the-N thrusting. In
421 the north of the Pingwu city, greywacke and siltstone were extensively folded verging
422 to the north with associated meter-scale reverse fault cutting fold limbs (Figures 3a and
423 7e). West to the Pingwu city, bedding (depicted by compositional banding) completely

transposed to SE-dipping foliations in response to the northward shearing and thrusting (Figures 3a and 7f).

In the southern Northwestern Zone, axial planar cleavage developed in the thin-bedded limestone mostly dips to the SE but strikes roughly along the margin of the Xuelongbao complex (Figure 2). On the foliation, lineation plunges to the SE at ~65–70° (Figures 2 and 3a). Along the lineation, asymmetric pressure shadow of calcite tail around garnet pseudomorph indicates a top-to-the-NW thrusting (Figures 3a and 7g). Asymmetric NW-verging folds are developed at different scales. The late Triassic turbidite was involved in a hundred-meter scale chevron fold verging to the NW that is consistent with the top-to-the-NW thrusting (Figures 3a and 7h). In the whole Northwestern Zone, the top-to-the-NW/N thrusting is indicated by the pervasive NW/N-overtaken bedding as well (Figure 2)(BGMRS, 1991).

3.3.3 Syntectonic (D_{1W}) Mupi pluton in the Northwestern Zone

In the Northwestern Zone, north of the Pingwu city, the Mupi pluton is a coarse-grained diorite and granodiorite pluton with prominent syntectonic features (Figure 2). At its southern margin, the contact between the pluton and the country rock is rather ambiguous. Strong mylonitisation with S-dipping foliation deformed both the country rocks and the south margin of the pluton (Figures 3a and 8c). On the S-dipping foliation (S_1), aligned biotite and sheared feldspar define a mylonitic lineation plunging south at ~45–75°. Kinematic indicators including sigmoidal feldspar porphyroclasts and shear bands both indicate a top-to-the-N shearing related to thrusting (Figures 3a and 8d). About 10–15 meters northwards from the south margin of the pluton, a set of quartz + feldspar veins parallel to the shear foliation (Figures 3a and 8b), in which the quartz and feldspar did not record ductile deformation. Further north ~30 meters from the south margin of the pluton, the pluton rocks show magmatic fabric with euhedral feldspar randomly distributed (Figures 3a and 8a). Cracks in feldspar grains are sub-parallel to the shear planes and infilled with quartz (Figures 8e and S1). These cracks in conjunction with the foliation-parallel quartz + feldspar veins indicate that a brittle deformation of the feldspar occurred in the presence of residual melt. Therefore, based on the textural criteria proposed by Paterson et al. (1989), we prefer to interpret the Mupi pluton as a syntectonic one relating to the D_{1W} deformation.

3.4 Top-to-the-NW normal faulting along the northwest boundary of the basement complexes (D₂)

Most of the Neoproterozoic basement complexes in the Southeastern Zone are devoid of ductile deformation but characterized by widespread subhorizontal joints (Xue et al., 2017). Mylonite is localized along the northwest boundary of the Xuelongbao and Pengguan complexes. Along the western boundary of the Xuelongbao complex, foliation dips to the NW at 45–80° with lineation plunging to the west (Figure 2). These structural foliations and lineations are difficult to distinguish from those of the D_{1E} deformation structures. However, sigmoidal feldspar clasts and S/C fabric indicate different kinematics of top-to-the-NW normal faulting (Figures 3a and 9a). The Silurian rock directly overlies the Neoproterozoic Pengguan complex also supports NW-ward normal faulting (Figure 2).

Along the northwest boundary of the Pengguan complex, mylonitic foliation dips to the NW at high angles of ~80° (Figures 3a and 9b) with lineation steeply plunging to the NW, except for a few SE- or N-dipping foliations that probably resulted from tilting of the original foliations by the Cenozoic deformation (Figure 2). Photomicrographs including asymmetric mica fishes (Figures 3a and 9c), sigmoidal feldspar porphyroclasts (Figures 3a and 9d), and S/C fabric of white mica (Figures S2c and e) indicate a top-to-the-NW normal faulting. The deformed rocks experienced syntectonic greenschist-facies metamorphism with a mineral assemblage of quartz + feldspar + biotite + white mica + chlorite (Figures S2e–f).

Microscopic structures from the northwest Pengguan complex show sigma-shaped porphyroclast, quartz ribbon (Figures 3a and 9f), and white mica shear-band fabric (Figures 3a and 9g), indicating a top-to-the-SE shearing related to the D_{1E} deformation. However, all these structures are cut by NW-dipping shear-bands (Figures 3a and 9e). In the western flank of the Wulong complex, about 60 km southwest to the Pengguan complex, a similar top-to-the-NW normal fault with greenschist-facies metamorphism that cuts D_{1E}-related thrust sheet has been previously reported as well (Figure 1b)(Burchfiel et al., 1995; Tian et al., 2016; Airaghi et al., 2018b). Thus, we interpret the top-to-the-NW normal faulting in the northwest Pengguan complex and the Xuelongbao complex as the D₂ event after the D_{1E} top-to-the-SE thrusting in the Southeastern Zone.

4. Geochronological constraints

In order to constrain the timing of the D₁ double-vergent thrusting and the D₂ top-to-the-NW normal faulting, SIMS U–Pb dating on zircons from the synkinematic Mupi pluton and ⁴⁰Ar–³⁹Ar dating on metamorphic white mica, biotite and fine-grained white mica were carried out.

4.1 Analytical methods

4.1.1 Zircon SIMS U-Pb dating

Zircon concentrates were extracted using conventional crushing and separation techniques. After set in epoxy resin mounts, mineral grains were polished to their cores. Cathodoluminescence (CL) imaging was performed to observe their internal structure and find suitable positions for spot analyses. Geochronological data were acquired at the State Key Laboratory of Lithospheric Evolution, Institute of Geology and Geophysics, Chinese Academy of Sciences (IGGCAS).

U–Pb zircon dating was conducted on a Cameca IMS–1280HR SIMS following the analytical procedures described by Li et al. (2009). The primary O^{2−} ion beam was focused to elliptical spots with the sizes of 20 μm×30 μm. Measured Pb/U ratios were calibrated relative to the zircon standard Plešovice (Sláma et al., 2008). Non-radiogenic Pb was subtracted from the measured Pb isotopic composition using the measured ²⁰⁴Pb and the present-day average terrestrial Pb isotopic composition in the model of (Stacey and Kramers, 1975). U–Pb data were plotted and calculated using the program Isoplot 3.00 (Ludwig, 2003), and age uncertainties are quoted at the 95% confidence level except for those noted otherwise. Measurements on a second zircon standard Qinghu were interspersed with unknowns to monitor the external uncertainties, and yielded concordia ages of 159.6 ± 1.6 Ma (MSWD = 0.28, N = 6), which are identical within errors to the recommended age (159.5 ± 0.2 Ma)(Li et al., 2013b).

4.1.2 ⁴⁰Ar–³⁹Ar dating

Micas were separated from eight micaschist, gneiss, and mylonite samples for ⁴⁰Ar/³⁹Ar dating by step-heating technique. Minerals were irradiated and measured in the Ar–Ar Laboratory, School of Earth Sciences of the University of Melbourne. Hand-picked grains were cleaned in de-ionized water and acetone. Samples were weighted and loaded into aluminum packets for irradiation. The packets were placed in a silicate glass tube (Can UM#80), interleaved with packets containing the flux monitor Mount Dromedary Biotite (Age = 99.125 ± 0.076 Ma)(Phillips et al., 2017). The canister was

irradiated for 40 MWhr in the CLICIT facility of the Oregon State University TRIGA reactor, USA. The measured isotopic ratios were corrected for system blanks, mass discrimination, and irradiation-induced interference. Plateau and/or total gas ages with 2σ uncertainties were calculated and calibrated based on methods proposed by (Lee et al., 2006). Analytical data are presented in Table S2 (Xue et al., 2020), and the age spectra are presented in Figure 11. Argon age plateaus are defined by contiguous gas fractions that together represent more than 50% of the total ^{39}Ar released from the sample and for which no difference in age can be detected between any two fractions at the 95% confidence level (Fleck et al., 1997). According to this definition, all argon age spectra in this work, except for sample CX33, do not yield plateau ages (Figure 11). We choose contiguous steps within 95% confidence close to 50% of the total ^{39}Ar released (CX6A white mica), or contiguous steps close to 95% confidence more than 50% of the total ^{39}Ar released (CX2, CX6B, CX25, CX44 white mica), to calculate weighted mean ages (Figure 11).

4.2 Results

4.2.1 U–Pb dating on synkinematic Mupi pluton

Zircons from the Mupi pluton are pink to colorless, 80 to 200 μm long with length/width ratios from 2 to 3. They show well-developed crystal morphologies. CL images of zircons show that most of the zircons have well-developed oscillatory zoning of igneous origin and some grains have an inherited core (Figure 10a). Under guidance of the zircon CL images, analyzed spots were positioned on the oscillatory zircon mantle/margin when there are exist inherited zircon cores. SIMS U–Pb zircon results are listed in Table S1. The SIMS U–Pb dating of the Mupi pluton yields a concordant age of 219.1 ± 1.7 Ma (MSWD = 0.99, N = 15; Figure 10b).

4.2.2 ^{40}Ar – ^{39}Ar dating

Sample CX44 collected from the northern Northwestern Zone (Figure 12) is micaschist with penetrative SSE-dipping foliation and associated S-plunging lineation related to the D_{1W} top-to-the-N thrusting. The CX44 micaschist contains domains rich in fine-grained white mica and quartz that define the shear foliation (Figures S2a–b). Multi-grain white mica (211–152 μm) separated from CX44 yields a weighted mean age of 224.1 ± 1.7 Ma, accounting for 81% of released ^{39}Ar (Figures 11a and Table 1).

Sample CX12 collected from the southern Northwestern Zone (Figure 12) is micaschist involved in the D_{1W} top-to-the-NW thrusting. Shear foliation is defined by aligned fine-grained white mica (Figure 7g). Two aliquots of fine-grained white mica (104–66 µm) of CX12A & CX12B show discordant stair-shaped spectra increasing from ~65–70 Ma to ~150 Ma and ~120 Ma, respectively (Figures 11b–c and Table 1).

Samples CX6 and CX33 (Figure 12), collected from the Inner Metamorphic Core within the Southeastern Zone, are characterized by penetrative NW-dipping foliation and NW-plunging lineation that was involved in D_{1E} top-to-the-SE thrusting. In these two samples, the foliations and associated lineations are defined by white mica (Figures 6d and S2g). Two aliquots of white mica (152–211 µm) of CX6A and CX6B yield equivalent weighted mean ages of 124 ± 1.9 Ma and 126.3 ± 3.3 Ma, respectively (Figure 11d–e). Multi-grain white mica (152–211 µm) from CX33 yields a plateau age of 148.2 ± 1.2 Ma (>60% of the total ³⁹Ar; Figure 11f and Table 1).

Samples CX25 from micaschist of the Xuelongbao complex and CX18, CX19 from gneiss/mylonite of the Pengguan complex (Figures 12) show well-developed NW-dipping foliations related to the D₂ top-to-the-NW normal faulting (Figures S2c–f). White mica (211–152 µm) in CX25 that defines the shear planes (Figures S2e–f) was selected for ⁴⁰Ar/³⁹Ar dating. Five steps of sample CX25 with ~80% released ³⁹Ar yields a weighted mean age of 79.7 ± 2.7 Ma (Figure 11h). Samples of CX18 and CX19 have stair-shaped spectra, in which ages increase from ~50–60 Ma to ~110–135 Ma from low- to high-temperature steps (Figure 11i–k and Table 1).

Sample CX2 from the northwest boundary of the Pengguan complex (in a structural position equivalent to the samples CX18 and CX19; Figure 12) was involved in the D₂ top-to-the-NW normal faulting (Figures 3 and 9g–i). Multi-grain white mica (211–152 µm) from CX2 yields a weighted mean age of 26.23 ± 0.99 Ma at low- to medium-temperature steps (>50% of the total ³⁹Ar released; Figure 11g and Table 1).

4.3 Interpretation of the geochronological results

4.3.1 Zircon U-Pb data interpretation

In the northern Northwestern Zone, the Mupi pluton shows a continuous transition from mylonitic fabrics along the pluton margin to magmatic fabrics in the pluton interior, arguing for its synkinematic emplacement. This deduction is supported by the fact that quartz + feldspar veins parallel to the mylonitic foliation, and feldspar records

587 brittle deformation occurred in the presence of residual melt (see details in Section
588 3.3.3). Thus, the D_{1W} top-to-the-N/NW thrusting is closely related to the emplacement
589 of the Mupi pluton. Our zircon U–Pb ages reveal that the Mupi pluton emplaced at *ca.*
590 219 Ma, therefore place age constraint on the D_{1W} top-to-the-N/NW thrusting at *ca.*
591 219 Ma (Figure 12).

592 4.3.2 ⁴⁰Ar–³⁹Ar data interpretation

593 All samples show high radiogenic ³⁹Ar and almost all steps are clustered at the
594 horizontal ³⁹Ar/⁴⁰Ar axis that does not define a real inverse isochron line (Table S2;
595 Figure S3). Most of the step-ages calculated both by using initial ⁴⁰Ar/³⁶Ar of 298.56
596 (Lee et al., 2006) and from the so-called inverse isochron show nearly no difference
597 (Table S2). Thus, the inverse isochron ages (though identical to their weighted mean
598 ages or plateau ages) and the fake extraneous atmosphere argon are meaningless and
599 will not be further discussed (Figure S3; Table S2).

600 Concordant white mica ⁴⁰Ar–³⁹Ar spectra, such as those yielded by samples of CX44
601 (81.6% released ³⁹Ar), CX6A (49.5% released ³⁹Ar), CX6B (52.4% released ³⁹Ar),
602 CX33 (60.1% released ³⁹Ar), CX2 (53.6% released ³⁹Ar) and CX25 (79.7% released
603 ³⁹Ar), are typical of samples that have remained a closed system since
604 thermal/mylonitic resetting or crystallization (Figures 11a and d–h). Similar rocks to
605 sample CX44 recorded greenschist facies metamorphism under temperatures of 260–
606 310 °C (chlorite + white mica multi-equilibrium thermometer; Liu et al., 2008), which
607 is lower than the white mica argon closure temperature of *ca.* 425 °C (100 μm
608 grain)(Harrison et al., 2009). Considering that the dated mica grains are strongly
609 deformed by the top-to-the-N/NW thrusting, we, therefore, interpret that the 224.1 ±
610 1.7 Ma argon age of CX44 recorded the timing of the white mica recrystallization
611 related to the D_{1W} thrusting event.

612 Samples of CX6A & CX6B, and CX33 from the Inner Metamorphic Core, CX2 and
613 CX25 from the mylonite zone along the northwest boundary of the Pengguan and
614 Xuelongbao complexes have been deformed and metamorphosed at temperatures
615 of >530°C (Raman Spectroscopy on Carbonaceous Material; Robert et al., 2010b).
616 These temperatures are higher than the closure temperature (425 °C) of the *ca.* 100 μm
617 radius muscovite grain (Harrison et al., 2009). There is no evidence of any post late
618 Cretaceous to Cenozoic magmatism along the LMTB and its adjacent eastern Songpan–
619 Ganzi Terrane. Thus, the flattened spectra of CX6A (124 ± 1.9 Ma), CX6B (126.3 ±

3.3), CX33 (148.2 ± 1.2 Ma), CX2 (26.2 ± 0.9 Ma), and CX25 (79.7 ± 2.7 Ma) are probably cooling ages when deformation rapidly exhumed those rocks through the ca. 425 °C geothermal depth. These cooling ages, therefore, can represent the timing of deformations. In this scenario, considering samples of CX6A, CX6B and CX33 were involved in top-to-the-SE thrusting, we interpret the dates of 148 Ma and 124 Ma as ages to SE-ward thrusting and rapid exhumation. Samples of CX2 and CX25 collected from the northwest boundary of the Pengguan and Xuelongbao complexes were involved in top-to-the-NW normal faulting, which may imply that the top-to-the-NW normal faulting occurred at *ca.* 27 Ma along the northwest boundary of the Pengguan complex, and at *ca.* 80 Ma along the northwest boundary of the Xuelongbao complex, respectively.

Discordant stair-shaped spectra, such as those observed from the samples of CX12, CX18, and CX19 (Figures 11b–c, i–k), are commonly observed in fine-grained mineral mixes (Tian et al., 2016). Those staircase spectra indicate ^{40}Ar diffusive loss that is possibly related to chemical alteration, a subsequent thermal event disturbance, or argon loss during the late stage mylonitisation (Costa and Maluski, 1988; Lovera et al., 1989; West and Lux, 1993). The oldest age of the stair-shaped spectra is a minimum one for the fine-grained white mica closure age (McDougall and Harrison, 1988). Samples of CX12, CX18, and CX19 have been deformed at temperatures $>530^\circ\text{C}$ as well (Raman Spectroscopy on Carbonaceous Material; Robert et al., 2010). Considering the size of white mica (104–66 μm), the argon-argon system of fine-grained mica should be completely reset (Harrison et al., 2009). According to the stair-shaped spectra, samples of CX12A, CX12B, CX18, CX19A, CX19B may have recrystallized or been reset before 146.5 ± 2.7 Ma, 119 ± 1.3 Ma, 111.1 ± 0.2 Ma, 133.7 ± 3.3 Ma, and 139.3 ± 1.1 Ma, respectively (Figures 11b–c, i–k). These oldest ages of the stair-shaped spectra possibly resulted from the 148–124 Ma mylonitisation as implied by ^{40}Ar - ^{39}Ar cooling ages of CX6A, CX6B, and CX33.

Chemical alteration should have led to a more profound influence on the fine-grained white mica of CX12, CX18, and CX19 that may result in ^{40}Ar diffusive loss and flattened spectra (Harrison et al., 2009). In this way, considering samples of CX2, CX18, CX19, and CX25 located at the same structural position along the northwest boundary of the Neoproterozoic complexes, we may expect that CX2 and CX25, showing stair-shaped spectrum and CX18, CX19 and presenting flattened spectra, yield

younger ages, however, it is not the case (Figures 11g–k). Thus, chemical alteration and related ^{40}Ar diffusive loss may not be the main factor to account for the stair-shaped spectra.

An alternative interpretation may suggest that the youngest age of the stair-shaped spectra gives a maximum age of the thermal overprint or late-stage mylonitisation (Costa and Maluski, 1988; Lovera et al., 1989; West and Lux, 1993). As stated above, no magmatism of late Mesozoic–Cenozoic ages has been reported along the LMTB. Thus, the post-cooling thermal event may not be responsible for the minimum ages of the stair-shaped spectra. The youngest ages in the stair-shaped spectra therefore most likely indicate late-stage mylonitisation after ~60–50 Ma (Figures 11i–k), which possibly related to the *ca.* 27 Ma top-to-the-NW faulting along the northwest boundary of the Pengguan complex. It is worth to note that the CX12A and CX12B fine-grained white mica from the Northwestern Zone (Figure 12), far away from the Pengguan complex, show older minimum ages of 64.7 ± 1.3 Ma and 72.2 ± 1.4 Ma comparing with the 50 ± 0.1 Ma, 59 ± 0.3 Ma, and 58.4 ± 0.4 Ma of the CX18, CX19A and CX19B (Figures 11b–c, j–l), which may suggest that the CX12 has been less affected by the top-to-the-NW normal faulting comparing to the CX18 and CX19.

5 Discussion

5.1 Summary of our newly obtained structural results

On the basis of detailed structural analyses, a Main Central Boundary and a D_{1W} top-to-the-N/NW thrusting in the Northwestern Zone are argued for the first time in the LMTB. The Main Central Boundary corresponds to a major tectonic boundary supported both by field structures and geophysical profile (Jia et al., 2006; Robert et al., 2010b), which divides the LMTB into a Northwestern Zone (part of the Songpan–Ganzi Terrane) to the northwest and a Southeastern Zone to the southeast. The Southeastern Zone is characterized by exclusively NW-dipping foliations and associated D_{1E} top-to-the-SE thrusting, while the Northwestern Zone is characterized by exclusively SE-dipping foliations and associated D_{1W} top-to-the-NW thrusting.

This study subdivides the Southeastern Zone of the LMTB into four units based on increasing deformation intensities and metamorphic grades from southeast to northwest, namely the Autochthon, Para-autochthon, Allochthon, and Inner Metamorphic Core. This subdivision pattern is revised from the previous one that divided the LMTB into

an allochthonous sheet thrust southeastwards on top of the autochthonous Sichuan Foreland Basin (Xu et al., 1992; Burchfiel et al., 1995; Worley and Wilson, 1996).

In the Southeastern Zone, a D₂ top-to-the-NW normal faulting that cuts the D_{1E} top-to-the-SE thrusting is recognized along the northwest boundary of the Pengguan and Xuelongbao complexes.

5.2 Polyphase deformations along the LMTB

According to the geometric and kinematic features and overprinting relationship, together with ⁴⁰Ar-³⁹Ar and zircon U-Pb data, which allow us to draw a polyphase deformation sequence of the LMTB.

5.2.1 Timing of the D₁ main deformation event

Timing of the D_{1W} top-to-the-N/NW thrusting is firstly constrained by the emplaced timing of the syntectonic Mupi pluton at *ca.* 219 Ma (Figure 12). Synkinematic recrystallized white mica of sample CX44 that is involved in the top-to-the-N/NW thrusting yields a weighted mean age of 224.1 ± 1.7 Ma, which places a second age constraint on the D_{1W} top-to-the-N/NW thrusting. This newly obtained argon age is consistent with previously published white mica ⁴⁰Ar-³⁹Ar ages at *ca.* 237–202 Ma from the northern Northwestern Zone (Figure 12)(Yan et al., 2011; Yan et al., 2018). The fact that the zircon U-Pb age of the Mupi pluton is *ca.* 5 Ma younger than the synkinematic white mica ⁴⁰Ar-³⁹Ar age may suggest the Mupi pluton emplaced during the late phase of the D_{1W} event. Based on the zircon U-Pb age of syntectonic Mupi pluton and synkinematic white mica ⁴⁰Ar-³⁹Ar age, we, therefore, suggest that the D_{1W} top-to-the-N/NW thrusting event occurred at *ca.* 224–219 Ma.

The D_{1E} top-to-the-SE thrusting within the Southeastern Zone has been constrained in the late Triassic for the following arguments: 1) Field observations and seismic reflection profile revealed that late Jurassic–Cenozoic rocks unconformably overly on intensively folded rocks as young as late Triassic (Figures 2 and 4a–d)(BGMRSRSP, 1991; Jia et al., 2006); 2) Detailed petrological analyses and in-situ allanite U-Pb dating and biotite ⁴⁰Ar/³⁹Ar dating revealed a *ca.* 204–190 Ma peak metamorphism in the Inner Metamorphic Core that postdates the D_{1E} event (Airaghi et al., 2017a; 2018a; 2018c); 3) The ages of the voluminous magmatism intruding the extensively folded Songpan–Ganzi Terrane range between 223 and 200 Ma, which postdates the folding in the Songpan–Ganzi Terrane and the top-to-the-SE thrusting of the LMTB (Figure 12)(Xu

et al., 1992; Roger et al., 2004; Harrowfield and Wilson, 2005; Hu et al., 2005; Zhang et al., 2007; de Sigoyer et al., 2014). Thus, the regional D_{1E} top-to-the-SE thrusting was at least activated during the late Triassic, together with the contemporary D_{1W} top-to-the-N/NW thrusting recorded within the Northwestern Zone, both of which compose the *ca.* 224–219 Ma main D₁ double-vergent event.

5.2.2 Timing of the D₂ top-to-the-NW normal faulting

According to the newly obtained ⁴⁰Ar–³⁹Ar ages, the timing of the D₂ top-to-the-NW normal faulting occurred along the northwest boundary of the Pengguan (*ca.* 27 Ma) and Xuelongbao (*ca.* 80 Ma) complexes are quite different. The top-to-the-NW/N normal faulting is widespread in an over 300 km long zone along the northern margins of the Danba and Kangding complexes, and the northwest margins of the Baoxing, Wulong, Xuelongbao, Moutuo, Pengguan, and Jiaoziding complexes (Figure 12)(Burchfiel et al., 1995; Xu et al., 2008; Tian et al., 2016; Xue et al., 2017). This belt of top-to-the-NW normal faulting has previously been entitled East Tibet Detachment (Figure 12)(Wang et al., 2012b). Timing of the East Tibet Detachment top-to-the-NW normal faulting is different at different localities. Hames and Burchfiel (1993) reported a top-to-the-NW normal faulting and syntectonic metamorphism at *ca.* 25 Ma along the northwest Pengguan complex that is in accordance with the *ca.* 27 Ma top-to-the-NW normal faulting as indicated by sample CX2 (Figure 12).

Along the northwest boundary of the Wulong complex and north boundary of the Danba complex, the top-to-the-NW/N normal faulting was constrained to *ca.* 81–47 Ma (⁴⁰Ar/³⁹Ar dating on metamorphic biotite and white mica; Figure 12)(Zhou et al., 2008; Tian et al., 2016). This phase of the top-to-the-NW normal faulting is recorded along the northwest boundary of the Xuelongbao complex (*ca.* 80 Ma ⁴⁰Ar–³⁹Ar age of sample CX25) as well.

A period of the top-to-the-NW/N normal faulting at *ca.* 166–159 Ma was reported at the northwest boundary of the Jiaoziding and Pengguan complexes, and the north boundary of the Danba complex (Figure 12)(Dirks et al., 1994; Zhou et al., 2008; Li, 2009). Whereas, younger ages of *ca.* 122–120 Ma related to the top-to-the-N normal faulting at the north boundary of the Danba complex were reported as well (Figure 12)(Xu et al., 2008).

In summary, the timing of the top-to-the-NW normal faulting of the East Tibet Detachment are scattered at *ca.* 166–159 Ma (north Danba, northwest Pengguan,

northwest Jiaoziding), *ca.* 122–120 Ma (north Danba), *ca.* 81–47 Ma (north Danba, northwest Wulong) and *ca.* 27–25 Ma (northwest Pengguan; Figure 12). Nonetheless, a process is required to explain these four groups of well-defined argon plateau ages that do not show an age-changing trend in map view (Figure 12). We, therefore, prefer to interpret that the top-to-the-NW/N normal faulting of the East Tibet Detachment was episodically and co-planarly activated at *ca.* 166–159 Ma, *ca.* 122–120 Ma, *ca.* 81–47 Ma, and *ca.* 27–25 Ma (Figure 12). The late-stage activations could partially reset the fine-grained white mica argon system that resulted in stair-shaped spectra yielded by samples of CX18 and CX19 (Figures 11 and 12).

5.3 Episodic basement-slice-imbrication and amphibolite facies metamorphic rock exhumation

The Western Sichuan Foreland Basin (WSFB) was formed by loading and subsidence due to the southeastward thrusting of the LMTB along the Yingxiu–Beichuan Fault and Xiaoguanzi Fault (T_2 in Figure 12)(Chen and Wilson, 1996; Li et al., 2003). Episodic terrestrial deposition in the WSFB from the late Triassic to the present day indicates that the LMTB continued to be tectonically active episodically during the late Triassic to early Jurassic, late Jurassic to early Cretaceous, and late Cretaceous to earliest Paleocene (Figure 12)(Xu et al., 2008; Li et al., 2013c; Tian et al., 2016).

The late Jurassic to early Cretaceous southeastward thrusting of the Yingxiu–Beichuan Fault and Xiaoguanzi Fault (T_2 in Figure 12), as the southeast boundary Fault of the Pengguan and Baoxing complexes, overlaps the *ca.* 166–159 Ma and *ca.* 122–120 Ma top-to-the-NW/N normal faulting along the NW boundary of the Jiaoziding complex and the north boundary of the Danba complex (Figure 12)(Xu et al., 2008; Zhou et al., 2008; Li, 2009). The late Cretaceous–earliest Paleocene southeastward thrusting of the Yingxiu–Beichuan Fault and Xiaoguanzi Fault is also coeval with the top-to-the-NW/N normal faulting along the NW boundary of the Xuelongbao (CX25 in this study) and Wulong (Tian et al., 2016) complexes (Figure 12). Furthermore, both the top-to-the-SE thrusting and top-to-the-NW normal faulting along the southeast and northwest boundaries of the complexes occurred under similar greenschist-facies metamorphic conditions (Airaghi et al., 2017b; Xue et al., 2017). Conjugated kinematics of boundary SE-thrusting and NW-normal faulting characterized by similar

metamorphic conditions and synchronous timing let us propose that the Xuelongbao, Pengguan, Baoxing and Wulong complexes were episodically imbricated to the southeast as basement-slices during late Jurassic–early Cretaceous and late Cretaceous–earliest Paleocene.

Numerical modeling of co-seismic slip vectors vs. fault angles indicates that the latest high-angle listric Yingxiu–Beichuan Fault might have facilitated the vertical imbrication of the Pengguan complex with no need of the SE-ward thrusting and loading (Feng et al., 2015). This feature explains the absence of a rejuvenated foreland basin contemporary to the 27–25 Ma basement imbrication. Nevertheless, the 27–25 Ma basement slice imbrication of the Pengguan complex is compatible with its 30–25 Ma fast cooling (Figure 12)(Wang et al., 2012a).

In the map view, two amphibolite facies metamorphic zones wrap the Xuelongbao and Danba complexes (Figure 13a). Both amphibolite facies metamorphic zones reached peak conditions at *ca.* 210–180 Ma post to the late Triassic orogeny (Figure 12)(Dirks et al., 1994; Huang et al., 2003a; Weller et al., 2013; Airaghi et al., 2018a). Gravity anomaly indicated that the Xuelongbao complex probably extends northeastwards beneath and parallel to the amphibolite facies metamorphic rocks (Xue et al., 2017). Coincidentally, the Wenchuan–Xuelongbao and Danba amphibolite facies metamorphic zones experienced rapid exhumation and cooling during the late Jurassic to early Cretaceous (CX12 and CX33 in this work) that is coeval with the aforementioned late Jurassic–early Cretaceous basement-slice-imbrication (Figure 13b)(Arne et al., 1997; Huang et al., 2003a; Tian et al., 2016; Airaghi et al., 2018a). Both gravity inversion (Xue et al., 2017) and seismic reflection profiles (Guo et al., 2013) revealed a belt of basement-slice hidden right beneath a Mesozoic Laojungou granite (224 ± 5 Ma; Zhao et al., 2007) within the east Songpan–Ganzi Terrane (Figure 13a). The Laojungou granite recorded rapid exhumation and cooling during the late Jurassic to early Cretaceous as well (Figure 13b)(Yuan et al., 1991).

Considering the spatial–cooling–imbricating relationships between the Laojungou granite, amphibolite facies metamorphic zones and basement-slices, we interpret that the Laojungou granite and amphibolite facies metamorphic rocks were popped up by a basement-slice-imbrication during the late Jurassic–early Cretaceous. The basement-slice-imbrication was responsible for localized fast cooling and exhumation of the amphibolite-facies metamorphic rocks. However, the Wenchuan–Xuelongbao

metamorphic sedimentary cover does not show fast cooling during the late Cretaceous–earliest Paleocene and at *ca.* 27–25 Ma comparing to the basement-slice-imbrication (Figure 13b), which needs further work.

5.4 Mesozoic tectonic evolution of the LMTB

The Mesozoic tectonic evolution of the LMTB can be summarized as follows without consideration of the top-to-the-N/NW thrusting in the Northwestern Zone. Namely, the Songpan–Ganzi Terrane subducted to the north along the Anyimaqen Suture Zone and/or to the southwest along the Jinshajiang Suture Zone, which resulted in the top-to-the-S shearing of the thick Triassic flysch cover along a basal decollement layer (Mattaueer et al., 1992; Calassou, 1994) and associated sinistral strike-slip to the southeastward thrusting in the LMTB (Xu et al., 1992; Dirks et al., 1994; Burchfiel et al., 1995; Worley and Wilson, 1996; Roger et al., 2004; Harrowfield and Wilson, 2005). The amphibolite facies metamorphism in the Wenchuan–Xuelongbao area reached the peak pressure of $> 11 \pm 2$ kbar at *ca.* 200 Ma (Airaghi et al., 2018b) required a sedimentary cover more than 30 km in thickness. However, high-resolution seismic reflection and gravity modeling profiles revealed a ~20 km thickness cover of the Songpan–Ganzi Terrane (Guo et al., 2013; Xue et al., 2017) that is far thinner than the required ~30 km thickness.

In contrast, Sengör (1984) and Luo (1991) regarded the LMTB as an early Mesozoic intracontinental subduction zone but lack details of structural observation. Considering the pervasive top-to-the-N/NW thrusting in the Northwestern Zone and top-to-the-SE thrusting in the Southeastern Zone, we thus prefer an early Mesozoic intracontinental subduction along the LMTB.

In this work, the Main Central Boundary, devoid of ophiolitic rocks separating the Southeastern Zone to the SE from the Northwestern Zone to the NW, does not represent an ophiolitic suture zone, but a tectonic “scar” corresponding to the place where the Yangtze basement was underthrust. Like the Chenzhou–Linwu Fault in the Xuefengshan Intracontinental Orogen (South China) where the South China Block underthrust (Chu et al., 2012a; Chu et al., 2012b). Along the LMTB, the underthrusting was coeval with important burial ($> 11 \pm 2$ kbar) and the double-vergent thrusting in both sides of the Main Central Boundary. The basement underthrusting is also supported by the blueschist with intraplate affinity reported along the Qingchuan–

Pingwu Fault (Wei, 1994; Xu et al., 2020). A new tectonic scenario is proposed here (Figure 14). In the pre-middle Triassic times, the Longmenshan area along the western Yangtze margin was a failed rift evidenced by radiolarian chert, abyssal deposits, slump structures (Long, 1991; Luo, 1991; Chang, 2000; Li, 2009) and seismic profile (Jia et al., 2006). Accompany with the pre-middle Triassic rifting, a series of NW-dipping syn-depositional normal faults developed in the Longmenshan area that dissected the western Yangtze basement to slices, and the lower crust thinned significantly that resulted in a local “weak zone” in the lower crust (Figure 14a).

During the late Triassic, the Yangtze basement underthrusting along the “weak zone” accommodated by inversion of the normal faults, which can be linked to the far-field effect of the Paleotethys’ closure (Figure 14b)(Luo, 1991; Chen and Wilson, 1996; Meng et al., 2005; Yan et al., 2019). The sedimentary cover was simultaneously shortened by a top-to-the-SE thrusting and a top-to-the-NW thrusting associated with the main D₁ event (Figure 14b). The domain of the top-to-the-SE thrusting in the Southeastern Zone is larger and more extensive than the top-to-the-NW thrusting in the Northwestern Zone. Similar structures were developed in the north Himalaya where a large domain experienced a top-to-the-S thrusting, compared to a small domain with a top-to-the-N thrusting due to the northward subduction of the India plate (Dewey, 1988; Harrison, 1992). We thus favor an NW-ward basement underthrusting along the LMTB (Figure 14b). The large amount of rift rocks transported southeastwards and overlain the autochthonous foreland basin. In addition, a large volume of rocks was underthrust to crustal depth deeper than 30 km and metamorphosed, as documented by the *ca.* 210–180 Ma amphibolite-facies metamorphic zone that metamorphosed at a peak pressure of $> 11 \pm 2$ kbar (Figure 12)(Weller et al., 2013; Airaghi et al., 2018a; Airaghi et al., 2018b; Airaghi et al., 2018c).

During periods of the late Jurassic–early Cretaceous, late Cretaceous–earliest Paleogene, the accretion of Gondwana-derived Terranes to the southern Eurasia plate had resulted in episodic compression in the Longmenshan area (Dewey, 1988; Xu et al., 1992). We cannot rule out the possible contribution of the 150–140 Ma and 120–105 Ma compression resulted from the westward subduction of the Paleo-Pacific plate subduction (Chu et al., 2019). Under the regional episodic compression, basement-slices represented by the Neoproterozoic complexes along the weak zone of the Yangtze Block, including the Xuelongbao, Moutuo, and Danba complexes,

episodically imbricated southeastwards along the inherited faults. Basement-slice-imbrication then exhumed the basement complexes and the amphibolite facies metamorphic rocks as documented by the synchronously fast cooling of the meta-sediments and the basement complexes (Figures 12 and 14d)(Yuan et al., 1991; Arne et al., 1997; Huang et al., 2003b; Weller et al., 2013; Tian et al., 2016). Basement-slice-imbrication also reworked the late Triassic allochthonous–autochthonous LMTB into four subdivisions (Figure 14d).

7 Conclusions

Based on structural analysis and previous geophysical profile, we have delineated a Main Central Boundary that separates the Longmenshan Thrust Belt into a Southeastern Zone to the southeast and a Northwestern Zone to the northwest.

On the basis of different lithologies, increasing strain intensity and metamorphic grade, the Southeastern Zone is revised into four subdivisions, namely, the Autochthon, Para-autochthon, Allochthon, and Inner Metamorphic Core.

The detailed structural analysis allows us recognizing the multiphase deformation of the LMTB. A main D_1 double-vergent thrusting event, characterized by the D_{1E} top-to-the-SE thrusting in the Southeastern Zone and the D_{1W} top-to-the-NW/N thrusting in the Northwestern Zone, has been recognized. The D_{1E} and D_{1W} events were synchronously occurred corresponding to a late Triassic intracontinental underthrusting along a “weak zone” of the LMTB. A D_2 top-to-the-NW normal faulting that cuts the D_{1E} top-to-the-SE thrusting has been recognized along the NW boundary of the basement complexes in the Southeastern Zone.

Geochronological constraints obtained from this study and the previously published ones indicate that the main D_1 event occurred at *ca.* 224–219 Ma and the D_2 was episodically activated at *ca.* 166–160 Ma, 141–120 Ma, 81–47 Ma, and 27–25 Ma. The episodically and synchronously activated top-to-the-NW normal faulting and top-to-the-SE thrusting along the northwest and SE boundaries of the basement complexes let us propose that the basement-slices episodically imbricated to the SE during the late Jurassic–early Cretaceous and late Cretaceous–earliest Paleocene. The basement-slice imbrication exhumed the D_1 -related amphibolite facies rocks to the subaerial and reworked allochthonous–autochthonous LMTB into a four subdivision one.

Acknowledgement

This work is supported by the National Nature Science Foundation of China (91855212, 91755205 and 41802219) and the National Key R&D Program of China (Grant Numbers 2016YFC0600401, 2016YFC0600102) as well as French LABEX VOLTAIRE (ANR-10-LABX-100-01), and EQUIPEX PLANET (ANR-11-EQPX-0036). Two anonymous reviewers are sincerely thanked for their elaborative reviews and pertinent comments. The geochronological data including SIMS U–Pb dating on zircons (Table S1) and $^{40}\text{Ar}/^{39}\text{Ar}$ dating on micas (Table S2) for this paper are available in the supplementary materials.

Figure captions

Figure 1. (a) Simplified tectonic sketch of China. QLDB: Qingling–Dabie Orogenic Belt; EKL–ANMQS: East Kunlun–Anyimaqen Suture Zone; JSZ: Jinshajiang Suture Zone; SGT: Songpan–Ganzi Terrane; LMTB: Longmenshan Thrust Belt; ICB: Indochina Block; (b) Simplified geological map of the Longmenshan Thrust Belt, modified after BGMRS (1991), Burchfiel et al. (1995) and Xu et al. (2008). T₁: Anxian–Guanxian Fault; T₂: Yingxiu–Beichuan Fault; T₃: Wenchuan–Maoxian Fault; T₄: Qingchuan–Pingwu Fault; JZD: Jiaoziding complex; PG: Pengguan complex; XLB: Xuelongbao complex; BX: Baoxing complex; KD: Kangding complex; DB: Danba complex; MT: Moutuo complex; WL: Wulong complex; RLG: Rilonguan granite; LJG: Laojungou granite.

Figure 2. Geological map of the north and middle parts of the LMTB and its adjacent area, modified after BGMRS (1991) and Xu et al. (2008). MP: Mupi pluton. Symbols and acronyms are the same as in Figure 1.

Figure 3. (a) Structural division, field elements including foliation, lineation, and kinematics of the LMTB. (b–d) π -circle projection of the bedding (S_0) in the Para-autochthon from south to north. All field elements are shown in a lower hemisphere equal-area projection. Symbols and acronyms are the same as in Figure 1.

Figure 4. (a–d) Geological cross-sections perpendicular to the LMTB. Purple solid lines represent the foliations attitude (Locations are marked in Figure 2). Subaerial structures of these cross-sections are based on our field mapping and the deep structures are based on seismic reflection data of Jia et al. (2006) and Guo et al. (2013). The black dashed lines represent speculated folds. (e) Flower structure revealed

by the seismic reflection profile processed by Jia et al. (2006)(Location is marked in Figure 4d). MCB: Main Central Boundary; AGF: Anxian–Guanxian Fault; YBF: Yingxiu–Beichuan Fault; WMF: Wenchuan–Maoxian Fault; QCF: Qingchuan–Pingwu Fault. Other symbols and acronyms are the same as in Figure 1

Figure 5. Rock characteristics and deformation style of the Main Central Boundary. (a) Intensely deformed argillaceous limestone characterized by a vertical foliation [31°34.833' N, 103°24.041' E]. (b) Graphite-bearing pelite was intensely foliated to shale with vertical attitude [31°35.916' N, 103°24.008' E]. (c) Foliated thin-bedded limestone and quartz ribbon interlayer and rootless intra-folial folds coexist with vertical foliation [31°48.403' N, 103°44.080' E]. Rootless folds in figures b–c implying significant shortening. Locations are presented in Figure 3a.

Figure 6. Structural features related to the Main Deformation (D_{IE}) in the Southeastern Zone. **In the Inner Metamorphic Core:** (a) Lineation formed by aligned biotite plunging N335° by 30° on the S-dipping foliation [31°30.359' N, 103°34.123' E]. (b) Asymmetric pyrite pressure shadow indicating a top-to-the-SE thrusting [31°42.210' N, 103°51.067' E]. (c) Sigma-shaped feldspar porphyroclast in paragneiss indicating a top-to-the-SE thrusting [31°05.193' N, 103°18.730' E]. (d) S/C fabric observed under Cross Polarized Light (CPL) indicating a top-to-the-SE thrusting. **In the Allochthon:** (e) Sigmoid pelitic lens and “cold” S/C structures developed in the Neoproterozoic Huangshuihe Group suggesting a top-to-the-SE thrusting [31°16.195' N, 103°46.747' E]. (f) Sigma-shaped sandy lens in Silurian pelitic schist within the nappe indicating a top-to-the-SE thrusting [32°04.577' N, 104°40.599' E]. (g) At the rear of the klippe, thick-bedded limestone thrust to the SE along an NW-dipping reverse fault [31°11.530' N, 103°45.929' E]. **In the Para-autochthon:** (h) Sigmoid lens and “cold” S/C fabrics indicating a top-to-the-SE thrusting [30°55.403' N, 103°20.495' E]. Figure locations are marked in Figure 3a. X and Z axes are marked on the figures. Mineral abbreviations are after Kretz (1983).

Figure 7. Structural features related to the Main Deformation (D_{IW}) in the Northwestern Zone. **In the northern Northwestern Zone:** (a) Aligned calcite, fine-grained white mica, and oxide-pods define lineation plunging to the south on the S-dipping foliation [32°30.892' N, 104°08.178' E]. (b) Sigma-shaped pebbles in a meta-conglomerate indicating a top-to-the-N sense of shear [32°53.562' N, 105°04.820' E]. (c) Deflected quartz ribbon observed under CPL indicating a top-to-the-N thrusting. (d) Asymmetric feldspar porphyroclast wrapped by deflected white mica suggesting a top-to-the-N thrusting. (e) Northward verging chevron fold associated with meter-scale reverse fault indicating a top-to-the-N

thrusting [32°25.219' N, 104°31.513' E]. (f) Bedding depicted by compositional banding showing an N-verging fold, and bedding to sub-parallel to the foliation in the limbs [32°228.591' N, 104°26.763' E]. **In the southern Northwestern Zone:** (g) Asymmetric garnet pseudomorph pressure shadow suggesting a top-to-the-NW thrusting. (h) Hundred-meter scale chevron fold verging to the NW indicating a top-to-the-NW thrusting [31°33.912' N, 103°16.680' E]. Figure locations are marked in Figure 3a. X and Z axes are marked on the figures. Mineral abbreviations are after Kretz (1983).

Figure 8. Features of the syn-tectonic Mupi pluton in the Northwestern Zone [32°31.445' N, 104°32.456' E]. (a) Magmatic fabric of the Mupi pluton with euhedral feldspar randomly distributed, about 30 meters north to the south margin of the Mupi pluton. (b) Quartz + feldspar veins parallel to the shear foliation, about 10-15 meters north to the pluton south margin. (c) Strong mylonitisation at the south margin of the pluton shares similar structural geometry and kinematics with the country rock. (d) Asymmetric feldspar porphyroclasts, synthetic offset of feldspar fragments along shear band in XZ plane indicating a top-to-the-N thrusting. (e) Panorama of thin section under CPL of the Mupi pluton, the feldspar brittle cracks (marked by red arrow) infilled by quartz subparallel to the shear foliation. X and Z axes are marked on the figures. Figure locations are marked in Figure 3a. Mineral abbreviations are after Kretz (1983).

Figure 9. Structural features related to the top-to-the-NW normal faulting along the northwest boundary of the Neoproterozoic complexes. (a) At the northwest boundary of the **Xuelongbao complex**, sigmoid feldspar phenocrysts associated with S/C fabric indicating a top-to-the-NW normal faulting [31°19.683' N, 103°23.874' E]. At the northwest boundary of the **Pengguan complex**: (b) Granitic mylonite having foliation dips to the NW [31°22.931' N, 103°30.364' E]. (c–d) Mica fish in the quartz schist (PPL) and σ -type porphyroclast (CPL) wrapped by deflected micas indicating a top-to-the-NW normal faulting. (e) Shear-band (white dashed line) distributes from the upper-right corner to the lower-left corner that cuts the whole thin section indicate a top-to-the-NW normal faulting, CPL. Sigmoid porphyroclast (e), incipient shear band inclining to the SE (f), and white mica shear-band fabric (g) indicate a top-to-the-SE thrusting related to D_{1E}. All of them are cut by the white shear band related to the D₂ top-to-the-NW normal faulting, CPL. Locations are marked in Figure 3a. X and Z axes are marked on the figures. Mineral abbreviations are after Kretz (1983).

Figure 10. (a) CL images of the representative zircons from the Mupi pluton. The scale bar represents 100 μ m. (b) SIMS U–Pb Concordia diagram of zircons of the syntectonic Mupi pluton.

Figure 11. ^{40}Ar - ^{39}Ar age spectra of the dated samples. We carried out a two-aliquot analysis of micas from CX6, CX12, and CX19 to exclude the influence of occasional deviation on the stair-case age spectrum, and the two-aliquot results are labeled with the suffix of A and B (eg., CX6A, CX6B, CX12A, CX12B, CX19A, CX19B). Inverse isochron diagrams are presented in Figure S3. ^{40}Ar - ^{39}Ar calculation results are summarized in Table 1, and detailed step-heating results are presented in the supporting information Table S2.

Figure 12. Geochronological constraints compiled from published and this study. Ages are shown by the method used to determine the age (A = Ar/Ar; K = K/Ar; U = U/Pb; F = Fission track) and mineral used (Am = Amphibole; B = biotite; M = Muscovite; Z = zircon; Mn = monazite; I = illite; S = Sericite; Wm = white mica). 1. Li (2009); 2. Yan et al. (2018); 3. Yan et al. (2011); 4. Arne et al. (1997); 5. Airaghi et al. (2017b); 6. Airaghi et al. (2018a); 7. Airaghi et al. (2018b); 8. Airaghi et al. (2018c); 9. Zheng et al. (2016); 10. Yan et al. (2008); 11. Zheng et al. (2014); 12. Tian et al. (2016); 13. Huang et al. (2003a); 14. Weller et al. (2013); 15. Xu et al. (2008); 16. Zhou et al. (2008); 17. Wallis et al. (2003); 18. Zhou et al. (2002); 19. Xia (1993); 20. Xu et al. (1992); 21. Yuan et al. (1991); 22. Hu et al. (2005); 23. Zhao (2007); 24. Kirby et al. (2002); 25. Roger et al. (2004); 26. Zhang et al. (2006a); 27. Yuan et al. (2010); 28. BGMRS (1991); 29. Li et al. (2013c); 30. Wang et al. (2012a). 31. Hames and Burchfiel (1993). Symbols and acronyms are the same as in Figure 2.

Figure 13. (a) Spatial relationship between the amphibolite facies metamorphic zones and the basement complexes along the peripheries of the Songpan–Ganzi Terrane. Isograd lines are presented in color dashed lines accompany relating mineral abbreviations. Mineral abbreviations are after Kretz (1983) except Liq representing the magmatic zone. Symbols and acronyms are the same as in Figure 12. (b) Compiled cooling histories for the Wenchuan–Xuelongbao amphibolite facies metamorphic belt (WXAB, bold dashed line)(Airaghi et al., 2018a), the Danba amphibolite facies metamorphic belt (DBAB), and migmatite zone (DBM, thin dashed line)(Huang et al., 2003a), and the Mesozoic Laojougou granite (LJG, bold gray line)(Yuan et al., 1991). Gray frames show the three episodes of basement-slice-imbrication at late Jurassic–early Cretaceous, late Cretaceous–Earliest Paleocene, ca. 27–25 Ma. The Wenchuan–Xuelongbao and the Danba amphibolite facies metamorphic belt and the Laojougou granite experienced a rapid exhumation and cooling during the late Jurassic to the late Cretaceous, and 27–25 Ma.

Figure 14. A possible Mesozoic tectonic evolutionary scenario of the LMTB and adjacent area. (a) Pre-middle Triassic rifting of the Yangtze Block resulted in crustal thinning and a potential “weak zone” in the future Longmenshan area. (b) During the late Triassic, NW-ward Yangtze basement underthrusting along the “weak zone” resulted in extensive shortening, inversion of syn-depositional normal Fault. The sedimentary rocks were buried to a crustal depth deeper than 30 km. NW- and SE-ward thrusting, and associated nappe belt formed at this time. (c) During the late Triassic to early Jurassic, the underthrustured rocks were metamorphosed. Gradient color from green to dark blue in the Inner Metamorphic Core represents show the low- to high-metamorphic grade. (d) late Jurassic–Miocene, basement-slice episodically imbricated to the SE that exhumed the amphibolite facies metamorphic rocks to the subaerial and reworked the LMTB. Symbols and acronyms are the same as in Figure 2.

Table 1. Summary of ^{40}Ar - ^{39}Ar dating results

References

- Airaghi, L., Janots, E., Lanari, P., de Sigoyer, J., and Magnin, V., 2018a, Allanite petrochronology in fresh and retrogressed garnet-biotite metapelites from the Longmen Shan (eastern Tibet): *Journal of Petrology*, p. egy109-egy109, <https://doi.org/10.1093/petrology/egy109>.
- Airaghi, L., Lanari, P., de Sigoyer, J., and Guillot, S., 2017a, Microstructural vs compositional preservation and pseudomorphic replacement of muscovite in deformed metapelites from the Longmen Shan (Sichuan, China): *Lithos*, v. 282-283, p. 262-280, <https://doi.org/10.1016/j.lithos.2017.03.013>.
- Airaghi, L., Sigoyer, J., Guillot, S., Robert, A., Warren, C. J., and Deldicque, D., 2018b, The Mesozoic along-strike tectono-metamorphic segmentation of Longmen Shan (eastern Tibetan plateau): *Tectonics*, v. 37, p. 4655-4678, <https://doi.org/10.1029/2018TC005005>.
- Airaghi, L., Sigoyer, J. d., Lanari, P., Guillot, S., Vidal, O., Monié, P., Sautter, B., and Tan, X., 2017b, Total exhumation across the Beichuan fault in the Longmen Shan (eastern Tibetan plateau, China): constraints from petrology and thermobarometry: *Journal of Asian Earth Sciences*, <https://doi.org/10.1016/j.jseaes.2017.04.003>.
- Airaghi, L., Warren, C. J., de Sigoyer, J., Lanari, P., and Magnin, V., 2018c, Influence of dissolution/reprecipitation reactions on metamorphic greenschist to

1082 amphibolite facies mica $^{40}\text{Ar}/^{39}\text{Ar}$ ages in the Longmen Shan (eastern Tibet):
 1083 Journal of Metamorphic Geology, v. 36, no. 7, p. 933-958,
 1084 <https://doi.org/10.1111/jmg.12420>.
 1085 Arne, D., Worley, B., Wilson, C., Chen, S. F., Foster, D., Luo, Z. L., Liu, S. G., and
 1086 Dirks, P., 1997, Differential exhumation in response to episodic thrusting along
 1087 the eastern margin of the Tibetan Plateau: tectonophysics, v. 280, no. 3, p. 239-
 1088 256, [https://doi.org/10.1016/S0040-1951\(97\)00040-1](https://doi.org/10.1016/S0040-1951(97)00040-1).
 1089 BGMRSP, 1991, Regional geology of Sichuan Province, Beijing, Geological
 1090 Publishing House, 730 p.:
 1091 Billerot, A., Duchene, S., Vanderhaeghe, O., and de Sigoyer, J., 2017, Gneiss domes of
 1092 the Danba Metamorphic Complex, Songpan Ganze, eastern Tibet: Journal of
 1093 Asian Earth Sciences, v. 140, p. 48-74,
 1094 <https://doi.org/10.1016/j.jseaes.2017.03.006>.
 1095 Burchfiel, B. C., Chen, Z., Liu, Y., and Royden, L. H., 1995, Tectonics of the Longmen
 1096 Shan and Adjacent Regions, Central China: International Geology Review, v.
 1097 37, no. 8, p. 661-735, <https://doi.org/10.1080/00206819509465424>.
 1098 Burchfiel, B. C., Royden, L. H., van der Hilst, R. D., Hager, B. H., Chen, Z., King, R.
 1099 W., Li, C., Lü, J., Yao, H., and Kirby, E., 2008, A geological and geophysical
 1100 context for the Wenchuan earthquake of 12 May 2008, Sichuan, People's
 1101 Republic of China: GSA Today, v. 18, no. 7, p. 4-11,
 1102 <https://doi.org/10.1130/gsatg18a.1>.
 1103 Calassou, S., 1994, Etude tectonique d'une chaîne de décollement. A- Tectonique
 1104 triasique et tertiaire de la chaîne de Songpan Garzê(Est Tibet). B- Géométrie et
 1105 cinématique des déformations dans les prismes d'accrétion sédimentaire:
 1106 Modélisation analogique.
 1107 Chang, E. Z., 2000, Geology and Tectonics of the Songpan-Ganzi Fold Belt,
 1108 Southwestern China: International Geology Review, v. 42, no. 9, p. 813-831,
 1109 <https://doi.org/10.1080/00206810009465113>.
 1110 Chen, S., and Wilson, C. J. K., 1996, Emplacement of the Longmen Shan Thrust-Nappe
 1111 Belt along the eastern margin of the Tibetan Plateau: Journal of Structural
 1112 Geology, v. 18, no. 4, p. 413-429. (In Chinese with english abstract)
 1113 Chen, S., Wilson, C. J. L., Luo, Z., and Deng, Q., 1994, The evolution of the Western
 1114 Sichuan Foreland Basin, southwestern China: Journal of Asian Earth Sciences,

1115 v. 10, no. 94, p. 159–168, [https://doi.org/10.1016/0743-9547\(94\)90016-7](https://doi.org/10.1016/0743-9547(94)90016-7).

1116 Chen, S., Wilson, C. J. L., and Worley, B. A., 1995, Tectonic transition from the
 1117 Songpan-Garzê Fold Belt to the Sichuan Basin, south-western China: Basin
 1118 Research, v. 7, no. 3, p. 235-253, [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2117.1995.tb00108.x)
 1119 [2117.1995.tb00108.x](https://doi.org/10.1111/j.1365-2117.1995.tb00108.x).

1120 Chu, Y., Faure, M., Lin, W., and Wang, Q., 2012a, Early Mesozoic tectonics of the
 1121 South China block: Insights from the Xuefengshan intracontinental orogen:
 1122 Journal of Asian Earth Sciences, v. 61, p. 199-220,
 1123 <https://doi.org/10.1016/j.jseaes.2012.09.029>.

1124 Chu, Y., Faure, M., Lin, W., Wang, Q., and Ji, W., 2012b, Tectonics of the Middle
 1125 Triassic intracontinental Xuefengshan Belt, South China: new insights from
 1126 structural and chronological constraints on the basal décollement zone:
 1127 International Journal of Earth Sciences, v. 101, no. 8, p. 2125-2150,
 1128 <https://doi.org/10.1007/s00531-012-0780-5>.

1129 Chu, Y., Lin, W., Faure, M., Xue, Z., Ji, W., and Feng, Z., 2019, Cretaceous Episodic
 1130 Extension in the South China Block, East Asia: Evidence From the
 1131 Yuechengling Massif of Central South China: Tectonics, v. 38, no. 10, p. 3675-
 1132 3702, <https://doi.org/10.1029/2019tc005516>.

1133 Clark, M. K., and Royden, L. H., 2000, Topographic ooze: Building the eastern margin
 1134 of Tibet by lower crustal flow: Geology, v. 28, no. 8, p. 703-706,
 1135 <https://doi.org/10.1130/0091-7613>.

1136 Costa, and Maluski, 1988, Use of the ⁴⁰Ar-³⁹Ar stepwise heating method for dating
 1137 mylonite zones: An example from the St. Barth616my Massif (North Pyrenees,
 1138 France). Chemical Geology, v. 72, no. 2, p. 127-144,
 1139 [https://doi.org/10.1016/0168-9622\(88\)90061-9](https://doi.org/10.1016/0168-9622(88)90061-9).

1140 de Sigoyer, J., Vanderhaeghe, O., Duchêne, S., and Billerot, A., 2014, Generation and
 1141 emplacement of Triassic granitoids within the Songpan Ganze accretionary-
 1142 orogenic wedge in a context of slab retreat accommodated by tear faulting,
 1143 Eastern Tibetan plateau, China: Journal of Asian Earth Sciences, v. 88, p. 192-
 1144 216, <https://doi.org/10.1016/j.jseaes.2014.01.010>.

1145 Dewey, J. F., 1988, The Tectonic Evolution of the Tibetan Plateau: Philosophical
 1146 Transactions of the Royal Society of London A Mathematical Physical &
 1147 Engineering Sciences, v. 327, no. 1594, p. 379-413,

1148 <https://doi.org/10.1098/rsta.1988.0135>.

1149 Dirks, P., Wilson, C., Chen, S., Luo, Z., and Liu, S., 1994, Tectonic evolution of the NE
 1150 margin of the Tibetan Plateau; evidence from the central Longmen Mountains,
 1151 Sichuan Province, China: *Journal of Southeast Asian Earth Sciences*, v. 9, no. 1,
 1152 p. 181-192, [https://doi.org/10.1016/0743-9547\(94\)90074-4](https://doi.org/10.1016/0743-9547(94)90074-4).

1153 Druschke, P. A., and Wang, T., 2006, Stratigraphic and UPb SHRIMP Detrital Zircon
 1154 Evidence for a Neoproterozoic Continental Arc, Central China: Rodinia
 1155 Implications: *Journal of Geology*, v. 114, no. 5, p. 627-636.

1156 Enkelmann, E., Weislogel, A., Ratschbacher, L., Eide, E., Renno, A., and Wooden, J.,
 1157 2007, How was the Triassic Songpan-Ganzi basin filled? A provenance study:
 1158 *Tectonics*, v. 26, no. 4, <https://doi.org/10.1029/2006TC002078>.

1159 Feng, S., Zhang, P., Liu, B., Wang, M., Zhu, S., Ran, Y., Wang, W., Zhang, Z., Zheng,
 1160 W., and Zheng, D., 2015, Deep crustal deformation of the Longmen Shan,
 1161 eastern margin of the Tibetan Plateau, from seismic reflection survey and finite
 1162 element modelling: *Journal of Geophysical Research: Solid Earth*, v. 121, p.
 1163 767-787, <https://doi.org/10.1002/2015JB012352>.

1164 Fleck, R. J., Sutter, J. F., and Elliot, D. H., 1997, Interpretation of discordant $^{40}\text{Ar}/^{39}\text{Ar}$
 1165 age-spectra of mesozoic tholeiites from antarctica: *Geochimica Et*
 1166 *Cosmochimica Acta*, v. 41, no. 1, p. 15-32, [https://doi.org/10.1016/0016-](https://doi.org/10.1016/0016-7037(77)90184-3)
 1167 [7037\(77\)90184-3](https://doi.org/10.1016/0016-7037(77)90184-3).

1168 Guo, X., Gao, R., Randy Keller, G., Xu, X., Wang, H., and Li, W., 2013, Imaging the
 1169 crustal structure beneath the eastern Tibetan Plateau and implications for the
 1170 uplift of the Longmen Shan range: *Earth and Planetary Science Letters*, v. 379,
 1171 p. 72-80, <https://doi.org/10.1016/j.epsl.2013.08.005>.

1172 Hames, W. E., and Burchfiel, B. C., 1993, Laser $^{40}\text{Ar}/^{39}\text{Ar}$ dating of Cenozoic,
 1173 greenschist-facies shear zones, Longmen Shan, China: *Geological Society of*
 1174 *America*, v. 25, no. 6, p. A-118, [https://doi.org/10.1016/0012-821X\(94\)00079-](https://doi.org/10.1016/0012-821X(94)00079-4)
 1175 [4](https://doi.org/10.1016/0012-821X(94)00079-4).

1176 Harrison, T. M., 1992, Raising Tibet: *Science*, v. 255, no. 5052, p. 1663-1670,
 1177 <https://doi.org/doi:10.1126/science.55.5052.1663>.

1178 Harrison, T. M., Clerier, J., Aikman, A. B., Hermann, J., and Heizler, M. T., 2009,
 1179 Diffusion of ^{40}Ar in muscovite: *Geochimica Et Cosmochimica Acta*, v. 73, no.
 1180 4, p. 1039-1051, <https://doi.org/https://doi.org/10.1016/j.gca.2008.09.038>.

1181 Harrowfield, M. J., and Wilson, C. J. L., 2005, Indosinian deformation of the Songpan
1182 Garze Fold Belt, northeast Tibetan Plateau: *Journal of Structural Geology*, v. 27,
1183 no. 1, p. 101-117, <https://doi.org/10.1016/j.jsg.2004.06.010>.

1184 Hu, J., Meng, Q., Shi, Y., and Qu, H., 2005, SHRIMP U-Pb dating of zircons from
1185 granitoid bodies in the Songpan-Ganzi terrane and its implications: *Acta*
1186 *Petrologica Sinica*, v. 21, no. 3, p. 867-880. (In Chinese with english abstract)

1187 Huang, M., Maas, R., Buick, I. S., and Williams, I. S., 2003a, Crustal response to
1188 continental collisions between the Tibet, Indian, South China and North China
1189 Blocks: geochronological constraints from the Songpan-Garzê Orogenic Belt,
1190 western China: *Journal of Metamorphic Geology*, v. 21, no. 3, p. 223-240,
1191 <https://doi.org/10.1046/j.1525-1314.2003.00438.x>.

1192 Huang, M. H., Buick, I. S., Hou, L. W., Huang, M. H., and Hou, L. W., 2003b,
1193 Tectonometamorphic evolution of the eastern Tibet plateau: Evidence from the
1194 central Songpan-Garze orogenic belt, western China: *Journal of Petrology*, v.
1195 44, no. 2, p. 255-278, <https://doi.org/10.1093/petrology/44.2.255>.

1196 Hubbard, J., and Shaw, J. H., 2009, Uplift of the Longmen Shan and Tibetan plateau,
1197 and the 2008 Wenchuan (M = 7.9) earthquake: *Nature*, v. 458, no. 7235, p. 194-
1198 197, <https://doi.org/10.1038/nature07837>.

1199 Jia, D., Wei, G., Chen, Z., Li, B., Zeng, Q., and Yang, G., 2006, Longmen Shan fold-
1200 thrust belt and its relation to the western Sichuan Basin in central China: New
1201 insights from hydrocarbon exploration: *AAPG bulletin*, v. 90, no. 9, p. 1425-
1202 1447, <https://doi.org/10.1306/03230605076>.

1203 Kirby, E., Reiners, P. W., Krol, M. A., Whipple, K. X., Hodges, K. V., Farley, K. A.,
1204 Tang, W., and Chen, Z., 2002, Late Cenozoic evolution of the eastern margin of
1205 the Tibetan Plateau: Inferences from $^{40}\text{Ar}/^{39}\text{Ar}$ and (U - Th)/He
1206 thermochronology: *Tectonics*, v. 21, no. 1, p. 1-1-20,
1207 <https://doi.org/10.1029/2000TC001246>.

1208 Kretz, R., 1983, Symbols for rock-forming minerals: *Amer Miner*, v. 68, no. 1, p. 277-
1209 279, [https://doi.org/10.1016/0040-1951\(84\)90122-7](https://doi.org/10.1016/0040-1951(84)90122-7).

1210 Lee, J.-Y., Marti, K., Severinghaus, J. P., Kawamura, K., Yoo, H.-S., Lee, J. B., and
1211 Kim, J. S., 2006, A redetermination of the isotopic abundances of atmospheric
1212 Ar: *Geochimica et Cosmochimica Acta*, v. 70, no. 17, p. 4507-4512,
1213 <https://doi.org/10.1016/j.gca.2006.06.1563>.

- 1214 Li, H., Wang, H., Xu, Z., Si, J., Pei, J., Li, T., Huang, Y., Song, S.-R., Kuo, L.-W., Sun,
1215 Z., Chevalier, M.-L., and Liu, D., 2013a, Characteristics of the fault-related
1216 rocks, fault zones and the principal slip zone in the Wenchuan Earthquake Fault
1217 Scientific Drilling Project Hole-1 (WFSD-1): Tectonophysics, v. 584, p. 23-42,
1218 <https://doi.org/10.1016/j.tecto.2012.08.021>.
- 1219 Li, H., Wang, H., Yang, G., Xu, Z., Li, T., Si, J., Sun, Z., Huang, Y., Chevalier, M.-L.,
1220 Zhang, W., and Zhang, J., 2015, Lithological and structural characterization of
1221 the Longmen Shan fault belt from the 3rd hole of the Wenchuan Earthquake
1222 Fault Scientific Drilling project (WFSD-3): International Journal of Earth
1223 Sciences, <https://doi.org/10.1007/s00531-015-1285-9>.
- 1224 Li, X. H., Liu, Y., Li, Q. L., Guo, C. H., and Chamberlain, K. R., 2009, Precise
1225 determination of Phanerozoic zircon Pb/Pb age by multicollector SIMS without
1226 external standardization: Geochemistry, Geophysics, Geosystems, v. 10, no. 4,
1227 <https://doi.org/10.1029/2009gc002400>.
- 1228 Li, X. H., Tang, G. Q., Gong, B., Yang, Y. H., Hou, K. J., Hu, Z. C., Li, Q. L., Liu, Y.,
1229 and Li, W. X., 2013b, Qinghu zircon: A working reference for microbeam
1230 analysis of U-Pb age and Hf and O isotopes: Chinese Science Bulletin, v. 58,
1231 no. 36, p. 4647-4654, <https://doi.org/10.1007/s11434-013-5932-x>.
- 1232 Li, Y., Allen, P. A., Densmore, A. L., and Xu, Q., 2003, Evolution of the Longmen Shan
1233 foreland basin (western Sichuan, China) during the Late Triassic Indosinian
1234 orogeny: Basin Research, v. 15, no. 1, p. 117-138,
1235 <https://doi.org/10.1046/j.1365-2117.2003.00197.x>.
- 1236 Li, Y., Su, D., Zhou, R., Li, H., Densmore, A. L., Yan, L., and Yan, Z., 2013c, Episodic
1237 orogeny deduced from coeval sedimentary sequences in the foreland basin and
1238 its implication for uplift process of Longmen Mountain, China: Journal of
1239 Mountain Science, v. 10, no. 1, p. 29-42, [https://doi.org/10.1007/s11629-013-](https://doi.org/10.1007/s11629-013-2238-z)
1240 [2238-z](https://doi.org/10.1007/s11629-013-2238-z).
- 1241 Li, Z., 2009, Composition, structural characteristics and evolution of back-
1242 Longmenshan orogen (north section) in the northwest margin of Yangtze block
1243 [Doctoral thesis: Chang'an University, 211 p.
- 1244 Lin, A., Rao, G., and Yan, B., 2014, Structural analysis of the right-lateral strike-slip
1245 Qingchuan fault, northeastern segment of the Longmen Shan thrust belt, central
1246 China: Journal of Structural Geology, v. 68, p. 227-244,

- 1247 <https://doi.org/10.1016/j.jsg.2014.09.014>.
- 1248 Liu, H., Yan, D., and Wei, G., 2008, Deformation and metamorphism sequence of bikou
1249 terrane in the northwest margin of yangtze plate : implications for extension
1250 collapse and transition of songpan—garze orogenic belt: *Acta Geologica Sinica*,
1251 no. 04, p. 464-474+578-579. (In Chinese with english abstract)
- 1252 Long, X., 1991, Several questions of geochronic evolution in the mid-northeast
1253 segment of Longmenshan Mountains: *Journal of Chengdu College of geology*,
1254 v. 18, no. 1, p. 8-16. (In Chinese with english abstract)
- 1255 Lovera, O. M., Richter, F. M., and Harrison, T. M., 1989, $^{40}\text{Ar}/^{39}\text{Ar}$ geothermometry
1256 for slowly cooled samples having a distribution of diffusion domain size:
1257 *Journal of Geophysical Research*, v. 94, no. 17, p. 917-936,
1258 <https://doi.org/10.1029/JB094iB12p17917>.
- 1259 Ludwig, K. R., 2003, User's manual for Isoplot 3.00: a geochronological toolkit for
1260 Microsoft Excel: Berkeley Geochronol. Center Special Publication, v. 4, p. 47–
1261 93.
- 1262 Luo, Z., 1991, The dynamical model of the lithospheric evolution in Longmenshan
1263 Orogenic Belt: *Journal of Chengdu College of geology*, no. 1, p. 1-7. (In Chinese
1264 with english abstract)
- 1265 Ma, Y., Wang, G., and Hu, X., 1996, Tectonic deformation of Pengguan complex: *Acta*
1266 *Geologica Sichuan*, no. 02, p. 110-114. (In Chinese with english abstract)
- 1267 Mattauer, M., Malavieille, J., Calassou, S., Lancelot, J., Roger, R., Hao, Z., Xu, Z. Q.,
1268 and Hou, L. W., 1992, Le chaine Triassique de Songpan-Garze (Ouest Sichuan
1269 et est Tibet): Une chaine de plissement-decollement sur marge passive: *Comptes*
1270 *Rendus de L'Academie Des Sciences*, p. 619-626.
- 1271 McDougall, I., and Harrison, T. M., 1988, Geochronology and thermo- chronology by
1272 the $^{40}\text{Ar}/^{39}\text{Ar}$ method, New York, Oxford University Press.
- 1273 Meng, Q.-R., Wang, E., and Hu, J.-M., 2005, Mesozoic sedimentary evolution of the
1274 northwest Sichuan basin: Implication for continued clockwise rotation of the
1275 South China block: *Geological Society of America Bulletin*, v. 117, no. 3-4, p.
1276 396-410, <https://doi.org/10.1130/B25407.1>.
- 1277 Paterson, S. R., Tobisch, O. T., and Bhattacharyya, T., 1989, Regional, structural and
1278 strain analyses of terranes in the Western Metamorphic Belt, Central Sierra
1279 Nevada, California: *Journal of structural geology*, v. 11, no. 3, p. 255-273,

1280 [https://doi.org/10.1016/0191-8141\(89\)90066-7](https://doi.org/10.1016/0191-8141(89)90066-7).

1281 Phillips, D., Matchan, E. L., Honda, M., and Kuiper, K. F., 2017, Astronomical
 1282 calibration of 40 Ar/ 39 Ar reference minerals using high-precision, multi-
 1283 collector (ARGUSVI) mass spectrometry: *Geochimica et Cosmochimica Acta*,
 1284 v. 196, p. 351-369, <https://doi.org/10.1016/j.gca.2016.09.027>.

1285 Richardson, N. J., Densmore, A. L., Seward, D., Fowler, A., Wipf, M., Ellis, M. A.,
 1286 Yong, L., and Zhang, Y., 2008, Extraordinary denudation in the Sichuan Basin:
 1287 Insights from low-temperature thermochronology adjacent to the eastern margin
 1288 of the Tibetan Plateau: *Journal of Geophysical Research*, v. 113, no. B4,
 1289 <https://doi.org/10.1029/2006jb004739>.

1290 Robert, A., Pubellier, M., De Sigoyer, J., Vergne, J., Lahfid, A., Cattin, R., Findling, N.,
 1291 and Zhu, J., 2010a, Structural and thermal characters of the Longmen Shan
 1292 (Sichuan, China): *tectonophysics*, v. 491, no. 1, p. 165-173,
 1293 <https://doi.org/10.1016/j.tecto.2010.03.018>.

1294 Robert, A., Zhu, J., Vergne, J., Cattin, R., Chan, L. S., Wittlinger, G., Herquel, G., de
 1295 Sigoyer, J., Pubellier, M., and Zhu, L. D., 2010b, Crustal structures in the area
 1296 of the 2008 Sichuan earthquake from seismologic and gravimetric data:
 1297 *Tectonophysics*, v. 491, no. 1-4, p. 205-210,
 1298 <https://doi.org/10.1016/j.tecto.2009.11.010>.

1299 Roger, F., Jolivet, M., Cattin, R., and Malavieille, J., 2011, Mesozoic-Cenozoic
 1300 tectonothermal evolution of the eastern part of the Tibetan Plateau (Songpan-
 1301 Garze, Longmen Shan area): insights from thermochronological data and
 1302 simple thermal modelling: *Geological Society, London, Special Publications*, v.
 1303 353, no. 1, p. 9-25, <https://doi.org/10.1144/sp353.2>.

1304 Roger, F., Malavieille, J., Leloup, P. H., Calassou, S., and Xu, Z., 2004, Timing of
 1305 granite emplacement and cooling in the Songpan–Garzê Fold Belt (eastern
 1306 Tibetan Plateau) with tectonic implications: *Journal of Asian Earth Sciences*, v.
 1307 22, no. 5, p. 465-481, [https://doi.org/10.1016/s1367-9120\(03\)00089-0](https://doi.org/10.1016/s1367-9120(03)00089-0).

1308 Royden, L. H., Burchfiel, B. C., and van der Hilst, R. D., 2008, The geological
 1309 evolution of the Tibetan Plateau: *science*, v. 321, no. 5892, p. 1054-1058,
 1310 <https://doi.org/10.1126/science.1155371>.

1311 Sengör, A. M. C., 1984, The Cimmeride Orogenic System and the Tectonics of Eurasia:
 1312 *Geological Society of America Special Papers*, v. 195, p. 1-74,

- 1313 <https://doi.org/10.1130/SPE195-p1>.
- 1314 Shen, Z.-K., Sun, J., Zhang, P., Wan, Y., Wang, M., Bürgmann, R., Zeng, Y., Gan, W.,
1315 Liao, H., and Wang, Q., 2009, Slip maxima at fault junctions and rupturing of
1316 barriers during the 2008 Wenchuan earthquake: *Nature Geoscience*, v. 2, no. 10,
1317 p. 718-724, <https://doi.org/10.1038/ngeo636>.
- 1318 Sláma, J., Košler, J., Condon, D. J., Crowley, J. L., Gerdes, A., Hanchar, J. M.,
1319 Horstwood, M. S. A., Morris, G. A., Nasdala, L., Norberg, N., Schaltegger, U.,
1320 Schoene, B., Tubrett, M. N., and Whitehouse, M. J., 2008, Plešovice zircon —
1321 A new natural reference material for U–Pb and Hf isotopic microanalysis:
1322 *Chemical Geology*, v. 249, no. 1, p. 1-35,
1323 <https://doi.org/10.1016/j.chemgeo.2007.11.005>.
- 1324 Stacey, J. S., and Kramers, J. D., 1975, Approximation of terrestrial lead isotope
1325 evolution by a two-stage model: *Earth and Planetary Science Letters*, v. 26, no.
1326 2, p. 207-221, [https://doi.org/10.1016/0012-821X\(75\)90088-6](https://doi.org/10.1016/0012-821X(75)90088-6).
- 1327 Tian, Y., Kohn, B. P., Phillips, D., Hu, S., Gleadow, A. J., and Carter, A., 2016, Late
1328 Cretaceous–earliest Paleogene deformation in the Longmen Shan fold-and-
1329 thrust belt, eastern Tibetan Plateau margin: Pre-Cenozoic thickened crust?:
1330 *Tectonics*, p. 1-20, <https://doi.org/10.1002/2016TC004182>.
- 1331 Wallis, S. R., Tsujimori, T., Aoya, M., Kawakami, T., Terada, K., Suzuki, K., and Hyodo,
1332 H., Cenozoic and Mesozoic metamorphism in the Longmenshan orogen:
1333 Implications for geodynamic models of eastern Tibet, *in* *Proceedings AGU Fall*
1334 *Meeting Abstracts*2003, p. 745-748.
- 1335 Wang, E., Kirby, E., Furlong, K. P., van Soest, M., Xu, G., Shi, X., Kamp, P. J. J., and
1336 Hodges, K. V., 2012a, Two-phase growth of high topography in eastern Tibet
1337 during the Cenozoic: *Nature Geosci*, v. 5, no. 9, p. 640-645,
1338 <https://doi.org/10.1038/ngeo1538>.
- 1339 Wang, E., Meng, K., Chu, J. J., Wang, G., and Chen, Z., East Tibet Detachment:
1340 Evidence for the uplift mechanism of the Longmen Shan fault belt2012b.
- 1341 Wang, Q., Liang, B., Xie, Q., and He, W., 2000, Research on microstructures and
1342 deformation conditions of the Qingchuan fault zone: *Journal of Mineral*
1343 *Petrology*, v. 87, no. 01, p. 87-90. (In Chinese with english abstract)
- 1344 Wei, C., 1994, Metamorphism and its geological significance of the Bikou Group in the
1345 Shanxi–Gansu–Sichuan border region: *Acta Geologica Sinica*, v. 68, no. 03, p.

1346 241-254. (In Chinese with english abstract)

1347 Weislogel, A. L., Graham, S. A., Chang, E. Z., Wooden, J. L., and Gehrels, G. E., 2010,
 1348 Detrital zircon provenance from three turbidite depocenters of the Middle-
 1349 Upper Triassic Songpan-Ganzi complex, central China: Record of collisional
 1350 tectonics, erosional exhumation, and sediment production: Geological Society
 1351 of America Bulletin, v. 122, no. 11-12, p. 2041-2062,
 1352 <https://doi.org/10.1130/b26606.1>.

1353 Weller, O. M., St-Onge, M. R., Waters, D. J., Rayner, N., Searle, M. P., Chung, S. L.,
 1354 Palin, R. M., Lee, Y. H., and Xu, X., 2013, Quantifying Barrovian
 1355 metamorphism in the Danba Structural Culmination of eastern Tibet: Journal of
 1356 Metamorphic Geology, v. 31, no. 9, p. 909-935,
 1357 <https://doi.org/10.1111/jmg.12050>.

1358 West, D. P., and Lux, D. R., 1993, Dating mylonitic deformation by the ^{40}Ar - ^{39}Ar
 1359 method: An example from the Norumbega Fault Zone, Maine: Earth and
 1360 Planetary Science Letters, v. 120, no. 3-4, p. 221-237,
 1361 [https://doi.org/10.1016/0012-821X\(93\)90241-Z](https://doi.org/10.1016/0012-821X(93)90241-Z).

1362 Worley, B. A., and Wilson, C. J., 1996, Deformation partitioning and foliation
 1363 reactivation during transpressional orogenesis, an example from the Central
 1364 Longmen Shan, China: Journal of Structural Geology, v. 18, no. 4, p. 395-411,
 1365 [https://doi.org/10.1016/0191-8141\(95\)00095-U](https://doi.org/10.1016/0191-8141(95)00095-U).

1366 Xia, Z., 1993, Syntectonic regional epimetamorphism of the Triassic terrigenous
 1367 detrital rocks in the Songpan-Ganze orogenic belt during the early Yanshanian
 1368 orogeny: Acta Geologica Sichuan, no. 03, p. 189-192. (In Chinese with english
 1369 abstract)

1370 Xu, C., Wells, M. L., Yan, D.-P., Qiu, L., Zhou, Y., and Zhang, Y., 2020, Phase equilibria
 1371 and geochronology of Triassic blueschists in the Bikou terrane and Mesozoic
 1372 tectonic evolution of the northwestern margin of the Yangtze Block (SW China):
 1373 Journal of Asian Earth Sciences, v. 201, p. 104513,
 1374 <https://doi.org/https://doi.org/10.1016/j.jseaes.2020.104513>.

1375 Xu, Z., Hou, L., and Wang, Z., 1992, Orogenic process of the Songpan-Ganzé Orogenic
 1376 belt of China., Geological Publishing House, Beijing, 1-189 p.:

1377 Xu, Z., Ji, S., Li, H., Hou, L., Fu, X., and Cai, Z., 2008, Uplift of the Longmen Shan
 1378 range and the Wenchuan earthquake: Episodes, v. 31, no. 3, p. 291-301,

1379 <https://doi.org/10.1007/s00254-008-1393-y>.

1380 Xue, Z., Martelet, G., Lin, W., Faure, M., Chen, Y., Wei, W., Li, S., and Wang, Q., 2017,
 1381 Mesozoic crustal thickening of the Longmenshan belt (NE Tibet, China) by
 1382 imbrication of basement slices: insights from structural analysis, petrofabric and
 1383 magnetic fabric studies, and gravity modeling: *Tectonics*, p. 3110-3134,
 1384 <https://doi.org/10.1002/2017TC004754>.

1385 Xue, Z., Shang, Q., Jiang, W., Wang, Q., and Li, S., 2016, Emplacement age and
 1386 tectonic implications of the brecciated limestone at the edge of the
 1387 Longmenshan klippe: *Science China Earth Sciences*, v. 59, no. 3, p. 590-600,
 1388 <https://doi.org/10.1007/s11430-015-5209-0>.

1389 Yan, D., Zhou, M., Li, S., and Wei, G., 2011, Structural and geochronological
 1390 constraints on the Mesozoic-Cenozoic tectonic evolution of the Longmen Shan
 1391 thrust belt, eastern Tibetan Plateau: *Tectonics*, v. 30, no. 6, p. 1-24,
 1392 <https://doi.org/10.1029/2011TC002867>.

1393 Yan, D., Zhou, M., Wei, G., Gao, J., Liu, S., Xu, P., and Shi, X., 2008, The Pengguan
 1394 tectonic dome of Longmen Mountains, Sichuan Province: Mesozoic denudation
 1395 of a Neoproterozoic magmatic arc-basin system: *Science in China Series D:*
 1396 *Earth Sciences*, v. 51, no. 11, p. 1545-1559, [https://doi.org/10.1007/s11430-](https://doi.org/10.1007/s11430-008-0126-0)
 1397 [008-0126-0](https://doi.org/10.1007/s11430-008-0126-0).

1398 Yan, D.-P., Qiu, L., Wells, M. L., Zhou, M.-F., Meng, X., Lu, S., Zhang, S., Wang, Y.,
 1399 and Li, S.-B., 2018, Structural and Geochronological Constraints on the Early
 1400 Mesozoic North Longmen Shan Thrust Belt: Foreland Fold-Thrust Propagation
 1401 of the SW Qinling Orogenic Belt, Northeastern Tibetan Plateau: *Tectonics*, v.
 1402 37, no. 12, p. 4595-4624, <https://doi.org/10.1029/2018tc004986>.

1403 Yan, Q., Wang, Z., Hanson, A., Druschke, P., Yan, Z., Liu, D., Jian, P., Song, B., Wang,
 1404 T., and Jiang, C., 2003, SHRIMP age and geochemistry of the Bikou volcanic
 1405 terrane: implications for Neoproterozoic tectonics on the northern margin of the
 1406 Yangtze Craton: *Acta Geologica Sinica (English Edition)*, v. 77, no. 4, p. 479-
 1407 490.

1408 Yan, Z., Tian, Y., Li, R., Vermeesch, P., Sun, X., Li, Y., Rittner, M., Carter, A., Shao, C.,
 1409 Huang, H., and Ji, X., 2019, Late Triassic tectonic inversion in the upper
 1410 Yangtze Block: Insights from detrital zircon U–Pb geochronology from south-
 1411 western Sichuan Basin: *Basin Research*, v. 31, no. 1, p. 92-113,

1412 <https://doi.org/https://doi.org/10.1111/bre.12310>.

1413 Yin, A., The Songpan-ganzi terrane of Tibet: was it a part of South China since the
 1414 Neoproterozoic, *in* Proceedings 2015 GSA Annual Meeting in Baltimore,
 1415 Maryland, USA, 2015, Volume 47, Geological Society of America, p. 38.

1416 Yin, A., and Nie, S., 1993, An indentation model for the North and South China
 1417 collision and the development of the Tan-Lu and Honam fault systems, Eastern
 1418 Asia: Tectonics, v. 12, no. 4, p. 801-813.

1419 Yuan, C., Zhou, M.-F., Sun, M., Zhao, Y., Wilde, S., Long, X., and Yan, D., 2010,
 1420 Triassic granitoids in the eastern Songpan Ganzi Fold Belt, SW China:
 1421 Magmatic response to geodynamics of the deep lithosphere: Earth and Planetary
 1422 Science Letters, v. 290, no. 3-4, p. 481-492,
 1423 <https://doi.org/10.1016/j.epsl.2010.01.005>.

1424 Yuan, H., Zhang, Z., and Zhang, P., 1991, The uplift and cooling histories of Laojungou
 1425 granite in the western margin of the central Longmen Mountain: Journal of
 1426 Chengdu College of geology, v. 18, no. 1, p. 17-22. (In Chinese with english
 1427 abstract)

1428 Zhang, H., Parrish, R., Zhang, L., Xu, W., Yuan, H., Gao, S., and Crowley, Q. G., 2007,
 1429 A-type granite and adakitic magmatism association in Songpan–Garze fold belt,
 1430 eastern Tibetan Plateau: Implication for lithospheric delamination: Lithos, v. 97,
 1431 no. 3-4, p. 323-335, <https://doi.org/10.1016/j.lithos.2007.01.002>.

1432 Zhang, H., Zhang, L., Harris, N., Jin, L., and Yuan, H., 2006a, U–Pb zircon ages,
 1433 geochemical and isotopic compositions of granitoids in Songpan-Garze fold
 1434 belt, eastern Tibetan Plateau: constraints on petrogenesis and tectonic evolution
 1435 of the basement: Contributions to Mineralogy and Petrology, v. 152, no. 1, p.
 1436 75-88, <https://doi.org/10.1007/s00410-006-0095-2>.

1437 Zhang, H.-P., Liu, S.-F., Yang, N., Zhang, Y.-Q., and Zhang, G.-W., 2006b, Geomorphic
 1438 characteristics of the Minjiang drainage basin (eastern Tibetan Plateau) and its
 1439 tectonic implications: New insights from a digital elevation model study: The
 1440 Island Arc, v. 15, no. 2, p. 239-250, <https://doi.org/10.1111/j.1440-1738.2006.00524.x>.

1442 Zhang, K.-J., Li, B., Wei, Q.-G., Cai, J.-X., and Zhang, Y.-X., 2008, Proximal
 1443 provenance of the western Songpan–Ganzi turbidite complex (Late Triassic,
 1444 eastern Tibetan plateau): Implications for the tectonic amalgamation of China:

1445 Sedimentary Geology, v. 208, no. 1-2, p. 36-44,
 1446 <https://doi.org/10.1016/j.sedgeo.2008.04.008>.

1447 Zhang, P., Shen, Z., Wang, M., Gan, W., Bürgmann, R., Molnar, P., Wang, Q., Niu, Z.,
 1448 Sun, J., Wu, J., Hanrong, S., and Xinzhaoy, Y., 2004, Continuous deformation of
 1449 the Tibetan Plateau from global positioning system data: Geology, v. 32, no. 9,
 1450 p. 809, <https://doi.org/10.1130/g20554.1>.

1451 Zhang, Y. X., Li, B., Wei, Q. G., Wang, Y., and Zhang, K. J., 2006c, Detrital zircon
 1452 provenance of the Late Triassic Songpan-Ganzi complex: Sedimentary record
 1453 of collision of the North and South China blocks: Comment and Reply:
 1454 COMMENT: Geology, v. 34, no. 1, p. e107-e107,
 1455 <https://doi.org/10.1130/g22944.1>.

1456 Zhang, Z., Wang, Y., Chen, Y., Houseman, G. A., Tian, X., Wang, E., and Teng, J., 2009,
 1457 Crustal structure across Longmenshan fault belt from passive source seismic
 1458 profiling: Geophysical Research Letters, v. 36, no. 17, p. L17310-,
 1459 <https://doi.org/10.1029/2009gl039580>.

1460 Zhang, Z., Yuan, X., Chen, Y., Tian, X., Kind, R., Li, X., and Teng, J., 2010, Seismic
 1461 signature of the collision between the east Tibetan escape flow and the Sichuan
 1462 Basin: Earth and Planetary Science Letters, v. 292, no. 3-4, p. 254-264,
 1463 <https://doi.org/10.1016/j.epsl.2010.01.046>.

1464 Zhao, Y., 2007, Mesozoic granitoids in Eastern Songpan-Garze: Geochemistry,
 1465 petrogenesis and tectonic implications [Doctor: Graduate school of the Chinese
 1466 Academy of Sciences, 111 p.

1467 Zheng, Y., Kong, P., and Fu, B., 2014, Time constraints on the emplacement of klippen
 1468 in the Longmen Shan thrust belt and tectonic implications: Tectonophysics, v.
 1469 634, p. 44-54, <https://doi.org/10.1016/j.tecto.2014.07.018>.

1470 Zheng, Y., Li, H., Sun, Z., Wang, H., Zhang, J., Li, C., and Cao, Y., 2016, New
 1471 geochronology constraints on timing and depth of the ancient earthquakes along
 1472 the Longmen Shan fault belt, eastern Tibet: Tectonics, v. 35, no. 12, p. 2781-
 1473 2806, <https://doi.org/10.1002/2016TC004210>.

1474 Zhou, M., Yan, D., Kennedy, A. K., Li, Y., and Ding, J., 2002, SHRIMP U–Pb zircon
 1475 geochronological and geochemical evidence for Neoproterozoic arc-
 1476 magmatism along the western margin of the Yangtze Block, South China: Earth
 1477 and Planetary Science Letters, v. 196, no. 1, p. 51-67,

1478 [https://doi.org/10.1016/S0012-821X\(01\)00595-7](https://doi.org/10.1016/S0012-821X(01)00595-7).
1479 Zhou, M., Yan, D., Vasconcelos, P. M., Li, J., and Hu, R., 2008, Structural and
1480 geochronological constraints on the tectono-thermal evolution of the Danba
1481 domal terrane, eastern margin of the Tibetan plateau: Journal of Asian Earth
1482 Sciences, v. 33, no. 5, p. 414-427, <https://doi.org/10.1002/2016TC004210>.
1483

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

~30 meters north to the south margin of the Mupi pluton

~10–15 meters north to the south margin of the Mupi pluton

South margin of the Mupi pluton

Figure 9

Figure 10

Northwestern Zone

Southeastern Zone -- Inner metamorphic core

Basement Complex

Figure 11

Figure 12

Figure 13

Figure 14

✓ Pre T₂ normal fault / Reverse fault during T₃–aJ₁ ✓ / Episodic and synchronous NW-normal faulting and SE-thrusting during J₃–K₁, K₂–E₁

J–Q T₃ D–T₂ E–S Pt basement Meta-unit

Supplemental Material

Figure S1. (a) Reflection image show location of the crack in feldspar, same crack in lower middle of Figure 8e from the Mupi pluton. (b) Same crack in Figure a, cross-polarized light. (c) Back scattered electron image show points selected for energy-dispersive X-ray spectrometer measurement. (d and h) Energy-dispersive X-ray spectra of point 1 and point 6 in the crack show typical feature of quartz. (e–g) Energy dispersive X-ray spectra of point 2, point 3, and point 5 indicate component of feldspar with different Ca content.

Figures S2. Features of the samples selected for ^{40}Ar - ^{39}Ar dating. Shear plane is defined by fine-grained white mica (c–d) or white mica (a–b, e–f, and g). (h) Field observation of sample CX6, where sigma-shaped quartz lens indicate a top-to-the-SE thrusting. Mineral abbreviations after Kretz (1983). CPL—cross-polarized light; PPL—plain-polarized light.

Figure S3. All samples show high radiogenic ^{40}Ar (Table S2) and almost all steps are clustered at the $^{39}\text{Ar}/^{40}\text{Ar}$ axis that do not define a real inverse isochron line. The initial $^{40}\text{Ar}/^{36}\text{Ar}$ ratio originate from the inverse isochron line is meaningless as well. Most of the step-ages calculated both by using initial $^{40}\text{Ar}/^{36}\text{Ar}$ of 298.56 (Lee et al., 2006) and from the “inverse isochron”, show nearly no differences (Table S2). Thus, the inverse isochron ages, though identical to their weighted mean ages or plateau ages, are meaningless and have not been discussed in the main text.

Table S1. Detailed SIMS U-Pb zircon results of sample CX48

Table S2. $^{40}\text{Ar}/^{39}\text{Ar}$ VG3600 Furnace Step-Heating analytical results

Figure S1

Figure S2

Figure S3
Northwestern Zone

Southeastern Zone --
Inner metamorphic core

Basement Complex