First green lacewing (Insecta: Neuroptera: Chrysopidae) from the Paleocene Sanshui Basin of Guangdong, South China

Diyang Huang, Xinneng Lian, Corentin Jouault, Jian Gao, André Nel

► To cite this version:

Diyang Huang, Xinneng Lian, Corentin Jouault, Jian Gao, André Nel. First green lacewing (Insecta: Neuroptera: Chrysopidae) from the Paleocene Sanshui Basin of Guangdong, South China. *Historical Biology*, 2022, 34 (4), pp.704-712. 10.1080/08912963.2021.1943381 . insu-03272764

HAL Id: insu-03272764

<https://hal-insu.archives-ouvertes.fr/insu-03272764>

Submitted on 28 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Historical Biology

First green lacewing (Insecta: Neuroptera: Chrysopidae) from the Paleocene Sanshui Basin of Guangdong, South China

Diyang Huang^{a, *}, Xinneng Lian^a, Corentin Jouault^{b, c}, Jian Gao^a and André Nel^{b, *}

^aState Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of Geology and Palaeontology and Center for Excellence in Life and Paleoenvironment, Chinese Academy of Sciences, 39 East Beijing Road, Nanjing 210008, PR China; ^bInstitut de Systématique, Évolution, Biodiversité (ISYEB) Muséum national d'Histoire naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles, CP50, 57 rue Cuvier 75005 Paris, France; ^cUniv. Rennes, CNRS, Géosciences Rennes, UMR 6118, 35000, Rennes, France

*Corresponding authors: dyhuang@nigpas.ac.cn; anel@mnhn.fr

ABSTRACT

Sinonothochrysa zhang Huang and Nel gen. et sp. nov., a new genus and species of nothochrysine lacewing, is described and illustrated from the Paleocene Buxin Formation, Sanshui Basina, Foshan City, Guangdong Province, South China. The new taxon is the oldest described representative and first report this subfamily in China. It is also the first formally described fossil insects from the Buxin Formation of the Sanshui Basin.

KEYWORDS

Neuroptera; Nothochrysinae; fossil insects; Buxin Formation; middle Paleocene

Introduction

The family Chrysopidae, also called green lacewings due to the color of their extant representatives, is a widespread group of Neuroptera, which has a quite rich fossil record, comprising more than 20 fossil genera and more than 60 fossil species (compared to other Neuroptera as the Myrmeleontidae) (Table. 1; <http://fossilworks.org>). Their oldest representatives are dated back from the Jurassic of China and Kazakhstan (e.g. Martynov 1927; Panfilov, 1980; Khramov et al. 2015), but their highest species diversity, in the fossil record, is known during the Paleogene. Representatives of the stem-group appear early in the evolutionary history of the family as the subfamily Limaiinae is known since the Jurassic period (Table. 1), suggesting that the family may be older.

Currently, the family is divided in four subfamilies (viz. Apochrysinae + Chrysopinae + Nothochrysinae + Limaiinae), with the Limaiinae is a strictly fossil subfamily. Each subfamilies present huge disparities in their fossil species diversity (Table. 1) since the subfamily Apochrysinae is completely absent from the fossil record while the two other extant subfamilies possess fossil representatives. Interestingly, while the majority of the fossil record of the Limaiinae is Mesozoic, this group survived the Cretaceous-Cenozoic crisis with some Paleocene and the Eocene representatives, viz. three species in the genus *Protochrysa* Willmann and Brooks, 1991 from the Paleocene-Eocene of Denmark and the Early Eocene of British Columbia (Nel et al. 2005; Makarkin and Archibald 2013). This scenario is very similar to the fate of the Mammalia Multituberculata that were very diverse during the Mesozoic and became extinct during the Paleocene, well after the K-T crisis. The fossil record of the crown group Chrysopidae is currently only represented by Cenozoic Nothochrysinae and Chrysopinae (Table 1).

The position of the family Chrysopidae is not fully elucidated and alternative hypotheses are proposed depending on the methods or the analyzed data (e.g. Engel et al. 2018; Vasilikopoulos et al. 2020). Winterton et al. (2019) proposed a molecular phylogenetic analysis of the Chrysopidae, dating the separation of their stem group from their putative sister group Myrmeleontoidea from the late Triassic and their crown group from the early Cretaceous. However, it seems that the Chrysopidae would closely related to the clade (Mantispidae + Berothidae) according to recent molecular analyses (Vasilikopoulos et al. 2020: fig. 1) with a divergence estimate to the early Jurassic. However, the calibration used in this most recent work seem to provide questionable age for the group. In fact, Vasilikopoulos et al. (2020) used the chrysopid fossil species *Mesypochrysa cf. intermedia* to calibrate the crown-group (Chrysopidae + (Mantispidae + Berothidae)) using a soft minimum constrain set to 158.1 Ma (Vasilikopoulos et al., 2020: additional file 1: table S10), which fits with the datation of the Karabastau Formation of Kazakhstan, dated back to the Callovian/Kimmeridgian (Doludenko and Orlovskaya 1976a, Doludenko and Orlovskaya 1976b) or possibly Oxfordian–Kimmeridgian (Doludenko et al. 1990). Recent study on Yanliao Biota late assemblage i.e. Linglongta biota at the Guancaishan locality was considered correlate to Karatau biota of Kazakhstan (Huang 2015, 2016) that yielded some very similar insects such as *Cicadomorpha* (Fu et al. 2020). An ash layer appeared less one meter up the fossil insect layer at the Guancaishan locality providing a SIMS zircon U-Pb dating as 157.3 ± 1.5 Ma that support Karabastau Formation is more likely an Oxfordian–Kimmeridgian age (Huang 2016). However, the Toarcian genus *Protoaristenymphes* Nel and Henrotay, 1994 is currently placed in the Mesozoic family Mesochrysopidae and the Chrysopoidea (a superfamily that also comprises the Mesozoic Ascalochrysidae), and would be the oldest known representative of this clade (Nel and Henrotay 1994; Yang et al. 2012; Makarkin et al. 2013), supporting a possible older age for the superfamily, more in accordance with Winterton et al. (2019), even if these last

1
2
3 authors did not used *Protoaristenymphes* to calibrate theirs datations. Vasilikopoulos et al.
4
5 (2020) also ignored the Mesochrysopidae and the genus *Protoaristenymphes* in their list of
6
7 fossils used as calibration points. We are aware that the choice of fossil taxa in calibrations is
8
9 difficult and that applying too much constraints on trees may results in dubious results, but, at
10
11 least the putative oldest records of the concerned clades should be discussed. Therefore, we
12
13 wonder if the clade or at least the Chrysopidae could be older than previously thought, and
14
15 suggest that it would be interesting to explore the Neuroptera phylogeny testing different
16
17 calibrations of the stem- and crown-groups for each family.
18
19

20
21 Herein we described a new genus and species of Nothochrysinæ from the Paleocene,
22
23 confirming the important diversity of this subfamily at that time.
24
25

26 27 28 **Geological setting**

29
30 The Paleogene deposit of the Sanshui Basin of South Guangdong Province, South China is
31
32 developed and detailed hereafter (Fig. 1). This strata is divided into the Xinzhuangcun, the
33
34 Buxin, the Baoyue, and the Huachong formations from bottom to top. Buxin Formation was
35
36 established at 1964 for a set of lacustrine deposit comprised dark gray oil shale, mud stone,
37
38 chalky clay, and sand stone. It was firstly considered as an Eocene or Early Oligocene strata.
39
40 The age of the Buxin Formation was argued by authors based on different biochronometer with
41
42 various opinions, summarized by Zhang et al. (1993): 1) Early Eocene by ostracodes,
43
44 conchostracans, gastropods, and fishes; 2) Late Paleocene-Early Eocene by bivalves; 3) Early-
45
46 Middle Eocene by plant fossils and spore-pollen studies; 4) Middle Eocene by stoneworts. Later,
47
48 a systematical biostratigraphic study on ostracods suggested that the Xinzhuangcun, Buxin, and
49
50 Baoyue formations would be assigned to the Paleocene and the Huachong Formation would be
51
52 assigned to the Early Eocene. Therefore, the above authors indicated that the age of Buxin
53
54 Formation could be assigned to the Middle Paleocene (Zhang et al. 2008). A recent integrative
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

study on China’s Paleogene stratigraphy and timescale indicated that the age of Buxin Formation would range from 62.2 Ma to 59.2 Ma, i.e. late Danian to Selandian-Thanetian boundary, mainly based on a paleomagnetism study (Wang et al. 2019). According to this opinion, the age of the present fossil material could be even as old as 62 Ma.

Most isotopic dating of volcanic rocks in the Sanshui Basin used K-Ar method (Guangdong Provincial Bureau of Geology and Mineral Resources 1997) could not provide convincing geochronological ages. A Paleocene age of the Buxin Formation was also supported by a geochronological study that indicated that the volcanic eruption started at approximately 60 Ma in the Sanshui Basin. These authors got two set of ages by zircon U-Pb dating as 59.3 ± 0.2 Ma and 56.2 ± 0.3 Ma. The Buxin Formation is subdivided into four members as Honggang Member, Dagang Member, Fenggang Member, and Gaogang Member from bottom to top. The herein described fossil insect was collected from the lower part of the Honggang Formation. The volcanic rocks in the Paleogene of Sanshui Basin initially appeared in the Honggang Member and particularly occurred at the bottom of the Gaogang Member, and then were widely distributed in the lower parts of the Huachong Formation (Zhang et al. 1993). Therefore, the above isotopic age 59.3 ± 0.2 Ma and 56.2 ± 0.3 Ma more likely corresponds to the volcanic events of the bottom Gaogang Member of Buxin Formation and the Lower part of the Huachong Formation respectively. This opinion would indicate that the age of the present insect layer is about 61–62 Ma approximately. A correspondence of these two volcanic rock layers with the lower and upper parts of the Buxin Formation cannot be excluded. Then, the age of this fossil insect layer would be 60–59 Ma approximately. To briefly summarize, the age of the lower part of the Buxin Formation that yielded numerous fossil insects would belong to the Middle Paleocene, very close to the famous French volcano-sedimentary maar deposit of Menat in age (60–61 Ma; e.g. Nel and Auvray 2006; Wappler et al. 2009). Nevertheless, both localities with rich fossil entomofaunas need future better accurate isotopic dating.

The fossil insects from the Sanshui Basin have been found from the Buxin and the Baoyue formations but without a systematically collecting and study. These fossil insects were identified by Qibin Lin (NIGPAS) including *Laodiscis melanis*, *Nisocercopis* sp., *Platokoleos* sp. etc., all taxa that firstly described from the Paleogene of Mengla County, Yunnan Province, Southwestern China (Zhang et al. 1993). The so-called Mengyejing Formation at Shangyong of Mengla County yielded a very rich fossil entomofauna (e.g. Lin 1977; Lin et al. 2010a, b), and its age was also discussed by authors. Analyses of spore-pollen assemblages suggested a Paleogene age for the ‘Mengyejing’ Formation at Shangyong, but a Late Cretaceous age was not excluded (Nanjing Institute of Geology and Palaeontology et al. 1975). Study of conchostracans suggested an early Eocene age i.e. Ypresian (Chen and Shen 1980; Shen and Chen 1984; Shen et al. 2006). This entomofauna was considered as Eocene (Lin 1977) or Paleocene (Hong 1984) respectively. For the general analysis, the insect-rich layer of the Buxin Formation at the Sanshui Basin and ‘Mengyejing’ Formation at Shangyong are correlated and both would be Paleocene.

Material and method

The specimen describe herein derives from the chalky clay of lower part of Honggang Member, Buxin Formation at western of Sanshui District, Foshan City, Guangdong Province, about 40 km west of Guangzhou, the capital of the province (Fig. 1).

The holotype of *Sinonothochrysa zhang* Huang and Nel gen. et sp. nov. is an imprint/compression fossil with three-dimensional preservation. It is a nearly complete insect with well-preserved wing venation but the body structures are damaged by crystallization. The specimen was examined and photographed with a LeicaMZ APO and a Canon EOS 5DMark II camera. All images are digitally stacked photomicrographic composites of several individual focal planes, which were obtained using a Zeiss Discovery V20 stereo microscope. The figures

and drawings were composed with Adobe Illustrator CC2019 and Photoshop CC2019 softwares. The holotype specimen presented herein is housed in NIGPAS, Nanjing Institute of Geology and Palaeontology, Chinese Academy of Science, China under the accession number NIGP175154. Wing venation nomenclature follows Breitkreuz et al. (2017) and in part Brooks and Barnard (1990).

Abbreviations are as follows: A anal vein; bsx basal subcostal crossvein; C costa; C1 and C2 first and second cubital cells; CuA cubitus anterior; CuP cubitus posterior; dcc discal cubital cell; ig inner gradate crossveins; im intramedian cell; MA media anterior; MP media posterior; M1 and M2 first and second median cells; og outer gradate crossveins; Psc Pseudo-cubital vein; Psm pseudo-median vein RA radius anterior; RP radius posterior; ScP subcostal posterior.

Published work and nomenclatural acts are registered in ZooBank urn:lsid:zoobank.org:pub:16ED73C0-9126-4313-837E-D2A76B8738C8.

Systematic palaeontology

Order Neuroptera Linnaeus, 1758

Family Chrysopidae Schneider, 1851

Subfamily ?Nothochrysinæ Navás, 1910

Sinonothochrysa Huang and Nel, gen. nov.

urn:lsid:zoobank.org:act:176B7F87-BA8C-4CCF-80A5-2121B79C9FEF

Type species

Sinonothochrysa zhangii n. sp.

Diagnosis

Forewing characters. Two series of gradate crossveins; cell *im* nearly quadrangular and very broad, twice as long as broad, with its anterior side curved; vein *Psm* with three branches of *RP* entering it, *Psm* continued by inner gradates; crossvein *2m-cu* at basal third of cell *im*; series of cells between *Psm* and *Psc* distinctly broader than long; inner gradate series anteriorly curved.

Etymology

Named after the Latin name 'Sina' for China, and the extant genus name *Nothochrysa* McLachlan, 1868.

Sinonhothochrysa zhangi Huang and Nel sp. nov.

Figs 2–3

urn:lsid:zoobank.org:act:0238777A-68AE-453D-A83B-2A793B4F06DC

Etymology

Named after Prof. Xianqiu Zhang, Chinese senior palaeontologist who made a systematically biostratigraphic study of the Sanshui Basin. The specific epithet is to be treated as a noun in the genitive case.

Type material

Holotype NIGP175154 (crystallized body with complete but superposed right fore- and hind wings and fragments of the left fore- and hind wings).

Type locality

Sanshui District, Foshan City, Guangdong Province, China.

Stratigraphic occurrence

Middle Paleocene, Buxin Formation at Western of Sanshui District.

Diagnosis

As for the genus.

Description

Body ca. 5.0 mm long; thorax ca. 1.0 mm wide.

Forewing (Figs 2–3), 9.0 mm long, ca. 2.2 mm wide; apex rounded; costal space poorly preserved, with simple subcostal veinlets relatively widely spaced; ScP very long, with apex not preserved but probably entering margin near wing apex; subcostal space moderately broad; proximal subcostal crossvein (bsx or 1scP-r) present; no other crossvein between ScP and RA detected; tympanum not detectable; RA long, entering margin at wing apex; origin of RP relatively close to wing base, at 3.0 mm of it; length of RP proximad 1r–m 10 times as long as 1r–m; RA space (between RA and RP) broad, with 10 crossveins, the most basal one being well distal to 1r–m; RP weakly zigzagged, with 10 branches, two of them entering Psm; basal crossvein 1r–m very short, perpendicular to M, connecting RP with MA; M forked at an open angle opposite origin of RP; intramedian cell very broad, only two times as long as broad, broadly rectangular with anterior side strongly curved; MA not proximally fused with RP1; Psm well-developed, slightly zigzagged and distally prolonged by inner row of gradates; MP distally fused with CuA; Psc with three-four simple branches between it and posterior wing margin; cells between Psm and Psc rather short but very broad; Psc distally prolonged by outer row of gradates; basal crossvein 1m–cu short, apparently located basal to origin of CuP; 2m–cu present between MP and CuA at basal third of cell im; anal and CuP veins incompletely;

two complete gradate series of crossveins, outer series not parallel to posterior wing margin; inner series not parallel to outer series; wing membrane apparently hyaline, only darkened in distal part of ScP (pterostigma).

Hind wing (Figs 2–3) ca. 8.8 mm long, ca. 2.1 mm wide; apex acute; costal space not preserved; only basal part of ScP visible, subcostal space as broad as that of forewing; proximal subcostal crossvein (bsx or 1scp-r) present; no other crossvein between ScP and RA detected; RA long, entering margin at wing apex; origin of RP relatively close to wing base, at ca. 1.1 mm of it; length of basal part of RP as long as part fused with MA; RA space (between RA and RP) broad, with 11 crossveins; RP weakly zigzagged, with eighth branches, three of them entering Psm; M forked at an open angle slightly basal to origin of RP; MA distally fused with RP1 and separating again; Psm well-developed, slightly zigzagged and distally prolonged by inner row of gradates; MP distally fused with CuA; Psc with four simple branches between it and posterior wing margin; cells between Psm and Psc large, rather short but very broad; Psc distally prolonged by outer row of gradates; intracubital crossveins C1 and C2 and anal veins not preserved except for forked first anal vein; two complete gradate series of crossveins, wing membrane apparently hyaline.

Discussion

After the key to the extant chrysopid subfamilies of Brooks and Barnard (1990: 132), *Sinonothochrysa* gen. nov. falls in the Nothochrysinae because of the character ‘forewing Psm continuous with inner row of gradates’. Tauber (2014) proposed a list of putative apomorphies of the Nothochrysinae and two groups of genera, but none of these are preserved in our specimen. Makarkin and Archibald (2013) considered the character ‘Psm not extending to the outer row of gradates’ as possibly characteristic of the Nothochrysinae. They also have the

crossvein 2m-cu located in the proximal half of im. This character is also present in some Chrysopinae, but not in the Cretaceous-Paleogene Limaiinae.

Garzón-Orduña et al. (2019) listed nine extant genera in the Nothochrysinae, viz. *Nothochrysa* McLachlan, 1868, *Pimachrysa* Adams, 1957, and *Hypochrysa* Hagen, 1866, *Asthenochrysa* Adams and Penny, 1992, *Leptochrysa* Adams and Penny, 1992, *Pamochrysa* Tjeder, 1966, *Kimochrysa* Tjeder, 1966 (considered as a subjective synonym of *Hypochrysa* by Tauber, 2014), *Triplochrysa* Kimmins, and *Dictyochrysa* Esben-Petersen (Tjeder 1966; Adams 1967; Adams and Penny 1992). Several fossil genera are currently put in this subfamily (Table 1), viz. *Adamsochrysa* Makarkin and Archibald, 2013 (Eocene), *Archaeochrysa* Adams, 1967 (Eocene to Oligocene), *Asiachrysa* Makarkin, 2014 (Eocene), *Dyspetochrysa* Adams, 1967 (Oligocene), *Lithochrysa* Carpenter, 1935 (Oligocene), *Okanaganochrysa* Makarkin and Archibald, 2013 (Eocene), *Palaeochrysa* Scudder, 1890 (Oligocene), *Pronothochrysa* Peñalver et al., 1995 (Miocene), *Pseudochrysopa* Makarkin and Archibald, 2013 (Eocene), and *Tribochrysa* Scudder, 1890 (Oligocene). As Winterton et al. (2019) found the Nothochrysinae as paraphyletic in their phylogenetic analysis, the exact positions of all these fossils remain questionable.

Dictyochrysa and *Triplochrysa* are excluded from consideration because the gradate crossveins in *Sinonothochrysa* gen. nov. are arranged in two series only (vs. more than two in these genera, see respectively Brooks and Barnard 1990: fig. 527; New 1980: fig. 13). *Pamochrysa* has the forewing cell im rhomboidal, while it is nearly quadrangular in *Sinonothochrysa* gen. nov. (Brooks and Barnard 1990: fig. 554).

Sinonothochrysa gen. nov. shares with *Kimochrysa* a forewing cell im ca. twice as long as broad, unlike the genus *Nothochrysa* in which im is much narrower (Adams 1967: fig. 3; Kovanci and Canbulat 2007; Tauber 2019). But the forewing cells im of the *Kimochrysa* spp. are triangular or pentagonal (Tjeder 1966; Adams 1978; Brooks and Barnard 1990: fig. 539).

Hypochrysa elegans (Burmeister, 1839) and *Asthenochrysa* (type species *A. viridula* (Adams, 1978)) have also a triangular cell im (Adams 1967: fig. 5, 1978: fig. 1; Brooks and Barnard 1990: fig. 533). In *Sinonothochrysa* gen. nov., the anterior side of the cell im is curved, while in *Kimochrysa impar* Tjeder, 1966, it is slightly angular, rendering im pentagonal (Tjeder 1966: fig. 785), the difference being rather weak. Nevertheless, the vein 'distal part of MA – Psm' is much shorter in *K. impar* and *K. africana* (Kimmins, 1937) than in *Sinonothochrysa* gen. nov., with only two short cells above it basal to the first i.g. vein instead of three large ones.

While *Pimachrysa grata* Adams, 1957, *P. fusca* Adams, 1967 and *P. intermedia* Adams, 1967 have rather narrow cells im, *P. nigra* Adams, 1967 has one nearly ca. twice as long as broad as in *Sinonothochrysa* gen. nov. (Adams 1957: fig. 1, 1967: figs 1-2, 4). They all differ from *Sinonothochrysa* gen. nov. in the crossvein 2m-cu basal or very close to basal angle of cell im and the series of cells between Psm and Psc nearly as long as broad.

Leptochrysa has a very particular cell im, quadrangular but with the cell below it strongly reduced (Adams and Penny 1992: fig. 10; Tauber 2019), strongly differing from the configuration of im in *Sinonothochrysa* gen. nov. (pentagonal without reduction of the cell below it).

Okanaganochrysa has two series of cells between Psm and Psc (Makarkin and Archibald 2013: fig. 4). The *Adamsochrysa* spp. have a hexagonal cell im, but broadly similar to that of *Sinonothochrysa* gen. nov. (Makarkin and Archibald 2013: fig. 6), but they also have four rows of gradate crossveins. The *Archaeochrysa* spp. have a quadrangular cell im, but much narrower than in *Sinonothochrysa* gen. nov. (Adams 1967; Makarkin and Archibald 2013). Makarkin and Archibald (2013) provisionally put *Pseudochrysopa* in the Nothochrysinae because of the two following characters: 'Psm does not extend to the outer gradate series'; and '2m-cu is located in the proximal half of im'. The exact position of this genus needs to be re-

evaluated. Nevertheless, it strongly differs from *Sinonothochrysa* gen. nov. in the narrower cell im and vein Psm not continued by the inner gradates.

Dyspetochrysa vetuscula (Scudder, 1890), *Tribochrysa inaequalis* Scudder, 1885, *Tribochrysa firmata* Scudder, 1890, *Lithochrysa wickhami* (Cockerell, 1914), *Palaeochrysa stricta* Scudder, 1890 and *P. creedei* Carpenter, 1935 differ from *Sinonothochrysa* gen. nov. in the elongate cell im, a course of the inner gradates parallel to posterior wing margin, instead of being anteriorly curved (Carpenter 1935: figs 1, 3-4, 6-7). *Palaeochrysa fracta* Cockerell, 1914 has also a narrow cell im and an irregular series of inner gradates (maybe an aberrant feature) (Carpenter 1935: fig. 2).

Pronothochrysa vivesi Peñalver et al., 1995 has also a narrow cell im, plus a much longer vein 'distal part of MA – Psm' than in *Sinonothochrysa* gen. nov. with six cells between it and RP instead of three (Peñalver et al. 1995: fig. 4).

Asiachrysa tadushiella Makarkin, 2014 has a cell im similar to that of *Sinonothochrysa* gen. nov., although slightly narrower. Nevertheless, its cells between Psm and Psc are longer and narrower than in *Sinonothochrysa* gen. nov. and overall, the crossvein 2m-cu is situated below mid part of cell im instead of being in the basal third of this cell (Makarkin 2014: fig. 3).

Conclusion

Except for a Miocene genus, the fossil record of the subfamily Nothochrysinæ is restricted to the Eocene – Oligocene, with an impressive diversity during the Eocene to the earliest Oligocene with ten described extinct genera contra only one (*Nothochrysa*) during the Middle – Late Oligocene, confirmed by the newly described genus. Possibly this subfamily collapsed during the Oligocene in relation to the general cooling and/or bat diversification, as already suggested by Archibald et al. (2014) and Ngô-Muller et al. (2019). The Paleocene

Sinonothochrysa zhang Huang and Nel gen. nov. et sp. nov. is the oldest record of the Nothochrysinae.

Acknowledgements

We thank two anonymous reviewers for their constructive comments. Financial support was provided by the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB18000000 and XDB26000000), and the National Natural Science Foundation of China (41925008 and 41688103).

References

- Adams PhA. 1957. A new genus and new species of Chrysopidae from the Western United States, with remarks on the wing venation of the family (Neuroptera). *Psyche* 63:67–74. doi.org/10.1155/1956/47287
- Adams PhA. 1967. A review of the Mesochrysinae and Nothochrysinae (Neuroptera: Chrysopidae). *Bulletin of the Museum of Comparative Zoology* 135:215–238. creativecommons.org/licenses/by-nc-sa/3.0/
- Adams PhA. 1978. New species of *Hypochrysa* and a new subgenus and species of *Mallada* (Neuroptera, Chrysopidae). *Pan-Pacific Entomologist* 54:292–296.
- Adams PhA, Penny ND. 1992. New genera of Nothochysinae from South America. *Pan-Pacific Entomologist* 68:216–221.
- Ansorge J, Schlüter T. 1990. The earliest chrysopid: *Liassochrysa stigmatica* n. g., n. sp. from the Lower Jurassic of Dobbertin, Germany. *Neuroptera International* 6:87–93.
- Archibald SB, Makarkin VN. 2015. A new species of *Archaeochrysa* Adams (Neuroptera: Chrysopidae) from the early Eocene of Driftwood Canyon, British Columbia, Canada. *The Canadian Entomologist* 147:359–369. doi.org/10.4039/tce.2014.53

- Archibald SB, Makarkin VN. 2017. A new fossil green lacewing (Neuroptera: Chrysopidae) from the early Eocene Driftwood Canyon, Canada. *Zootaxa* 4324:397–400. doi.org/10.11646/zootaxa.4324.2.13
- Archibald SB, Makarkin VN, Greenwood DR, Gunnell GF. 2014. The Red Queen and court jester in green lacewing evolution: bat predation and global climate change. *Palaios* 29:185–191. doi.org/10.2110/palo.2013.089
- Breitkreuz LCV, Winterton SL, Engel MS. 2017. Wing tracheation in Chrysopidae and other Neuropterida (Insecta): a resolution of the confusion about vein fusion. *American Museum Novitates* 3890:1–44. doi.org/10.1206/3890.1
- Brooks SJ, Barnard PC. 1990. The green lacewings of the World: a generic review (Neuroptera: Chrysopidae). *Bulletin of the British Museum, (Natural History), Entomology* 59:117–286. creativecommons.org/licenses/by-nc-sa/4.0/
- Burmeister H. 1839. *Handbuch der Entomologie*, Reimer (publ.), Berlin, 1:1–400.
- Carpenter FM. 1935. Tertiary insects of the family Chrysopidae. *Journal of Paleontology* 9:259–271. jstor.org/stable/1298167
- Chen P-J, Shen Y-B. 1980. On the *Paraleptestheria menglaensis* fauna with reference to the age of the Lofochai Group. *Acta Palaeontologica Sinica* 19:182–189 (In Chinese with English Summary).
- Cockerell TDA. 1909. Two fossil Chrysopidae. *The Canadian Entomologist* 41:218–219. doi.org/10.4039/Ent41218-7
- Cockerell TDA. 1914. New and little-known insects from the Miocene of Florissant, Colorado. *Journal of Geology* 22:714–724. https://doi.org/10.1086/622186
- Doludenko, MP, Orlovskaya, E.R. 1976a. Jurassic floras of the Karatau Range, southern Kazakhstan. *Palaeontology* 19:627–640. go.palass.org/3zu
- Doludenko MP, Orlovskaya ER. 1976b. Jurassic flora of Karatau. Nauka, Moscow (In Russian).

- Doludenko MP, Sakulina GV, Ponomarenko AG. 1990. The geology and Late Jurassic fauna and flora of a unique locality Aulie (Karatau Mountains, Southern Kazakhstan) Geologicheskii Institut AN SSSR, Moscow (In Russian)
- Engel MS, Grimaldi DA. (2007). The neuropterid fauna of Dominican and Mexican amber (Neuropterida: Megaloptera, Neuroptera). American Museum Novitates 3587:1–58. doi.org/10.1206/3587.1
- Fu Y-z, Cai C-y, Huang D-y. 2019. A new palaeontinid (Insecta, Hemiptera, Cicadomorpha) from the Upper Jurassic Tiaojishan Formation of northeastern China and its biogeographic significance. *Journal of Paleontology* 94:513–520. doi: 10.1017/jpa.2019.95.
- Garzón-Orduña IJ, Winterton SL, Jiang Y-l, Breitzkreuz LCV, Duelli P, Engel MS, Penny ND, Tauber CA, Mochizuki A, Liu X-y. 2019. Evolution of green lacewings (Neuroptera: Chrysopidae): a molecular supermatrix approach. *Systematic Entomology* 44:499–513. doi.org/10.1111/syen.12339
- Guangdong Provincial Bureau of Geology and Mineral Resources. 1997 Lithologic Stratigraphy of Guangdong Province. Wuhan: China University of Geosciences Press:264 pp.
- Handschin E. 1937. Fossile Insekten aus Siebenbürgen. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 17:25–29.
- Hong Y-c. 1984. Tertiary insects of China. In: Li Y-t. (Ed.), *The Tertiary System of China*. Beijing: Geological Publishing House:308–309.
- Huang D-y. 2015. Yanliao biota and Yanshan movement. *Acta Palaeontologica Sinica* 54:501–546 [In Chinese].
- Huang D-y. 2016. *The Daohugou Biota*. Shanghai Scientific and Technical Publishers, Shanghai:332 pp. [In Chinese].

- Jepson JE, Makarkin VN, Coram RA. 2012. Lacewings (Insecta: Neuroptera) from the Lower Cretaceous Purbeck Limestone Group of southern England. *Cretaceous Research* 34:31–47. doi.org/10.1016/j.cretres.2011.10.001
- Khramov AV, Liu Q, Zhang H-C, Jarzembowski EA. 2015. Early green lacewings (Insecta: Neuroptera: Chrysopidae) from the Jurassic of China and Kazakhstan. *Papers in Palaeontology* 2:25–39. doi.org/10.1002/spp2.1024
- Kimmins DE. 1937. A new African *Hypochrysa* (Neuroptera). *Annals and Magazine of Natural History* (10) 19:307–308. doi.org/10.1080/00222933708655265
- Khramov AV. 2018. A new assemblage of Early Cretaceous green lacewings (Chrysopidae: Neuroptera) from Transbaikalia. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh* 107:195–202. doi.org/10.1017/S1755691017000342
- Kovanci B, Canbulat S. 2007. A new species of the genus *Nothochrysa* McLachlan 1868 from northwestern Turkey (Neuroptera: Chrysopidae) with a key to western Palaearctic species. *Annales de la Société Entomologique de France* (N.S.) 43:165–168. doi.org/10.1080/00379271.2007.10697507
- Lin Q-b. 1977. Fossil insects of Yunnan. In: Nanjing Institute of Geology and Palaeontology (Ed.), *Mesozoic fossils of Yunnan*. Science Press, Beijing: 373–384.
- Lin Q-b, Petrusevičius JF, Huang D-y, Nel A, Engel MS. 2010a. First fossil Calopterygoidea from Southeastern Asia (Odonata: Zygoptera): a new genus and species from the Paleogene of China. *Geobios* 43:349–353. doi.org/10.1016/j.geobios.2009.11.005
- Lin Q-b, Szwedo J, Huang D-y, Stroiński A. 2010b. *Weioboidae* fam. nov. of ‘higher’ Fulgoroidae (Hemiptera: Fulgoromorpha) from the Eocene deposits of Yunnan, China. *Acta Geologica Sinica* (English Edition) 84:751–755. doi.org/10.1111/j.1755-6724.2010.00263.x

- Lu X-m, Wang B, Ohl M, Liu X-y. 2018. The first green lacewing (Insecta: Neuroptera: Chrysopidae) from the mid-Cretaceous amber of Myanmar. *Zootaxa* 4399:563–570. doi.org/10.11646/zootaxa.4399.4.6
- Makarkin VN. 1991. Miocene insects (Neuroptera) from the northern Caucasus and Sikhote-Alin. *Paleontological Journal* 25:55–65.
- Makarkin VN. 1994. Upper Cretaceous Neuroptera from Russia and Kazakhstan. *Annales de la Société Entomologique de France (N.S.)* 30:283–292.
- Makarkin VN. 1997. Fossil Neuroptera of the Lower Cretaceous of Baisa, east Siberia. Part 3. Chrysopidae (Insecta). *Spixiana* 20:107–118.
- Makarkin VN. 2014. A new fossil green lacewing (Neuroptera: Chrysopidae) from the Eocene Tadushi Formation, eastern Sikhote-Alin. *Far Eastern Entomologist* 272:1–7.
- Makarkin VN, Archibald SB. 2013. A diverse new assemblage of green lacewings (Insecta, Neuroptera, Chrysopidae) from the Early Eocene Okanagan Highlands, Western North America. *Journal of Paleontology* 87:123–146. doi.org/10.1666/12-052R.1
- Makarkin VN, Yang Q, Shi C-f, Ren D. 2013. The presence of the recurrent veinlet in the Middle Jurassic Nymphidae (Neuroptera): a unique character condition in Myrmeleontoidea. *ZooKeys* 325:1–20. doi.org/10.3897/zookeys.325.5453
- Makarkin VN, Wedmann S, Weiterschan T. 2018. The first green lacewings from the late Eocene Baltic amber. *Acta Palaeontologica Polonica* 63:527–537. doi.org/10.4202/app.00504.2018
- Martins-Neto RG. 1997. Neurópteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. X - descrição de novos táxons (Chrysopidae, Babinskaiidae, Myrmeleontidae, Ascalaphidae e Psychopsidae). *Revista da Universidade de Guarulhos, Ciências Exatas e Tecnológicas* 2:68–83.

- Martins-Neto RG, Vulcano MA. 1988. Neurópteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, nordeste do Brasil. I - Família Chrysopidae. Anais da Academia Brasileira de Ciências 60:189–201.
- Martynov AV. 1927. Jurassic fossil Insects from Turkestan. 7. Some Odonata, Neuroptera, Thysanoptera. Izvestiya Akademii Nauk SSSR 21:757–768.
- Nanjing Institute of Geology and Palaeontology, Bureau of Geology and Mineral Resources of Yunnan Province and Bureau of Metallurgy of Yunnan Province Geological Company. 1975. Mesozoic Red Beds of Yunnan. Science Press, Beijing:1–201 (In Chinese).
- Nanjing Institute of Geology and Palaeontology, Bureau of Geology and Mineral Resources of Yunnan Province and Bureau of Metallurgy of Yunnan Province Geological Company. 1975. Mesozoic Red Beds of Yunnan. Science Press, Beijing: 1–201 (In Chinese).
- Nel A, Delclòs X, Hutin A. 2005. Mesozoic chrysopid-like Planipennia: a phylogenetic approach (Insecta: Neuroptera). Annales de la Société Entomologique de France (N.S.) 41:29–69. doi.org/10.1080/00379271.2005.10697440
- Nel. A, Auvray F. 2006. The oldest Vespinae from the Paleocene of Menat (France) (Hymenoptera: Vespidae). Zootaxa 1344:59–62. doi.org/10.11646/zootaxa.1344.1.5
- Nel A, Séméria Y. 1986. Une nouvelle espèce de chrysopide fossile du Stampien supérieur (Oligocène) d'Aix-en-Provence *Notochrysa stampieni* sp. nov. (Neuroptera, Chrysopidae, Notochrysinæ). Neuroptera International 4:23–30.
- New TR. 1980 A revision of the Australian Chrysopidae (Insecta: Neuroptera) Australian Journal of Zoology, Supplementary Series 77:1–143. <https://doi.org/10.1071/AJZS077>
- Ngô-Muller V, Blot K, Nel A. 2019. A new species of *Nothochrysa* from the latest Oligocene of France (Insecta: Neuroptera: Chrysopidae). Palaeoentomology 2:111–114. doi.org/10.11646/palaeoentomology.2.2.2

- Panfilov DV. 1980. Novye predstaviteli setcharokrylykh (Neuroptera) iz yury Karatau [New representative of Neuroptera from the Jurassic in the Karatau region.] pp. 82–111. In: Dolin VG, Panfilov DV, Ponomarenko AG, Pritykina LN. (eds). Iskopajemye nasekomyje mezozoja. [Fossil insects of the Mesozoic.] Akademiya nauk Ukrainskoj SSR, Institut Zoologii, Naukova Dumka, Kiev, 135 pp. [in Russian.]
- Peñalver E, Nel A, Martínez-Delclòs X. 1995. New Nothochrysinae from the Spanish Miocene (Neuroptera: Chrysopidae). Bulletin de la Société Entomologique de France 100:481–487. www.persee.fr/doc/bsef_0037-928x_1995_num_100_5_17194
- Ponomarenko AG. 1992. Novye setchatokrylye (Insecta: Neuroptera) iz mezozoya Mongolii. Novye Taksony Iskopaemykh Bespozvonochnykh Mongolii. Sovmestnaya Rossiisko-Mongol'skaya Paleontologicheskaya Ekspeditsiya 41:101–111.
- Ren D, Guo Z-G. 1996 On the new fossil genera and species of Neuroptera (Insecta) from the Late Jurassic of northeast China. Acta Zootaxonomica Sinica 21:461–479.
- Schlüter T. 1982. *Cimbrochrysa moleriensis* n. g. sp. nov. und *Hypochrysa hercyniensis* sp. nov., zwei fossile Chrysopidae-Arten (Insecta: Planipennia) aus dem europäischen Tertiär. Neues Jahrbuch für Geologie und Paläontologie, Monatshefte 1982:257–264.
- Scudder SH. 1885. 4. Classe Insecta. Insecten. Handbuch der Palaeontologie, I Abtheilung. Palaeozoologie 2:747–831.
- Scudder SH. 1890. The Tertiary insects of North America. Report of the United States Geological Survey of the Territories 13:1–734.
- Séméria Y, Nel A 1990. *Paleochrysopa monteilsensis* gen. et sp. nov., a new fossil of Chrysopidae from the Upper Eocene Formation of Monteils (France), with a review of known chrysopid fossils (Insecta: Neuroptera). Advances in Neuropterology. Proceedings of the Third International Symposium on Neuropterology: 27–32.

- Shen Y-b, Chen P-j. 1984. Tertiary conchostracans of China. In: Li Y-t. (ed.), The Tertiary System of China. Beijing: Geological Publishing House: 306–308.
- Shen Y-b., Gallego OF, Buchheim HP, Biaggi RE. 2006. Eocene conchostracans from the Laney Member of the Green River Formation, Wyoming, USA. *Journal of Paleontology* 80:447–454. doi.org/10.1666/0022-3360(2006)80[447:ECFTLM]2.0.CO;2
- Statz S. 1936. Über neue Funde von Neuropteren, Panorpaten und Trichopteren aus dem Tertiären schiefen von Rott am Siebengebirge. *Decheniana* 93:208–255.
- Tauber CA. 2014. Nothochrysinæ (Neuroptera: Chrysopidae): new larval description and generic synonymy, with a consideration of generic relationships. *Psyche* 2014 (ID 839261):1–10. doi.org/10.1155/2014/839261
- Tauber CA. 2019. South American Nothochrysinæ (Neuroptera, Chrysopidae): I. Description of *Nothochrysa ehrenbergi* sp. nov. *ZooKeys* 86:1–18. doi.org/10.3897/zookeys.866.35394
- Tauber CA. 2019. South American Nothochrysinæ (Neuroptera, Chrysopidae): II. Redescription of *Leptochrysa prisca* Adams & Penny. *ZooKeys* 866:19–38. doi.org/10.3897/zookeys.866.35396
- Tjeder B. 1966. Neuroptera-Planipennia. The lace-wings of Southern Africa. 5. Family Chrysopidae. pp. 228–534. In: Hanström B, Brinck P, Rudebec G. (eds.). *South African Animal Life*, Swedish Natural Science Research Council, Stockholm, 12.
- Vasilikopoulos, A., Misof, B., Meusemann, K., Lieberz, D., Flouri, T., Beutel, R.G., Niehuis, O., Wappler, T., Rust, J., Peters, R.S., Donath, A., Podsiadlowski, L., Mayer, C., Bartel, D., Böhm A, Liu, S-l, Kapli P, Greve C, Jepson JE, Liu X-y, Zhou, X, Aspöck H, Aspöck U. 2020. An integrative phylogenomic approach to elucidate the evolutionary history and divergence times of Neuropterida (Insecta: Holometabola). *BMC Evolutionary Biology* 20:1–24. doi.org/10.1186/s12862-020-01631-6

- Wang Y-q, Li Q, Bai B, Jin X, Mao F-y, Meng J. 2019. Paleogene integrative stratigraphy and timescale of China. *Science China Earth Science* 62:287–309.
- Wappler T, Currano ED, Wilf P, Rust J, Labandeira CC. 2009. No post-Cretaceous ecosystem depression in European forests? Rich insect-feeding damage on diverse middle Palaeocene plants, Menat, France. *Proceedings of Royal Society of London B* 276:4271–4277. doi.org/10.1098/rspb.2009.1255
- Willmann R. 1993. Insekten aus der Fur-Formation von Dänemark (Moler, ob. Paleozän/unt. Eozän? 8. Zwei neue Vertreter der Chrysopidae (Neuroptera). *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 1993:239–245.
- Willmann R, Brooks SJ. 1991. Insekten aus der Fur-Formation von Dänemark (Moler, ob. Paleozän/unt. Eozän ?). 6. Chrysopidae (Neuroptera). *Meyniana* 43:125–135.
- Winterton SL, Gillung JP, Garzón-Orduña IJ, Badano D, Breitkreuz LCV, Duelli P, Engel MS, Liu X-g, Machado RJP, Mansell M, Mochizuki A, Penny ND, Tauber CA, Oswald JD. 2019. Evolution of green lacewings (Neuroptera: Chrysopidae): an anchored phylogenomics approach. *Systematic Entomology* 44:514–526. https://doi.org/10.1111/syen.12347
- Yang J-K, Hong Y-c. 1990. *Drakochrysa*, an Early Cretaceous new genus of Chrysopidae (Insecta: Neuroptera) from Laiyang basin, Shandong Province. *Geoscience - Journal of Graduate School, China University of Geosciences* 4:15–26.
- Yang Q, Makarkin VN, Ren D 2012. New fossil Mesochrysopidae (Neuroptera) from the Mesozoic of China. *Zootaxa* 3597:1–14. doi.org/10.11646/zootaxa.3597.1.1
- Zhang X-q, Li G, Yang R-l, Li H-m. 2008. Palaeogene ostracods from the Sanshui Basin of Guangdong. *Acta Micropalaeontologica Sinica* 25:235–265.
- Zhang X-q, Zhou X-p, Chen X-y. 1993. Atlas of Cretaceous to Tertiary strata classification and correlation in the Sanshui Basin of South China. Marine Press, Beijing: 183 pp.

Zhou H-m, Xiao L, Dong Y-x, Wang C-z, Wang F-z, Ni P-z. 2009. Geochemical and geochronological study of the Sanshui basin bimodal volcanic rock suite, China: Implications for basin dynamics in southeastern China. Journal of Asian Earth Sciences 34:178–189. doi.org/10.1016/j.jseaes.2008.05.001

Table 1. Fossil record of the family Chrysopidae

Figure 1. Fossil locality map.

Figure 2. *Sinonhothochrysa zhang*i Huang and Nel gen. et sp. nov., holotype NIGP175154. (A) photograph of habitus; (B) forewing reconstruction; arrows: course of forewing CuP; (C) hind wing reconstruction. Scale bars = 1 mm.

Figure 3. *Sinonhothochrysa zhang*i Huang and Nel gen. et sp. nov., holotype NIGP175154. Photographs. (A) thorax with bases of wings; (B) detail of bases of wings. Scale bars = 1 mm (A), 0.5 mm (B)

figure 1

figure 2

figure 3

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Subfamily / Tribe / Genus	Species	Distribution	Age
Subfamily uncertain			
<i>Cimbrochrysa</i> Schlüter, 1982			
	<i>C. moleriensis</i> Schlüter, 1982	Denmark	Ypresian
<i>Cretachrysa</i> Makarkin, 1994			
	<i>C. martynovi</i> Makarkin, 1994	Russian Federation	Campanian
<i>Danochrysa</i> Willmann, 1993			
	<i>D. madseni</i> Willmann, 1993	Denmark	Ypresian
<i>Paralembochrysa</i> Nel et al., 2005			
	<i>P. splendida</i> Nel et al., 2005	China	Aptian
<i>Stephenbrooksia</i> Willmann, 1993			
	<i>S. multifurcata</i> Willmann, 1993	Denmark	Ypresian
Chrysopinae Schneider, 1851			
<i>Paleochrysopa</i> Séméria & Nel, 1990			
	<i>P. monteilsensis</i> Séméria & Nel, 1990	France	Priabonian
<i>Pseudosencera</i> Makarkin et al., 2018			
	<i>P. baltica</i> Makarkin et al., 2018	Russian Federation	Priabonian
Chrysopini Schneider 1851			
<i>Chrysopa</i> Leach, 1815			
	<i>C. glaesaria</i> Engel & Grimaldi, 2007	Dominican Republic	Burdigalian/Langhian
	<i>C. martynovae</i> Makarkin, 1991	Russian Federation	Miocene
	<i>C. miocenea</i> Makarkin, 1991	Russian Federation	Miocene
	<i>C. sarmatica</i> Handschin, 1937	Romania	middle Serravallian
	<i>C. stavropolitana</i> Makarkin, 1991	Russian Federation	Miocene
	<i>C. vetula</i> Engel & Grimaldi 2007	Dominican Republic	Burdigalian/Langhian
Leucochrysini Adams 1978			
<i>Leucochrysa</i> McLachlan, 1868			
	<i>L. (Nodita) prisca</i> Engel & Grimaldi, 2007	Dominican Republic	Burdigalian/Langhian
Limaiinae Martins-Neto and Vulcano, 1988			
<i>Aberrantochrysa</i> Khramov, 2018			
	<i>A. buryatica</i> Khramov, 2018	Russian Federation	Aptian
	<i>A. pulchella</i> Khramov, 2018	Russian Federation	Aptian
<i>Araripechrysa</i> Martins-Neto & Vulcano, 1988			
	<i>A. magnifica</i> Martins-Neto & Vulcano, 1988	Brazil	Aptian
<i>Baisochrysa</i> Makarkin, 1997			
	<i>B. multinervis</i> Makarkin, 1997	Russian Federation	Aptian
	<i>B. pumila</i> Khramov et al., 2015	Kazakhstan	Callovian/Oxfordian
<i>Drakochrysa</i> Yang & Hong, 1990			
	<i>D. sinica</i> Yang & Hong, 1990	China	Aptian
<i>Limaia</i> Martins-Neto & Vulcano, 1988			
	<i>L. adicotomica</i> Martins-Neto, 1997	Brazil	Aptian

	<i>L. conspicua</i> Martins-Neto & Vulcano, 1999	Brazil	Aptian
	<i>Mesypochrysa</i> Martynov, 1927		
	<i>M. angustialata</i> Makarkin, 1997	Russian Federation	Aptian
	<i>M. cannabina</i> Khramov, 2018	Russian Federation	Aptian
	<i>M. chrysopa</i> Makarkin, 1997	Russian Federation	Aptian
	<i>M. chrysopoides</i> Ponomarenko, 1992	Mongolia	Aptian
	<i>M. curvimedia</i> Makarkin, 1997	Russian Federation	Aptian
	<i>M. falcata</i> Makarkin, 1997	Russian Federation	Aptian
	<i>M. intermedia</i> Panfilov, 1980	Kazakhstan	Callovian/Oxfordian
	<i>M. latipennis</i> Martynov, 1927	Kazakhstan	Callovian/Oxfordian
	<i>M. magna</i> Makarkin, 1997	Russian Federation	Aptian
	<i>M. minima</i> Makarkin, 1997	Russian Federation	Aptian
	<i>M. miniscula</i> Ren & Guo, 1996	China	Aptian
	<i>M. minuta</i> Jepson et al., 2012	United Kingdom	Berriasian
	<i>M. naranica</i> Khramov, 2018	Russian Federation	Aptian
	<i>M. polyclada</i> Panfilov, 1980	Kazakhstan	Callovian/Oxfordian
	<i>M. polyneura</i> Ren & Guo, 1996	China	Aptian
	<i>M. reducta</i> Panfilov, 1980	Kazakhstan	Callovian/Oxfordian
	<i>M. sinica</i> Khramov et al., 2015	China	Callovian/Oxfordian
	<i>Parabaisochrysa</i> Lu et al., 2018		
	<i>P. xingkei</i> Lu et al., 2018	Myanmar	Cenomanian
	<i>Protochrysa</i> Willmann & Brooks, 1991		
	<i>P. aphrodite</i> Willmann & Brooks, 1991	Denmark	Ypresian
	<i>P. fuscobasalis</i> Makarkin & Archibald, 2013	Canada	Ypresian
	Nothochrysinæ Navas 1910		
	<i>Adamschrysa</i> Makarkin & Archibald, 2013		
	<i>A. aspera</i> Makarkin & Archibald, 2013	Canada	Ypresian
	<i>A. wilsoni</i> Makarkin & Archibald, 2013	Canada	Ypresian
	<i>Archaeochrysa</i> Adams, 1967		
	<i>A. creedei</i> Carpenter, 1935	USA	Oligocene
	<i>A. fracta</i> Cockerell, 1914	USA	earliest Oligocene
	<i>A. paranervis</i> Adams, 1967	USA	earliest Oligocene
	<i>A. profracta</i> Makarkin & Archibald, 2013	Canada	Ypresian
	<i>A. sanikwa</i> Archibald & Makarkin, 2015	Canada	Ypresian
	<i>Asiachrysa</i> Makarkin, 2014		
	<i>A. tadushiella</i> Makarkin, 2014	Russian Federation	Ypresian
	<i>Dyspetochrysa</i> Adams, 1967		
	<i>D. vetuscula</i> Scudder, 1890	USA	earliest Oligocene
	<i>Hypochrysa</i> Gould, 1861		
	<i>H. hercyniensis</i> Schlüter, 1982	Germany	Pliocene
	<i>Lithochrysa</i> Carpenter, 1935		
	<i>L. borealis</i> Archibald & Makarkin, 2017	Canada	Ypresian
	<i>L. wickhami</i> Cockerell, 1914	USA	earliest Oligocene
	<i>Nothochrysa</i> MacLachlan, 1868		

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	<i>N. praeclara</i> Statz, 1936	Germany	Chattian
	<i>N. stampieni</i> Nel & Séméria, 1986	France	Chattian
	<i>N. praeclara</i> Statz, 1936	Germany	Chattian
<i>Okanaganochrysa</i> Makarkin & Archibald, 2013			
	<i>O. coltsunae</i> Makarkin & Archibald, 2013	Canada	Ypresian
<i>Palaeochrysa</i> Scudder, 1890			
	<i>P. concinnula</i> Cockerell, 1909	USA	Chadronian
	<i>P. stricta</i> Scudder, 1890	USA	Chadronian
<i>Pronothochrysa</i> Peñalver et al., 1995			
	<i>P. vivesi</i> Peñalver et al., 1995	Spain	Miocene
<i>Pseudochrysopa</i> Makarkin & Archibald, 2013			
	<i>P. harveyi</i> Makarkin & Archibald, 2013	Canada	Ypresian
<i>Sinonothochrysa</i> gen. nov.			
	<i>Sinonothochrysa eocenica</i> sp. nov.	China	middle Paleocene
<i>Tribochrysa</i> Scudder, 1885			
	<i>T. firmata</i> Scudder, 1890	USA	Chadronian
	<i>T. inequalis</i> Scudder, 1885	USA	Chadronian

Locality / Formation	Reference
Gullerup / Fur Formation	Schlüter, 1982
Obeshchayushchiy / Ola Formation	Makarkin, 1994
Sundby Klint, Mors / Fur Formation	Willmann, 1993
Huangbanjigou, Chaomidian Village / Yixian Formation	Nel et al., 2005
Ejerslev Industrial Pit / Fur Formation	Willmann, 1993
Alès-Monteils / Monteils Formation	Séméria and Nel, 1990
Baltic amber	Makarkin et al., 2018
Dominican amber	Engel and Grimaldi, 2007
Vishnevaya Balka creek, outcrop 4, Stavropol	Makarkin, 1991
Vishnevaya Balka creek, outcrop 4, Stavropol	Makarkin, 1991
Magyar Sáros	Handschin, 1937
Vishnevaya Balka creek, outcrop 3 bed IX, Stavropol	Makarkin, 1991
Dominican amber	Engel and Grimaldi, 2007
Dominican amber	Engel and Grimaldi, 2007
Khasurty	Khramov, 2018
Khasurty	Khramov, 2018
Crato / Crato Formation	Martins-Neto and Vulcano, 1999
Baissa / Zaza Formation	Makarkin, 1997
Karatau-Mikhailovka / Karabastau Formation	Khramov et al., 2015
84GT1, west Tuanwang Village / Laiyang Formation	Yang and Hong, 1990
Crato / Crato Formation	Martins-Neto, 1997

Crato / Crato Formation	Martins-Neto, 1988
Baissa / Zaza Formation	Makarkin, 1997
Khasurty	Khramov, 2018
Baissa / Zaza Formation	Makarkin, 1997
Bon Tsagaan / Dzun-Bain Formation	Ponomarenko, 1992
Baissa / Zaza Formation	Makarkin, 1997
Baissa / Zaza Formation	Makarkin, 1997
Karatau-Mikhailovka / Karabastau Formation	Panfilov, 1980
Galkino / Karabastau Formation	Martynov, 1927
Baissa / Zaza Formation	Makarkin, 1997
Baissa / Zaza Formation	Makarkin, 1997
Near Chaomidian Village / Yixian Formation	Ren and Guo, 1996
Durlston Bay, Bed DB175, Swanage (Manchester Mus) / Durlston Formation	Jepson et al., 2012
Khasurty	Khramov, 2018
Karatau-Mikhailovka / Karabastau Formation	Panfilov, 1980
Near Chaomidian Village / Yixian Formation	Ren and Guo, 1996
Karatau-Mikhailovka / Karabastau Formation	Panfilov, 1980
Daohugou / Daohugou Formation	Khramov et al., 2015
Burmese amber	Lu et al., 2018
Fur Formation	Willmann and Brooks, 1991
McAbee	Makarkin and Archibald, 2013
McAbee	Makarkin and Archibald, 2013
McAbee	Makarkin and Archibald, 2013
Creede / Creede Formation	Carpenter, 1935
Wilson Ranch, Florissant	Cockerell, 1914
Florissant, Station 14 / Florissant Formation	Adams, 1967
McAbee	Makarkin and Archibald, 2013
Driftwood Canyon	Archibald and Makarkin, 2015
Tadushi locality / Tadushi Formation	Makarkin, 2014
Florissant / Florissant Formation	Scudder, 1890
Willershausen clay pit	Schlüter, 1982
Driftwood Canyon	Archibald and Makarkin, 2017
Florissant / Florissant Formation	Cockerell, 1914

Rott / Rott Formation	Statz, 1936
Plâtrières, Aix-en-Provence	Nel and Séméria, 1986
Rott / Rott Formation	Statz, 1936
McAbee	Makarkin and Archibald, 2013
Florissant / Florissant Formation	Cockerell, 1909
Florissant / Florissant Formation	Scudder, 1890
La Rinconada, Ribesalbes	Peñalver et al., 1995
Driftwood Canyon	Makarkin and Archibald, 2013
Sanshui District / Buxin Formation	This paper
Florissant / Florissant Formation	Scudder, 1890
Florissant / Florissant Formation	Scudder, 1885