

## Mass-Radius relationships of small, highly irradiated exoplanets with small water mass fractions

Hugo Vivien, Artem Aguichine, Olivier Mousis, M. Deleuil, Emmanuel Marcq

► **To cite this version:**

Hugo Vivien, Artem Aguichine, Olivier Mousis, M. Deleuil, Emmanuel Marcq. Mass-Radius relationships of small, highly irradiated exoplanets with small water mass fractions. European Planetary Science Congress. EPSC 2021, Sep 2021, Virtual Meeting, France. 10.5194/epsc2021-249 . insu-03317763

**HAL Id: insu-03317763**

**<https://hal-insu.archives-ouvertes.fr/insu-03317763>**

Submitted on 7 Aug 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



EPSC Abstracts

Vol. 15, EPSC2021-249, 2021, updated on 01 Aug 2021

<https://doi.org/10.5194/epsc2021-249>

European Planetary Science Congress 2021

© Author(s) 2021. This work is distributed under the Creative Commons Attribution 4.0 License.



## Mass-Radius relationships of small, highly irradiated exoplanets with small water mass fractions

Hugo Vivien<sup>1</sup>, Artem Agüichine<sup>1</sup>, Olivier Mousis<sup>1</sup>, Magali Deleuil<sup>1</sup>, and Emmanuel Marcq<sup>2</sup>

<sup>1</sup>Aix Marseille Univ, CNRS, CNES, LAM, Marseille, France

<sup>2</sup>LATMOS/IPSL, UVSQ Université Paris-Saclay, Sorbonne Université, CNRS, Guyancourt, France

This work aims to determine the mass-radius relationships of highly irradiated ( $500 < T_{\text{irr}} < 2000 \text{ K}$ ) small planets ( $0.2 < M < 2.3 M_{\oplus}$ ) with water contents up to 5%. To do so, we coupled an internal model of small terrestrial planets (Brugger et al., 2017) to the atmosphere model elaborated by Marcq et al. (2017, 2019), following the approach depicted in Agüichine et al. (2021) and Mousis et al. (2020).

We show that these planets, even with small water contents, can become strongly inflated and produce large radii for small masses. We also show that strongly irradiated small planets cannot sustain their atmospheres due to the lack of hydrostatic stability, implying they cannot preserve any hydrosphere. The temperature and the water mass fraction are the key parameters controlling the extent of inflation and the thickness of the supercritical layer. An important amount of water also leads to the contraction of the rocky interior. However, the composition of the rocky interior only has a limited impact on the final mass-radius relationship, and barely impacts the behavior of the hydrosphere.