Mass-Radius relationships of small, highly irradiated exoplanets with small water mass fractions

Hugo Vivien, Artem Aguichine, Olivier Mousis, M. Deleuil, Emmanuel Marcq

► To cite this version:

Hugo Vivien, Artem Aguichine, Olivier Mousis, M. Deleuil, Emmanuel Marcq. Mass-Radius relationships of small, highly irradiated exoplanets with small water mass fractions. European Planetary Science Congress. EPSC 2021, Sep 2021, Virtual Meeting, France. 10.5194/epsc2021-249 . insu-03317763

HAL Id: insu-03317763

<https://hal-insu.archives-ouvertes.fr/insu-03317763>

Submitted on 7 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

Mass-Radius relationships of small, highly irradiated exoplanets with small water mass fractions

Hugo Vivien¹, Artem Aguichine¹, Olivier Mouis¹, Magali Deleuil¹, and Emmanuel Marcq²

¹Aix Marseille Univ, CNRS, CNES, LAM, Marseille, France

²LATMOS/IPSL, UVSQ Université Paris-Saclay, Sorbonne Université, CNRS, Guyancourt, France

This work aims to determine the mass-radius relationships of highly irradiated ($500 < T_{\text{irr}} < 2000 \text{ K}$) small planets ($0.2 < M < 2.3 M_{\oplus}$) with water contents up to 5%. To do so, we coupled an internal model of small terrestrial planets (Brugger et al., 2017) to the atmosphere model elaborated by Marcq et al. (2017, 2019), following the approach depicted in Aguichine et al. (2021) and Mouis et al. (2020).

We show that these planets, even with small water contents, can become strongly inflated and produce large radii for small masses. We also show that strongly irradiated small planets cannot sustain their atmospheres due to the lack of hydrostatic stability, implying they cannot preserve any hydrosphere. The temperature and the water mass fraction are the key parameters controlling the extent of inflation and the thickness of the supercritical layer. An important amount of water also leads to the contraction of the rocky interior. However, the composition of the rocky interior only has a limited impact on the final mass-radius relationship, and barely impacts the behavior of the hydrosphere.