

Molecular hydrogen in oxidized atmospheres of terrestrial exoplanets: Implications for water and oxygen formation

Thomas Drant, Nathalie Carrasco, Zoé Perrin, Ludovic Vettier, Thomas Gautier, Shang-Min Tsai, Kevin Heng

► **To cite this version:**

Thomas Drant, Nathalie Carrasco, Zoé Perrin, Ludovic Vettier, Thomas Gautier, et al.. Molecular hydrogen in oxidized atmospheres of terrestrial exoplanets: Implications for water and oxygen formation . European Planetary Science Congress. EPSC 2021, Sep 2021, Virtual Meeting, France. 10.5194/epsc2021-503 . insu-03317872

HAL Id: insu-03317872

<https://hal-insu.archives-ouvertes.fr/insu-03317872>

Submitted on 8 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Molecular hydrogen in oxidized atmospheres of terrestrial exoplanets : Implications for water and oxygen formation

Thomas Drant¹, Nathalie Carrasco¹, Zoe Perrin¹, Ludovic Vettier¹, Thomas Gautier¹, Shang-Min Tsai², and Kevin Heng³

¹IPSL, University of Paris Saclay, LATMOS, France (thomas.drant@latmos.ipsl.fr)

²Department of Physics, University of Oxford, Oxford, England

³Center for Space and Habitability, Bern University, Bern, Switzerland

1. Introduction

Hydrogen emission by degassing magma is likely to have a significant impact on the H₂ mixing ratio in early atmospheres of terrestrial planets depending on the surface pressure, volcanic flux and oxygen fugacity of the magma [1]. This process can account for the presence of a few percent of hydrogen in the early atmospheres of terrestrial exoplanets.

The formation of water using a N₂-CO₂-H₂ mixture was identified in a previous experimental study focusing on the early Earth with nitrogen as a dominant gas and an initial H₂ mixing ratio fixed at 4% [2]. We now explore various amounts of hydrogen in the mixture to cover the unknown parameters regarding surface emission and to highlight the role played by hydrogen in the chemistry of these exoplanetary atmospheres.

2. Experimental method

The PAMPRE (french for "aerosol production in microgravity using a reactive plasma") experimental setup [3] is a plasma reactor used to simulate the photochemistry at low pressure (around 1 hPa) in planetary atmospheres. The CO₂-N₂-H₂ mixture is injected in the reactor with various N₂ to H₂ abundance ratio and a fixed abundance of 70% for carbon dioxide. The molecular hydrogen mixing ratio is varied from 5% down to 0.5% in volume. The gas phase in the PAMPRE reactor is analyzed using a quadrupole mass spectrometer (Hiden Analyticals) to identify the species formed once the plasma is turned on.

3. Experimental results




Figure 1. Production of water and oxygen with 0.5% H₂







Figure 2. Production of water and oxygen with 5% H₂

The significant oxygen production associated to a hydrogen ratio of 0.5% is reduced drastically when increasing the hydrogen mixing ratio by a factor of 10. The formation of oxygen via the HOx cycle well known in the case of Mars becomes secondary when the hydrogen mixing ratio increases. As a result, the production rate of the OH radical formed by (1) increases.



The radical OH then reacts by (2) and (3) which explains the significant formation of water with an initial hydrogen mixing ratio of 5%.



These new experiments put forward the presence of a hydrogen mixing ratio threshold over which the formation of water dominates the formation of oxygen. Two regimes can therefore be identified :

O₂-dominant production regime at low amounts of H₂ and H₂O-dominant production regime at higher amounts of H₂.

4. Conclusions and perspectives

For terrestrial exoplanets known to be in the habitable zone of their host star, the present study shows that molecular hydrogen in the atmosphere is critical to characterize their habitability as this particular molecule leads to an important formation of water in oxydized conditions. It is also shown that based on the hydrogen abundance in the atmosphere, two regimes can be identified and this particular process suggests that high amounts of hydrogen could lead to a significant abundance of water in these atmospheres even without liquid water at the surface. Quantification of relative concentrations using quadrupole mass spectrometry by a recently developed tool [4] is in progress and would allow to quantify the H₂ mixing ratio which separates the two regimes in our experimental conditions. Based on these experimental results, the impact of this process is now being studied using the chemical kinetics model VULCAN [5] taking into account vertical mixing and using a known star spectrum. The predictions obtained from the model will be used to predict future observations of these specific environments in the frame of JWST and ARIEL space missions.

Acknowledgements

NC acknowledges the financial support of the European Research Council (ERC Starting Grant PRIMCHEM, Grant agreement no. 636829).

References

- [1] Liggins, P., Shorttle, O., Rimmer, P., 2020. Can volcanism build hydrogen-rich early atmospheres ? *Earth and Planetary Science Letters* 550
- [2] Fleury, B., Carrasco, N., Marcq, E., Vettier, L., Määttänen, A., 2015. *The Astrophysical Journal Letters* 807
- [3] Szopa, C., Cernogora, G., Boufendi, L., Correia, J., Coll, P., 2006. *Planetary and Space Science* 54
- [4] Gautier, T., Serigano, J., Bourgalais, J., Hörst, S.M., Trainer, M.G., 2020. Decomposition of electron ionization mass spectra for space application using a Monte-Carlo approach. *Rapid Communications in Mass Spectrometry*
- [5] Tsai, S-M., Lyons, J.R., Grosheintz, L., Rimmer, P.B., Kitzmann, D., Heng, K., 2017. VULCAN : An Open-Source, Validated Chemical Kinetics Python Code for Exoplanetary Atmospheres. *The Astrophysical Journal Supplement Series* 228