



HAL
open science

The French Massif Central: a witness of successive weathering periods since the Early Cretaceous in the Alpine foreland

Augustin Dekoninck, Gilles Ruffet, Julien Baptiste, Robert Wyns, Eric Lasseur, Jean-Yves Roig, Johan Yans

► To cite this version:

Augustin Dekoninck, Gilles Ruffet, Julien Baptiste, Robert Wyns, Eric Lasseur, et al.. The French Massif Central: a witness of successive weathering periods since the Early Cretaceous in the Alpine foreland. *Geologica Belgica Meeting 2021, Sep 2021, Tervuren, Belgium*. pp.159-160. insu-03381898

HAL Id: insu-03381898

<https://insu.hal.science/insu-03381898>

Submitted on 18 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/355059637>

The French Massif Central: a witness of successive weathering periods since the Early Cretaceous in the Alpine foreland

Conference Paper · September 2021

CITATIONS

0

READS

33

7 authors, including:



Augustin Dekoninck

Royal Museum for Central Africa

52 PUBLICATIONS 119 CITATIONS

[SEE PROFILE](#)



Gilles Ruffet

French National Centre for Scientific Research

200 PUBLICATIONS 5,007 CITATIONS

[SEE PROFILE](#)



Julien Baptiste

French Geological Survey

7 PUBLICATIONS 22 CITATIONS

[SEE PROFILE](#)



Eric Lasseur

French Geological Survey

98 PUBLICATIONS 607 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Geochronology, petrology and source characteristics of Cenozoic magmatism in NE Turkey [View project](#)



Manganese occurrences in the Vosges Massif [View project](#)

The French Massif Central: a witness of successive weathering periods since the Early Cretaceous in the Alpine foreland

Augustin DEKONINCK^{*1}, Gilles RUFFET², Julien BAPTISTE³, Robert WYNS³, Eric LASSEUR³, Jean-Yves ROIG³, Johan YANS¹

1. *Université de Namur, ILEE, Département de géologie – Belgium*
(augustin.dekoninck@unamur.be)(johan.yans@unamur.be)

2. *CNRS (CNRS/INSU) et Université de Rennes 1, Géosciences (UMR6118) – France*
(gilles.ruffet@univ-rennes1.fr)

3. *Bureau de Recherches Géologiques et Minières (BRGM) – France*
(j.baptiste@brgm.fr)(e.lasseur@brgm.fr)(jy.roig@brgm.fr)(r.wyns@free.fr)

Many European uplands have remained topographic highs for protracted periods of weathering during the Mesozoic and Cenozoic. The long-term elevation position as low relief of these massifs acted as a source for adjacent sedimentary basins, their weathering playing a major role in the production of sediments available for erosion. Weathering profiles on these uplands are mostly incomplete/truncated since weathering products have often been washed away during repeated periods of erosions, leading to patchworks of remaining weathered rocks. In order to better address the events that have shaped these source areas, a complementary approach of both the source (upland) and sink (sedimentary basin) areas is crucial, but the record in the hinterland often remains partial due to lack of dating material and incomplete sequence.

The study presented here after is part of the BRGM-TOTAL *Source-to-Sink* project and aims at identifying key periods of exposure of the Massif Central (France). We used the combination of a careful petrogenesis and ⁴⁰Ar/³⁹Ar dating of K-bearing Mn oxides (coronadite group and romanechite). The French Massif Central hosts several Mn occurrences sporadically mined until the first half of the 20th Century. We have targeted five of these sites that contain dating minerals: (1) “Nontronais” (Haute-Dordogne; Mn-Fe; south-west), (2) Villerembert (Montagne Noire; Mn; south), (3) Vieussan (Montagne Noire; Mn; south), (4) Auxilhac (Lozère; Mn-Fe; south), and (5) Romanèche (Mn-F-Ba; east). The combined petrography, mineralogy and geochemistry of these Mn ores show that Mn deposits 1, 2 and 3 belong to a weathering crust model (“laterite”), sampling site 4 follows a karst-hosted model whereas sampling site 5 follows an epithermal model. In any of these cases, the dated K-Mn oxides were formed at the end of the weathering paragenetic sequence, except maybe for Romanèche which encompasses epithermal formation. This and their oxidizing precipitation environment support a formation under surface or sub-surface conditions. Therefore, their ages are instructive of periods of meteoric waters circulation under near-surface conditions. For the first time in these areas, the ⁴⁰Ar/³⁹Ar ages define at least four weathering periods: (i) Albian and older, (ii) Campanian, (iii) Oligocene-Early Miocene and (iv) Late Miocene to Quaternary. The French Massif Central is a key place in the West European landscape that better constrain the long-term evolution of weathering that has affected basement rocks and their cover. The Early Cretaceous period is well known in Western Europe for having provided various and thick weathering products (bauxite, kaolin, siderolithic). The Campanian

age fits well with the first slow Pyrenean compression that exposed the Montagne Noire (south Massif Central), whereas Oligocene-Quaternary weathering events correspond to widespread weathering phases and compressional phases of the Alpine and Pyrenees. The lack of any Eocene period in the age record might be due to erosional conditions that overcome weathering, as it corresponds to the paroxysmal phase of the Pyrenean compression and strong siliciclastic discharge from the Massif Central to the Aquitaine and Paris basins.

This work is funded and carried out in the framework of the BRGM-TOTAL project Source-to-Sink.