Journal of Paleontology

CAMBRIDGE UNIVERSITY PRESS

The termite genus *Glyptotermes* (Isoptera: Kalotermitidae) in Miocene amber from Ethiopia

Journal:	*Journal of Paleontology*
Manuscript ID	JPA-RA-2021-0060.R2
Manuscript Type:	Regular Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Bouju, Valentine; Université de Rennes 1, Géosciences Jouault, Corentin; Université de Rennes 1, Géosciences; Muséum National d'Histoire Naturelle, Institut de Systématique, Evolution, Biodiversité (ISYEB) Perrichot, Vincent; Université de Rennes 1, Géosciences
Taxonomy:	Insecta, Blattodea < Insecta, Isoptera
Broad Geologic Time:	Neogene < Cenozoic
Detailed Geologic Time:	Miocene < Neogene
Subject area Geographic Location:	Africa, Ethiopia
Abstract:	A new species of drywood termite (Isoptera: Kalotermitidae) is described from a nearly complete alate specimen preserved in early Miocene Ethiopian amber. *Glyptotermes abyssinicus* n. sp., is distinguished by its U-shaped head with 12-segmented antennae, the ocelli separated from eye margin, the right mandible with obtuse angle between apical and first marginal teeth, the left mandible with obtuse angle between apical and first + second marginal teeth, and its wing venation. This is the first termite reported from Ethiopian amber, and the fourth Miocene species of the extant genus *Glyptotermes*, together with species previously described from diatomites of China and amber from the Dominican

	Republic. As the oldest report of the genus known from Africa, *G. abyssinicus* n. sp. constitutes an interesting new record for the biogeographical history of the kalotermitid lineage.

SCHOLARONE™
Manuscripts

The termite genus *Glyptotermes* (Isoptera: Kalotermitidae) in Miocene amber from Ethiopia

Valentine Bouju[1], Corentin Jouault[1,2] and Vincent Perrichot[1*]

[1]Univ Rennes, CNRS, Géosciences Rennes, UMR 6118, 35000, Rennes, France.

[2]Muséum national d'Histoire naturelle, Institut de Systématique, Evolution, Biodiversité (ISYEB), CNRS, Sorbonne Université, EPHE, Université des Antilles, CP50, 57 rue Cuvier, 75005 Paris, France.

<valentine.bouju@univ-rennes1.fr> ; orcid.org/0000-0003-3185-5047

<jouaultc0@gmail.com>; orcid.org/0000-0002-3680-5172

<vincent.perrichot@univ-rennes1.fr>; orcid.org/0000-0002-7973-0430

Running Header: New Miocene drywood termite

Abstract.—A new species of drywood termite (Isoptera: Kalotermitidae) is described from a nearly complete alate specimen preserved in early Miocene Ethiopian amber. *Glyptotermes abyssinicus* n. sp., is distinguished by its U-shaped head with 12-segmented antennae, the ocelli separated from eye margin, the right mandible with obtuse angle between apical and first marginal teeth, the left mandible with obtuse angle between apical and first + second marginal teeth, and its wing venation. This is the first termite reported from Ethiopian amber, and the fourth Miocene species of the extant genus *Glyptotermes*, together with species previously described from diatomites of China and amber from the

For Review Only

Dominican Republic. As the oldest report of the genus known from Africa, *G. abyssinicus* n. sp. constitutes an interesting new record for the biogeographical history of the kalotermitid lineage.

UUID: http://zoobank.org/7670b045-fb31-4809-8116-4d14c4dd275b

Introduction

The Kalotermitidae, also named drywood termites for their high tolerance to dry conditions, is a monophyletic family consistently retrieved as sister to the Neoisoptera in recent molecular and morphological phylogenetic analyses (Engel et al., 2009; Bourguignon et al., 2015; Legendre et al., 2015; Bucek et al., 2019; Zhao et al., 2019; Jouault et al., 2021). While some discrepancies are observed regarding the divergence time estimates of several isopteran constitutive families, depending on the method or dataset used, molecular- and morphology-based analyses agree for an Early Cretaceous appearance of the kalotermitids (Bourguignon et al., 2015; Bucek et al., 2019; Jouault et al., 2021).

The comprehension of the Kalotermitidae have greatly benefited from the outstanding work of Kumar Krishna who proposed the first phylogenetic hypothesis and described tens of new species and genera, e.g., *Bifiditermes* Krishna, 1961, *Incisitermes* Krishna, 1961, *Postelectrotermes* Krishna, 1961, and the fossil species *Incisitermes peritus* Engel and Krishna, 2007, or *Cryptotermes glaesarius* Engel and Krishna, 2007.

The kalotermitids have a rather extensive fossil record, with 74 registered occurrences ranging from the Cretaceous to Quaternary periods (http://fossilworks.org/). The oldest representative of the family was described from the Brazilian Crato Formation, as *Cratokalotermes santanensis* Bechly, 2007, and the family seems to have diversified during the mid-Cretaceous, with various species

For Review Only

reported from Cenomanian Burmese amber (Cockerell, 1916, 1917; Engel et al., 2007; Poinar, 2009; Zhao et al., 2021).

Glyptotermes Froggatt, 1897 is a cosmopolitan genus principally represented in tropical environments and is the most speciose of kalotermitid genera, with at least 128 living species (Krishna et al., 2013; Scheffrahn, 2021). *Glyptotermes* is commonly found infesting sound or rotting dry wood, as well as wood scars in live trees (Scheffrahn et al., 2001; Engel and Krishna, 2007). Like all kalotermitids, *Glyptotermes* reproductives are likely to be trapped in resin during their swarming flight since they commonly land on tree trunks and rarely reach the ground, and in fact the genus is recorded by the numerous wings and winged specimens mentioned from many deposits worldwide (Cowie et al., 1990; Nel and Paicheler, 1993; Engel and Krishna, 2007; Krishna et al., 2013; Engel and Kaulfuss, 2017). As for many extant genera, the fossil record of *Glyptotermes* remains relatively scarce, with only four species described: *G. grimaldii* Engel and Krishna, 2007, and *G. paleoliberatus* Engel and Krishna, 2007, from early Miocene Dominican amber; *G. shandongianus* (Zhang, 1989), from Miocene diatomitic deposits of Shanwang, China (Zhang, 1989; Zhang et al., 1994); and *G. pusillus* (Heer, 1849), from Pleistocene or younger East African copal (Heer, 1849; Snyder, 1949; Krishna et al., 2013). The new specimen described herein represents the first fossil record of the genus, and one of the very few fossil kalotermitids, from Africa.

Material and methods

The studied specimen is preserved in a piece of amber originating from a deposit located in the North Shewa Zone of Ethiopia. The exact locality is unknown but is undoubtedly among the four localities reported by Bouju et al. (2021, fig. 2), as the material was accessed through an amber trader from Addis Ababa who obtains amber exclusively from those four localities. Apparently, the amber is dug

For Review Only

from the same geological layer in the four deposits. The amber bed corresponds to a siltstone situated within a series of Pre-Oligocene to Miocene basalts and ignimbrite (Coulié et al., 2003; Kieffer et al., 2004; Belay et al., 2009) that are exposed down the gorges of the Jema and Wenchit Rivers or their tributaries (Bouju and Perrichot, 2020). Unfortunately, no radiometric age could be obtained from the adjacent basaltic rocks sampled by two of us in 2019 from the locality of Woll (see Bouju et al., 2021: fig. 2). However, the geological data, combined with studies of the amber chemistry, organismic inclusions, and palynomorphs of the amber-bearing sediment, support an early Miocene age for the amber (Bouju and Perrichot, 2020; Bouju et al., 2021).

The specimen was found preserved in syninclusion with a tiny chalcid wasp (Pteromalidae?) in a piece of greenish amber. The raw amber piece was ground and polished using thin silicon carbide papers for an optimal observation of the insect specimens in frontal, dorsal, and ventral views. The amber piece was ultimately embedded in a block of epoxy resin (Araldite 2020) following the method in Sadowski et al. (2021, fig. 8) for consolidation and long-term conservation. Photographs and examination of the specimen were made using a Leica M205 C stereomicroscope equipped with a Leica DMC4500 digital camera. The series of photographs taken at multiple focal planes were stacked using Helicon Focus 6.7 software. Illustrations were made using Adobe Illustrator and Photoshop CC 2019 softwares. We follow the morphological terminology and classification of termites as presented in Krishna et al. (2013), except for the wing venation that follows Schubnel et al. (2019).

Repository and institutional abbreviations.—The type specimen examined in this study is deposited in the amber collection of the Geological Department and Museum of the University of Rennes (IGR), France, under the registration number IGR.ET2020/017.

Systematic paleontology

For Review Only

Order Blattodea Brunner von Wattenwyl, 1882

Infraorder Isoptera Brullé, 1832

Family Kalotermitidae Froggatt, 1897

Genus *Glyptotermes* Froggatt, 1897

Type species.—*Glyptotermes tuberculatus* Froggatt, 1897 by subsequent designation in Snyder, 1949: 45.

Glyptotermes abyssinicus new species

Figures 1–3

Holotype.—One alate specimen, IGR.ET2020/017, in amber from the lower Miocene (16–23 Ma) of the North Shewa Zone, Amhara and Oromia regions, Northwestern Plateau of Ethiopia.

Diagnosis.—Head U-shaped in posterodorsal view; eyes small, sub-circular and feebly protruding from head laterally; ocelli distinctly separated from eye margin, longest obliquely; antenna 12-segmented, AII longer than AIII, AIII subequal to slightly shorter than AIV; left mandible with apical tooth stout, angle between apical and first + second marginal teeth right to nearly obtuse; right mandible with long, stout apical tooth, angle between apical and first marginal teeth obtuse; fore wing scale not fully overlapping hind wing scale; fore wing with Sc and RA short, RP long with two anterior branches, M sclerotized, running very close to and parallel to RP to reach wing apex, with faint cross-veins between M and radial sector, CuA with at least 8–9 posterior branches.

For Review Only

Description.—Body 3.8 mm long (from tip of labrum to abdomen apex), 7.0 mm with wings (Fig. 1). Head (Fig. 2.1, 2.2) prognathous, 1.1 mm long and 0.65 mm wide excluding compound eyes, with straight sides, anterolateral corners slightly angulate, posterolateral corners rounded; compound eye sub-circular, slightly wider than long, ca. 0.25 mm wide, protruding from head, separated from posterior head margin by more than its length; ocelli white, ovoid (about 0.06 mm long and 0.05 mm wide), located above eye mid-length, very close to but not touching eye margin; fontanelle absent; antenna (Fig. 2.2) moniliform, with 12 antennomeres, AII longer than AIII, AIII subequal to slightly shorter than AIV, AIV shorter than AV, following antennomeres progressively increasing in length toward apex; left mandible with slightly obtuse to nearly right angle between apical and first + second marginal teeth, third marginal tooth hidden; right mandible with obtuse angle between apical and first marginal teeth; maxillary palps hidden by labial palps in frontal view; labial palps with 3 segments (Fig. 2.3); labrum about 0.18 mm long and 0.24 mm wide, posterior margin convex, anterior margin straight or nearly so; postclypeus slightly raised medially (Fig. 2.1, 2.3), rectangular and conspicuously wider than long (about 0.24 mm wide and 0.06 mm long).

Pronotum (Fig. 2.5) flat to slightly arched in lateral view, anterior margin slightly concave, sides rounded, slightly narrower than head. Legs slender; profemur *ca.* 0.5 mm long, protibia *ca.* 0.4 mm long, protarsus at least 0.2 mm long; protibia with three spurs f1, f2, f3; mesofemur *ca.* 0.5 mm long, mesotibia *ca.* 0.5 mm long, mesotarsus at least 0.4 mm long; mesotibia with three spurs m1, m2, m3 (Fig. 2.6); metafemur *ca.* 0.5 mm long, metatibia *ca.* 0.8 mm long, metatarsus at least 0.4 mm long; metatibia with three spurs h1, h2, h3; no additional spine on meso- and metatibiae; tarsi tetramerous, with apical tarsomere as long as or slightly longer than combined length of preceding ones (Fig. 2.7); arolium present (Fig. 2.8).

Wings (Fig. 3.1–3.5) 4.75 mm long (measured from suture to apex), membrane reticulate; fore wing scale 0.75 mm long, not covering hind wing scale, with humeral margin nearly straight, claval suture (CuP) straight, slightly curved distally, basal suture straight; fore wing with M and CuA veins

For Review Only

arising independently from inside the wing scale; Sc short and simple; RA and RP fused basally, RA short, RP long with 2 anterior branches or fusion point with C; M sclerotized, running to wing apex, close and parallel to RP, with at least 1 anterior, short branch located near wing apex, and maybe two unsclerotized anterior branches located respectively in basal and apical wing third; CuA unsclerotized, with 8–9 posterior branches and maybe two unsclerotized anterior branches; hind wing not describable; anal lobe absent on both wings.

Abdomen not fully preserved, ca. 1.4 mm long, with at least 6 to 7 observable segments (Fig. 1.1); abdominal segments apparently widest at mid-length. Cerci and styli not visible.

Etymology.—The specific epithet *abyssinicus* refers to the Abyssinian origin of the amber piece.

Remarks.—The new fossil does not fit in the early diverged termite families, including the extinct Cratomastotermitidae and Termopsidae, since it combines the following series of derived characters: a reduced wing venation (vs. more complete, RP with at least several branches and no costalisation of veins in early diverged families), 4-segmented tarsi (vs. 5-segmented, plesiomorphic character), a reduced tibial spur formula 3-3-3 (vs. enriched with additional spurs along tibiae), and a head wider than pronotum (vs. smaller than pronotum). Compared to most of the other families, the fossil lacks a fontanelle (apomorphic character of the Neoisoptera; Krishna et al. 2013). Following the key to the families based on imago characters by Krishna et al. (2013, p. 69), the new fossil keys out in the Kalotermitidae for its hind wing without anal lobe, tarsi 4-segmented, antenna 12-segmented, fontanelle absent, fore wing scale not overlapping hind wing scale, ocelli present, and pronotum flat. Using the key to the kalotermitid genera of Krishna et al. (2013: p. 75–76), the new fossil species keys out either in *Glyptotermes*, *Calcaritermes* Snyder, 1925, or *Proneotermes* Holmgren, 1911, depending on character states detailed in the sixth and thirteenth couplets but invisible on the fossil, i.e., the left mandibular portion posterior to the first + second marginal tooth. However, the specimen differs from *Calcaritermes* and *Proneotermes* by the 12 antennomeres (vs. 13–14 in *Calcaritermes*, 17–18 in

For Review Only

Proneotermes). Additionally, it differs from *Proneotermes* but concords with *Glyptotermes* by the small eye diameter (0.25 mm vs. 0.39–0.43 mm in *Proneotermes*) and the fore wing with M as heavily sclerotized as the radial sector, running very close to and parallel to the radial sector to reach apex of wing, with faint branches between M and radial sector, and radial sector very close to costal vein.

Glyptotermes abyssinicus n. sp. can be distinguished from the extant Neotropical and African *Glyptotermes* species, by the combination of characters stated in the diagnosis. More precisely, *G. abyssinicus* n. sp. differs from *G. adamsoni* Krishna and Emerson, 1962*, G. sicki* Krishna and Emerson, 1962, or the African *G. sinomalatus* Krishna and Emerson, 1962, by a U-shaped head (vs. circular or semi-circular in the other species). It differs from other species by the 12-segmented antennae (vs. 11-segmented in *G. marlatti* (Snyder, 1926) and *G. planus* (Snyder, 1925); more than 12-segmented in *G. amplus* Scheffrahn et al., 2001, *G. kawandae* Wilkinson, 1954, or *G. longuisculus* Krishna and Emerson, 1962.

Glyptotermes abyssinicus n. sp. can also be distinguished from other species by the antennae with AII longer than AIII (vs. AII shorter than AIII *in G. asperatus* (Snyder, 1926), *G. ignotus* Wilkinson, 1959, or *G. tuberifer* Krishna and Emerson, 1962; by measurements such as the body, head and wing length (*G. rotundifrons* Krishna and Emerson, 1962 and *G. suturis* (Snyder, 1925) are larger species with wider head and pronotum, and longer fore wings; *G. contracticornis* (Snyder, 1925) and *G. longipennis* Krishna and Emerson, 1962 are about twice as large as *G. abyssinicus* n. sp.); by the ocelli separated from the eye margin (vs. ocelli touching eye margin in *G. jurioni* Krishna and Emerson, 1962, *G. parki* Krishna and Emerson, 1962, *G. parvoculatus* Krishna and Emerson, 1962 or *G. truncatus* Krishna and Emerson, 1962.

Finally, *Glyptotermes abyssinicus* n. sp. differs from the fossil species *G. grimaldii* Engel and Krishna, 2007 and *G. paleoliberatus* Engel and Krishna, 2007 by a smooth head capsule (vs. granulose) and the ocelli separated from the eye margin (vs. touching). The old description of *G. pusillus* (Heer, 1849), from East African copal, challenges its comparison with the other *Glyptotermes* species.

For Review Only

However, it is clear that our specimen does not belong to *G. pusillus* owing to the differential of temporal range (Pleistocene vs. Early Miocene).

Discussion

The genus *Glyptotermes* has a worldwide distribution with a higher diversity in tropical areas (Scheffrahn et al., 2001; Krishna et al., 2013; Debelo and Degaga, 2014; Yashiro et al., 2019; Scheffrahn, 2021). Our current knowledge on the *Glyptotermes* fossil record, and its correlation with the distribution of the extant representatives of the genus, seems to indicate that *Glyptotermes* was also widely distributed in the tropical region during the Miocene period and may have experimented successive dispersal events during older geological periods (Krishna and Emerson, 1962; Emerson, 1969; Nel and Paicheler, 1993; Engel and Krishna, 2007).

The kalotermitids are generally of particular importance in extant and past ecosystems due to their biology. In fact, they are the only termites that do not need to construct their nest in contact with soil (a direct adaptation from their high tolerance to dryness). The kalotermitids can live exclusively within the wood and participate in its active degradation by consuming the trunk in which they construct their nest. Extant *Glyptotermes* species, however, are very tolerant to environmental conditions and can live in swamp forests or mangroves (Johnson et al., 1980; Tho, 1992). And according to Scheffrahn (2021), extant, Neotropical '*Glyptotermes* species have a rather high wood moisture requirement and, therefore, are not found in arid parts of the Neotropics'. The same was likely true for the Miocene species newly described here, as this accords with indications of a generally humid, tropical, forest environment with freshwater suggested by the palynomorphs associated with Ethiopian amber as well as by other organismal inclusions in amber (Bouju and Perrichot, 2020; Bouju et al., 2021).

For Review Only

Conclusion

Although Isoptera are relatively well known from the fossil record, numerous fossil species have been described based on compression or imprint fossils of isolated wings, challenging placements and questioning some systematic attributions (Krishna et al., 2013). The present description of a nearly complete, three-dimensional, *Glyptotermes* specimen provides elements for more comprehensive, morphological-based phylogenetic studies. Additionally, *Glyptotermes abyssinicus* n. sp. can be considered as the oldest and first « true » *Glyptotermes* fossil from Africa. This discovery confirms the presence of the genus since the Miocene period and will allow further inference on the biogeographical history of the genus. The genus *Chilgatermes* Engel et al., 2013 (Stolotermitidae), described from an Oligocene imprint fossil, was hitherto the only fossil termite known from Ethiopia. Therefore, the description of *Glyptotermes abyssinicus* n. sp. extends our knowledge of the Isoptera fossil diversity in Ethiopia.

Acknowledgments

We are grateful to Yale Goldman, Bloomfield, Connecticut, who provided the Ethiopian amber piece studied here. We thank Amde Zewdalem, Jacksonville, Florida, and Benyam Teferi, Addis Ababa, Ethiopia, for information and access to the amber deposits of the North Shewa Zone, Ethiopia; and the reviewers and editor for their helpful comments. Partial support for this work was provided by the Tellus-INTERRVIE programme of CNRS INSU (project AMBRAFRICA to V. Perrichot, 2019), enabling field work in Ethiopia for V. Bouju and V. Perrichot.

References

For Review Only

240 Bechly, G., 2007, Isoptera: termites, *in* Martill, D.M., Bechly, G., and Loveridge, R.F., eds., The Crato
241 Fossil Beds of Brazil: Window into an Ancient World: Cambridge, Cambridge University Press, p.
242 249–262.
243 Belay, T., Tesfay, I., Ayalew, A., Yohannes, G., Zewdie, T., Bekele, H., Tadesse, M., Demisse, T., and
244 Alemu, T., 2009, Geology of the Were-Ilu Area: Geological Survey of Ethiopia Memoir 25, 56 pp.
245 Bouju, V., and Perrichot, V., 2020, A review of amber and copal occurrences in Africa and their
246 paleontological significance: BSGF-Earth Sciences Bulletin, v. 191, 17, doi:10.1051/bsgf/2020018
247 Bouju, V., Feldberg, K., Kaasalainen, U., Schaefer-Verwimp, A., Hedenäs, L., Buck, W.R., Wang, B.,
248 Perrichot, V., and Schmidt, A.R., 2021, Miocene Ethiopian amber: a new source of fossil
249 cryptogams: Journal of Systematics and Evolution, doi:10.1111/jse.12796.
250 Bourguignon, T., Lo, N., Cameron, S.L., Šobotník, J., Hayashi, Y., Shigenobu, S., Watanabe, D.,
251 Roisin, Y., Miura, T., and Evans, T.A., 2015, The evolutionary history of termites as inferred from
252 66 mitochondrial genomes: Molecular Biology and Evolution, v. 32, p. 406–421.
253 Brunner von Wattenwyl, K., 1882, Prodromus der europäischen Orthopteren: Leipzig, W. Engelmann,
254 466 pp.
255 Brullé, G.A., 1832, Insectes. Expédition scientifique de Morée. Section des sciences physiques, Tome
256 III, Partie 1. Zoologie. Sect. 2. Des animaux articulés: Paris, F.G. Levrault, p. 1–29 + 64–395.
257 Bucek, A., Šobotník, J., He, S., Shi, M., McMahon, D.P., Holmes, E.C., Roisin, Y., Lo, N., and
258 Bourguignon, T., 2019, Evolution of termite symbiosis informed by transcriptome-based: Current
259 Biology, v. 29, p. 3728–3734.
260 Cockerell, T.D.A., 1916, Insects in Burmese amber: American Journal of Science, Series 4, v. 42, p.
261 135–138.
262 Cockerell, T.D.A., 1917, Insects in Burmese amber: Annals of the Entomological Society of America,
263 v. 10, p. 323–329.

For Review Only

Coulié, E., Quidelleur, X., Gillot, P.-Y., Courtillot, V., Lefèvre, J.-C., and Chiesa, S., 2003, Comparative K–Ar and Ar/Ar dating of Ethiopian and Yemenite Oligocene volcanism: implications for timing and duration of the Ethiopian traps: Earth and Planetary Science Letters, v. 206, p. 477–492.

Cowie, R.H., Wood, T.G., Barnett, E.A., Sands, W.A., and Black, H.I.J., 1990, A checklist of the termites of Ethiopia with a review of their biology, distribution and pest status: African Journal of Ecology, v. 28, p. 21–33.

Debelo, D.G., and Degaga, E.G., 2014, Termite species composition in the central rift valley of Ethiopia: Agriculture and Biology Journal of North America, v. 5, no. 3, p. 123–134.

Emerson, A.E., 1969, A revision of the Tertiary fossil species of the Kalotermitidae (Isoptera): American Museum Novitates, v. 2359, p. 1–57.

Engel, M.S., and Kaulfuss, U., 2017, Diverse, primitive termites (Isoptera: Kalotermitidae, incertae sedis) from the Early Miocene of New Zealand: Austral Entomology, v. 56, p. 94–103.

Engel, M.S., and Krishna, K., 2007, Drywood termites in Dominican amber (Isoptera: Kalotermitidae): Beiträge zur Entomologie, v. 57, p. 263–275.

Engel, M.S., Grimaldi, D.A., and Krishna, K., 2007, Primitive termites from the Early Cretaceous of Asia (Isoptera): Stuttgarter Beiträge zur Naturkunde B, v. 371, p. 1–32.

Engel, M.S., Grimaldi, D.A., and Krishna, K., 2009, Termites (Isoptera): their phylogeny, classification, and rise to ecological dominance: American Museum Novitates, v. 3650, p. 1–27.

Engel, M., Pan, A.D., and Jacobs, B.F., 2013, A termite from the Late Oligocene of northern Ethiopia: Acta Palaeontologica Polonica, v. 58, p. 331–334.

Froggatt, W.W., 1897, Australian Termitidae, Part II: Proceedings of the Linnean Society of New South Wales, v. 21, p. 510–552.

For Review Only

Heer, O., 1849, Die Insektenfauna der Tertiärgebilde von Oeningen und von Radoboj in Croatien. Zweiter Teil: Heuschrecken, Florfliegen, Aderflügler, Schmetterlinge und Fliegen. Leipzig, W. Engelmann, iv + 264 p.

Holmgren, N., 1911, Termitenstudien. 2. Systematik der Termiten. Die Familien Mastotermitidae, Protermitidae und Mesotermitidae: Kungliga Svenska Vetenskaps-Akademiens Handlingar, v. 46, no. 6, p. 1–86 + 6 pl.

Johnson, R.A., Lamb, R.W., Sands, W.A., Shittu, M.O., Williams, R.M.C., and Wood, T.G., 1980, A checklist of Nigerian termites (Isoptera) with brief notes on their biology and distribution: Nigerian Field, v. 45, p. 50–64.

Jouault, C., Legendre, F., Grandcolas, P., and Nel, A., 2021, Revising dating estimates and the antiquity of eusociality in termites using the fossilized birth-death process: Systematic Entomology, v. 46, p. 592–610.

Kieffer, B., Arndt, N., Lapierre, H., Bastien, F., Bosch, D., Pecher, A., Yirgu, G., Ayalew, D., Weis, D., Jerram, D.A., Keller, F., and Meugniot, C., 2004, Flood and shield basalts from Ethiopia: magmas from the African superswell: Journal of Petrology, v. 45, p. 793–834.

Krishna, K., 1961, A generic revision and phylogenetic study of the family Kalotermitidae (Isoptera): Bulletin of the American Museum of Natural History, v. 122, p. 303–408.

Krishna, K., and Emerson, A.E., 1962, New species of the genus *Glyptotermes* Froggatt from the Papuan, Oriental, Ethiopian, and Neotropical regions (Isoptera, Kalotermitidae): American Museum Novitates, v. 2089, p. 1–65.

Krishna, K., Grimaldi, D.A., Krishna, V., and Engel, M.S., 2013, Treatise on the Isoptera of the world: Bulletin of the American Museum of Natural History, v. 377, p. 1–2704.

For Review Only

309 Legendre, F., Nel, A., Svenson, G.J., Robillard, T., Pellens, R., and Grandcolas, P., 2015, Phylogeny of
310 Dictyoptera: dating the origin of cockroaches, praying mantises and termites with molecular data
311 and controlled fossil evidence: PLoS ONE, v. 10, e0130127, doi:10.1371/journal.pone.0130127
312 Nel, A., and Paicheler, J.C., 1993, Les Isoptera fossiles. Etat actuel des connaissances, implications
313 paléoécologiques et paléoclimatologiques (Insecta, Dictyoptera): Cahiers de Paléontologie, Paris,
314 CNRS Editions, p. 101–179.
315 Poinar, G., Jr., 2009, Description of an Early Cretaceous termite (Isoptera: Kalotermitidae) and its
316 associated intestinal protozoa, with comments on their co-evolution: Parasites & Vectors, v. 2, no.
317 12, doi:10.1186/1756-3305-2-12
318 Sadowski, E.M., Schmidt, A.R., Seyfullah, L.J., Solórzano-Kraemer, M.M., Neumann, C., Perrichot,
319 V., Hamann, C., Milke, R., and Nascimbene, P.C., 2021, Conservation, preparation and imaging of
320 diverse ambers and their inclusions: Earth-Science Reviews, 103653,
321 doi:10.1016/j.earscirev.2021.103653
322 Scheffrahn, R.H., 2021, The termite genus *Glyptotermes* (Isoptera: Kalotermitidae) from Paraguay:
323 ZooKeys, v. 1059, p. 23–33.
324 Scheffrahn, R.H., Su N.Y., and Křeček, J., 2001, *Glyptotermes amplus*, a new dampwood termite
325 (Isoptera: Kalotermitidae) from St. Lucia: Florida Entomologist, v. 84, p. 426–430.
326 Schubnel, T., Desutter-Grandcolas, L., Legendre, F., Prokop, J., Mazurier, A., Garrouste, R.,
327 Grandcolas, P., and Nel., A., 2019, To be or not to be: postcubital vein in insects revealed by
328 microtomography: Systematic Entomology, v. 45, p. 327–336.
329 Snyder, T.E., 1925, New American termites, including a new subgenus: Journal of the Washington
330 Academy of Sciences, v. 15, p. 152–162.
331 Snyder, T.E., 1926, New termites from Guatemala, Costa Rica, and Colombia: Journal of the
332 Washington Academy of Sciences, v. 16, p. 18–28.

For Review Only

333 Snyder, T.E., 1949, Catalogue of termites (Isoptera) of the world: Smithsonian Miscellaneous
334 Collections, v. 112, p. 1–490.
335 Tho, Y.P., 1992, Termites of peninsular Malaysia: Kuala Lumpur, Malaysia, Forest Research Institute
336 Malaysia (Malayan Forest Records, v. 36), 224 p.
337 Wilkinson, W., 1954, Three new species of Kalotermitidae (Isoptera) from East Africa: Proceedings of
338 the Royal Entomological Society of London, Series B, Taxonomy, v. 23, p. 75–82.
339 Wilkinson, W., 1959, Four new species of Kalotermitidae from East Africa (Isoptera): Proceedings of
340 the Royal Entomological Society of London, Series B, Taxonomy, v. 28, p. 61–72.
341 Yashiro, T., Takematsu, Y., Ogawa, N., and Matsuura, K., 2019, Taxonomic assessment of the termite
342 genus Neotermes (Isoptera: Kalotermitidae) in the Ryukyu-Taiwan Island arc, with description of a
343 new species: Zootaxa, v. 4604, p. 549–561.
344 Zhang, J.F., 1989, Fossil Insects from Shanwang, Shandong, China: Jinan, Shandong Science and
345 Technology Publishing House, 459 p. [in Chinese, with English summary]
346 Zhang, J.F., Sun, B., and Zhang, X., 1994, Miocene Insects and Spiders from Shanwang, Shandong:
347 Beijing, Science Press, 298 p. [in Chinese, with English summary]
348 Zhao, Z., Eggleton, P., Yin, X., Gao, T., Shih, C., and Ren, D., 2019, The oldest known mastotermitids
349 (Blattodea : Termitoidae) and phylogeny of basal termites: Systematic Entomology, v. 44, p. 612–
350 623.
351 Zhao, Z., Shih, C., Gao, T., and Ren, D., 2021, Termite communities and their early evolution and
352 ecology trapped in Cretaceous amber: Cretaceous Research, v. 117, 104612,
353 doi:10.1016/j.cretres.2020.104612
354
355

For Review Only

Figure captions

Figure 1. *Glyptotermes abyssinicus* n. sp., holotype IGR.ET2020/017, from early Miocene Ethiopian amber. (**1**) Habitus in dorsal view; (**2**) habitus in ventral view. Scale bars = 1 mm.

Figure 2. *Glyptotermes abyssinicus* n. sp., holotype IGR.ET2020/017, from early Miocene Ethiopian amber. (**1**) Head in dorsal view; (**2**) left antenna in dorsal view; (**3**) mouthparts in frontal view; (**4**) line drawing of mandibles in frontal view; (**5**) pronotum in anterodorsal view; (**6**) right mesotibia in ventral view; (**7**) left mesotarsus in ventral view; (**8**) right protarsus in ventral view. I–IV = tarsomeres 1–4; AII–AV = antennomeres 2–5; ar = arolium; la = left mandible apical tooth; lbr = labrum; lm1+2 = left mandible 1st + 2nd marginal tooth; lp = labial palp; m1, m2, m3 = mesotibial spurs 1–3; mp = maxillary palp; oc = ocellus; pc = postclypeus; ra = right mandible apical tooth; rm1 = right mandible 1st marginal tooth. Scale bars = 25 µm (**4**), or 100 µm (**6–8**), or 125 µm (**1–3, 5**).

Figure 3. *Glyptotermes abyssinicus* n. sp., holotype IGR.ET2020/017, from early Miocene Ethiopian amber. (**1**) Right hind and fore wing bases in dorsal view; (**2**) line drawing of right wing scales in dorsal view; (**3**) left fore wing scale in dorsal view; (**4**) left fore wing in dorsal view; (**5**) explanatory drawing of fore wing venation. BS = basal suture; CuA = anterior cubital vein; CuP = claval suture; fsc = fore wing scale; hsc = hind wing scale; M = medial vein; RA = anterior radial vein; RP = posterior radial vein; Sc = subcostal vein. Scale bars = 250 µm (**1–3**), or 750 µm (**4, 5**).

For Review Only




Figure 1. *Glyptotermes abyssinicus* n. sp., holotype IGR.ET2020/017, from early Miocene Ethiopian amber. (1) Habitus in dorsal view; (2) habitus in ventral view. Scale bars = 1 mm.

170x218mm (300 x 300 DPI)




Figure 2. *Glyptotermes abyssinicus* n. sp., holotype IGR.ET2020/017, from early Miocene Ethiopian amber. (1) Head in dorsal view; (2) left antenna in dorsal view; (3) mouthparts in frontal view; (4) line drawing of mandibles in frontal view; (5) pronotum in anterodorsal view; (6) right mesotibia in ventral view; (7) left mesotarsus in ventral view; (8) right protarsus in ventral view. I–IV = tarsomeres 1–4; AII–AV = antennomeres 2–5; ar = arolium; la = left mandible apical tooth; lbr = labrum; lm1+2 = left mandible 1st + 2nd marginal tooth; lp = labial palp; m1, m2, m3 = mesotibial spurs 1–3; mp = maxillary palp; oc = ocellus; pc = postclypeus; ra = right mandible apical tooth; rm1 = right mandible 1st marginal tooth. Scale bars = 125 µm (1–3, 5), or 100 µm (6–8).

170x225mm (300 x 300 DPI)




Figure 3. *Glyptotermes abyssinicus* n. sp., holotype IGR.ET2020/017, from early Miocene Ethiopian amber. (1) Right hind and fore wing bases in dorsal view; (2) same as 1, but colour-marked to highlight the wing scales; (3) left fore wing scale in dorsal view; (4) left fore wing in dorsal view; (5) explanatory drawing of fore wing venation. BS = basal suture; CuA = anterior cubital vein; CuP = claval suture; fsc = fore wing scale; hsc = hind wing scale; M = medial vein; RA = anterior radial vein; RP = posterior radial vein; Sc = subcostal vein. Scale bars = 250 μm (1–3), or 750 μm (4).

179x133mm (600 x 600 DPI)