EGU21-15707

<https://doi.org/10.5194/egusphere-egu21-15707>

EGU General Assembly 2021

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.

Why switchbacks may be related to solar granulation

Naïs Fargette¹, Benoit Lavraud^{1,2}, Alexis Rouillard¹, Victor Réville¹, Tai Phan³, Stuart D. Bale³, Thierry Dudok De Wit⁴, Clara Froment⁴, Justin Kasper⁵, Jasper S. Halekas⁶, Philippe Louarn¹, Anthony W. Case⁷, Kelly E. Korreck⁷, Davin E. Larson³, David Malaspina^{8,9}, Marc Pulupa³, Michael L. Stevens⁷, Phyllis L. Whittlesey³, and Matthieu Berthomier¹⁰

¹Institut de Recherche en Astrophysique et Planétologie, CNRS, UPS, CNES, Toulouse, France (nais.fargette@irap.omp.eu)

²Laboratoire d'Astrophysique de Bordeaux, CNRS, Bordeaux, France

³Space Sciences Laboratory, University of California, Berkeley, Berkeley, CA, USA

⁴LPC2E, CNRS and University of Orléans, Orléans, France

⁵Climate and Space Sciences and Engineering, University of Michigan, Ann Arbor, MI, US

⁶Department of Physics and Astronomy, University of Iowa, Iowa City, Iowa, United States

⁷Smithsonian Astrophysical Observatory, Cambridge, Massachusetts, US

⁸Department of Astrophysical and Planetary Sciences, University of Colorado, Boulder, CO, USA

⁹Laboratory for Atmospheric and Space Physics, University of Colorado, Boulder, CO, USA

¹⁰Laboratoire de Physique des Plasmas, CNRS, Sorbonne Université, Ecole Polytechnique, Observatoire de Paris, Université Paris-Saclay, Paris, France

Parker Solar Probe data below 0.3 AU have revealed a near-Sun magnetic field dominated by Alfvénic structures that display back and forth reversals of the radial magnetic field. They are called magnetic switchbacks, they display no electron strahl variation consistent with magnetic field foldings within the same magnetic sector, and are associated with velocity spikes during an otherwise calmer background. They are thought to originate either at the photosphere through magnetic reconnection processes, or higher up in the corona and solar wind through turbulent processes.

In this work, we analyze the spatial and temporal characteristic scales of these magnetic switchbacks. We define switchbacks as a deviation from the Parker spiral direction and detect them automatically through perihelia encounters 1 to 6. We analyze the solid angle between the magnetic field and the Parker spiral both over time and space. We perform a fast Fourier transformation to the obtained angle and find a periodical spatial variation with scales consistent with solar granulation. This suggests that switchbacks form near the photosphere and may be caused, or at least modulated, by solar convection.