



HAL
open science

Using the 3D MOCAGE CTM to simulate the chemistry of halogens in the volcanic plume of Etna's eruption in December 2018 at the regional scale

Herizo Narivelo, Virginie Marécal, Paul David Hamer, Luke Surl, Tjarda J Roberts, Mickaël Bacles, Simon Warnach, Thomas Wagner

► To cite this version:

Herizo Narivelo, Virginie Marécal, Paul David Hamer, Luke Surl, Tjarda J Roberts, et al.. Using the 3D MOCAGE CTM to simulate the chemistry of halogens in the volcanic plume of Etna's eruption in December 2018 at the regional scale. vEGU21 - EGU General Assembly 2021, Apr 2021, Online, France. 10.5194/egusphere-egu21-12205 . insu-03559302

HAL Id: insu-03559302

<https://insu.hal.science/insu-03559302>

Submitted on 7 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

EGU21-12205

<https://doi.org/10.5194/egusphere-egu21-12205>

EGU General Assembly 2021

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



Using the 3D MOCAGE CTM to simulate the chemistry of halogens in the volcanic plume of Etna's eruption in December 2018 at the regional scale

Herizo Narivelo¹, Virginie Marécal¹, Paul David Hamer², Luke Surl^{3,4}, Tjarda Roberts^{3,4}, Mickaël Bacles¹, Simon Warnach⁵, and Thomas Wagner⁵

¹Centre National de Recherches Météorologiques (CNRM), Toulouse, France

²Norsk institutt for luftforskning (NILU), Kjeller, Norway

³Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E), Orléans, France

⁴Laboratoire Atmosphères, Milieux, Observations Spatiales (LATMOS), Paris, France

⁵Max-Planck-Institut für Chemie (MPI-C), Mainz, Germany

Volcanoes emit different gaseous species, SO₂ and in particular halogen species especially bromine and chlorine compounds. In general, halogens play an important role in the atmosphere by contributing to ozone depletion in the stratosphere (WMO Ozone assessment, 2018) and by modifying air composition and oxidizing capacity in the troposphere (Von Glasow et al. 2004). The halogen species emitted by volcanoes are halides. The chemical processing occurring within the plume leads to the formation of BrO from HBr following the 'bromine explosion' mechanism as evidenced from both observations and modelling (e.g., Bobrowski et al. Nature, 2003; Roberts et al., Chem. Geol. 2009). Oxidized forms of chlorine and bromine are modelled to be formed within the plume due to the heterogeneous reaction of HOBr with HCl and HBr, forming BrCl and Br₂ that photolyses and produces Br and Cl radicals. So far, modelling studies were mainly focused on the very local scale and processes occurring within a few hours after eruption.

In this study, the objective is to go a step further by analyzing the impact at the regional scale over the Mediterranean basin of a Mt Etna eruption event. For this, we use the MOCAGE model (Guth et al., GMD, 2016), a chemistry transport model run with a resolution of 0.2°x 0.2°, to quantify the impacts of the halogens species emitted by the volcano on the tropospheric composition. We have selected here the case of the eruption of Mount Etna around Christmas 2018 characterised by large amounts of emissions over several days (Calvari et al., remote sensing 2020; Corradini et al., remote sensing 2020). The results show that MOCAGE represents rather well the chemistry of the halogens in the volcanic plume because it established theory of plume chemistry. The bromine explosion process takes place on the first day of the eruption and even more strongly the day after, with a rapid increase of the in-plume BrO concentrations and a corresponding strong reduction of ozone and NO₂ concentrations.

We also compared MOCAGE results with the WRF-CHEM model simulations for the same case study. We note that the tropospheric column of BrO and SO₂ in the two models have the same

order of magnitude with more rapid bromine explosion occurring in WRF-CHEM simulations. Finally, we compared the MOCAGE results to tropospheric columns of BrO and SO₂ retrieved from TROPOMI spaceborne instrument.