EGU21-14686

<https://doi.org/10.5194/egusphere-egu21-14686>

EGU General Assembly 2021

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.

Encounter of Parker Solar Probe and a Comet-like Object During Their Perihelia: Simulations and Measurements

Jiansen He¹, Bo Cui¹, Liping Yang², Chuanpeng Hou¹, Lei Zhang¹, Wing-Huen Ip³, Yingdong Jia⁴, Chuanfei Dong⁵, Die Duan¹, Qiugang Zong¹, Stuart Bale^{6,7,8}, Marc Pulupa⁶, John Bonnell⁶, Thierry Dudok de Wit¹⁰, Keith Goetz¹¹, Peter Harvey⁶, Robert MacDowall¹², and David Malaspina¹³

¹School of Earth and Space Sciences, Peking University, Beijing, 100871, P. R. China

²State Key Laboratory of Space Weather, National Space Science Center, Chinese Academy of Sciences, Beijing, 100190, P. R. China

³Institute of Astronomy, National Central University, Taoyuan, 32001, Taiwan, R.O. China

⁴IGPP and EPSS, University of California, Los Angeles, CA 90095, USA

⁵Department of Astrophysical Sciences, Princeton University, Princeton, NJ 08544, USA

⁶Space Sciences Laboratory, University of California, Berkeley, CA 94720-7450, USA

⁷Physics Department, University of California, Berkeley, CA 94720-7300, USA

⁸The Blackett Laboratory, Imperial College London, London, SW7 2AZ, UK

¹⁰LPC2E, CNRS and University of Orleans, Orleans, France

¹¹School of Physics and Astronomy, University of Minnesota, Minneapolis, MN 55455, USA

¹²Solar System Exploration Division, NASA/Goddard Space Flight Center, Greenbelt, MD, 20771

¹³Laboratory for Atmospheric and Space Physics, University of Colorado, Boulder, CO 80303, USA

Parker Solar Probe (PSP) aims at exploring the nascent solar wind close to the Sun. Meanwhile, PSP is also expected to encounter small objects like comets and asteroids. In this work, we survey the ephemerides to find a chance of recent encounter, and then model the interaction between released dusty plasmas and solar wind plasmas. On 2019 September 2, a comet-like object 322P/SOHO just passed its perihelion flying to a heliocentric distance of 0.12 au, and swept by PSP at a relative distance as close as 0.025 au. We present the dynamics of dust particles released from 322P, forming a curved dust tail. Along the PSP path in the simulated inner heliosphere, the states of plasma and magnetic field are sampled and illustrated, with the magnetic field sequences from simulation results being compared directly with the in-situ measurements from PSP. Through comparison, we suggest that 322P might be at a deficient activity level releasing limited dusty plasmas during its way to becoming a “rock comet”. We also present images of solar wind streamers as recorded by WISPR, showing an indication of dust bombardment for the images superposed with messy trails. We observe from LASCO coronagraph that 322P was transiting from a dimming region to a relatively bright streamer during its perihelion passage, and simulate to confirm that 322P was flying from relatively faster to slower solar wind streams, modifying local plasma states of the streams.