



HAL
open science

Secondary bremsstrahlung X-rays emitted during TGF propagation

Nini Lykke Berge, Sebastien Celestin

► **To cite this version:**

Nini Lykke Berge, Sebastien Celestin. Secondary bremsstrahlung X-rays emitted during TGF propagation. 20th EGU General Assembly, Apr 2018, Vienne, Austria. pp.15089. insu-03566321

HAL Id: insu-03566321

<https://hal-insu.archives-ouvertes.fr/insu-03566321>

Submitted on 11 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License



Secondary bremsstrahlung X-rays emitted during TGF propagation

Nini Lykke Berge and Sebastien Celestin

LPC2E, University of Orleans, CNRS, France (nini.berge@cnsr-orleans.fr)

Terrestrial gamma ray flashes (TGFs) are high-energy bursts of photons bright enough to be routinely observed from space. They originate in thunderstorms in association with lightning activity. As the gamma rays propagate through the atmosphere, interactions of photons with air molecules produce secondary electrons. When trapped around geomagnetic field lines, some of these electrons can be observed as terrestrial electron beams (TEBs) at satellite altitude. Most of secondary electrons do not make it as far as space; rather they lose their energy through interactions with the atmosphere. Through bremsstrahlung, the secondary electrons also produce another population of photons in the X-ray energy range.

The satellite Taranis (CNES) and the observatory ASIM (ESA) have been specifically designed to study TGFs and other lightning-related phenomena, and will have X- and gamma ray detection capabilities.

Through Monte Carlo simulations of particle propagation and interaction with the atmosphere, we model the production and properties of secondary electrons and the induced second generation of photons. We aim to predict how secondary X-rays might be detected by space-based instruments. We find that especially at the lower end of the TGF energy range, the secondary X-rays make a noticeable contribution to the spectrum.