



**HAL**  
open science

## Solar wind turbulence at kinetic scales inside 1 AU: HELIOS revisited

Vamsee Krishna Jagarlamudi, Olga Alexandrova, Thierry Dudok de Wit,  
Milan Maksimovic

► **To cite this version:**

Vamsee Krishna Jagarlamudi, Olga Alexandrova, Thierry Dudok de Wit, Milan Maksimovic. Solar wind turbulence at kinetic scales inside 1 AU: HELIOS revisited. 20th EGU General Assembly, 2018, Vienne, Austria. pp.9438. insu-03566332

**HAL Id: insu-03566332**

**<https://hal-insu.archives-ouvertes.fr/insu-03566332>**

Submitted on 11 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License



## **Solar wind turbulence at kinetic scales inside 1 AU: HELIOS revisited**

Vamsee krishna Jagarlamudi (1), Olga Alexandrova (2), Thierry Dudok de Wit (3), and Milan Maksimovic (4)

(1) LPC2E, LESIA - Observatoire de Paris, France (vjagarla@cnsr-orleans.fr), (2) LESIA - Observatoire de Paris (olga.alexandrova@obspm.fr), (3) University of Orléans, LPC2E (ddwit@cnsr-orleans.fr), (4) LESIA - Observatoire de Paris (milan.maksimovic@obspm.fr)

We use Search Coil Magnetometer data on Helios 1 in 1975 to study the radial evolution of solar wind magnetic field turbulence in the kinetic range, at scales smaller than the ion characteristic scales. Helios being a closest mission to Sun till date offers a unique opportunity to study turbulence in different regions of the inner Heliosphere. The magnetic spectra from 7Hz and up to  $\sim 680$  Hz has been studied. We see the signatures of quasi-parallel whistlers waves in the magnetic spectra at a fraction of the local electron cyclotron frequency. We relate the presence of whistlers and their evolution with radial distance to different physical parameters such as solar wind type, plasma beta and features of the electron distribution functions.