



**HAL**  
open science

## Time of flight dispersed and repetitive ion structures in the diffuse auroral zone originating from 1-2 Re altitudes

Jean-André Sauvaud, Dominique Delcourt, Michel Parrot, Denis Payan,  
Emmanuel Penou

### ► To cite this version:

Jean-André Sauvaud, Dominique Delcourt, Michel Parrot, Denis Payan, Emmanuel Penou. Time of flight dispersed and repetitive ion structures in the diffuse auroral zone originating from 1-2 Re altitudes. 20th EGU General Assembly, 2018, Vienne, Austria. pp.2062. insu-03566339

**HAL Id: insu-03566339**

**<https://hal-insu.archives-ouvertes.fr/insu-03566339>**

Submitted on 11 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License



## **Time of flight dispersed and repetitive ion structures in the diffuse auroral zone originating from 1-2 Re altitudes**

Jean-André Sauvaud (1), Dominique Delcourt (2), Michel Parrot (3), Denis Payan (4), and Emmanuel Penou (1)

(1) IRAP, GPPS, Toulouse Cedex 4, France (jsauvaud@irap.omp.eu), (2) LPP, école polytechnique, Palaiseau, France, (3) LPC2E, Orléans, France, (4) CNES, Toulouse, France

The AMBRE experiment onboard the ocean topography mapper JASON-3 aims at measuring auroral particle precipitation using two top-hat analyzers for electrons and ions in the 20 eV-28 keV energy range. The JASON-3 spacecraft that has a nearly circular orbit at an altitude of 1336 km with an inclination of  $66^\circ$ , at times probes the equatorward part of the auroral oval in a nearly tangentially manner upon leaving the outer radiation belt. In this region of space, during periods of enhanced geomagnetic activity with small or moderate storms, AMBRE detected recurrent ion bands/micro-injections with energies in the 200 eV-28 keV range and which exhibit clear time of flight dispersions. Ray tracing using single trajectory computations suggests that these ions are launched from a source located in the 8000-12000 km altitudinal range and subsequently propagate downward toward the ionosphere. More radial orbits show that the ion bands are detected inside the diffuse auroral zone up to the encounter of auroral arcs. Such observations of dispersed downflowing ions are new and we argue that these structures are produced by localized wave-particle interactions.