



HAL
open science

Estimation of Physical Properties of Streamers in Transient Luminous Events from Non-Steady State Optical Emissions

Kevin Mohand A. Ihaddadene, Sebastien Celestin

► **To cite this version:**

Kevin Mohand A. Ihaddadene, Sebastien Celestin. Estimation of Physical Properties of Streamers in Transient Luminous Events from Non-Steady State Optical Emissions. EGU General Assembly 2016, Apr 2016, Vienne, Austria. insu-03573509

HAL Id: insu-03573509

<https://hal-insu.archives-ouvertes.fr/insu-03573509>

Submitted on 15 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License



Estimation of Physical Properties of Streamers in Transient Luminous Events from Non-Steady State Optical Emissions

Kevin Mohand A. Ihaddadene and Sebastien Celestin

LPC2E, University of Orleans, CNRS, Orleans, France (mohand.ihaddadene@cnsr-orleans.fr)

Optical emissions from sprite streamers are used to estimate peak electric fields and electron energies [e.g., Kuo et al., *GRL*, 32, L19103, 2005; Adachi et al., *GRL*, 33, L17803, 2006]. It has been shown that significant correction factors need to be used to account for the spatial shift between distributions of optical emissions in streamers and peak electric fields in their heads [Celestin and Pasko, *GRL*, 37, L07804, 2010]. The latter study involved the excited species $N_2(C_3\Pi_u)$ and $N_2^+(B_2\Sigma_u^+)$, whose populations are considered to be in steady state. The species $N_2(C_3\Pi_u)$ and $N_2^+(B_2\Sigma_u^+)$ are responsible for the second positive ($2PN_2$) and first negative ($1NN_2^+$) band systems of N_2 and N_2^+ , respectively.

In this work, we show how this technique can be extended to non-steady state optical emissions, such as those produced by $N_2(a^1\Pi_g)$ and $N_2(B^3\Pi_g)$ in the form of Lyman-Birge-Hopfield (LBH) and first positive ($1PN_2$) band systems, respectively. Additionally, we simulate numerically downward propagating sprite streamers and their optical emissions for the following band systems: $1PN_2$, $2PN_2$, LBH, and $1NN_2^+$, and show how they relate to specific physical properties. This study particularly focuses on improving analysis of observational results from the future missions ASIM (ESA) and TARANIS (CNES) that will detect various optical emissions produced by transient luminous events in the nadir.