IASI observations at the city scale

Sarah Safieddine, Maya George, Cathy Clerbaux, Ana Claudia Parracho,
Anne Boynard, Juliette Hadji-Lazaro, Lieven Clarisse, Simon Whitburn,
Pierre-François Coheur

► To cite this version:

Sarah Safieddine, Maya George, Cathy Clerbaux, Ana Claudia Parracho, Anne Boynard, et al.. IASI observations at the city scale. EGU General Assembly 2020, May 2020, Online, Unknown Region. 10.5194/egusphere-egu2020-18569 . insu-03573583

HAL Id: insu-03573583

<https://hal-insu.archives-ouvertes.fr/insu-03573583>

Submitted on 14 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EGU2020-18569

<https://doi.org/10.5194/egusphere-egu2020-18569>

EGU General Assembly 2020

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.

IASI observations at the city scale

Sarah Safieddine¹, Maya George¹, Cathy Clerbaux^{1,2}, Ana Paracho¹, Anne Boynard¹, Juliette Hadji-Lazaro¹, Lieven Clarisse², Simon Whitburn², and Pierre-Francois Coheur²

¹LATMOS/IPSL, Sorbonne Université, UVSQ, CNRS, Paris, France (sarah.safieddine@latmos.ipsl.fr)

²Université libre de Bruxelles (ULB), Spectroscopy, Quantum Chemistry and Atmospheric Remote Sensing (SQUARES), Brussels, Belgium

IASI is a versatile mission, allowing the measurement of both meteorological parameters such as temperature and atmospheric composition for infrared absorbing species. With its long observation record and frequent overpasses, IASI is able to follow changes at different spatial scales. We studied IASI's capability to track the anthropogenic signature associated with large cities, both in terms of temperature fingerprint (urban heat islands) and carbon monoxide (CO) content, a good tracer of human activity (transport, heating, and industrial activities). For this study we averaged the IASI data available since the launch of the first IASI, in order to increase the signal to noise, and allow discriminating the city from its surroundings. For skin temperatures we show that some cities experience much warmer temperatures than nearby rural areas, with day and night differences, whereas other urban areas appear as cold urban islands when surrounded by deserts. Examples will be shown and compared with MODIS observations. For CO emitted by human activities, we identified some cities that stand out from their background, and were able to compare their CO associated signatures with measurements provided by other available spaceborne instruments such as Mopitt and TROPOMI.