



**HAL**  
open science

## Herschel/HIFI deepens the circumstellar NH<sub>3</sub> enigma

K. M. Menten, F. Wyrowski, J. Alcolea, E. de Beck, L. Decin, A. P. Marston,  
V. Bujarrabal, J. Cernicharo, C. Dominik, K. Justtanont, et al.

► **To cite this version:**

K. M. Menten, F. Wyrowski, J. Alcolea, E. de Beck, L. Decin, et al.. Herschel/HIFI deepens the circumstellar NH<sub>3</sub> enigma. *Astronomy and Astrophysics*, EDP Sciences, 2010, 521, 10.1051/0004-6361/201015108 . insu-03625638

**HAL Id: insu-03625638**

**<https://hal-insu.archives-ouvertes.fr/insu-03625638>**

Submitted on 31 Mar 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License

LETTER TO THE EDITOR

## ***Herschel/HIFI deepens the circumstellar NH<sub>3</sub> enigma*<sup>\*,\*\*</sup>**

K. M. Menten<sup>1</sup>, F. Wyrowski<sup>1</sup>, J. Alcolea<sup>2</sup>, E. De Beck<sup>3</sup>, L. Decin<sup>3,4</sup>, A. P. Marston<sup>5</sup>, V. Bujarrabal<sup>6</sup>, J. Cernicharo<sup>7</sup>, C. Dominik<sup>5,8</sup>, K. Justtanont<sup>9</sup>, A. de Koter<sup>5,10</sup>, G. Melnick<sup>11</sup>, D. A. Neufeld<sup>12</sup>, H. Olofsson<sup>9,13</sup>, P. Planesas<sup>6,15</sup>, M. Schmidt<sup>14</sup>, F. L. Schöier<sup>9</sup>, R. Szczerba<sup>14</sup>, D. Teyssier<sup>5</sup>, L. B. F. M. Waters<sup>4,3</sup>, K. Edwards<sup>16,17</sup>, M. Olberg<sup>9,17</sup>, T. G. Phillips<sup>18</sup>, P. Morris<sup>19</sup>, M. Salez<sup>20,21</sup>, and E. Caux<sup>22,23</sup>

(Affiliations are available on page 5 of the online edition)

Received 31 May 2010 / Accepted 29 June 2010

### ABSTRACT

**Context.** Circumstellar envelopes (CSEs) of a variety of evolved stars have been found to contain ammonia (NH<sub>3</sub>) in amounts that exceed predictions from conventional chemical models by many orders of magnitude.

**Aims.** The observations reported here were performed in order to better constrain the NH<sub>3</sub> abundance in the CSEs of four, quite diverse, oxygen-rich stars using the NH<sub>3</sub> ortho  $J_K = 1_0-0_0$  ground-state line.

**Methods.** We used the Heterodyne Instrument for the Far Infrared aboard *Herschel* to observe the NH<sub>3</sub>  $J_K = 1_0-0_0$  transition near 572.5 GHz, simultaneously with the ortho-H<sub>2</sub>O  $J_{K_a,K_c} = 1_{1,0}-1_{0,1}$  transition, toward VY CMa, OH 26.5+0.6, IRC+10420, and IK Tau. We conducted non-LTE radiative transfer modeling with the goal to derive the NH<sub>3</sub> abundance in these objects' CSEs. For the last two stars, Very Large Array imaging of NH<sub>3</sub> radio-wavelength inversion lines were used to provide further constraints, particularly on the spatial extent of the NH<sub>3</sub>-emitting regions.

**Results.** We find remarkably strong NH<sub>3</sub> emission in all of our objects with the NH<sub>3</sub> line intensities rivaling those of the ground state H<sub>2</sub>O line. The NH<sub>3</sub> abundances relative to H<sub>2</sub> are very high and range from  $2 \times 10^{-7}$  to  $3 \times 10^{-6}$  for the objects we have studied.

**Conclusions.** Our observations confirm and even deepen the circumstellar NH<sub>3</sub> enigma. While our radiative transfer modeling does not yield satisfactory fits to the observed line profiles, it does lead to abundance estimates that confirm the very high values found in earlier studies. New ways to tackle this mystery will include further *Herschel* observations of more NH<sub>3</sub> lines and imaging with the Expanded Very Large Array.

**Key words.** stars: AGB and post-AGB – supergiants – circumstellar matter

### 1. Introduction

Ammonia (NH<sub>3</sub>) was the first polyatomic molecule detected in an astronomical object (Cheung et al. 1968). It is ubiquitous in dark, dense interstellar cloud cores and an eminently useful thermometer of these regions (Walmsley & Ungerechts 1983; Danby et al. 1988). This, aided by the easy observability of its inversion lines – many of the astronomically most important ones crowd around 1.3 cm wavelength (24 GHz frequency) – make NH<sub>3</sub> one of the most frequently observed interstellar molecules (Ho & Townes 1983).

NH<sub>3</sub> has also been detected toward a still limited, but diverse number of CSEs around evolved stars, first using infrared (IR) heterodyne absorption spectroscopy toward the high mass-loss asymptotic giant branch (AGB), extreme carbon star CW Leo (= IRC+10216; Betz et al. 1979). In addition, absorption was found toward a number of oxygen-rich objects that included the long-period variable (LPV) *o* Ceti (Betz & Goldhaber 1985) and the super- or even hyper-luminous objects VY CMa and IRC+10420 (McLaren & Betz 1980; Monnier et al. 2000). The last study finds that around VY CMa the NH<sub>3</sub> is forming at  $\geq 40$  stellar radii away from the star, where dust formation has well started.

Contemporaneously, several radio inversion lines were detected also toward IRC+10216 (Kwok et al. 1981;

Bell et al. 1982; Nguyen-Q-Rieu et al. 1984). Later on, high-velocity cm-wavelength NH<sub>3</sub> emission plus absorption was found toward the bipolar protoplanetary nebulae (PPNe) CRL 2688 and CRL 618 (Truong-Bach et al. 1988; Martin-Pintado & Bachiller 1992; Truong-Bach et al. 1996). Toward CRL 618, P Cygni profiles are observed with a full width at zero power (FWZP) of  $\approx 100$  km s<sup>-1</sup>. Menten & Alcolea (1995) detected high-velocity NH<sub>3</sub> radio emission toward IRC+10420, the high mass-loss rate LPV IK Tau, and the PPN OH 231.8+4.2. In the last case, they find high-velocity emission over  $\approx 70$  km s<sup>-1</sup> FWZP. Recently, Hasegawa et al. (2006) report and discuss observations of the NH<sub>3</sub> line central to the present study, the ortho-NH<sub>3</sub>  $1_0-0_0$  transition, toward IRC+10216 made with the Odin satellite.

One common, surprising result of *all* the above studies is the exceedingly high NH<sub>3</sub> abundances they report. Most of them cite values of several times  $10^{-7}$  or even  $10^{-6}$  relative to molecular hydrogen. These numbers are in stark contrast to the results of thermodynamical equilibrium calculations for the atmospheres of cool stars, which predict the production of only negligible amounts of NH<sub>3</sub>, of order  $10^{-12}$  (Tsuji 1964). The pioneering study of Tsuji has been confirmed by more recent work (see, e.g., Lafont et al. 1982; Cherchneff & Barker 1992, both for C-rich CSEs). Somewhat ad hoc approaches to bringing observations and theory closer together involved injecting a significant amount of NH<sub>3</sub> in the inner envelope (Nejad & Millar 1988; Nercessian et al. 1989). Willacy & Cherchneff (1998) include shock chemistry in their model of IRC+10216, but still only produce an abundance of  $4 \times 10^{-11}$ , at least three orders of

\* *Herschel* is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA.

\*\* Appendix A (page 5) is only available in electronic form at <http://www.aanda.org>

magnitude below the value implied by observations. As to bona fide shocked regions like PPN outflows, [Morris et al. \(1987\)](#) suggested that, for OH 231.8+4.2, N<sub>2</sub>, which binds most of the nitrogen, might be dissociated in the high-velocity gas and that the high NH<sub>3</sub> abundance might be the result of a series of hydrogenation reactions. Whether this can be confirmed by detailed chemical models remains to be explored.

For the present study, as described in Sect. 2, we observed the NH<sub>3</sub>  $J_K = 1_0-0_0$  ortho ground-state transition in O-rich stars of widely different natures and mass loss rates: the high mass-loss LPV IK Tau, the peculiar red supergiant VY CMa, the archetypical OH/IR star OH 26.5+0.6, and the hypergiant IRC+10420. We chose a receiver setting that allowed simultaneous observations of the  $J_{K_a, K_c} = 1_{1,0}-1_{0,1}$  transition of ortho-H<sub>2</sub>O. All of these objects have dense CSEs, and NH<sub>3</sub> has been previously detected toward all of them but OH 26.5+0.6. In particular, for IK Tau and IRC+10420, single-dish observations of the  $(J, K) = (1, 1)$  and  $(2, 2)$  inversion lines have been reported by [Menten & Alcolea \(1995\)](#). Moreover, the emission in these lines has subsequently been imaged with the NRAO Very Large Array (VLA) with a resolution of a few arcseconds ([Menten et al.](#), in prep.; see Sect. 2.3).

The critical density of the cm-wavelength inversion lines is on the order of  $10^4 \text{ cm}^{-3}$ , while that of the sub-mm  $1_0-0_0$  transition has a value  $\sim 4$  orders of magnitude higher. Thus, both types of lines should provide complementary information on different regions of the envelope. In Sect. 3.1, we give a general description of the sub-mm spectra we obtained with HIFI aboard *Herschel* ([Pilbratt et al. 2010](#)). Thereafter, in Sect. 3.2, we present radiative transfer calculations conducted to model the observed line profiles, taking advantage of the constraints from the VLA imaging. These lead to NH<sub>3</sub> abundance determinations.

## 2. Observations

### 2.1. Herschel/HIFI submillimeter observations

The observations were made with the two orthogonal HIFI receivers available for each band, which in all cases work in double side-band (DSB) mode (see [de Graauw et al. 2010](#)). This effectively doubles the instantaneous intermediate frequency (IF) coverage. We observed the four stars described above with a tuning that, in the upper sideband, covers the frequency of the  $J_K = 1_0-0_0$  ground state transition of ortho-NH<sub>3</sub> at 572.4981 GHz. The tuning was chosen to also cover the frequency of the  $J_{K_a, K_c} = 1_{1,0}-1_{0,1}$  ground state line of ortho-H<sub>2</sub>O at 556.9360 GHz in HIFI's lower sideband. The observations were obtained using the dual-beam-switching (DBS) mode. In this mode, the HIFI internal steering mirror chops between the source position and a position believed to be free of emission, which was certainly the case for our observations. The telescope then alternately locates the source in either of the chopped beams, providing a double-difference calibration scheme, which allows a more efficient cancellation of the residual standing waves in the spectra. Additional details on this observing mode can be found in [de Graauw et al. \(2010\)](#). The double sideband system temperature was  $\approx 100$  K, and the calibration uncertainty is estimated to be 10%. Spectral baselines were excellent. *Herschel*'s beam had a size of  $37''$  FWHM at the observing frequency, which is much larger than the NH<sub>3</sub>-emitting regions of all our sources.

The HIFI data shown here were obtained using the wide band spectrometer (WBS), which is an acousto-optical spectrometer, providing a simultaneous coverage of the full instantaneous IF band in the two available orthogonal receivers, with a (oversampled) channel spacing of 0.5 MHz ( $0.27 \text{ km s}^{-1}$ ), about half the

effective resolution. Spectra in the figures have been resampled and smoothed to a channel spacing of  $\approx 1.1 \text{ km s}^{-1}$ .

The data were processed with the standard HIFI pipeline using HIPE, and nonstitched Level-2 data were exported using the HiClass tool available in HIPE. Further processing, i.e. blanking spurious signals, first order polynomial baseline removal, stitching of the spectrometer subbands and averaging, was performed in CLASS. Since the quality of the spectra measured in both horizontal and vertical polarization was good, these were averaged to lower the final noise in the spectrum. This approach is justified since polarization is not a concern for the presented molecular-line analysis. All HIFI data were originally calibrated in units of antenna temperature ( $T_A^*$ ) and were converted to the main-beam temperature ( $T_{\text{MB}}$ ) scale according to  $T_{\text{MB}} = T_A^*/\eta_{\text{MB}}$ , with the main-beam efficiency  $\eta_{\text{MB}} = 0.68$ . In all cases we have assumed a side-band gain ratio of one.

### 2.2. Ammonia spectroscopy and astrophysics


The main focus of this letter is on the  $J_K = 1_0-0_0$  line of ortho-NH<sub>3</sub>. Ammonia microwave spectroscopy has a long history (see, e.g., [Townes & Schawlow 1955](#), [Kukolich 1967](#)). Very briefly, because of the possible orientations of the hydrogen spins, two different species of NH<sub>3</sub> exist that do not interconvert, ortho-NH<sub>3</sub> and para-NH<sub>3</sub>. Ortho-NH<sub>3</sub> assumes states,  $J_K$ , with  $K = 0$  or  $3n$ , where  $n$  is an integer (all H spins parallel), whereas  $K \neq 3$  for para-NH<sub>3</sub> (not all H spins parallel). The principal quantum numbers  $J$  and  $K$  correspond to the total angular momentum and its projection on the symmetry axis of the pyramidal molecule.

The temperature corresponding to the energy of the lowest para level ( $J_K = 1_1$ ) is 22 K above that of the lowest ortho level ( $J_K = 0_0$ ). Therefore, for formation in the interstellar gas phase, which involves reactions with high exothermicities, the ortho-to-para-NH<sub>3</sub> ratio is expected to attain its equilibrium value of unity ([Umamoto et al. 1999](#)). This situation is also expected to hold for CSEs, given that the IR studies cited above place the NH<sub>3</sub> they observe in the hot medium close to the star.

A high-resolution study of the NH<sub>3</sub>  $1_0-0_0$  transition has very recently been presented by [Cazzoli et al. \(2009\)](#). (Only) its upper state is split into several hyperfine structure (hfs) components with the  $\sim 1-2$  MHz splitting resulting from the coupling of the quadrupole moment of the N nucleus with the electric field of the electrons. Two of these components are further split by magnetic interactions. The mean frequency is 572498.1 MHz and the centroid frequencies of the three main hfs groups are all within 2 MHz, corresponding to  $\approx 1 \text{ km s}^{-1}$ , much less than the line widths observed for the targets of this study (see Fig. 1).

### 2.3. VLA observations of inversion lines

In Sect. 3.2 we use data of the  $(J, K) = (1, 1)$  and  $(2, 2)$  inversion lines to constrain our models for IK Tau and IRC+10420. For these stars, single-dish observations of those lines made with the Effelsberg 100 m telescope were reported by [Menten & Alcolea \(1995\)](#). In addition, to place constraints on the spatial distribution of the NH<sub>3</sub> molecules, we have used data obtained with the VLA that will be published separately ([Menten et al.](#) in prep.). The hfs splitting in the inversion lines in velocity units is much wider than for the rotation line. However, because both the  $(1, 1)$  and  $(2, 2)$  lines are very optically thin, as indicated by the spectra and supported by our modeling (see Sect. 3.2), any contribution of the hfs components will be factors of several weaker than the main hfs component and neglected in the modeling. The intensities of the spectra produced from the VLA images, which were restored with a circular beam of  $3.7''$  FWHM, used in that section, are consistent with the published 100 m telescope values.



**Fig. 1.** HIFI spectra compared to radiative transfer model results: The thick black lines show spectra of the  $J_K = 1_0-0_0$  transition of ortho- $\text{NH}_3$  for (top to bottom) IK Tau, VY CMa, OH 26.5+0.6, and IRC+10420. The blue lines represent the ortho- $\text{H}_2\text{O}$   $J_{K_a,K_c} = 1_{1,0}-1_{0,1}$  spectra for the same stars in the same order. The dashed red lines represent the predictions for the  $\text{NH}_3$  line resulting from our radiative transfer modeling. The left- and right hand ordinates give the main-beam brightness temperature scales for the  $\text{NH}_3$  and the  $\text{H}_2\text{O}$  lines, respectively. Except for IRC+10420, they have different ranges.

### 3. Results and analysis

#### 3.1. $\text{NH}_3$ versus $\text{H}_2\text{O}$ emission

In Figs. 1, A.1 and Table 1 we present the results of our HIFI observations of both the  $\text{NH}_3$  and the  $\text{H}_2\text{O}$  ortho ground state lines, together with the results of our  $\text{NH}_3$  modeling. All our observed positions agree to within  $2''$  with the stars' 2MASS positions, which themselves have an absolute accuracy of better than  $0''.1$  (Cutri et al. 2003). The determination of the LSR ranges is somewhat subjective and the upper and lower velocities are uncertain by  $\sim$  a few  $\text{km s}^{-1}$  for weaker lines. For all entries, the formal error in  $\int T_{\text{MB}} dv$  is smaller than  $0.1 \text{ K km s}^{-1}$ . For VY CMa we used the higher of the literature mass-loss rate values scaled to the recently measured trigonometric parallax distance, 1100 pc (Choi et al. 2008). For both IK Tau and IRC+10420 the lower and the higher values of  $X_{\text{NH}_3}$  are implied by the cm lines and the submm line, respectively.

Inspecting Fig. 1 and the table, it is striking to see that the luminosity (integrated intensity) in the  $\text{NH}_3$  and  $\text{H}_2\text{O}$  ground-state lines is of comparable magnitude for all of our objects. The  $\text{NH}_3/\text{H}_2\text{O}$  line ratios are  $\approx 0.50, 0.30, 0.28,$  and  $0.42$  for IK Tau, VY CMa, OH 26.5+0.6, and IRC+10420, respectively. One has to keep in mind that  $\text{H}_2\text{O}$  is a major molecular constituent of our CSEs, while even the presence of observable  $\text{NH}_3$  emission is completely unexplained!

Another remarkable result is that the velocity ranges covered by the two lines are almost identical, which suggests that the bulk of the material producing the emission for both is similar. Moreover, for all our targets, both lines' FWZP values are lower, but comparable to twice the terminal velocity, implying that both molecules are present in the outer layers of the envelope, where the material has almost been fully accelerated. Furthermore, we point out the clear self absorption in the blue wing of the  $\text{H}_2\text{O}$  lines toward IK Tau and VY CMa, which proves that the line emitting region covers the *whole* envelope. Whether this is also true for the  $\text{NH}_3$  line is a priori not clear.

#### 3.2. Radiative transfer modeling and constraints on abundances

The  $\text{NH}_3$  emission of the sources has been modeled with the Monte Carlo radiative transfer code RATRAN developed and described by Hogerheijde & van der Tak (2000). For  $\text{NH}_3$ , RATRAN uses collision rates calculated by Danby et al. (1988). Power laws for the density and temperature were used to describe the physical structure of the envelope, using as input published values for the mass loss rate and expansion velocity (see Table 1). For IRC+10420, envelope parameters from Dinh-V.-Trung et al. (2009) were used and for IK Tau, VY CMa, and OH26.5+0.6 we refer to the modeling of Decin et al. (2010), Decin et al. (2006), and Justanont et al. (2006), respectively. The VLA data yield an extent of  $\approx 4''$  for the  $\text{NH}_3$  emitting region around IRC+10420 and  $\approx 2''.6$  for that around IK Tau, numbers we use for our modeling.

To fit the submm  $\text{NH}_3$  lines observed with HIFI, the  $\text{NH}_3$  abundance was varied in a first iteration. For the two sources with additional data from the cm inversion lines, those (para) lines were modeled as well, using an ortho-to-para ratio of 1, appropriate for formation of  $\text{NH}_3$  under high temperatures (see Sect. 2.2). Interestingly, this does not lead to a satisfying fit for both the cm  $\text{NH}_3$  inversion lines *and* the submm ground state line. With a fit adjusted to reproduce the cm lines, the submm line is underestimated by a factor of 10 for IK Tau.

Since high densities are needed to excite the submm line, its emission must arise from the inner part of the envelope, further inward than the cm-line emitting region. Our modeling suggests densities above a few times  $10^6 \text{ cm}^{-3}$  and temperatures in the 10–100 K range. This discrepancy between the physical conditions required to produce the observed cm emission, on the one hand, and the submm emission, on the other, is even greater for the very extended but relatively low-density shell of IRC+10420. To reach densities high enough to excite the  $\text{NH}_3$  sub-mm line, the mass-loss rate had to be increased to  $2 \times 10^{-3} M_{\odot} \text{ yr}^{-1}$ , i.e., a factor 3 higher than the value derived by Dinh-V.-Trung et al. (2009) and the inner radius reduced to  $5 \times 10^{15} \text{ cm}$ , which is twice the value of the hot inner shell proposed by these authors. Then agreement between the abundances obtained from the cm and submm lines can be reached within a factor of 2. Interestingly, De Beck et al. (2010, accepted for publication), derive a value of  $3.6 \times 10^{-3} M_{\odot} \text{ yr}^{-1}$  for IRC+10420's mass loss rate based on multi-transition modeling of CO.

To reproduce the strong submm  $\text{NH}_3$  line from VY CMa, the highest mass-loss rate and largest radius from the various shells discussed by Decin et al. (2006) had to be used, scaled to  $D = 1100 \text{ pc}$  (see Sect. 3.1). The outer radius that led to a best fit for OH26.5+0.6 is  $2.5 \times 10^{16} \text{ cm}$ . For this source, the same temperature profile as for IK Tau was used.

The line profiles produced by our  $\text{NH}_3$  model calculations are shown in Figs. 1 and A.1 overlaid on the measured spectra.

**Table 1.** Results of HIFI NH<sub>3</sub> and H<sub>2</sub>O observations and NH<sub>3</sub> modeling.

Object	$\alpha_{J2000}$	$\delta_{J2000}$	$D$	$\dot{M}^{\text{Lit}}$	$v_{\infty}$	$v_{\text{LSR}}$ range	$\int T_{\text{MB}} dv$	$\dot{M}^{\text{Mod}}$	$r_{\text{out}}$	$\theta_{\text{out}}$	$X_{\text{NH}_3}$
			(kpc)	( $M_{\odot} \text{ yr}^{-1}$ )	( $\text{km s}^{-1}$ )	( $\text{km s}^{-1}$ )	( $\text{K km s}^{-1}$ )	( $M_{\odot} \text{ yr}^{-1}$ )	(cm)	( $''$ )	
IK Tau	03 <sup>h</sup> 53 <sup>m</sup> 28 <sup>s</sup> .8	+11°24′23″	0.25 {1}	$8 \times 10^{-6}$ {2}	18 {2}	[11.5, 57.2] [10.4, 59.5]	5.1 10.3	$8 \times 10^{-6}$	$5.0 \times 10^{15}$	1.3	(1–3) $10^{-6}$
VY CMa	07 22 58.3	-25 46 03	1.1 {3}	$(0.8\text{--}3.4) 10^{-4}$ {4}	35 {4}	[-23.7, 64.3] [-34, 79]	15.5 51.2	$1.8 \times 10^{-4}$	$3.5 \times 10^{16}$	2.1	$3 \times 10^{-6}$
OH26.5+0.6	18 37 32.5	-05 23 59	1.37 {5}	$2.4 \times 10^{-4}$ {5}	16 {6}	[14.6, 43.2] [15.0, 43.2]	0.96 3.4	$2.4 \times 10^{-4}$	$2.5 \times 10^{16}$	1.2	$3 \times 10^{-7}$
IRC+10420	19 26 48.1	+11 21 17	5 {7}	$8 \times 10^{-4}$ {7}	38 {7}	[39, 112] [42.8, 116.4]	3.1 7.4	$2 \times 10^{-3}$	$1.5 \times 10^{17}$	2.0	(2–5) $10^{-7}$

**Notes.** Columns are (from left to right) the object, J2000 right ascension and declination of observed position, distance, mass-loss rate from the literature, terminal expansion velocity (from CO data), FWZP LSR velocity range with observed emission, integrated main-beam brightness temperature, as well as mass-loss, outer radius of the NH<sub>3</sub> distribution (in cm and in arcsec), and [NH<sub>3</sub>/H<sub>2</sub>] abundance ratio derived from our modeling. For the  $v_{\text{LSR}}$  range and  $\int T_{\text{MB}} dv$  for each source, the first and second rows give the values for the NH<sub>3</sub> and the H<sub>2</sub>O lines, respectively. Numbers in braces relate to the following references for distances and mass loss rates: {1} Hale et al. (1997); {2} Decin et al. (2010); {3} Choi et al. (2008); {4} Decin et al. (2006); {5} Justtanont et al. (2006); {6} Kemper et al. (2003); {7} Dinh-V-Trung et al. (2009).

The resulting NH<sub>3</sub> abundances for the four observed stars range from  $2 \times 10^{-7}$  to  $3 \times 10^{-6}$  (see Table 1). While they are in line with circumstellar NH<sub>3</sub> abundances derived from the inversion lines alone and from IR spectroscopy, we note, as a caveat, that our model calculations did not consider the possibility of IR pumping of the  $1_0\text{--}0_0$  transition. IR pumping of the H<sub>2</sub>O  $1_{1,0}\text{--}1_{0,1}$  line in IRC+10216’s CSE via various vibrational bands has been investigated by Agúndez & Cernicharo (2006), who found it to be the dominant source of excitation of this high critical density line over much of the star’s outer envelope. Consequently, their modeling suggests an order of magnitude lower H<sub>2</sub>O abundance than invoked earlier from SWAS and Odin observations (Melnick et al. 2001; Hasegawa et al. 2006).

#### 4. Summary and outlook

The high critical density of the  $J_K = 1_0\text{--}0_0$  ortho-NH<sub>3</sub> line allows investigations of NH<sub>3</sub> in a new density regime of circumstellar envelopes. However, the remarkably high abundances we determine for all our objects confirm and significantly strengthen the finding that this molecule exists in a variety of CSEs at levels not explained by current chemical models. Better constraints on the emitting regions will come from HIFI observations of a number of NH<sub>3</sub> lines, which are actually evident in the (heavily spectrally diluted) PACS spectrum of VY CMa (Royer et al. 2010). Given their intensities, such observations appear eminently feasible. They would place tight constraints on future radiative transfer modeling that includes infrared excitation. Additionally, with its extremely wideband new generation digital correlator, the Expanded Very Large Array (EVLA) will allow simultaneous imaging of many NH<sub>3</sub> inversion lines. Nevertheless, given the weakness of these lines toward ordinary AGB stars, PPNe being an exception, EVLA observations will be challenging.

*Acknowledgements.* HIFI has been designed and built by a consortium of institutes and university departments from across Europe, Canada, and the United States under the leadership of SRON Netherlands Institute for Space Research, Groningen, The Netherlands and with major contributions from Germany, France, and the US. Consortium members are: Canada: CSA, U. Waterloo; France: CESR, LAB, LERMA, IRAM; Germany: KOSMA, MPIfR, MPS; Ireland, NUI Maynooth; Italy: ASI, IFSI-INAF, Osservatorio Astrofisico di Arcetri-INAF; Netherlands: SRON, TUD; Poland: CAMK, CBK; Spain: Observatorio Astronómico Nacional (IGN), Centro de Astrobiología (CSIC-INTA); Sweden: Chalmers University of Technology – MC2, RSS & GARD; Onsala Space Observatory; Swedish National Space Board, Stockholm University – Stockholm Observatory; Switzerland: ETH Zurich, FHNW; USA: Caltech, JPL, NHSC. HCSS / HSpot / HIPE is a joint development by the *Herschel* Science Ground Segment Consortium, consisting of ESA, the NASA *Herschel* Science Center, and the HIFI, PACS and SPIRE consortia. This work has been partially supported by the Spanish MICINN, within the program CONSOLIDER INGENIO 2010, under grant “Molecular Astrophysics: The


*Herschel* and ALMA Era – ASTROMOL” (ref.: CSD2009-00038). R. Sz. and M. Sch. acknowledge support from grant N 203 393334 from the Polish MNiSW. K. J. acknowledges the funding from SNSB. J. C. thanks funding from MICINN, grant AYA2009-07304.

#### References

- Agúndez, M., & Cernicharo, J. 2006, ApJ, 650, 374  
 Bell, M. B., Kwok, S., Matthews, H. E., & Feldman, P. A. 1982, AJ, 87, 404  
 Betz, A. L., & Goldhaber, D. M. 1985, in Mass Loss from Red Giants, ed. M. Morris, & B. Zuckerman, Astrophys. Space Sci. Libr., 117, 83  
 Betz, A. L., McLaren, R. A., & Spears, D. L. 1979, ApJ, 229, L97  
 Cazoli, G., Dore, L., & Puzzarini, C. 2009, A&A, 507, 1707  
 Cherchneff, I., & Barker, J. R. 1992, ApJ, 394, 703  
 Cheung, A. C., Rank, D. M., Townes, C. H., Thornton, D. D., & Welch, W. J. 1968, Phys. Rev. Lett., 21, 1701  
 Choi, Y. K., Hirota, T., Honma, M., et al. 2008, PASJ, 60, 1007  
 Cutri, R. M., Skrutskie, M. F., van Dyk, S., et al. 2003, 2MASS All Sky Catalog of point sources. (NASA/IPAC Infrared Science Archive)  
 Danby, G., Flower, D. R., Valiron, P., Schilke, P., & Walmsley, C. M. 1988, MNRAS, 235, 229  
 Decin, L., De Beck, E., Brunken, S., et al. 2010, A&A, 516, A69  
 Decin, L., Hony, S., de Koter, A., et al. 2006, A&A, 456, 549  
 de Graauw, Th., Helmich, F. P., Phillips, T. G., et al. 2010, A&A, 518, L6  
 Dinh-V-Trung, Muller, S., Lim, J., Kwok, S., & Muthu, C. 2009, ApJ, 697, 409  
 Hale, D. D. S., Bester, M., Danchi, W. C., et al. 1997, ApJ, 490, 407  
 Hasegawa, T. I., Kwok, S., Koning, N., et al. 2006, ApJ, 637, 791  
 Ho, P. T. P., & Townes, C. H. 1983, ARA&A, 21, 239  
 Hogerheijde, M. R., & van der Tak, F. F. S. 2000, A&A, 362, 697  
 Justtanont, K., Olofsson, G., Dijkstra, C., & Meyer, A. W. 2006, A&A, 450, 1051  
 Kemper, F., Stark, R., Justtanont, K., et al. 2003, A&A, 407, 609  
 Kukulich, S. G. 1967, Phys. Rev., 156, 83  
 Kwok, S., Bell, M. B., & Feldman, P. A. 1981, ApJ, 247, 125  
 Lafont, S., Lucas, R., & Omont, A. 1982, A&A, 106, 201  
 Martin-Pintado, J., & Bachiller, R. 1992, ApJ, 391, L93  
 McLaren, R. A., & Betz, A. L. 1980, ApJ, 240, L159  
 Melnick, G. J., Neufeld, D. A., Ford, K. E. S., Hollenbach, D. J., & Ashby, M. L. N. 2001, Nature, 412, 160  
 Menten, K. M., & Alcolea, J. 1995, ApJ, 448, 416  
 Monnier, J. D., Danchi, W. C., Hale, D. S., Tuthill, P. G., & Townes, C. H. 2000, ApJ, 543, 868  
 Morris, M., Guilloteau, S., Lucas, R., & Omont, A. 1987, ApJ, 321, 888  
 Nejad, L. A. M., & Millar, T. J. 1988, MNRAS, 230, 79  
 Nercessian, E., Omont, A., Benayoun, J. J., & Guilloteau, S. 1989, A&A, 210, 225  
 Nguyen-Q-Rieu, Graham, D., & Bujarrabal, V. 1984, A&A, 138, L5  
 Pilbratt, G. L., Riedinger, J. R., Passvogel, T., et al. 2010, A&A, 518, L1  
 Royer, P., Decin, L., Wesson, R., et al. 2010, A&A, 518, L145  
 Townes, C. H., & Schawlow, A. L. 1955, Microwave Spectroscopy (New York: McGraw-Hill)  
 Truong-Bach, Graham, D., & Nguyen-Q-Rieu. 1996, A&A, 312, 565  
 Truong-Bach, Nguyen-Q-Rieu, & Graham, D. 1988, A&A, 199, 291  
 Tsuji, T. 1964, Annals of the Tokyo Astronomical Observatory, 9  
 Umemoto, T., Mikami, H., Yamamoto, S., & Hirano, N. 1999, ApJ, 525, L105  
 Walmsley, C. M., & Ungerechts, H. 1983, A&A, 122, 164  
 Willacy, K., & Cherchneff, I. 1998, A&A, 330, 676

## Appendix A: NH<sub>3</sub> rotational and inversion line spectra and fits for IK Tau and IRC+10420

Figure A.1 shows, for IK Tau and IRC+10420, the HIFI spectra of the  $J_K = 1_0-0_0$  ortho-NH<sub>3</sub> line and the spectra of the  $(J, K) = (1, 1)$  and  $(2, 2)$  para-NH<sub>3</sub> lines produced from our VLA data together with our best fit model predictions.



**Fig. A.1.** Spectra of NH<sub>3</sub> transitions for IK Tau (left column) and IRC+10420 (right column). Top row: HIFI spectra of the  $J_K = 1_0-0_0$  ortho-NH<sub>3</sub> line. Middle row: VLA spectra of the  $(J, K) = (1, 1)$  para-NH<sub>3</sub> line. Bottom row: VLA spectra of the  $(J, K) = (2, 2)$  para-NH<sub>3</sub> line. The intensity scales apply for both sources. The lefthand ordinate gives main-beam brightness temperature in a  $40''$  FWHM beam, while the righthand ordinate gives flux density in Jy units. In all panels the red dashed line represents our best-fit model prediction. The blue bars give the spacings and theoretical relative intensities of the main groups of hfs components determined by Cazzoli et al. (2009) for the  $1_0-0_0$  transition and by Kukolich (1967) for the  $(1, 1)$  and  $(2, 2)$  inversion lines, respectively.

- <sup>1</sup> Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany  
e-mail: kmenten@mpi.fr.de
- <sup>2</sup> Observatorio Astronómico Nacional (IGN), Alfonso XII No 3, 28014 Madrid, Spain
- <sup>3</sup> Instituut voor Sterrenkunde, Katholieke Universiteit Leuven, Celestijnenlaan 200D, 3001 Leuven, Belgium
- <sup>4</sup> Sterrenkundig Instituut Anton Pannekoek, University of Amsterdam, Science Park 904, 1098 Amsterdam, The Netherlands
- <sup>5</sup> European Space Astronomy Centre, ESA, PO Box 78, 28691 Villanueva de la Cañada, Madrid, Spain
- <sup>6</sup> Observatorio Astronómico Nacional (IGN), Ap 112, 28803 Alcalá de Henares, Spain
- <sup>7</sup> CAB, INTA-CSIC, Ctra de Torrejón a Ajalvir, km 4, 28850 Torrejón de Ardoz, Madrid, Spain
- <sup>8</sup> Department of Astrophysics/IMAPP, Radboud University, Nijmegen, The Netherlands
- <sup>9</sup> Onsala Space Observatory, Dept. of Radio and Space Science, Chalmers University of Technology, 43992 Onsala, Sweden
- <sup>10</sup> Astronomical Institute, Utrecht University, Princetonplein 5, 3584 CC Utrecht, The Netherlands
- <sup>11</sup> Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA
- <sup>12</sup> The Johns Hopkins University, 3400 North Charles St, Baltimore, MD 21218, USA
- <sup>13</sup> Department of Astronomy, AlbaNova University Center, Stockholm University, 10691 Stockholm, Sweden
- <sup>14</sup> N. Copernicus Astronomical Center, Rabiańska 8, 87-100 Toruń, Poland
- <sup>15</sup> Joint ALMA Observatory, El Golf 40, Las Condes, Santiago, Chile
- <sup>16</sup> Department of Physics and Astronomy, University of Waterloo, Waterloo, ON Canada N2L 3G1, Canada
- <sup>17</sup> SRON Netherlands Institute for Space Research, Landleven 12, 9747 AD Groningen, The Netherlands
- <sup>18</sup> California Institute of Technology, Cahill Center for Astronomy and Astrophysics 301-17, Pasadena, CA 91125, USA
- <sup>19</sup> Infrared Processing and Analysis Center, California Institute of Technology, MS 100-22, Pasadena, CA 91125, USA
- <sup>20</sup> Laboratoire d'Études du Rayonnement et de la Matière en Astrophysique, UMR 8112 CNRS/INSU, OP, ENS, UPMC, UCP, Paris, France
- <sup>21</sup> LERMA, Observatoire de Paris, 61 avenue de l'Observatoire, 75014 Paris, France
- <sup>22</sup> Institute Centre d'Étude Spatiale des Rayonnements, Université de Toulouse [UPS], 31062 Toulouse Cedex 9, France
- <sup>23</sup> CNRS/INSU, UMR 5187, 9 avenue du Colonel Roche, 31028 Toulouse Cedex 4, France